

EPP0527

Deinstitutionalization in Georgia-why it is so slowN. Zavrashvili^{1*} and G. Matiashvili¹¹School of health sciences, The University of Georgia, Tbilisi, Georgia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.651

Introduction: Mental health reform represents one of the most transformative changes in the field of healthcare, as it not only changes the forms of services but also the nature of services offered. While many countries have successfully implemented such reforms, others, including Georgia, have struggled with a protracted and inconsistent process. Despite decades of advocacy by professionals for deinstitutionalization and the development of community-based services, psychiatric hospital treatment continues to dominate in Georgia.

Objectives: The purpose of the review is to explore the concept of deinstitutionalization within the mental health landscape and assess its status in the context of Georgia. It aims to study the lessons learned from successful deinstitutionalization and illuminate achievements and challenges surrounding deinstitutionalization in Georgia's reality.

Methods: A qualitative analysis including desk review, in-depth interviews and focus group discussions was conducted. Proceeding from the research objectives we analyzed the existing legislation, strategic documents and clinical practices concerning individuals with mental disorders; Interviews were also conducted with key informants on the shortcomings and problems in deinstitutionalization practices

Results: The review findings reveal, that despite recent progress such as the development of community mobile teams and increased funding allocated for community services within mental health budget, several issues persist: there is no agreement among stakeholders on how to restructure existing hospital beds and financial provisions remain unresolved. The field of mental health in Georgia suffers from a lack of human resources. Attracting new personnel, ensuring regional distribution, and enhancing qualifications are necessary components of deinstitutionalization that require the involvement of all stakeholders, coordinated and time-planned action. The current mental healthcare system in Georgia is characterized by a lack of coordination and collaboration among its various components. Establishing patient care pathways with clear referral criteria is crucial for improving the efficiency of mental health services. This research highlights that successful deinstitutionalization requires additional funds, time, and trained people. Institutions should have a long-term (3-5) year development plan, detailing the source of funding, activities to be implemented, and expected outcomes. In the absence of such a plan, progress remains sporadic, intermittent, uncoordinated, and less effective.

Conclusions: In conclusion, the protracted nature of deinstitutionalization in Georgia requires a multifaceted approach involving various stakeholders. By addressing identified challenges and promoting coordination among mental health components, Georgia can guide a more effective course toward a community-based, patient-centered mental healthcare system.

Disclosure of Interest: None Declared

EPP0528

A Cost-Effectiveness Analysis of a Ketamine-assisted Psychotherapy Program Compared to Online Group Psychotherapy in British Columbia, Canada

V. Tsang

UBC, Vancouver, Canada

doi: 10.1192/j.eurpsy.2024.652

Introduction: Depression continues to present significant economic burdens to the Canadian healthcare system. Novel therapies, including those that incorporate psychoactive substances such as ketamine, present an opportunity to evaluate both clinical and economic effectiveness against current standards of care, which may be repeatedly proving ineffective in treating depression for some individuals.

Objectives: This paper evaluates the cost-effectiveness of the Roots to Thrive ketamine program compared to group psychotherapy covered through the medical services plan in British Columbia, Canada.

Methods: A discrete-time Markov-model is used to estimate depressive states over five cycles for a treatment cohort and a synthetic control cohort. The transition probabilities for the treatment cohort are calculated from Roots to Thrive program data (n = 62) over the past 3 years, with the control cohort using published values from the literature. Both cohorts use the same starting state distribution, excess healthcare utilization rates for each severity level of depression, and utility outcomes based on depression state severity.

Results: Compared to the control cohort, the Roots to Thrive program was less expensive and produced better outcomes as measured by PHQ-9 scores and Quality-Adjusted life years over 5 treatment cycles. On average, the Roots to Thrive program would save \$14,481 and produce 0.94 additional QALY's per individual compared to group psychotherapy of three patients per provider in the current standard of care.

Conclusions: From an economic perspective, incorporating the Roots to Thrive program - or a program like it - into care in British Columbia would provide both an improvement in health outcomes and reduce expenditure by the ministry of health. These funds could be reinvested into other areas of the healthcare system to improve the lives of all British Columbians, even those that do not engage in psychedelic-assisted psychotherapy.

Disclosure of Interest: None Declared

Neuroscience in Psychiatry

EPP0530

Gamma activity in Autism Spectrum Disorder: Enhanced response to visual inputB. Kakuszi^{1*}, B. Szuromi², M. Tóth¹, I. Bitter¹ and P. Czobor¹¹Psychiatry and Psychotherapy, Semmelweis University and ²OMIII-Nyíró Gyula Hospital, Budapest, Hungary

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.653