


RESEARCH ARTICLE

Calculating premium principles from the mode of a unimodal weighted distribution

Georgios Psarrakos 

Department of Statistics and Insurance Science, University of Piraeus, Piraeus, Greece
Email: gpsarr@unipi.gr

Received: 15 October 2023; **Revised:** 17 March 2024; **Accepted:** 17 April 2024; **First published online:** 15 May 2024

Keywords: Weighted premium calculation principle; exponential premium; Esscher premium; Kamps premium; stochastic ordering; mode ordering; weighted distribution; unimodal distribution

2020 Mathematics Subject Classification: 91G05; 62P05

Abstract

The theory of utility is a well-known method of constructing insurance premiums (see e.g., Newton *et al.* (1986) *Actuarial Mathematics*. Itasca, Illinois: The Society of Actuaries.). Furman and Zitikis ((2008) *Insurance: Mathematics and Economics*, 42, 459–465.) proposed an alternative method using the mean value of a weighted random variable. According to this approach, for various choices of weighting, popular premiums such as net premium, modified variance premium, Esscher premium, and Kamps premium are obtained. On the other hand, some premiums cannot be obtained with this method, such as the premium of the exponential principle. In this paper, we provide a complementary theory by introducing a family of unimodal weighted distributions for which the mode is a premium principle.

1. Introduction

Let $X \geq 0$ be a loss random variable with probability density function (pdf) $f(x)$, cumulative distribution function (cdf) $F(x) = \mathbb{P}(X \leq x)$, and tail function $\bar{F}(x) = 1 - F(x) = \mathbb{P}(X > x)$. Moreover, let $\mathbb{E}(X) = \int_0^\infty x f(x) dx = \int_0^\infty \bar{F}(x) dx$ be the mean value and m_x a mode of the distribution of X satisfying the condition $f'(m_x) = 0$. The weighted premium principle introduced by Furman and Zitikis (2008) is defined and denoted by:

$$H_w(X) = \frac{\mathbb{E}[X w(X)]}{\mathbb{E}[w(X)]}, \quad (1.1)$$

where $w: [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing function such that $0 < \mathbb{E}[w(X)] < \infty$. One option to obtain a weighted premium principle is to take the mean value of a weighted loss random variable X_w with pdf:

$$f_w(x) = \frac{w(x)}{\mathbb{E}[w(X)]} f(x), \quad x > 0.$$

Since w is nondecreasing by assumption, it holds that $H_w(X) = \mathbb{E}(X_w) \geq \mathbb{E}(X)$. We recall that, for different choices of w , some well-known premiums follow directly. For example, we can get the net premium $\mathbb{E}(X)$ (when $w(x) = c$, $c > 0$ is a constant), the modified variance premium $\mathbb{E}(X) + \text{Var}(X)/\mathbb{E}(X)$ (when $w(x) = x$), the Esscher premium $\mathbb{E}(X e^{\lambda X})/\mathbb{E}(e^{\lambda X})$ (when $w(x) = e^{\lambda x}$, $\lambda > 0$), and the Kamps premium $\mathbb{E}[X(1 - e^{-\lambda X})]/\mathbb{E}[1 - e^{-\lambda X}]$ (when $w(x) = 1 - e^{-\lambda x}$, $\lambda > 0$); see Furman and Zitikis (2008). Furthermore, for $w(x) = g'(\bar{F}(x))$, we obtain a large class of premium principles based on distorted expectation theory (when $g: [0, 1] \rightarrow [0, 1]$ is a continuous and differentiable function such as: (i) g

is increasing and concave function, (ii) $g(0) = 0$ and $g(1) = 1$). On the other hand, there are premium principles such as the exponential premium principle $\ln \mathbb{E}(e^{\lambda X})/\lambda$, where $\lambda > 0$ and ‘ln’ means natural logarithm, which are not obtained for some function w . For more details on premium principles, see Gerber (1979), Bühlmann (1980), Goovaerts *et al.* (1984), Newton *et al.* (1986), Gerber and Shiu (1994), Denneberg (1994), Wang (1995, 1996, 2003), Kamps (1998), Denuit *et al.* (2006), Laeven and Goovaerts (2008), Castaño-Martínez *et al.* (2020), and references therein.

The ordering is important in studying premium principles. Next, we recall two well-known stochastic orders, which imply the mean ordering; see the classical books of Müller and Stoyan (2002), Shaked and Shanthikumar (2007), and Belzunce *et al.* (2016). Let X and Y be two continuous random variables with pdf's $f_X(x)$ and $f_Y(x)$, and tail functions $\bar{F}_X(x)$ and $\bar{F}_Y(x)$, respectively. A random variable X is said to be smaller than Y in the usual stochastic order, denoted by $X \leq_{st} Y$, if $\bar{F}_X(x) \leq \bar{F}_Y(x)$ for all x . Moreover, $X \leq_{st} Y$, if $\mathbb{E}[\Phi(X)] \leq \mathbb{E}[\Phi(Y)]$ for all nondecreasing functions Φ such that the expectations exist. Furthermore, we say that X is smaller than Y in the likelihood ratio order, denoted by $X \leq_{lr} Y$, if $f_Y(x)/f_X(x)$ is nondecreasing with respect to x . The connection between these stochastic orders is given by the following diagram:

$$X \leq_{lr} Y \Rightarrow X \leq_{st} Y \Rightarrow \mathbb{E}(X) \leq \mathbb{E}(Y).$$

Calculating weighted premium principles, Furman and Zitikis (2008), applied the ordering $X \leq_{lr} X_w$, where w is a nondecreasing function, and introduced the premium given in (1.1) by using the mean of X_w , that is, $H_w(X) = \mathbb{E}(X_w) \geq \mathbb{E}(X)$. An important advantage of this method is that given the distribution of risk X , one can determine the distribution of X_w , namely, an actuary has more information from just the knowledge of premium principle $\mathbb{E}(X_w)$. A natural question here is: Can we use a different statistical measure instead of mean value on the distribution of a weighted random variable to obtain a premium principle?

The contribution of this work is to highlight the role of mode of a weighted risk in calculating premium principles. Our method is not only a complementary theory but also a generalization calculating weighted premiums. A challenging problem here is the mode ordering for two unimodal distributions. For this aim, we study how we can use the usual stochastic order to construct a family of unimodal weighted distributions, into which we can order the modes in the strict sense (i.e., for two unimodal random variables X and Y with modes m_X and m_Y , we say that $X <_{mode} Y$, if $m_X < m_Y$).

This paper is organized as follows. In Section 2, we calculate the mean value premium principle by using the mode of a weighted risk, and we provide some distributional properties. In Section 3, we give a generalization of the mean value premium principle, which also generalizes the method of calculating weighted premium principles proposed by Furman and Zitikis (2008). Furthermore, we give a generalization of the exponential premium principle and we provide a two-side bound for it in terms of the Esscher premium. In Section 4, we study the comparison of modes between two unimodal distributions, by introducing the concept of mode ordering.

Throughout the paper, we give some examples to illustrate the theoretical results and discuss how the mechanism of our method works. Furthermore, whenever we use an expectation, we are tacitly assuming that it exists. Moreover, we use the terms *nondecreasing* and *nonincreasing* in a wide sense, that is, a function g is nondecreasing (resp., nonincreasing) if $g(x) \leq g(y)$ (\geq) for all $x < y$. We also use the terms *increasing* and *decreasing* in a strict sense, that is, a function g is increasing (resp., decreasing) if $g(x) < g(y)$ ($>$) for all $x < y$.

2. Calculating the mean value premium principle from the mode of a unimodal weighted distribution

Let X be a continuous nonnegative random variable with pdf f , distribution F , and tail $\bar{F} = 1 - F$. Let also $w : [0, \infty) \rightarrow [0, \infty)$ is a continuous, increasing, and convex function such that $0 < \mathbb{E}[w(X)] < \infty$.

We consider the weighted random variable $X_{\tilde{w}}$ with mode $m_{X_{\tilde{w}}} = \tilde{m}_w$ and pdf:

$$f_{\tilde{w}}(x) = \frac{\tilde{w}(x)}{\mathbb{E}[\tilde{w}(X)]} f(x) = \frac{\bar{F}_w(x) - \bar{F}(x)}{\mathbb{E}(X_w) - \mathbb{E}(X)}, \quad x > 0, \tag{2.1}$$

provided that $\mathbb{E}(X_w) > \mathbb{E}(X) > 0$. Here, the weight $\tilde{w}(x)$ is

$$\tilde{w}(x) = \frac{\bar{F}_w(x) - \bar{F}(x)}{f(x)}.$$

Furthermore, $X_{\tilde{w}}$ is unimodal, with mode $\tilde{m}_w = w^{-1}(\mathbb{E}[w(X)])$. Therefore, the premium principle

$$T_{\tilde{w}}(X) := \tilde{m}_w = w^{-1}(\mathbb{E}[w(X)]) \tag{2.2}$$

is the well-known mean value principle; see for more details Goovaerts *et al.* (1984). Since, by assumption, $w(x)$ is convex, we have (by Jensen’s inequality) the property $\tilde{m}_w \geq \mathbb{E}(X)$.

Remark 2.1.

- (i) An interpretation of the density $f_{\tilde{w}}$ in applied probability is the following. Let X and Y be two nonnegative continuous random variables with $X \leq_{st} Y$ and $\mathbb{E}(X) < \mathbb{E}(Y)$. Let also $\bar{F}_X(x)$ and $\bar{F}_Y(x)$ be the tail functions of X and Y , respectively. Then, by Theorem 4.1 of Di Crescenzo (1999), there exists a suitable nonnegative random variable U such that

$$\mathbb{E}[g(Y)] - \mathbb{E}[g(X)] = \mathbb{E}[g'(U)] [\mathbb{E}(Y) - \mathbb{E}(X)], \tag{2.3}$$

provided that g is a measurable and differentiable function such that $\mathbb{E}[g(X)]$ and $\mathbb{E}[g(Y)]$ are finite, and that its derivative g' is measurable and Riemann-integrable. Here, the pdf of the random variable U is

$$f_U(x) = \frac{\bar{F}_Y(x) - \bar{F}_X(x)}{\mathbb{E}(Y) - \mathbb{E}(X)}.$$

Recalling that the mean value theorem states that if g is a function continuous in $[x, y]$ and differentiable in (x, y) , then there exists a point $u \in (x, y)$ such that

$$g(y) - g(x) = g'(u) (y - x),$$

formula (2.3) is a probabilistic analogue of the mean value theorem. Hence, applying this probabilistic analogue of the mean value theorem between X and X_w (i.e., X_w plays the role of Y), we obtain the new random variable $X_{\tilde{w}}$ (i.e., $X_{\tilde{w}}$ plays the role of U) with pdf $f_{\tilde{w}}$. For more details, see Di Crescenzo (1999), Psarrakos (2022), and reference therein. It is worth mentioning that the relation $\mathbb{E}(X) \leq \mathbb{E}(X_{\tilde{w}}) \leq \mathbb{E}(X_w)$ does not hold in general (see Proposition 4.1 in Di Crescenzo, 1999), which implies that the relation $X \leq_{st} X_{\tilde{w}} \leq_{st} X_w$ does not hold in general.

- (ii) The premium $T_{\tilde{w}}(X)$ for $w(x) = e^{\lambda x}$ gives the exponential premium $\ln \mathbb{E}(e^{\lambda X})/\lambda$ ($\lambda > 0$). Furthermore, for $w(x) = x^a$ ($a \geq 1$), $T_{\tilde{w}}(X)$ yields the premium $(\mathbb{E}[X^a])^{1/a}$; see Goovaerts *et al.* (1984, p. 123).

It is well known that if the mean is greater than the mode, then the distribution is positively skewed. We recall that Pearson (1895) proposed $(\mu - m)/\sigma$ as a measure of skewness for a univariate distribution with mean μ , mode m , and variance σ^2 . Next, we compare the mode and the mean of $X_{\tilde{w}}$, under the hypothesis that $X \leq_{st} X_{\tilde{w}}$.

Proposition 2.1. *If $X \leq_{st} X_{\tilde{w}}$, then $\tilde{m}_w \leq \mathbb{E}(X_{\tilde{w}})$.*

Proof. Let $X \leq_{st} X_{\tilde{w}}$. Recalling that w is an increasing and convex function, we have

$$\mathbb{E}[w(X)] \leq \mathbb{E}[w(X_{\tilde{w}})] \leq w[\mathbb{E}(X_{\tilde{w}})],$$

where the second inequality follows by Jensen’s inequality. Therefore,

$$w^{-1}(\mathbb{E}[w(X)]) \leq \mathbb{E}(X_{\tilde{w}}),$$

or equivalently,

$$\tilde{m}_w \leq \mathbb{E}(X_{\tilde{w}}),$$

which completes the proof. □

In the following theorem, for $w(x) = x$, we provide a characterization result for the case where $X_{\tilde{w}} =_d X_w$, where $=_d$ means the equality in distribution. This condition gives that $\tilde{w}(x) = w(x)$ is increasing in x . Hence, (keeping in mind that likelihood order \leq_{lr} implies the usual stochastic order \leq_{st}) the assumption $X \leq_{st} X_{\tilde{w}}$ in Proposition 2.1 is satisfied.

Theorem 2.1. *Consider the weighted risks X_w and $X_{\tilde{w}}$, where $w(x) = x, x > 0$. Then, $X_{\tilde{w}} =_d X_w$ if and only if X follows a Gamma distribution.*

Proof. (\Rightarrow) Let $X_{\tilde{w}} =_d X_w$. Then, we have

$$f_{\tilde{w}}(x) = \bar{f}_w(x),$$

for all $x > 0$, and using formula (2.1) we get

$$\frac{\bar{F}_w(x) - \bar{F}(x)}{\frac{\mathbb{E}(X^2)}{\mathbb{E}(X)} - \mathbb{E}(X)} = \frac{x}{\mathbb{E}(X)} f(x).$$

By differentiating and after some computation, we conclude that

$$\frac{f'(x)}{f(x)} = \frac{\mathbb{E}(X) - x - \frac{\text{Var}(X)}{\mathbb{E}(X)}}{\frac{\text{Var}(X)}{\mathbb{E}(X)} x}.$$

An alternative representation of the latter expression is

$$\frac{f'(x)}{f(x)} = \frac{k - x - q'(x)}{q(x)},$$

where $k = \mathbb{E}(X)$ and $q(x) = \frac{\text{Var}(X)}{\mathbb{E}(X)} x$. Therefore, by Ruiz and Navarro (1994), X follows a Gamma distribution. □

(\Leftarrow) Suppose that X follows a $\text{Gamma}(a, b)$ distribution with parameters $a > 0, b > 0$, and pdf

$$f(x) = \frac{b^a}{\Gamma(a)} x^{a-1} e^{-bx}, \quad x > 0, \tag{2.4}$$

where $\Gamma(a) = \int_0^\infty x^{a-1} e^{-x} dx$ is the Gamma function, and tail

$$\bar{F}(x) = \frac{\Gamma(a, bx)}{\Gamma(a)},$$

where $\Gamma(a, x) = \int_x^\infty y^{a-1} e^{-y} dy$ is the upper incomplete Gamma function. Then, X_w follows a $\text{Gamma}(a + 1, b)$ distribution. Recalling that

$$\Gamma(a + 1, x) = a \Gamma(a, x) + x^a e^{-x},$$

formula (2.1) yields

$$f_{\tilde{w}}(x) = \frac{\frac{\Gamma(a+1, bx)}{\Gamma(a+1)} - \frac{\Gamma(a, bx)}{\Gamma(a)}}{\frac{a+1}{b} - \frac{a}{b}} = \frac{b^{a+1}}{\Gamma(a+1)} x^a e^{-bx} = f_w(x),$$

and the result follows. □

Table 1. The values of $\mathbb{E}(X)$, $\mathbb{E}(X_w)$, $\mathbb{E}(X_{\tilde{w}})$, and \tilde{m}_w of Gamma distribution in Example 2.1, for $a = 3$, $b = 2$, and $c = 1, 2, 3, 4$.

c	$\mathbb{E}(X)$	$\mathbb{E}(X_w)$	$\mathbb{E}(X_{\tilde{w}})$	\tilde{m}_w
1	1.5	2	2	1.5
2	1.5	2.5	2.25	1.73205
3	1.5	3	2.5	1.95743
4	1.5	3.5	2.75	2.17794




Figure 1. The pdf $f_w(x)$ given in Example 2.1 for $a = 3$, $b = 2$ and $0 \leq x \leq 7$, when (i) $c = 1$ (left part), and (ii) $c = 2$ (right part).

Example 2.1. Let X follows a Gamma(a, b) distribution with parameters $a > 0$, $b > 0$, and pdf $f(x)$ given in (2.4). The r -order moment of X is

$$\mathbb{E}(X^r) = \frac{\Gamma(a+r)}{b^r \Gamma(a)}.$$

Let also $w(x) = x^c$, where $c \geq 1$. Then, X_w follows a Gamma($a + c, b$) with mean $\mathbb{E}(X_w) = \frac{a+c}{b}$. Furthermore, we have

$$f_w(x) = \frac{\frac{\Gamma(a+c, bx)}{\Gamma(a+c)} - \frac{\Gamma(a, bx)}{\Gamma(a)}}{\frac{c}{b}}$$

and

$$\mathbb{E}(X_{\tilde{w}}) = \frac{\mathbb{E}(X_w^2) - \mathbb{E}(X^2)}{2(\mathbb{E}(X_w) - \mathbb{E}(X))} = \frac{\frac{\Gamma(a+c+2)}{b^2 \Gamma(a+c)} - \frac{\Gamma(a+2)}{b^2 \Gamma(a)}}{2(\frac{a+c}{b} - \frac{a}{b})} = \frac{2a + c + 1}{2b}.$$

It is clear that for $c = 1$, we have $\mathbb{E}(X_w) = \mathbb{E}(X_{\tilde{w}})$ (see also Theorem 2.1), and for $c > 1$, we have $\mathbb{E}(X_w) > \mathbb{E}(X_{\tilde{w}})$. Furthermore, the mode of $X_{\tilde{w}}$ (see (2.2)) is

$$\tilde{m}_w = \left(\frac{\Gamma(a+c)}{\Gamma(a)} \right)^{1/c} \frac{1}{b}.$$

In Table 1, we give the values of $\mathbb{E}(X)$, $\mathbb{E}(X_w)$, $\mathbb{E}(X_{\tilde{w}})$, and \tilde{m}_w for $a = 3$, $b = 2$, and $c = 1, 2, 3, 4$. Furthermore, in Figures 1 and 2, we drew the pdf $f_w(x)$, giving also the values of mode and mean of $X_{\tilde{w}}$ (see the vertical lines).

Example 2.2. Suppose that X follows an Exponential(b) distribution with parameter $b > 0$, and pdf

$$f(x) = b e^{-bx}, \quad x > 0. \tag{2.5}$$




Figure 2. The pdf $f_w(x)$ given in Example 2.1 for $a = 3$, $b = 2$ and $0 \leq x \leq 7$, when (i) $c = 3$ (left part), and (ii) $c = 4$ (right part).

For $b > \lambda > 0$, let also $w(x) = e^{\lambda x}$. Then, X_w follows an Exponential($b - \lambda$) distribution, and the pdf of X_w is

$$f_w(x) = \frac{e^{-(b-\lambda)x} - e^{-bx}}{\frac{1}{b-\lambda} - \frac{1}{b}} = \frac{b}{\lambda} (b - \lambda) e^{-(b-\lambda)x} (1 - e^{-\lambda x}), \quad x > 0.$$

It is worth mentioning that X_w belongs to a family of weighed exponential distribution introduced by Gupta and Kundu (2009). We recall, see Gupta and Kundu (2009), that a nonnegative random variable X is said to have a weighted exponential distribution, with the shape and scale parameters as $\alpha > 0$ and $\beta > 0$, respectively, if the PDF of X is

$$f_x(x; \alpha, \beta) = \frac{\alpha + 1}{\alpha} \lambda e^{-\beta x} (1 - e^{-\alpha \beta x}), \quad x > 0.$$

Therefore, X_w follows a weighed exponential distribution with parameters $\alpha = \lambda / (b - \lambda)$ and $\beta = b - \lambda$. A probabilistic interpretation of the $f_w(x)$ is that $X_w =_d X + X_w$; see Di Crescenzo (1999); for further interpretations and properties, see Gupta and Kundu (2009). The mode and the mean value of X_w are

$$\tilde{m}_w = \frac{1}{\lambda} \ln \mathbb{E}(e^{\lambda X}) = \frac{1}{\lambda} \ln \frac{b}{b - \lambda}$$

and

$$\mathbb{E}(X_w) = \mathbb{E}(X) + \mathbb{E}(X_w) = \frac{1}{b} + \frac{1}{b - \lambda}.$$

One can also verify that the ratios

$$\frac{f_w(x)}{f(x)} = \frac{b - \lambda}{\lambda} (e^{\lambda x} - 1)$$

and

$$\frac{f_w(x)}{f_w(x)} = \frac{b}{\lambda} (1 - e^{-\lambda x})$$

are increasing functions. Hence, $X \leq_{lr} X_w \leq_{lr} X_w$. In Figure 3, we drew the pdf $f_w(x)$ for $b = 2$ and $\lambda = 1$, giving also the values $\tilde{m}_w = 0.693147$ and $\mathbb{E}(X_w) = 1.5$.

Example 2.3. Suppose that X follows a Pareto(a, b) type I distribution with parameters $a > 3$ and $b > 0$, pdf

$$f(x) = \frac{ab^a}{x^{a+1}}, \quad x > b,$$




Figure 3. The pdf $f_w(x)$ given in Example 2.2 for $b = 2, \lambda = 1$ and $0 \leq x \leq 5$.

and r -order moment $\mathbb{E}(X^r) = ab^r / (a - r)$ for $0 < r < a$. We also consider the weight function $w(x) = x^c$, for $1 \leq c < a - 1$. Then, X_w follows a Pareto($a - c, b$) type I distribution, and the pdf of X_w is

$$f_w(x) = \frac{\frac{b^{a-c}}{x^{a-c}} - \frac{b^a}{x^a}}{\frac{(a-c)b}{a-c-1} - \frac{ab}{a-1}} = \frac{a-1}{c} \frac{(a-c-1)b^{a-c-1}}{x^{a-c}} \left(1 - \frac{b^c}{x^c}\right), \quad x > b.$$

Keeping in mind the definition of weighed exponential distributions in the Example 2.2, an alternative representation of $f_w(x)$ is

$$f_w(x) = \frac{\frac{c}{a-c-1} + 1}{\frac{c}{a-c-1}} \frac{(a-c-1)b^{a-c-1}}{x^{a-c}} \left[1 - \left(\frac{b}{x}\right)^{\frac{c}{a-c-1}(a-c-1)}\right], \quad x > b,$$

and hence, in a similar way as Gupta and Kundu (2009), we can say that X_w follows a weighed Pareto type I distribution with parameters $c/(a - c - 1)$ and $a - c - 1$. The mode and the mean of X_w are

$$\tilde{m}_w = [\mathbb{E}(X^c)]^{1/c} = \left(\frac{a}{a-c}\right)^{1/c} b$$

and

$$\mathbb{E}(X_w) = \frac{(a-1)(a-c-1)}{(a-2)(a-c-2)} b,$$

respectively. Furthermore, the ratios

$$\frac{f_w(x)}{f(x)} = \frac{(a-c-1)(a-1)}{acb^{a+1}} (b^{a-c} x^{a+1} - b^a x)$$

and

$$\frac{f_w(x)}{f_w(x)} = \frac{(a-c-1)(a-1)}{(a-c)cb^{a-c+1}} (b^{a-c} x - b^a x^{-c+1})$$




Figure 4. The pdf $f_{\tilde{w}}(x)$ given in Example 2.3 for $a = 5, b = 1, c = 2$ and $0 \leq x \leq 7$.

are increasing functions. Hence, $X \leq_{lr} X_w \leq_{lr} X_{\tilde{w}}$. In Figure 4, we drew the pdf $f_{\tilde{w}}(x)$ for $a = 5, b = 2$ and $c = 2$, giving also the values $\tilde{m}_w = \sqrt{5/3}$ and mean of $\mathbb{E}(X_{\tilde{w}}) = 8/3$.

3. A generalization

By using a generalization of (2.2), we obtain a large class of weighted premiums, which includes the weighted premiums proposed by Furman and Zitikis (2008). In fact, let $w : [0, \infty) \rightarrow [0, \infty)$ and $v : [0, \infty) \rightarrow [0, \infty)$ two continuous functions such that $0 < \mathbb{E}[w(X)] < \infty$ and $0 < \mathbb{E}[v(X)] < \infty$. Furthermore, assume that w is increasing, v is nondecreasing and $\xi(x) = w(x)/v(x)$ is increasing and convex function. Let also consider a nonnegative weighted random variable $X_{\tilde{w}}$ of X with pdf

$$f_{\tilde{w}v}(x) = \frac{\tilde{w}v(x)}{\mathbb{E}[\tilde{w}v(X)]} f(x) = \frac{\bar{F}_w(x) - \bar{F}_v(x)}{\mathbb{E}(X_w) - \mathbb{E}(X_v)},$$

provided that $\mathbb{E}(X_w) > \mathbb{E}(X_v) > 0$. Here, the weight $\tilde{w}v(x)$ is

$$\tilde{w}v(x) = \frac{\bar{F}_w(x) - \bar{F}_v(x)}{f(x)}.$$

The mode of $X_{\tilde{w}v}$ follows by solving the equation

$$\frac{w(x)}{\mathbb{E}[w(X)]} = \frac{v(x)}{\mathbb{E}[v(X)]},$$

or equivalently,

$$\xi(x) = \frac{\mathbb{E}[w(X)]}{\mathbb{E}[v(X)]}. \tag{3.1}$$

Since the function $\xi(x) = w(x)/v(x)$ is increasing (by assumption), the solution of Equation (3.1) is unique. Denote by $m_{X_{\tilde{w}v}} = \tilde{m}_\xi$ the unique mode of $X_{\tilde{w}v}$. Then the generalized mode premium is defined by:

$$T_\xi(X) := \tilde{m}_\xi = \xi^{-1}\left(\frac{\mathbb{E}[w(X)]}{\mathbb{E}[v(X)]}\right). \tag{3.2}$$

Proposition 3.1. Under the assumptions given above for the functions $w(x)$, $v(x)$ and $\xi(x) = w(x)/v(x)$, it holds that $\tilde{m}_\xi \geq \mathbb{E}(X)$.

Proof. Since $\xi(x)$ is increasing and $v(x)$ is nondecreasing, we have

$$\frac{\mathbb{E}[w(X)]}{\mathbb{E}[v(X)]} = \frac{\mathbb{E}[\xi(X) v(X)]}{\mathbb{E}[v(X)]} \geq \mathbb{E}[\xi(X)]. \tag{3.3}$$

Furthermore, recalling that $\xi(x)$ is convex, applying Jensen’s inequality, it follows

$$\mathbb{E}[\xi(X)] \geq \xi(\mathbb{E}(X)). \tag{3.4}$$

By (3.3) and (3.4), we get

$$\frac{\mathbb{E}[w(X)]}{\mathbb{E}[v(X)]} \geq \xi(\mathbb{E}(X)),$$

and keeping in mind that $\xi(x)$ is increasing, we obtain

$$\xi^{-1}\left(\frac{\mathbb{E}[w(X)]}{\mathbb{E}[v(X)]}\right) \geq \mathbb{E}(X),$$

which completes the proof. □

Remark 3.1.

- (i) For $v(x) = a$, where a is a positive constant, we have $\tilde{m}_\xi = \tilde{m}_w$.
- (ii) For $w(x) = x v(x)$, we have that \tilde{m}_ξ is the unique solution of the equation:

$$\frac{x v(x)}{\mathbb{E}[X v(X)]} = \frac{v(x)}{\mathbb{E}[v(X)]}.$$

Therefore,

$$\tilde{m}_\xi = \frac{\mathbb{E}[X v(X)]}{\mathbb{E}[v(X)]};$$

for more details, see Furman and Zitikis (2008).

- (iii) For $w(x) = e^{\lambda x} v(x)$, we have that \tilde{m}_ξ is the unique solution of the equation:

$$\frac{e^{\lambda x} v(x)}{\mathbb{E}[e^{\lambda X} v(X)]} = \frac{v(x)}{\mathbb{E}[v(X)]}.$$

Therefore, we obtain the premium:

$$\tilde{m}_\xi = \frac{1}{\lambda} \ln \frac{\mathbb{E}[e^{\lambda X} v(X)]}{\mathbb{E}[v(X)]}, \tag{3.5}$$

which is a generalization of the exponential premium principle. In the following section, we study further this generalization for $v(x) = e^{\gamma x}$ (where $\lambda > \gamma > 0$).

- (iv) An analogous interpretation as the one given in Remark 2.1 for the density $f_{\tilde{w}}$ is the following. Applying a probabilistic analogue of the mean value theorem between X_v and X_w , we obtain the new random variable $X_{\tilde{w}}$ with pdf $f_{\tilde{w}}$.

3.1. A generalization of exponential premium principle

For $\lambda > \gamma > 0$, we consider the functions $w(x) = e^{\lambda x}$ and $v(x) = e^{\gamma x}$, and hence, $\xi(x) = e^{(\lambda-\gamma)x}$. By (3.2) (or (3.5)), it follows that

$$\tilde{m}_\xi = \frac{1}{\lambda - \gamma} [\ln \mathbb{E}(e^{\lambda X}) - \ln \mathbb{E}(e^{\gamma X})], \tag{3.6}$$

where \tilde{m}_ξ is the (unique) mode of X_{wv} with pdf

$$f_{wv}(x) = \frac{\int_x^\infty \frac{e^{\lambda y}}{\mathbb{E}(e^{\lambda X})} f(y) dy - \int_x^\infty \frac{e^{\gamma y}}{\mathbb{E}(e^{\gamma X})} f(y) dy}{\frac{\mathbb{E}(X e^{\lambda X})}{\mathbb{E}(e^{\lambda X})} - \frac{\mathbb{E}(X e^{\gamma X})}{\mathbb{E}(e^{\gamma X})}}.$$

Theorem 3.1. *The mode \tilde{m}_ξ of X_{wv} given in (3.6) satisfies the following properties:*

(i) *A two-side bound is*

$$\frac{\mathbb{E}[X e^{\gamma X}]}{\mathbb{E}[e^{\gamma X}]} \leq \tilde{m}_\xi \leq \frac{\mathbb{E}[X e^{\lambda X}]}{\mathbb{E}[e^{\lambda X}]} \tag{3.7}$$

(ii) *It holds that*

$$\frac{d\tilde{m}_\xi}{d\gamma} \geq 0 \text{ and } \frac{d\tilde{m}_\xi}{d\lambda} \geq 0.$$

(iii) *For $\gamma \rightarrow 0$ and $\gamma \rightarrow \lambda$, we have*

$$\lim_{\gamma \rightarrow 0} \tilde{m}_\xi = \frac{1}{\lambda} \ln \mathbb{E}[e^{\lambda X}] \text{ and } \lim_{\gamma \rightarrow \lambda} \tilde{m}_\xi = \frac{\mathbb{E}[X e^{\lambda X}]}{\mathbb{E}[e^{\lambda X}]},$$

respectively.

Proof.

(i) We have

$$\begin{aligned} \tilde{m}_\xi &= \frac{1}{\lambda - \gamma} [\ln \mathbb{E}(e^{\lambda X}) - \ln \mathbb{E}(e^{\gamma X})] \\ &= \frac{\lambda}{\lambda - \gamma} \frac{1}{\lambda} \int_0^\lambda \frac{\mathbb{E}(X e^{sX})}{\mathbb{E}(e^{sX})} ds - \frac{\gamma}{\lambda - \gamma} \frac{1}{\gamma} \int_0^\gamma \frac{\mathbb{E}(X e^{sX})}{\mathbb{E}(e^{sX})} ds \\ &= \frac{1}{\lambda - \gamma} \int_\gamma^\lambda \frac{\mathbb{E}(X e^{sX})}{\mathbb{E}(e^{sX})} ds. \end{aligned}$$

Since, $\frac{\mathbb{E}(X e^{sX})}{\mathbb{E}(e^{sX})}$ is nondecreasing in s , the two-side bound in (3.7) for \tilde{m}_ξ follows.

(ii) After some computation, we get

$$\frac{d\tilde{m}_\xi}{d\gamma} = \frac{1}{\lambda - \gamma} \left[\tilde{m}_\xi - \frac{\mathbb{E}[X e^{\gamma X}]}{\mathbb{E}[e^{\gamma X}]} \right]$$

and

$$\frac{d\tilde{m}_\xi}{d\lambda} = \frac{1}{\lambda - \gamma} \left[\frac{\mathbb{E}[X e^{\lambda X}]}{\mathbb{E}[e^{\lambda X}]} - \tilde{m}_\xi \right].$$

Therefore, by using formula (3.7), the result of part (ii) follows.

(iii) The result follows directly. □

Example 3.1. *Suppose that X follows an exponential distribution with parameter $b > 0$ and pdf $f(x)$ given in (2.5). Consider the following options for the weights $w(x)$ and $v(x)$, when $b > \lambda > \gamma > 0$:*

- (i) $w_1(x) = x v_1(x)$ and $v_1(x) = e^{\gamma x}$,
- (ii) $w_2(x) = x v_2(x)$ and $v_2(x) = e^{\lambda x}$,
- (iii) $w_3(x) = e^{\lambda x}$ and $v_3(x) = e^{\gamma x}$.

Table 2. For $i = 1, 2, 3$, the distribution of X_{w_i} , X_{v_i} , $\tilde{X}_i = {}_d X_{\tilde{w}_i v_i}$ and the mode \tilde{m}_i of \tilde{X}_i .

Weighted risk	Distribution	$\tilde{m}_i, i = 1, 2, 3$
X_{w_1}	Gamma(2, $b - \gamma$)	
X_{v_1}	Exponential($b - \gamma$)	
\tilde{X}_1	Gamma(2, $b - \gamma$)	$\tilde{m}_1 = \frac{1}{b - \gamma}$
X_{w_2}	Gamma(2, $b - \lambda$)	
X_{v_2}	Exponential($b - \lambda$)	
\tilde{X}_2	Gamma(2, $b - \lambda$)	$\tilde{m}_2 = \frac{1}{b - \lambda}$
X_{w_3}	Exponential($b - \lambda$)	
X_{v_3}	Exponential($b - \gamma$)	
\tilde{X}_3	Weighted Exponential($\frac{\lambda - \gamma}{b - \lambda}, b - \lambda$)	$\tilde{m}_3 = \frac{1}{\lambda - \gamma} \ln \frac{b - \gamma}{b - \lambda}$




Figure 5. The pdf $\tilde{f}_1(x) = f_{w_1 v_1}(x)$ (left part – dashed line), $\tilde{f}_2(x) = f_{w_2 v_2}(x)$ (center part - dotted line), and $\tilde{f}_3(x) = f_{w_3 v_3}(x)$ (right part – solid line) given in Example 3.1 for $b = 2, \lambda = 1, \gamma = 0.5$, and $0 \leq x \leq 5$.

For each $i = 1, 2, 3$, we also consider the random variable $\tilde{X}_i = {}_d X_{\tilde{w}_i v_i}$ with pdf $\tilde{f}_i(x) = f_{\tilde{w}_i v_i}(x)$ and its mode \tilde{m}_i , respectively. In Table 2, for $i = 1, 2, 3$, we determine the distribution of the above random risk trough its pdf:

$$f_{w_i}(x) = \frac{w_i(x)}{\mathbb{E}[w_i(X)]} f(x), f_{v_i}(x) = \frac{v_i(x)}{\mathbb{E}[v_i(X)]} f(x) \text{ and } \tilde{f}_i(x) = \frac{\bar{F}_{w_i}(x) - \bar{F}_{v_i}(x)}{\mathbb{E}(X_{w_i}) - \mathbb{E}(X_{v_i})}.$$

We note that

$$\tilde{f}_3(x) = \frac{\bar{F}_{w_3}(x) - \bar{F}_{v_3}(x)}{\mathbb{E}(X_{w_3}) - \mathbb{E}(X_{v_3})} = \frac{b - \gamma}{\lambda - \gamma} (b - \lambda) e^{-(b-\lambda)x} [1 - e^{-(\lambda-\gamma)x}], \quad x > 0,$$

and hence, \tilde{X}_3 follows a Weighted Exponential($\frac{\lambda - \gamma}{b - \lambda}, b - \lambda$); see Example 2.2. Furthermore, we have $\tilde{X}_1 = {}_d X_{w_1}$, $\tilde{X}_2 = {}_d X_{w_2}$, and $\tilde{X}_3 = {}_d X_{w_3} + X_{v_3}$. For $i = 1, 2, 3$, in Table 2, we also provide the mode of \tilde{X}_i ,

$$\tilde{m}_1 = \frac{\mathbb{E}(X e^{\gamma X})}{\mathbb{E}(e^{\gamma X})} = \frac{1}{b - \gamma}, \quad \tilde{m}_2 = \frac{\mathbb{E}(X e^{\lambda X})}{\mathbb{E}(e^{\lambda X})} = \frac{1}{b - \lambda}$$

and

$$\tilde{m}_3 = \frac{1}{\lambda - \gamma} [\ln \mathbb{E}(e^{\lambda X}) - \ln \mathbb{E}(e^{\gamma X})] = \frac{1}{\lambda - \gamma} \ln \frac{b - \gamma}{b - \lambda}.$$

In Figures 5 and 6, we drew the pdf $\tilde{f}_i(x) = f_{\tilde{w}_i v_i}(x)$, for $i = 1, 2, 3, b = 2, \lambda = 1$, and $\gamma = 0.5$. We observe that $\tilde{m}_1 = 0.66667$ (the lower bound in (3.7)), $\tilde{m}_2 = 1$ (the upper bound in (3.7)), and $\tilde{m}_3 = 0.81093$ (the exact value of the generalized exponential premium).




Figure 6. The pdf $\tilde{f}_i(x) = f_{\tilde{w}_i}(x)$ given in Example 3.1 for $i = 1$ (dashed line), $i = 2$ (dotted line), $i = 3$ (solid line), $b = 2$, $\lambda = 1$, $\gamma = 0.5$, and $0 \leq x \leq 2$.

4. Mode ordering for a class of unimodal weighted distributions

Let X and Y be two continuous nonnegative random variables with pdf's f_X and f_Y , distributions F_X and F_Y , and tails $\bar{F}_X = 1 - F_X$ and $\bar{F}_Y = 1 - F_Y$, respectively. We assume that $X \leq_{st} Y$ and $\mathbb{E}(X) < \mathbb{E}(Y)$. Let us also assume that X and Y are unimodal with mode m_X and m_Y , respectively.

Definition 4.1. We say that the mode of X is smaller than the mode of Y , denoted by $X <_{mode} Y$, if $m_X < m_Y$.

Next, we present a class of unimodal weighted distributions for which the mode is the mean value premium principle.

Definition 4.2. Let X_w be a weighted random variable of X , where $w : [0, \infty) \rightarrow [0, \infty)$ is a continuous and increasing function such that $0 < \mathbb{E}[w(X)] < \infty$ and $\mathbb{E}(X) < \mathbb{E}(X_w)$. Consider also the nonnegative weight function

$$\tilde{w}(x) = \frac{\bar{F}_{X_w}(x) - \bar{F}_X(x)}{f_X(x)}.$$

We say that $X_w \in \mathcal{A}_w(F_X)$ if the pdf of X_w is

$$f_{X_w}(x) = \frac{\tilde{w}(x)}{\mathbb{E}[\tilde{w}(X)]} f_X(x) = \frac{\bar{F}_{X_w}(x) - \bar{F}_X(x)}{\mathbb{E}(X_w) - \mathbb{E}(X)}.$$

The mode $m_{X_w} = w^{-1}(\mathbb{E}[w(X)])$ of $X_w \in \mathcal{A}_w(F)$ is unique. Let also $Y_w \in \mathcal{A}_w(F_Y)$, that is, the pdf of Y_w is

$$f_{Y_w}(x) = \frac{\tilde{w}(x)}{\mathbb{E}[\tilde{w}(Y)]} f_Y(x) = \frac{\bar{F}_{Y_w}(x) - \bar{F}_Y(x)}{\mathbb{E}(Y_w) - \mathbb{E}(Y)},$$

where

$$\tilde{w}(x) = \frac{\bar{F}_{Y_w}(x) - \bar{F}_Y(x)}{f_Y(x)}.$$

In the following proposition, we prove (under some assumptions) that $X_w <_{mode} Y_w$.

Proposition 4.1. Let X and Y be two continuous nonnegative random variables with distributions F_X and F_Y , respectively. Let also $X \leq_{st} Y$, and w be a continuous and increasing function such that $0 < \mathbb{E}[w(X)] < \mathbb{E}[w(Y)]$. If $X_{\tilde{w}} \in \mathcal{A}_w(F_X)$ and $Y_{\tilde{w}} \in \mathcal{A}_w(F_Y)$, then $m_{X_{\tilde{w}}} < m_{Y_{\tilde{w}}}$.

Proof. By assumptions, we have $\mathbb{E}[w(X)] < \mathbb{E}[w(Y)]$. Hence, $w^{-1}(\mathbb{E}[w(X)]) < w^{-1}(\mathbb{E}[w(Y)])$, which completes the proof. \square

Remark 4.1. The mode order can be also applied when the nonnegative continuous function $w(x)$ is decreasing such that $\mathbb{E}(X) > \mathbb{E}(X_w)$. In this case, we say that $X_{\tilde{w}} \in \bar{\mathcal{A}}_w(F)$ if the pdf of $X_{\tilde{w}}$ is

$$f_{X_{\tilde{w}}}(x) = \frac{\tilde{w}(x)}{\mathbb{E}[\tilde{w}(X)]} f_X(x) = \frac{\bar{F}_X(x) - \bar{F}_{X_w}(x)}{\mathbb{E}(X) - \mathbb{E}(X_w)},$$

where

$$\tilde{w}(x) = \frac{\bar{F}_X(x) - \bar{F}_{X_w}(x)}{f_X(x)}.$$

By using similar arguments as in the proof of Proposition 4.1, we can obtain that if $X_{\tilde{w}} \in \bar{\mathcal{A}}_w(F_X)$ and $Y_{\tilde{w}} \in \bar{\mathcal{A}}_w(F_Y)$, then $m_{X_{\tilde{w}}} > m_{Y_{\tilde{w}}}$.

Acknowledgments. The author wishes to thank the referees and the editor for their comments, which improved the paper. This research was financially supported by the MSc program in Actuarial Science & Risk Management of the University of Piraeus through the grant C.908: Development of new methodologies in insurance and risk management.

References

Belzunce, F., Martínez-Riquelme, C. and Mulero, J. (2016) *An Introduction to Stochastic Orders*. Amsterdam: Elsevier-Academic Press.

Bühlmann, H. (1980) An economic premium principle. *ASTIN Bulletin*, **11**, 52–60.

Castaño-Martínez, A., López-Blázquez, F., Pigeiras, G. and Sordo, M. (2020) A method for constructing and interpreting some weighted premium principles. *ASTIN Bulletin*, **50**(3), 1037–1064.

Di Crescenzo, A. (1999) A probabilistic analogue of the mean value theorem and its applications to reliability. *Journal of Applied Probability*, **36**, 706–719.

Denneberg, D. (1994) *Non-Additive Measure and Integral*. Dordrecht: Kluwer.

Denuit, M., Dhaene, J., Goovaerts, M. and Kaas, R. (2006) *Actuarial Theory for Dependent Risks: Measures, Orders and Models*. Hoboken, New Jersey, USA: John Wiley & Sons.

Furman, E. and Zitikis, R. (2008) Weighted premium calculation principles. *Insurance: Mathematics and Economics*, **42**, 459–465.

Gerber H.U. (1979) *An Introduction to Mathematical Risk Theory*. Huebner Foundation Monograph 8, distributed by Richard D. Irwin, Homewood Illinois.

Gerber, H. and Shiu, E.S.W. (1994) Option pricing by Esscher transforms. *Transactions of the Society of Actuaries*, **46**, 99–191.

Goovaerts, M.J., de Vylder, F. and Haezendonck, J. (1984) *Insurance Premiums: Theory and Applications*. Amsterdam: North-Holland.

Gupta, R.D. and Kundu, D. (2009) A new class of weighted exponential distributions. *Statistics*, **43**(6), 621–634.

Kamps, U. (1998) On a class of premium principles including the Esscher premium. *Scandinavian Actuarial Journal*, **1**, 75–80.

Laeven, R.J.A. and Goovaerts, M.J. (2008) Premium calculation and insurance pricing. In *Encyclopedia of Quantitative Risk Analysis and Assessment*. Wiley & Sons.

Müller, A. and Stoyan, D. (2002) *Comparison Methods for Stochastic Models and Risks*. Chichester: John Wiley and Sons.

Newton, L., Bowers, Jr., Gerber H.U., Hickman, J.A., Jones, D.A. and Nesbitt, C.J. (1986) *Actuarial Mathematics*. Itasca, Illinois: The Society of Actuaries.

Pearson, K. (1895) Contributions to the mathematical theory of evolution. II. Skew variation in homogeneous material. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, **186**, 343–414.

Psarrakos, G. (2022) How a probabilistic analogue of the mean value theorem yields Stein-type covariance identities. *Journal of Applied Probability*, **59**, 350–365.

Ruiz, J.M. and Navarro, J. (1994) Characterization of distributions by relationships between failure rate and mean residual life. *IEEE Transactions on Reliability*, **43**(4), 640–644.

Shaked, M. and Shanthikumar, J.G. (2007) *Stochastic Orders*. New York: Springer.

Wang, S.S. (1995) Insurance pricing and increased limits ratemaking by proportional hazards transforms. *Insurance: Mathematics and Economics*, **17**, 43–54.

Wang, S.S. (1996) Premium calculation by transforming the layer premium density. *ASTIN Bulletin*, **26**(1), 71–92.

Wang, S.S. (2003) Equilibrium pricing transforms: New results using Bühlmann’s 1980 economic model. *ASTIN Bulletin*, **33**(1), 57–73.