

# The European sea bass *Dicentrarchus labrax* in the Dutch Wadden Sea: from visitor to resident species

J.F.M.F. CARDOSO<sup>1,2</sup>, V. FREITAS<sup>1,2</sup>, I. QUILEZ<sup>1</sup>, J. JOUTA<sup>1,3</sup>, J.IJ. WITTE<sup>1</sup> AND H.W. VAN DER VEER<sup>1</sup>

<sup>1</sup>NIOZ – Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands, <sup>2</sup>CIIMAR/CIMAR – Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Rua dos Bragas 289, P 4050-123 Porto, Portugal, <sup>3</sup>RUG – University of Groningen, P.O. Box 11103, 9700 CC Groningen, the Netherlands

*This paper analyses the population dynamics, growth and feeding ecology of *Dicentrarchus labrax* in order to gain a better understanding of its present role in the western Dutch Wadden Sea ecosystem. Otolith analysis showed that the population is mostly comprised of individuals aged 3–5 years old and between 20 and 45 cm in length. In autumn, 0-group juveniles are also an important part of the population. Both juveniles and adults use the area as a feeding ground exhibiting an opportunistic feeding strategy that relies on available prey, especially the brown shrimp *Crangon crangon*. Stomach content analysis and nitrogen stable isotope analysis showed an ontogenetic shift towards piscivory and a general decrease in the dominance of invertebrates with increasing size. Over the last 50 years, large between-year fluctuations in *D. labrax* abundance have been observed with an underlying increasing trend from about 1990 until 2007 followed by a subsequent decline. Spring abundance showed significant relationships with temperature and salinity while autumn abundance was only related to temperature. Spring and autumn *D. labrax* abundance were also strongly related to abundance of brown shrimp *C. crangon* prey. Long-term trends in temperature and salinity in the area suggest that environmental conditions for juvenile growth have become optimal, resulting in increased abundance since the mid-1980s. Continued monitoring of the dynamics of this species in the Dutch Wadden Sea is important to understand and anticipate the effects of climate change on the *D. labrax* population and its role in the local food web.*

**Keywords:** population dynamics, long-term abundance trend, growth, age, feeding ecology, sea bass

Submitted 27 June 2014; accepted 15 October 2014; first published online 24 November 2014

## INTRODUCTION

The European sea bass *Dicentrarchus labrax* (Linnaeus, 1758) (Moronidae) is a demersal species of high commercial and recreational value in north-east Atlantic waters and in the Mediterranean. *Dicentrarchus labrax* inhabits estuaries, lagoons, coastal waters and occasionally rivers (Smith, 1990; Kottelat & Freyhof, 2007), and is found from Senegal to Norway, including the Mediterranean and the Black Sea (Smith, 1990; FAO, 2013; Froese & Pauly, 2013). Larvae develop in continental shelf waters and then migrate to sheltered coastal or estuarine areas where juveniles spend the next few years of life (Holden & Williams, 1974; Kelley, 1988; Pickett & Pawson, 1994). Estuaries are important nursery areas for *D. labrax* due to their high productivity and favourable environmental conditions for juvenile growth (Boeuf & Lasserre, 1978; Costa & Bruxelas, 1989).

The biology and ecology of *D. labrax* have been extensively studied in several Atlantic coastal areas and the Mediterranean, especially in the UK (e.g. Kennedy & Fitzmaurice, 1972; Holden & Williams, 1974; Claridge & Potter, 1983;

Aprahamian & Barr, 1985; Kelley, 1988, 2002; Pickett & Pawson, 1994; Henderson & Corps, 1997), French coast (Lam Hoai, 1970; Stequert, 1972; Barnabé, 1976, 1978; Fritsch, 2005; Dufour *et al.*, 2009) and Portuguese coast (Costa & Bruxelas, 1989; Cabral & Costa, 2001; Martinho *et al.*, 2008; Vasconcelos *et al.*, 2009; Vinagre *et al.*, 2009, 2012c). Contrasting results regarding recruitment, growth and condition of *D. labrax* in these areas illustrate the importance of population- and area-specific research to understand the effects of biotic and environmental variables on the population dynamics of this species across its distributional range. In Dutch waters, *D. labrax* abundance has shown a continuous increase from the mid-1980s to mid-2000s, both in the coastal zone as well as in estuarine areas (Tulp *et al.*, 2008). However, information regarding its population structure and dynamics in these areas is scarce and limited to reports (Kroon, 2007; Bierman *et al.*, 2010).

We report here on the growth, feeding ecology and abundance of *D. labrax* in the western Dutch Wadden Sea. The Dutch Wadden Sea is part of an international wetland ecosystem that has been designated as a UNESCO World Heritage Site in 2009. With 4700 km<sup>2</sup> of intertidal flats it is one of the world's largest intertidal systems. The western Dutch Wadden Sea is recognized as an important nursery and feeding ground for several fish species (Zijlstra, 1972; Bergman *et al.*, 1988, 1989; Van der Veer *et al.*, 1991, 2001)

**Corresponding author:**  
J.F.M.F. Cardoso  
Email: joanafcardoso@portugalmail.com

and serves as a breeding and wintering area for millions of migratory birds (Reise *et al.*, 2010). Given the recent colonization of the western Dutch Wadden Sea by *D. labrax* (Tulp *et al.*, 2008), the need arose to study the position of this species in the local food web to gain a better understanding of its present role in the Wadden Sea ecosystem. Since growth and feeding are two important aspects influencing population dynamics, an integrative approach combining information from multiple sources dealing with long-term abundance patterns, growth rates, sex-specific size and age distribution, stomach content and stable isotope analysis was used. In addition, in order to understand the fluctuations in *D. labrax* abundance in the area over the course of time, the effect of a number of abiotic (salinity, temperature) and biotic (prey abundance) variables was examined using available time-series.

## MATERIALS AND METHODS

### Data acquisition

Data used by this study are from two sources: the NIOZ fyke net programme and the beach seine surveys carried out in the western Dutch Wadden Sea (Figure 1). The NIOZ fyke net programme started in 1960 and consists of a passive fishing kom-fyke net with a stretched mesh-size of 20 mm near the southern point of Texel, at the entrance of the Dutch Wadden Sea which is deployed annually. This trap is a combination of a pound net and a fyke with a 200 m leader running from above high water into the subtidal where two end chambers collect the fish. The tidal range is between

1–2 m depending on tidal phase and weather conditions. Fishing is usually undertaken from April to June, interrupted in summer because of jellyfish and weed clogging, and then restarted in September until the end of October. In winter, when ice forms, the fyke is removed. Until 1995, the fyke was emptied every Monday to Friday morning. From 1995 onwards the fyke was emptied every day. Occasionally, when catches are low, the net was emptied every other morning. All fyke net catches were sorted immediately, identified to species and counted. Between 1970 and 1978, fyke catches were measured irregularly, with a frequency of less than once a week. From 1979 onwards, all fish caught were measured to the nearest cm total length (TL). Since 2005, catch sub-samples were analysed in more detail including sex, reproductive stage, diet and age. In 2011, fish samples were also collected for stable isotope analysis. The validity of data from a single fish trap for the study of long-term trends in fish populations relies on the assumption that, on average, a constant fraction of the study populations is sampled. This assumption was validated for most species by comparing contemporaneous trends in a number of traps in the study area (Van der Veer *et al.*, 1992; Van der Meer *et al.*, 1995).

Beach seine surveys were carried out in 2010 and 2011 to collect information on young-of-the-year *D. labrax*. Sampling was done around low water in the Amsteldiep tidal gully surrounding the Balgzand intertidal (Figure 1). At each sampling date four hauls were made from the low water mark to a depth of about 1.2 m and covering a surface area of about 60 m<sup>2</sup>. All fish collected were sorted immediately, counted and measured.

### Age and growth

Annual growth increments on *D. labrax* otoliths collected from 2005 onwards were identified and counted in order to estimate individual age. The sagittal otolith pair was removed, cleaned and either the left or right otolith was chosen for analysis depending on the clarity of the increment pattern. The selected otolith was submerged in a black vat filled with water with the sulcus facing down, and observed under a stereomicroscope using reflective light. Annual rings (annuli) were identified as the combination of opaque (indicating spring/summer growth) and translucent (indicating autumn/winter growth) bands alternating around a central nucleus. Counts were made by two readers and whenever the age obtained from two readings differed, the otolith was re-examined by both readers to make a final decision. In cases of disagreement, the otolith was excluded from further analyses. The transition between age groups was set as the 1st of January.

Growth curves were fitted to length-at-age data with the Von Bertalanffy growth model (VBGM) using the traditional version of the model developed by Von Bertalanffy with modifications by Beverton (1954) and Beverton & Holt (1957) (Cailliet *et al.*, 2006):

$$L_t = L_\infty \times (1 - e^{-K \times (t - t_0)}) \quad (1)$$

whereby  $L_\infty$  is the estimated maximum total length (cm),  $K$  is the growth rate constant ( $y^{-1}$ ),  $t$  is age (y),  $t_0$  is the age at length zero and  $L_t$  is the observed length at age  $t$ .

Growth was examined separately for females, males and all individuals together (including both sexes and immature




Fig. 1. Sampling areas in the Western Dutch Wadden Sea. Location of the NIOZ fyke net (■) near the southern tip of Texel, tidal gullies where beach seine hauls (▲) were performed and, the jetty used for measuring water temperatures (●).

specimens). In all specimens the gonad was inspected and, if possible, sex was determined. For analysis of sex-specific growth differences in model parameters, different models were compared. Sex-specific models (i.e. separate parameter estimates for males and females) were compared with models with one parameter in common (either  $L_\infty$  or  $K$  were the same for both sexes) and with a common model (same parameter estimates for both groups). All growth models were fitted using the R package version 3.1.0 programming environment (R Core Team, 2014), with parameters estimated using nonlinear least-squares (nls). An attempt to iteratively estimate all parameters was done resulting in a very large  $L_\infty$  (214 cm) which was unrealistic according to reports on maximum length (Pickett & Pawson, 1994; Freitas *et al.*, 2010). Therefore,  $t_0$  was fixed to day 59 (0.16 years), corresponding to a birth date of 1st of March based on the spawning season of *D. labrax* described for the UK (Thompson & Harrop, 1987; Jennings & Pawson, 1992; Pawson & Pickett, 1996). For each growth model, the parameters, their corresponding standard error (SE) and the lower and upper limits of the 95% confidence intervals and 95% prediction intervals were estimated. Model goodness of fit, comparison and selection were based on the Akaike information criterion (AIC), with the best model defined as the one having the lowest AIC value (Katsanevakis, 2006).

Comparison of growth parameters of the Von Bertalanffy curve with those described for other *D. labrax* populations were performed using the growth performance index  $\Phi$  ( $\Phi = \log_{10}K + 2 \times \log_{10}L_\infty$ ) described by Pauly & Munro (1984).

## Feeding ecology

Feeding patterns were studied based on stomach content and stable isotope analyses. For diet analyses, a subset of the fish collected in the fyke net (from 2005 to 2011) and in the beach seine surveys was used. Stomachs were removed and each prey item was identified to the lowest possible taxon, counted and measured for total length (TL). No adjustment of prey TL due to partial digestion was performed. For subsequent analyses, prey taxa were pooled into higher taxonomic groups, as well as unidentified or digested material.

The relative importance of each prey item in *D. labrax* diet was assessed using standard indices including the numerical index (NI), calculated as the percentage of the number of individuals of a prey category divided by the total number of individuals of all prey items combined, and the occurrence index (OI) estimated as the percentage of each prey item in all non-empty stomachs (Hyslop, 1980). Diet composition in terms of weight was not assessed due to a lack of systematic weighting of each prey item. To evaluate ontogenetic changes in the diet composition of *D. labrax*, dietary indices were calculated in relation to fish length grouped in 5 cm size classes.

Predator-prey relationships were further examined for the most abundant prey using quantile regression. To assess the relationship between minimum, median and maximum prey size consumed with *D. labrax* length, regressions on the 5, 50 and 95% quantiles were performed (Scharf *et al.*, 2000). The analysis was implemented in R using the 'quantreg' package (Koenker, 2013).

For nitrogen stable isotope analysis (SIA), tissue samples from both fyke and beach seines catches were collected in 2011. Fifty individuals covering the entire size range captured

(4–53 cm TL) were selected and, from each fish, a portion of the white dorsal muscle tissue was removed, cleaned of bones and scales, and freeze-dried in a glass vial for more than 48 h. Dried samples were then ground with a mortar and pestle to a fine and uniform powder. Aliquots of homogenized tissue (between 0.4–0.8 mg dry sample) were weighed to the nearest 0.001 mg and packed in tin cups in preparation for SIA. Isotopic ratios were measured in duplicates using a Flash 2000 elemental analyser (EA) coupled to a DeltaV Advantage-irmMS (Thermo Scientific). Stable isotope ratios were expressed in the  $\delta$  unit notation in units per millilitre according to:

$$\delta^{15}\text{N}(\text{‰}) = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 1000 \quad (2)$$

where  $R_{\text{sample}}$  and  $R_{\text{standard}}$  are the ratios of heavy to light isotopes ( $^{15}\text{N}/^{14}\text{N}$ ) of sample or standard, respectively.  $\delta^{15}\text{N}$  is reported relative to atmospheric  $\text{N}_2$ . The analytical precision for  $\delta^{15}\text{N}$  based on repeated measurements of a laboratory standard (acetanilide) was  $\pm 0.26\text{‰}$  ( $n = 60$ ). In order to test for ontogenetic variability in isotopic composition, Gaussian linear and additive models were applied using *D. labrax* length as an explanatory variable. Statistical tests were performed in R with a significance threshold fixed at 5%.

## Annual trends in abundance

Daily catches of the NIOZ fyke were used to calculate the total numbers of *D. labrax* caught per year during spring (April–June) and autumn (September–October). Generalized additive modelling (GAM) (Hastie & Tibshirani, 1990; Faraway, 2006; Zuur *et al.*, 2007) was used to describe the temporal trends in abundance and to compare between seasons (spring and autumn). Poisson GAMs were fitted to the abundance data using the mgcv (Wood, 2006) package in R. A quasi-Poisson error structure was employed to account for overdispersion (c.f. McCullagh & Nelder, 1989), as the dataset comprised a large number of observations with very low abundances.

Two models were compared, whereby the explanatory variables considered were season (nominal) and year (fitted as a smoother):

$$\text{Abundance}_i = \alpha + \text{offset}(\text{LogF}_i) + f(\text{year}_i) + \text{factor}(\text{season}_i) + \varepsilon_i \quad (3)$$

$$\text{Abundance}_i = \alpha + \text{offset}(\text{LogF}_i) + f_1(\text{year}_i) + f_2(\text{year}_i) \times \text{factor}(\text{season}_{\text{autumn},i}) + \text{factor}(\text{season}_i) + \varepsilon_i \quad (4)$$

where  $\alpha$  is an intercept,  $f$  is the smoothing function and  $\varepsilon_i$  is the residual. Model (3) assumes that both seasons have the same abundance-year relationship, i.e. one smoothing curve is used for both seasons. In model (4), the first smoother ( $f_1(\text{year}_i)$ ) represents the overall year effect in both seasons while the second smoother ( $f_2(\text{year}_i)$ ) represents the deviation in autumn from the overall abundance-year relationship. To adjust for differences in fishing effort between years and seasons, the natural log-transformed number of days fished ( $\text{LogF}_i$ ) was used as an offset variable. Because in 1967 no fishing was done in autumn, this year was excluded from

the analysis. The smooth functions were constructed as penalized (thin plate) regression splines, with automatic selection of the effective degrees of freedom during the fitting process (Wood, 2006). The effects of temporal autocorrelation were investigated by analysing the residual autocorrelation function (ACF). No temporal autocorrelation was detected in any of the models. An *F*-test (Wood, 2006) was applied to compare the two models.

### Abiotic and biotic variables

Time series of abiotic and biotic variables were used to examine whether *D. labrax* abundance was related to changes in environmental variables and food availability. Daily water temperature and salinity were obtained from a long-term monitoring program at the NIOZ sampling jetty, located in the vicinity (<1 km) of the NIOZ fyke (Van Aken, 2003). From the daily temperature and salinity measurements, an index for winter and summer was calculated respectively as the mean of December, January and February, and the mean of July and August (Supplementary Material Figure S1). Indices of spring and autumn abundance of the main preys of *D. labrax* (based on diet analysis) were determined as described above for the NIOZ fyke net catches (Supplementary Material Figure S2).

The influence of temperature, salinity and prey abundance on *D. labrax* spring and autumn abundances, from 1961–2011, was analysed with generalized additive modelling, similarly as described above. The general model structure for each season was:

$$\text{Abundance}_i = \alpha + \text{offset}(\text{Log}F_i) + \sum_k f_k(Z_{ki}) + \varepsilon_i \quad (5)$$

where  $f_k$  are smooth functions of the explanatory variables, and  $Z_{ki}$  is the value of the explanatory variable  $k$  per observation  $i$ . Analysis for *D. labrax* spring abundances considered seawater temperature and salinity from the previous winter and the previous summer, and prey abundance in spring. Analysis for *D. labrax* autumn abundances considered seawater temperature and salinity from the summer and the previous winter, and prey abundance in autumn. Correlations between explanatory variables (collinearity) were assessed by multiple pair-wise scatter plots (pairplots). A forward selection procedure was used (following Zuur *et al.*, 2009), whereby the starting point was a GAM with only one explanatory variable, which was fitted  $x$  times depending on the number of different variables (e.g. six times for spring abundances). Different models were compared in terms of the generalized cross-validation (GCV) scores (Wood, 2006). The model with the lowest GCV was then used in a next GAM with two explanatory variables, by adding each of the remaining variables to the model (resulting in e.g. five models with two variables for spring). This selection procedure was continued until a combination was found which resulted in higher GCV than the previous (simpler) model, resulting on the selection of the simpler model. Because for spring the number of smoothers in the final model used more degrees of freedom than the number of observations, the basis dimension parameter  $k$  was set to 6 for all smoothers in the model. The effects of temporal autocorrelation were investigated as described above whereby no autocorrelation was detected in any of the selected models.

## RESULTS

### Age and growth

In total, 335 individuals ranging from 3.6 to 70 cm TL were aged by otolith analysis. The maximum age found was 11 years. Results showed that  $L_\infty$  and  $K$  differed significantly between sexes (the model with the lowest AIC was the general model). Females had a significantly higher  $L_\infty$  (d.f. = 149;  $L_\infty$  = 114.6 cm, 95% C.I.: 95.1–146.6 cm) and a lower  $K$  ( $0.08 \text{ y}^{-1}$ , 95% C.I.:  $0.06\text{--}0.10 \text{ y}^{-1}$ ) than males (d.f. = 154;  $L_\infty$  = 83.5 cm, 95% C.I.: 70.8–101.8 cm;  $K$  =  $0.11 \text{ y}^{-1}$ , 95% C.I.:  $0.09\text{--}0.14 \text{ y}^{-1}$ ) (Figure 2). For comparison with published literature, VBGM parameters were also estimated considering all individuals (males, females, immature and non-identified specimens) which resulted in:  $L_\infty$  = 102.4 cm (95% C.I.: 90.6–121.3 cm) and  $K$  =  $0.09 \text{ y}^{-1}$  (95% C.I.:  $0.07\text{--}0.11 \text{ y}^{-1}$ ) (d.f. = 333). The calculated growth performance index  $\Phi$  was 3.02 for females, 2.88 for males and 2.97 considering all individuals. The largest size caught in the long-term data series was 85 cm (in 1999) but unfortunately no otoliths were retrieved from this animal.

### Feeding ecology

In total, 408 stomachs were analysed from individuals ranging from 2.5 to 70 cm TL. Of all stomachs inspected, 138 (about 34%) were empty. The *D. labrax* diet consisted of 32 prey item types belonging to seven major taxonomic groups, as well as unidentified materials (Figure 3, Supplementary Material Table S1). Decapods were the dominant prey, according to both numerical and occurrence indices, with values of 42 and 59%, respectively. Within this group, the brown shrimp *Crangon crangon* (Linnaeus, 1758) was by far the most important prey (NI = 37%, OI = 49%; Supplementary Material Table S1). Fish (Teleostei) and mysids (Mysidae) were the second and third most important prey groups with similar numerical contributions to the diet although fish occurred more frequently (Figure 3). Of the nine fish species found, herring *Clupea harengus*




Fig. 2. Estimated Von Bertalanffy growth curves for females (back symbols and full line) and males (grey symbols and dashed line).




Fig. 3. Size variation in diet composition of *D. labrax* according to (a) numerical and (b) occurrence indices. Occurrence index values were rescaled to sum 100% across all prey items.

(Linnaeus, 1758) was the most common (NI = 14%, OI = 17%; Supplementary Material Table S1).

Size-related differences in diet were observed and similar patterns were found irrespective of the index considered. The diet of small juvenile *D. labrax* (<10 cm) was dominated by mysids, but this group was practically absent in larger fish. From this size onwards, the contribution of decapods, mainly *C. crangon*, increased reaching 85% of the numerical diet composition in 25–29 cm TL *D. labrax*, thereafter slightly decreasing but still present in about 50% of the stomachs of all larger size classes (Figure 3). The degree of piscivory increased with size. In fish longer than 50 cm TL, plaice *Pleuronectes platessa* (Linnaeus, 1758) was also an important prey (NI: 24–32%, OI: 29–33%).

Figure 4 shows the predation of *D. labrax* on *C. crangon* and *C. harengus* according to size for all stomachs examined. As *D. labrax* increased in size, larger *C. crangon* were preyed upon as indicated by a significant positive relationship between predator and prey sizes (Pearson correlation,  $n = 186$ ,  $r = 0.58$ ,  $P < 0.001$ ). Minimum, median and maximum prey size increased significantly with increasing *D. labrax* length (slope and intercepts of 5; 50; 95% quantile regressions: 0.06 and 0.48; 0.06 and 2.04; 0.08 and 3.25, respectively;  $P < 0.05$ ). No significant difference was found between the slopes

of the 5 and 95% quantiles indicating that both the minimum and maximum *C. crangon* sizes consumed increased at a same rate with predator length. In the case of *C. harengus*, 4–10 cm fish were eaten but no relationship between predator and prey length was found over the range of sizes examined.

The trophic position of *D. labrax* was assessed with  $\delta^{15}\text{N}$  individual signatures. Values ranged from 12.6 to 19.8‰ with an average of 16.5‰ ( $\pm 1.1\%$ ). Values of  $\delta^{15}\text{N}$  increased significantly with body size (ANOVA;  $F_{1,45} = 30.15$ ;  $r^2 = 0.40$ ;  $P < 0.001$ ) indicating higher trophic levels for larger *D. labrax*, consistent with the progressive increase in piscivory observed in the diet (Supplementary Material Figure S3).

### Annual trends in abundance

Abundances of *D. labrax* in spring and autumn are presented in Figure 5, together with model predictions. Maximum abundance in spring was observed in 2005 although high abundances were also observed in 1990, 1999 and from 2001 to 2009. In autumn, highest numbers were caught in 2003 with peaks in 1984, 2001–2004, 2007 and 2011. Minimum spring abundance occurred in 1960 (one individual caught) while in several years (1966, 1970, 1974–1976, 1982 and 1987) no individuals were caught in autumn.




Fig. 4. Relationship between prey length and *D. labrax* total length. For *C. crangon* the quantile regression lines are shown. Solid line: median prey length (50% quantile), dashed lines: minimum (5% quantile) and maximum (95% quantile) prey length. Each symbol represents a single prey eaten by *D. labrax*.

Generalized additive modelling revealed that abundances did not differ significantly between seasons (ANOVA; model (3):  $F_{1,94} = 0.07$ ,  $P = 0.79$ ; model (4):  $F_{1,88} = 0.22$ ,  $P = 0.64$ ). The overall year effect (represented by the same smoother for both seasons) indicated that there was a significant year effect (ANOVA; model (3):  $F_{6,94} = 11.49$ ,  $P < 0.001$ ; model (4):  $F_{7,88} = 8.56$ ,  $P < 0.001$ ) on the abundance of *D. labrax*. However, the yearly pattern of abundance was not significantly different between seasons (i.e. the abundance-year pattern in autumn did not significantly differ from the overall pattern in model (4): ANOVA,  $F_{7,88} = 0.63$ ,  $P = 0.72$ ). Overall, *D. labrax* abundances showed an increasing trend, although a decrease was observed from 2007 onwards (Figure 5). Comparison of models (3) and (4) found no significant differences




Fig. 5. *Dicentrarchus labrax* abundance in spring and autumn (total numbers of fish caught per season) in the western Dutch Wadden Sea. Full lines are GAM model predictions and dashed lines represent 95% confidence intervals.

(ANOVA,  $F_{7,88} = 1.93$ ,  $P = 0.10$ ), confirming that abundance-year patterns were not different between seasons.

### Abiotic and biotic variables

Based on diet analysis, the spring and autumn abundance of brown shrimp *C. crangon* and herring *C. harengus* were selected as biotic explanatory variables in the *D. labrax* abundance model. For *C. harengus*, a selection of size classes potentially relevant as prey was made based on the size spectrum found in the examined stomachs (up to 10 cm). In the case of *C. crangon*, all size classes present in fyke catches were included in the analyses.

Initial exploratory data analysis on the explanatory variables influencing *D. labrax* abundances in spring and autumn revealed no strong collinearity between the selected explanatory variables ( $r < 0.51$  for spring and  $r < 0.58$  for autumn). The final model for spring included the effects of *C. crangon* spring abundance, temperature in the previous winter and previous summer, and salinity in the previous winter and previous summer (Supplementary Material Table S2). This model explained 88% of the deviance (adjusted  $r^2 = 0.87$ ). Spring abundance showed a marked relationship with seawater temperature and salinity in the previous winter and with shrimp abundance in spring (Figure 6, Table 1). *Dicentrarchus labrax* spring abundance showed an overall increase with *C. crangon* spring abundance. The shape of the smooth function showing a decrease in *D. labrax* abundance between 150 and 400 *C. crangon* in spring followed by an increase is probably due to the fact that very high abundances of *C. crangon* were found in a few years. Winter mean temperatures above 2°C led to an increase in *D. labrax* spring abundance and a peak in *D. labrax* abundance was seen at winter salinities around 28.5. *Dicentrarchus labrax* spring abundance was also associated with seawater temperature and salinity in the previous summer. The relationship between *D. labrax* spring abundance and temperature in the previous summer yielded a wiggly curve showing a general increase in *D. labrax* abundance with increasing temperature. A negative linear relationship was found between *D. labrax* spring abundance and salinity.

The final model for autumn included the effects of *C. crangon* abundance in autumn and temperature in the previous winter (Supplementary Material Table S2). This model explained 79% of the deviance (adjusted  $r^2 = 0.82$ ). Autumn abundance was associated with temperature in the previous winter (Figure 7, Table 1), whereby highest autumn abundances occurred at winter temperatures above 4.5°C. *Dicentrarchus labrax* autumn abundance was also positively associated with high autumn *C. crangon* abundance.

## DISCUSSION

### Population structure

The *D. labrax* population in the western Dutch Wadden Sea is dominated by juveniles older than 1 year and by adults, although in autumn 0-group juveniles are also an important part of the population. Most individuals caught belonged to age classes 3 to 5 years old, comprising a size range between 20 and 45 cm. Although reproduction data has not been analysed in the present study, reports of size at first maturity from




Fig. 6. Partial plots showing smoothers, with 95% confidence limits, fitted to (partial) effects of the predictor variables retained in the final model of spring abundance of *D. labrax* in the western Dutch Wadden Sea. Tick marks in the x-axis represent samples.

UK waters (Pawson *et al.*, 2007) suggest that this size class mostly corresponds to immature individuals. In the UK, *D. labrax* remain in coastal and estuarine nursery habitats until about 36 cm in length (Pawson *et al.*, 1987; Pickett *et al.*, 2004), after which adolescent or mature fish either stay at inshore feeding areas or migrate offshore to pre-spawning and spawning grounds (Pawson *et al.*, 1987). It thus seems that the western Dutch Wadden Sea supports juvenile *D. labrax* growth and the presence of o-group suggests that the area acts as a nursery for this species. The fact that only two 1-year-old animals were sampled is likely a sampling artefact,

due to the large mesh size of the fyke net combined with avoidance of the beach seine. Future sampling should focus on early juvenile stages (o- and I-group). These are sensitive to environmental conditions (Person-Le Ruyet *et al.*, 2004; Vinagre *et al.*, 2012a, b, c) and could therefore give a better indication of changes in the nursery function of this area for the species.

The estimated  $L_{\infty}$  values in this study were in line with predictions for the species using bioenergetics modelling (103 cm

Table 1. GAM results for the selected spring and autumn *D. labrax* abundance models along with their estimated degrees of freedom (Edf), F-statistics and P-values.

Variable	Edf	F	P-value
<i>Spring</i>			
s(shr_ab_sp)	4.81	5.15	< 0.01
s(prev_winterT)	3.39	10.17	< 0.001
s(prev_winterS)	4.62	5.94	< 0.01
s(prev_summerT)	4.61	3.59	< 0.05
s(prev_summerS)	1	4.38	< 0.05
<i>Autumn</i>			
s(shr_ab_au)	6.82	7.69	< 0.01
s(prev_winterT)	7.9	3.65	< 0.01

shr\_ab\_sp: *C. crangon* abundance in spring; prev\_winterT: winter temperature in the previous year; prev\_winterS: winter salinity in the previous year; prev\_summerT: summer temperature in the previous year; prev\_summerS: summer salinity in the previous year; shr\_ab\_au: *C. crangon* abundance in autumn.




Fig. 7. Partial plots showing smoothers, with 95% confidence limits, fitted to (partial) effects of the predictor variables retained in the final model of autumn abundance of *D. labrax* in the western Dutch Wadden Sea. Tick marks in the x-axis represent samples.

**Table 2.** Von Bertalanffy growth parameters  $L_{\infty}$  and  $K$ , and the growth performance index ( $\Phi$ ) for *D. labrax* data from different locations (adapted from Pickett & Pawson, 1994). F refers to females and M to males; data from this paper are in bold.

Area	$L_{\infty}$	$K$	$\Phi$	References
Ireland (F)	69.3	0.14	2.84	Kennedy & Fitzmaurice (1972)
Ireland (M)	67.3	0.14	2.81	Kennedy & Fitzmaurice (1972)
Irish Sea	63.3	0.16	2.8	Fahy <i>et al.</i> (2000)
UK all areas (F)	54.9	0.24	2.87	Pickett & Pawson (1994)
UK all areas (M)	53.8	0.23	2.83	Pickett & Pawson (1994)
UK east coast	55.1	0.38	3.06	Pickett & Pawson (1994)
UK west coast (F)	68.5	0.14	2.82	Kelley (1988)
UK west coast (M)	69.9	0.13	2.79	Kelley (1988)
<b>Dutch Wadden Sea</b>	<b>102</b>	<b>0.09</b>	<b>2.97</b>	<b>This paper</b>
<b>Dutch Wadden Sea (F)</b>	<b>115</b>	<b>0.08</b>	<b>3.02</b>	<b>This paper</b>
<b>Dutch Wadden Sea (M)</b>	<b>84</b>	<b>0.11</b>	<b>2.88</b>	<b>This paper</b>
South Brittany (France) (F)	77.2	0.16	2.98	Bertignac (1988)
South Brittany (France) (M)	76.4	0.15	2.95	Bertignac (1988)
Vendée (France) (sea)	76.9	0.14	2.93	Lam Hoai (1970)
Vendée (France) (lagoons)	84.3	0.13	2.97	Lam Hoai (1970)
Arcachon (France)	45	0.34	2.84	Stequert (1972)
Sète (France) (M)	57	0.38	3.09	Barnabé (1976)
Sète (France) (F)	79	0.24	3.17	Barnabé (1976)
Cadiz (South Spain)	99.3	0.14	3.14	Arias (1980)
Morocco	52.3	0.3	2.92	Gravier (1961)
Egypt	39	0.32	2.69	Rafail (1971)

for females and 87 cm for males; Freitas *et al.*, 2010) and were not far from the maximum observed size in the area of 85 cm (this study). Estimated  $L_{\infty}$  values were higher than the ones reported for populations from other areas (Table 2). The estimated  $K$  value was, on the other hand, lower than described in previous studies but this was expected as the two parameters are inversely related (Pauly, 1984). The index of growth performance  $\Phi$  resulted in a relatively high value (2.97), larger than the ones reported for British and Irish coasts and similar to the one described in more southern Atlantic areas, such as south Brittany and the Bay of Biscay (Table 2). Since no recent data on growth of *D. labrax* are available from other areas it is not possible to assess whether conditions for growth in the western Dutch Wadden Sea are better than in other areas or whether growth conditions have generally improved over the last decades.

### The role of *D. labrax* in the system

As in other areas, *D. labrax* diet in the western Wadden Sea is largely composed of crustaceans. *Dicentrarchus labrax* has been described as an opportunistic feeder, adapting its diet to the most available food sources (Pickett & Pawson, 1994; Laffaille *et al.*, 2000). Such flexible feeding behaviour has been observed in other areas (Arias, 1980; Cabral & Costa, 2001; Martinho *et al.*, 2008). The high dominance of shrimp *C. crangon* in the diet of juvenile *D. labrax* may also be related to its general high abundance and all-year around availability as *C. crangon* is a major component of the mobile epibenthic fauna of the western Wadden Sea tidal flats (Kuipers & Dapper, 1984; Beukema, 1992). The same opportunistic feeding strategy probably explains the dominance of small herring in *D. labrax* diet, as the Wadden Sea is well known as a nursery and feeding ground for *C. harengus* (I. Tulp, personal communication).

Adult *D. labrax* are placed high in the food chain due to their piscivorous diet (Pasquaud *et al.*, 2010; Spitz *et al.*, 2013). An increase in piscivory with *D. labrax* size observed

in the Wadden Sea has been reported in other wild populations (Barnes *et al.*, 2008; Pasquaud *et al.*, 2008) emphasizing the need to consider fish size when evaluating function and trophic position in the food web (Pasquaud *et al.*, 2010).

In many fish species, the larger the predator, the larger the size spectrum of prey consumed (Keast & Webb, 1966; Juanes & Conover, 1994) which seems to be related to anatomical changes in predation mechanisms as well as increased swimming capabilities and better visual acuity (Keast & Webb, 1966; Webb, 1976; Beamish, 1978). Such a relationship was found for *C. crangon* prey where the size of shrimps consumed increased linearly with *D. labrax* length. However, contrary to the widespread pattern of asymmetrical predator size–prey size relationships observed in many fish species (Scharf *et al.*, 2000), both maximum and minimum prey sizes increased in parallel for larger *D. labrax*. This indicates a constant niche breadth, consistent with the traditional foraging theory according to which predators should maximize their net energetic return by selecting the largest available prey (Schoener, 1971; Stephens & Krebs, 1986). From the prey viewpoint, these results indicate that a large part of the *C. crangon* population in the Dutch Wadden Sea is vulnerable to *D. labrax* predation. Fish predation has been identified as the main factor affecting autumn recruitment of the *C. crangon* population in the region (Campos *et al.*, 2010) but the specific impact of *D. labrax* was not assessed.

### Long-term trends in abundance

Over the last 50 years, an overall increasing trend in *D. labrax* in the western Dutch Wadden Sea was found especially from 1990 until 2007, however there were large between-year fluctuations. A similar increasing trend in abundance has been observed along the Dutch coast and in the Western Scheldt estuary in the south (Tulp *et al.*, 2008). Water temperature is a critical factor determining abundance and distribution of fish species, including *D. labrax*, in estuaries (Marshall & Elliott, 1998; Power *et al.*, 2000; Cabral & Costa, 2001;



Attrill & Power, 2002; Henderson *et al.*, 2011; Pasquaud *et al.*, 2012). The increase in mean annual temperature observed since the 1980s in the western Dutch Wadden Sea, as well as along the whole Dutch coastal zone (Van Aken, 2008a, 2010), suggests that temperatures for growth of *D. labrax* may be more favourable in recent years than in the past. During the last decade, mean summer water temperature in the western Dutch Wadden Sea was around 18°C, and in some months even reached 21°C (Van Aken, 2008a; H. van Aken, personal communication), which are optimal for growth of juvenile *D. labrax* (Barnabé, 1990; Vinagre *et al.*, 2012b).

*Dicentrarchus labrax* spring abundances were usually highest when temperatures in the previous year were around or above the average of the whole dataset (4°C in winter and 18°C in summer). A positive relationship was indeed observed between spring abundance and summer and winter temperatures in the previous year. In south-west Britain, higher year-classes occurred in warmer years, whereby spring and summer temperature appeared to determine the winter survival of o-group *D. labrax* and the number of I-group observed the next spring (Henderson & Corps, 1997). Autumn abundances in the western Dutch Wadden Sea also showed a strong relationship with temperature in the previous winter. Higher winter temperatures are known to lead to higher recruitment success the following year (Holmes & Henderson, 1990) and to improve survival of larvae and juvenile *D. labrax* during their first winter (Kelley, 1988; Pawson, 1992).

Salinity is also an important factor structuring estuarine fish communities (e.g. Marshall & Elliott, 1998; Drake *et al.*, 2002; Costa *et al.*, 2007; Henderson *et al.*, 2011; Pasquaud *et al.*, 2012), as a result of different physiological tolerance and habitat preference of species and life stages. Salinity in the previous summer and winter was seen to influence *D. labrax* spring abundance with highest abundance at a salinity of about 28. This is consistent with evidence from laboratory experiments where this species showed consistently better growth at an intermediate salinity of about 26 compared with higher or lower salinities (Johnson & Katavic, 1986). From 1861 to 2003, annual mean salinity decreased in the Dutch Wadden Sea, mainly due to changes in rainfall, river flow and engineering works on inland rivers (Van Aken, 2008b). This suggests that, in addition to more favourable temperatures, also salinity may have become more optimal for growth of *D. labrax*.

A recent decreasing trend (since 2007) in *D. labrax* abundance has been observed in the Dutch Wadden Sea. Such a trend has also been reported for the Bristol Channel, contemporary with a decline in average temperatures (Henderson & Bird, 2010). In the Dutch Wadden Sea, a decrease in temperature has not been observed yet but mean annual salinity has increased since the early 2000s (H. van Aken, personal communication), although the available time series is presently too short to see a clear trend. Moreover, our modelling results suggest that long-term fluctuations in *D. labrax* abundance are also affected by *C. crangon* abundance, as observed for other areas (Kelley, 1987; Costa & Bruxelas, 1989). A decreasing trend in the stock size of *C. crangon* has also been observed after 2000 (Campos *et al.*, 2010; this study), suggesting that a combination of decrease in prey availability and changing environmental conditions may be responsible for the recent declining trend in *D. labrax* abundance. In addition, other factors such as wind speed and direction, depth

and cannibalism (Henderson & Corps, 1997; Cabral & Costa, 2001; Martinho *et al.*, 2009; Pasquaud *et al.*, 2012) have also been proposed to explain variability in *D. labrax* year-to-year recruitment and abundance.

## CONCLUSIONS

This study shows that the western Dutch Wadden Sea has become an important area for *D. labrax*. It is unclear how the *D. labrax* population will develop in the near future and how this will impact their epibenthic and nektonic prey. Ocean warming seems to be the reason for the northerly extension in distribution of several fish species, including *D. labrax* (Brander *et al.*, 2003; Rose, 2005; ICES, 2012). However, the consistent use of coastal and estuarine areas depends on the quality of local feeding and nursery grounds. Climate variability is known to affect fish recruitment, spawning and feeding distributions (Beaugrand *et al.*, 2003; Ottersen *et al.*, 2004; Rose, 2005). Warmer winter temperatures are thought to have lengthened the residence time of *D. labrax* in summer feeding areas and caused *D. labrax* to spawn further away from their main spawning areas (Pawson & Pickett, 1996; Pawson *et al.*, 2007). A warming climate could prolong the residence time of juvenile *D. labrax* in the Dutch Wadden Sea and/or provide suitable overwintering conditions causing higher abundance of adults in autumn and winter. In addition, climate variability is also expected to affect individual prey species with impacts on predator-prey interactions and on the food web (Freitas *et al.*, 2007). A long-term increase in the western Dutch Wadden Sea *C. crangon* population has been observed (Campos *et al.*, 2010; Tulp *et al.*, 2012), which seems to have positively affected food availability for *D. labrax*. Dynamics of both prey and predator populations have changed in recent years (Campos *et al.*, 2010; this study) but analysing cross-correlations requires longer time series for sound conclusions. Combined with variations in the extent of prospective fisheries, the future of the *D. labrax* population in the Dutch Wadden Sea remains largely unknown. However, continued monitoring is an important tool to understand and anticipate *D. labrax* responses to further environmental changes.

## ACKNOWLEDGMENTS

The authors thank Peter Henderson for providing valuable comments on an earlier version of the manuscript.

## FINANCIAL SUPPORT

This study was partly supported by the Netherlands Organization for Scientific Research (NWO) via Project 839.08.241 of the National program Sea and Coastal Research (ZKO). J. Cardoso and V. Freitas were partly funded by Fundação para a Ciência e a Tecnologia (FCT, Portugal) and Fundo Social Europeu (POP/FSE) (grant nos. SFRH/BPD/34773/2007 and SFRH/BPD/75858/2011, respectively). The stable isotope analysis was carried out within the framework of the Waddensleutels project (<http://www.waddensleutels.nl/>) and supported by the Waddenfonds.

## Supplementary materials and methods

The supplementary material for this article can be found at <http://www.journals.cambridge.org/MBI>

## REFERENCES

- Aprahamian M.W. and Barr C.D.** (1985) The growth, abundance and diet of 0-group sea bass, *Dicentrarchus labrax*, from the Severn Estuary. *Journal of the Marine Biological Association of the United Kingdom* 65, 169–180.
- Arias A.** (1980) Crecimiento, régimen alimentario y reproducción de la dorada (*Sparus aurata* L.) y del robalo (*Dicentrarchus labrax* L.) en los esteros de Cádiz. *Investigación Pesquera* 44, 59–83.
- Attrill M.J. and Power M.** (2002) Climatic influence on a marine fish assemblage. *Nature* 417, 275–278.
- Barnabé G.** (1976) *Contribution à la connaissance de la biologie du loup Dicentrarchus labrax (Poisson Serranidae) de la région de Sète*. PhD thesis. Université des Sciences et Techniques de Languedoc, Montpellier, France.
- Barnabé G.** (1978) Étude dans le milieu naturel et au captivité de l'écoéthologie du loup *Dicentrarchus labrax* (L.) (Poisson, Serranidae) à l'aide de nouvelles techniques. *Annales des Sciences Naturelles – Zoologie et Biologie Animale* 20, 423–502.
- Barnabé G.** (1990) Rearing bass and gilthead bream. *Aquaculture* 2, 647–686.
- Barnes C., Jennings S., Polunin N.V. and Lancaster J.E.** (2008) The importance of quantifying inherent variability when interpreting stable isotope field data. *Oecologia* 155, 227–235.
- Beamish F.W.H.** (1978) Swimming capacity of fish. In Hoar W.S. and Randall D.J. (eds) *Fish physiology*. New York, NY: Academic Press, pp. 101–187 [Fish Physiology series, vol. 7].
- Beaugrand G., Brander K.M., Lindley J.A., Souissi S. and Reid P.C.** (2003) Plankton effect on cod recruitment in the North Sea. *Nature* 426, 661–664.
- Bergman M.J.N., Van der Veer H.W., Stam A. and Zuidema J.J.** (1989) Transport mechanisms of larval plaice (*Pleuronectes platessa* L.) from the coastal zone into the Wadden Sea nursery area. *Rapport et Procès-verbaux des Réunions, Conseil international pour l'Exploration de la Mer* 191, 43–49.
- Bergman M.J.N., Van der Veer H.W. and Zijlstra J.J.** (1988) Plaice nurseries: effects on recruitment. *Journal of Fish Biology* 33, 201–218.
- Bertignac M.** (1988) *L'exploitation du bar (Dicentrarchus labrax) dans le Mor Bras (Bretagne Sud)*. Thesis. Publications du Département d'Halieutique no. 7, Ecole nationale supérieure agronomique, Rennes, France.
- Beukema J.J.** (1992) Dynamics of juvenile shrimp *Crangon crangon* in a tidal-flat nursery of the Wadden Sea after mild and cold winters. *Marine Ecology Progress Series* 83, 157–165.
- Beverton R.J.H.** (1954) Notes on the use of theoretical models in the study of the dynamics of exploited fish populations. Miscellaneous Contribution. U.S. Fishery Laboratory, USA, no. 2, 181 pp.
- Beverton R.J.H. and Holt S.J.** (1957) *On the dynamics of exploited fish populations. Fishery Investigations, series II*. London: Ministry of Agriculture, Fisheries and Food, 533 pp.
- Bierman S., Van Overzee H., Van der Hammen T. and Quirijns F.** (2010) Kennisdocument Zeebaars. *IMARES Report C007/10*. Wageningen UR, Wageningen, 23 pp.
- Boeuf G. and Lasserre P.** (1978) Aspects de la régulation osmotique chez le bar juvénile (*Dicentrarchus labrax*) en élevage et introduit dans les lagunes aménagées de certains bassins d'Arcachon (Gironde). *Colloque National Ecotron* 7, 673–688.
- Brander K.M., Blom G., Borges M.F., Erzini K., Henderson G., MacKenzie B.R., Mendes H., Santos A.M.P. and Tøresen P.** (2003) Changes in fish distribution in the eastern North Atlantic: are we seeing a coherent response to changing temperature? *ICES Marine Science Symposium* 219, 260–273.
- Cabral H.N. and Costa M.J.** (2001) Abundance, feeding ecology and growth of 0-group sea bass, *Dicentrarchus labrax*, within the nursery areas of the Tagus estuary. *Journal of the Marine Biological Association of the United Kingdom* 81, 679–682.
- Cailliet G.M., Smith W.D., Mollet H.F. and Goldman K.J.** (2006) Age and growth studies of chondrichthyan fishes: the need for consistency in terminology, verification, validation, and growth function fitting. *Environmental Biology of Fish* 77, 211–228.
- Campos J., Bio A., Cardoso J.F.M.F., Dapper R., Witte J.I. and Van der Veer H.W.** (2010) Fluctuations of brown shrimp *Crangon crangon* abundance in the western Dutch Wadden Sea. *Marine Ecology Progress Series* 405, 203–219.
- Claridge P.N. and Potter I.C.** (1983) Movements, abundance, age composition and growth of bass, *Dicentrarchus labrax*, in the Severn estuary and inner Bristol Channel. *Journal of the Marine Biological Association of the United Kingdom* 63, 871–879.
- Costa M.J. and Bruxelles A.** (1989) The structure of fish communities in the Tagus Estuary, Portugal, and its role as a nursery for commercial fish species. *Scientia Marina* 53, 561–566.
- Costa M.J., Vasconcelos R., Costa J.L. and Cabral H.N.** (2007) River flow influence on the fish community of the Tagus estuary (Portugal). *Hydrobiologia* 587, 113–123.
- Drake P., Arias A.M., Baldo F., Cuesta J.A., Rodríguez A., Sobrino I., Silva-García A., García-González D. and Fernández-Delgado C.** (2002) Spatial and temporal variation of the aquatic nekton and hyperbenthos from a temperate European estuary with a regulated freshwater inflow. *Estuaries* 25, 451–468.
- Dufour V., Cantou M. and Lecomte F.** (2009) Identification of sea bass (*Dicentrarchus labrax*) nursery areas in the north-western Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* 89, 1367–1374.
- FAO** (2013) *Species Fact Sheets: Dicentrarchus labrax (Linnaeus, 1758)*. Available at <http://www.fao.org/fishery/species/2291/en>.
- Fahy E., Forrest N., Shaw U. and Green P.** (2000) Observations on the status of bass *Dicentrarchus labrax* stocks in Ireland in the late 1990s. *Irish Fisheries Investigations*, 5. Dublin Marine Institute, 29 pp.
- Faraway J.J.** (2006) *Extending the linear model with R*, 1st edn. London: Chapman and Hall/CRC Press.
- Freitas V., Campos J., Fonds M. and Van der Veer H.W.** (2007) Potential impact of temperature change on epibenthic predator–bivalve prey interactions in temperate estuaries. *Journal of Thermal Biology* 32, 328–340.
- Freitas V., Cardoso J.F.M.F., Lika K., Peck M.A., Campos J., Kooijman S.A.L.M. and Van der Veer H.W.** (2010) Temperature tolerance and energetics: a dynamic energy budget-based comparison of North Atlantic marine species. *Philosophical Transactions of the Royal Society B* 365, 3553–3565.
- Fritsch M.** (2005) *Traits biologiques et exploitation du bar commun Dicentrarchus labrax (L.) dans les pêcheries françaises de la manche et du golfe de Gascogne*. PhD thesis. University of Bretagne Occidentale, Brest, France.
- Froese R. and Pauly D.** (2013) *FishBase*. Available at <http://www.fishbase.org>.
- Gravier R.** (1961) Les bars (loups) du Maroc atlantique, *Morone labrax* (L.) et *Morone punctatus* (Bloch). *Revue des Travaux de l'Institut des Pêches Maritimes* 25, 281–292.

- Hastie T. and Tibshirani R.** (1990) *Generalized additive models*, 1st edn. London: Chapman and Hall.
- Henderson P.A. and Bird D.J.** (2010) Fish and macro-crustacean communities and their dynamics in the Severn estuary. *Marine Pollution Bulletin* 61, 100–114.
- Henderson P.A. and Corps M.** (1997) The role of temperature and cannibalism in interannual recruitment variation of bass in British waters. *Journal of Fish Biology* 50, 280–295.
- Henderson P.A., Seaby R.M.H. and Somes J.R.** (2011) Community level response to climate change: the long-term study of the fish and crustacean community of the Bristol Channel. *Journal of Experimental Marine Biology and Ecology* 400, 78–89.
- Holden M.J. and Williams T.** (1974) The biology, movements and population dynamics of bass, *Dicentrarchus labrax*, in English waters. *Journal of the Marine Biological Association of the United Kingdom* 54, 91–107.
- Holmes R.H.A. and Henderson P.A.** (1990) High fish recruitment in the Severn Estuary: the effect of a warm year? *Journal of Fish Biology* 36, 961–963.
- Hyslop E.J.** (1980) Stomach contents analysis – a review of methods and their application. *Journal of Fish Biology* 17, 411–429.
- ICES** (2012) Widely distributed and migratory stocks: European seabass in the Northeast Atlantic. *ICES Advice 2012*, 9.4.23. ICES, no. 9, 14 pp.
- Jennings S. and Pawson M.G.** (1992) The origin and recruitment of bass, *Dicentrarchus labrax*, larvae to nursery areas. *Journal of the Marine Biological Association of the United Kingdom* 72, 199–212.
- Johnson D.W. and Katavic I.** (1986) Survival and growth of sea bass (*Dicentrarchus labrax*) larvae as influenced by temperature, salinity, and delayed initial feeding. *Aquaculture* 52, 11–19.
- Juanes F. and Conover D.O.** (1994) Piscivory and prey size selection in young-of-the-year bluefish: predator preference or size-dependent capture success? *Marine Ecology Progress Series* 114, 59–69.
- Katsanevakis S.** (2006) Modelling fish growth: model selection, multi-model inference and model selection uncertainty. *Fisheries Research* 81, 229–235.
- Keast A. and Webb D.** (1966) Mouth and body form relative to feeding ecology in the fish fauna of a small lake, Lake Opinicon, Ontario. *Journal of the Fisheries Research Board of Canada* 23, 1845–1874.
- Kelley D.** (2002) Abundance, growth and first-winter survival of young bass in nurseries of south-west England. *Journal of the Marine Biological Association of the United Kingdom* 82, 307–319.
- Kelley D.F.** (1987) Food of bass in U.K. waters. *Journal of the Marine Biological Association of the United Kingdom* 67, 275–286.
- Kelley D.F.** (1988) Age determination in bass and assessment of growth and year-class strength. *Journal of the Marine Biological Association of the United Kingdom* 68, 179–214.
- Kennedy M. and Fitzmaurice P.** (1972) The biology of the bass, *Dicentrarchus labrax*, in Irish Waters. *Journal of the Marine Biological Association of the United Kingdom* 52, 557–597.
- Koenker R.** (2013) *quantreg: Quantile Regression*. R package version 4.97. Available at <http://CRAN.R-project.org/package=quantreg>.
- Kottelat M. and Freyhof J.** (2007) *Handbook of European freshwater fishes*. Cornol: Publications Kottelat.
- Kroon J.W.** (2007) Kennisdocument zeebaars, *Dicentrarchus labrax* (Linnaeus, 1758). *Sportvisserij Nederland, Bilthoven*, no. 21, 52 pp.
- Kuipers B.R. and Dapper R.** (1984) Nursery function of Wadden Sea tidal flats for the brown shrimp *Crangon crangon*. *Marine Ecology Progress Series* 17, 171–181.
- Laffaille P., Lefeuvre J.C. and Feunteun E.** (2000) Impact of sheep grazing on juvenile sea bass, *Dicentrarchus labrax*, in tidal salt marshes. *Biological Conservation* 96, 271–277.
- Lam Hoai T.** (1970) Contribution a l'étude des bars de la region des Sables d'Olonne. *Travaux de la Faculté des Sciences de Rennes, Série Oceanographie Biologique* 3, 39–68.
- Marshall S. and Elliott M.** (1998) Environmental influences on the fish assemblage of the Humber estuary, UK. *Estuarine, Coastal and Shelf Science* 46, 175–184.
- Martinho F., Dolbeth M., Viegas I., Teixeira C.M., Cabral H.N. and Pardal M.A.** (2009) Environmental effects on the recruitment variability of nursery species. *Estuarine, Coastal and Shelf Science* 83, 460–468.
- Martinho F., Leitão R., Neto J.M., Cabral H., Lagardère F. and Pardal M.A.** (2008) Estuarine colonization, population structure and nursery functioning for 0-group sea bass (*Dicentrarchus labrax*), flounder (*Platichthys flesus*) and sole (*Solea solea*) in a mesotidal temperate estuary. *Journal of Applied Ichthyology* 24, 229–237.
- McCullagh P. and Nelder J.A.** (1989) *Generalized linear models*, 2nd edn. London: Chapman and Hall/CRC Press.
- Ottersen G., Alheit J., Drinkwater K., Friedland K., Hagen E. and Stenseth N.C.** (2004) The responses of fish populations to oceanic climate fluctuations. In Stenseth N.C., Ottersen G., Hurrell J.W. and Belgrano A. (eds) *Marine ecosystems and climate variation. The North Atlantic: a comparative perspective*. Oxford: Oxford University Press, pp. 73–94.
- Pasquaud S., Béguer M., Larsen M.H., Chaalali A., Cabral H. and Lobry J.** (2012) Increase of marine juvenile fish abundances in the middle Gironde estuary related to warmer and more saline waters, due to global changes. *Estuarine, Coastal and Shelf Science* 104–105, 46–53.
- Pasquaud S., Elie P., Jeantet C., Billy I., Martinez P. and Girardin M.** (2008) A preliminary investigation of the fish food web in the Gironde estuary, France, using dietary and stable isotope analyses. *Estuarine, Coastal and Shelf Science* 78, 267–279.
- Pasquaud S., Pillet M., David V., Sautour B. and Elie P.** (2010) Determination of fish trophic levels in an estuarine system. *Estuarine, Coastal and Shelf Science* 86, 237–246.
- Pauly D.** (1984) Fish population dynamics in tropical waters: a manual for use with programmable calculator. *ICLARM Studies and Reviews*, 8. Manila, Philippines, 325 pp.
- Pauly D. and Munro J.L.** (1984) Once more on the comparison of growth in fish and invertebrates. *Fishbyte* 2, 2.
- Pawson M.G.** (1992) Climatic influences on the spawning success, growth and recruitment of bass (*Dicentrarchus labrax* L.) in British waters. *ICES Marine Science Symposia* 195, 388–392.
- Pawson M.G., Kelley D.F. and Pickett G.D.** (1987) The distribution and migrations of bass *Dicentrarchus labrax* L. in waters around England and Wales as shown by tagging. *Journal of the Marine Biological Association of the United Kingdom* 67, 183–217.
- Pawson M.G., Kupschus S. and Pickett G.D.** (2007) The status of sea bass (*Dicentrarchus labrax*) stocks around England and Wales, derived using a separable catch-at-age model, and implications for fisheries management. *ICES Journal of Marine Science* 64, 346–356.
- Pawson M.G. and Pickett G.D.** (1996) The annual pattern of condition and maturity in bass (*Dicentrarchus labrax* L.) in waters around the UK. *Journal of the Marine Biological Association of the United Kingdom* 76, 107–126.
- Person-Le Ruyet J., Mahé K., Le Bayon N. and Le Delliou H.** (2004) Effects of temperature on growth and metabolism in a Mediterranean population of European sea bass, *Dicentrarchus labrax*. *Aquaculture* 237, 269–280.

- Pickett G.D., Kelley D.F. and Pawson M.G.** (2004) The patterns of recruitment of sea bass, *Dicentrarchus labrax* L. from nursery areas in England and Wales and implications for fisheries management. *Fisheries Research* 68, 329–342.
- Pickett G.D. and Pawson M.G.** (1994) *Sea bass. Biology, exploitation and conservation*. London: Chapman and Hall [Fish and Fisheries Series, vol. 12, 358 pp.].
- Power M., Attrill M.J. and Thomas R.M.** (2000) Environmental factors and interactions affecting the temporal abundance of juvenile flatfish in the Thames Estuary. *Journal of Sea Research* 4, 135–149.
- R Core Team** (2014) *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing. <http://www.R-project.org>.
- Rafail S.Z.** (1971) Investigation on Sciaenidae and Moronidae catch and on the total catch by beach seine on U.A.R. Mediterranean Coast. *Etudes et Revues, Conseil General des Peches pour la Mediterranee* 48, 1–26.
- Reise K., Baptist M., Burbridge P., Dankers N., Fischer L., Flemming B., Oost A. P. and Smit C.** (2010) The Wadden Sea – a universally outstanding tidal wetland. *Wadden Sea Ecosystem. Common Wadden Sea Secretariat, Wilhelmshaven*, no. 29, pp. 7–24.
- Rose G.A.** (2005) On distributional responses of North Atlantic fish to climate change. *ICES Journal of Marine Science* 62, 1360–1374.
- Scharf F.S., Juanes F. and Rountree R.A.** (2000) Predator size–prey size relationships of marine fish predators: interspecific variation and effects of ontogeny and body size on trophic-niche breadth. *Marine Ecology Progress Series* 208, 229–248.
- Schoener T.W.** (1971) Theory of feeding strategies. *Annual Review of Ecology and Systematics* 2, 369–404.
- Smith C.L.** (1990) Moronidae. In Quero J.C., Hureau J.C., Karrer C., Post A. and Saldanha L. (eds) *Check-list of the Fishes of the Eastern Tropical Atlantic*. Paris: UNESCO, pp. 692–694 [CLOFETA, vol. 2.].
- Spitz J., Chauvelon T., Cardinaud M., Kostecki C. and Lorange P.** (2013) Prey preferences of adult sea bass *Dicentrarchus labrax* in the northeastern Atlantic: implications for bycatch of common dolphin *Delphinus delphis*. *ICES Journal of Marine Science* 70, 452–461.
- Stephens D.W. and Krebs J.R.** (1986) *Foraging theory*. Princeton, NJ: Princeton University Press.
- Stequert B.** (1972) *Contribution a l'étude du bar (Dicentrarchus labrax) des réservoirs à poissons de la région d'Arcachon. Thèse 3ème cycle*. Sciences biologiques, Bordeaux.
- Thompson B.R. and Harrop R.T.** (1987) The distribution and abundance of bass (*Dicentrarchus labrax*) eggs and larvae in the English Channel and southern North Sea. *Journal of the Marine Biological Association of the United Kingdom* 67, 263–274.
- Tulp I., Bolle L.J., Meesters E. and Vries P.** (2012) Brown shrimp abundance in northwest European coastal waters from 1970 to 2010 and potential causes for contrasting trends. *Marine Ecology Progress Series* 458, 141–154.
- Tulp I., Bolle L.J. and Rijnsdorp A.D.** (2008) Signals from the shallows: in search of common patterns in long-term trends in Dutch estuarine and coastal fish. *Journal of Sea Research* 60, 54–73.
- Van Aken H.M.** (2003) 140 years of daily observations in a tidal inlet (Marsdiep). *ICES Marine Science Symposia* 219, 359–361.
- Van Aken H.M.** (2008a) Variability of the water temperature in the western Wadden Sea on tidal to centennial time scales. *Journal of Sea Research* 60, 227–234.
- Van Aken H.M.** (2008b) Variability of the salinity in the western Wadden Sea on tidal to centennial time scales. *Journal of Sea Research* 59, 121–132.
- Van Aken H.M.** (2010) Meteorological forcing of long-term temperature variations of the Dutch coastal waters. *Journal of Sea Research* 63, 143–151.
- Van der Meer J., Witte J.I.J. and Van der Veer H.W.** (1995) The suitability of a single intertidal fish trap for the assessment of long-term trends in fish and epibenthic invertebrate populations. *Environmental Monitoring and Assessment* 36, 139–148.
- Van der Veer H.W., Bergman M.J.N., Dapper R. and Witte J.I.J.** (1991) Population dynamics of an intertidal o-group flounder *Platichthys flesus* population in the western Dutch Wadden Sea. *Marine Ecology Progress Series* 73, 141–148.
- Van der Veer H.W., Dapper R. and Witte J.I.J.** (2001) The nursery function of the intertidal areas in the western Wadden Sea for o-group sole *Solea solea* (L.). *Journal of Sea Research* 45, 271–279.
- Van der Veer H.W., Witte J.I.J., Beumkes H.A., Dapper R., Jongejan W.P. and Van der Meer J.** (1992) Intertidal fish traps as a tool to study long-term trends in juvenile flatfish populations. *Netherlands Journal of Sea Research* 29, 119–126.
- Vasconcelos R.P., Reis-Santos P., Fonseca V., Ruano M., Tanner S., Costa M.J. and Cabral H.N.** (2009) Juvenile fish condition in estuarine nurseries along the Portuguese coast. *Estuarine, Coastal and Shelf Science* 82, 128–138.
- Vinagre C., Ferreira T., Matos L., Costa M.J. and Cabral H.N.** (2009) Latitudinal gradients in growth and spawning of sea bass, *Dicentrarchus labrax* (L.) and their relationship with temperature and photoperiod. *Estuarine, Coastal and Shelf Science* 81, 375–380.
- Vinagre C., Madeira D., Narciso L., Cabral H. and Diniz M.** (2012a) Effect of temperature on oxidative stress in fish: lipid peroxidation and catalase activity in the muscle of juvenile seabass, *Dicentrarchus labrax*. *Ecological Indicators* 23, 274–279.
- Vinagre C., Madeira D., Narciso L., Cabral H. and Diniz M.** (2012b) Impact of climate change on coastal versus estuarine nursery areas: cellular and whole-animal indicators in juvenile seabass *Dicentrarchus labrax*. *Marine Ecology Progress Series* 464, 237–243.
- Vinagre C., Narciso L., Cabral H., Costa M.J. and Rosa R.** (2012c) Coastal versus estuarine nursery grounds: effect of temperature on growth, condition and metabolism in seabass, *Dicentrarchus labrax*. *Estuarine, Coastal and Shelf Science* 109, 133–137.
- Webb P.W.** (1976) The effect of size on the fast-start performance of rainbow trout *Salmo gairdneri*, and a consideration of piscivorous predator–prey interactions. *Journal of Experimental Biology* 65, 157–177.
- Wood S.N.** (2006) *Generalized additive models: an introduction with R*. Boca Raton, FL: Chapman and Hall/CRC Press.
- Zijlstra J.J.** (1972) On the importance of the Wadden Sea as a nursery area in relation to the conservation of the southern North Sea fishery resources. *Symposium of the Zoological Society of London* 29, 233–258.
- Zuur A., Ieno E.N. and Smith G.M.** (2007) *Analyzing ecological data*. New York, NY: Springer.
- and
- Zuur A., Ieno E.N., Walker N., Saveliev A.A. and Smith G.M.** (2009) *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.

#### Correspondence should be addressed to:

J.F.M.F. Cardoso

NIOZ – Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands

email: [joanafcardoso@portugalmail.com](mailto:joanafcardoso@portugalmail.com)