



Review Article

Cite this article: Cashman KD (2025). Vitamin D and other micronutrient deficiency prevention: the role of data in informing national, regional and global policy. *Proceedings of the Nutrition Society*, page 1 of 11. doi: [10.1017/S0029665124007626](https://doi.org/10.1017/S0029665124007626)

Received: 2 September 2024

Revised: 14 November 2024

Accepted: 19 November 2024

Keywords:

Vitamin D; Micronutrient deficiencies; Hidden hunger; Prevalence; Data

Corresponding author:

Kevin D. Cashman; Email: k.cashman@ucc.ie

Vitamin D and other micronutrient deficiency prevention: the role of data in informing national, regional and global policy

Kevin D. Cashman

Cork Centre for Vitamin D and Nutrition Research, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland

Abstract

The WHO describes micronutrient deficiencies, or hidden hunger, as a form of malnutrition that occurs due to low intake and/or absorption of minerals and vitamins, putting human development and health at risk. In many cases, emphasis, effort and even policy revolve around the prevention of deficiency of one particular micronutrient in isolation. This is understandable as that micronutrient may be among a group of nutrients of public health concern. Vitamin D is a good exemplar. This review will highlight how the actions taken to tackle low vitamin D status have been highly dependent on the generation of new data and/or new approaches to analysis of existing data, to help develop the evidence-base, inform advice/guidelines, and in some cases, translate into policy. Beyond focus on individual micronutrients, there has also been increasing international attention around hidden hunger, or deficiencies of a range of micronutrients, which can exist unaccompanied by obvious clinical signs but can adversely affect human development and health. A widely quoted estimate of the global prevalence of hidden hunger is a staggering two billion people, but this is now over 30 years old. This review will outline how strategic data sharing and generation is seeking to address this key knowledge gap in relation to the true prevalence of hidden hunger in Europe, a key starting point towards defining sustainable and cost-effective, food-based strategies for its prevention. The availability of data on prevalence and food-based strategies can help inform public policy to eradicate micronutrient deficiency in Europe.

We are fast approaching the latter end of the UN's *Decade of Action on Nutrition - 2016–2025*⁽¹⁾. This Nutrition Decade, the implementation of which is being co-led by the UN FAO and the WHO, is intended to provide an enabling environment for all countries, regardless of their income, the nature of their malnutrition challenges or the characteristics of their food and health systems, to ensure that action is taken by governments and stakeholders to develop and implement inclusive policies aimed at ending all forms of malnutrition⁽¹⁾. Malnutrition, in all its forms, includes undernutrition (wasting, stunting and underweight), micronutrient (vitamins and/or minerals) deficiencies, overweight/obesity and resulting diet-related noncommunicable diseases. In adopting the draft resolution on the UN Decade of Action on Nutrition, the General Assembly expressed concern that more than 2 billion people were suffering from micronutrient deficiencies, as well as the other forms of malnutrition being evident around the globe⁽¹⁾. Deficiencies of micronutrients, such as iron, zinc, iodine, folate, vitamins A, B₁₂ and D, amongst others, compromise the immune system, disrupt childhood growth and brain development and accelerate multi-system aging and non-communicable diseases⁽²⁾. At a societal level, micronutrient deficiencies can adversely affect the development potential of individuals, reducing educational attainment, as well as work capacity and productivity, ultimately hindering the development of societies and nations⁽²⁾. The staggering numbers affected by micronutrient deficiencies globally stem from the fact that a high proportion of the population is at risk of these deficiencies. For example, children, adolescents, women of reproductive age and pregnant and lactating women are at greater risk of micronutrient deficiencies due to their higher nutritional requirements during these stages of life⁽²⁾, whereas micronutrient deficiencies can also disproportionately affect older adults, because of biological and environmental challenges and often coexist with and can be exacerbated by obesity⁽³⁾. Collectively, these population subgroups represent more than two-thirds of the European population, a statistic that does not include immigrant and ethnic minority groups, who are also at risk of micronutrient deficiencies⁽⁴⁾.

The UN Sustainable Development Goal 2 has a highly ambitious, but laudable, target of ending micronutrient deficiencies, as well as other all forms of malnutrition, by 2030⁽⁵⁾. This requires that inclusive policies need to be urgently built around sound data on aspects of micronutrient status and intake, prevalence of deficiencies, contributory factors, as well as evidence-based strategies for improving micronutrient status in the affected populations. Data is

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.






Figure 1. The number of studies/papers of vitamin D status (and intake) in Ireland by year within the PUBMED database. 1998 EC Report = Report for the European Parliament on the prevention of osteoporosis; VDSP, Vitamin D Standardization Program.

key to decision making. The mathematician Clive Humby said that ‘like oil, data is valuable, but if unrefined it cannot really be used’⁽⁶⁾. The present review seeks to illustrate the critical importance of data, including its careful interrogation and strategic use, in the generation of the evidence-base that can inform guidelines and policies for addressing micronutrient deficiency. It will do this firstly by presenting the case for data from the prism of tackling vitamin D deficiency, as an exemplar micronutrient from among the two dozen or more currently considered essential. It will then consider how lessons learned from the vitamin D experience, may benefit our attempts at tackling deficiencies of other vitamins as well as minerals, also known as hidden hunger. The review will also stress the importance of the uptake and use of such data by key expert groups, agencies, policy decision-makers and other relevant stakeholders in terms of tackling micronutrient deficiencies. Finally, it will also acknowledge the role funding agencies have had, and continue to have, in terms of the generation of data that is much needed to inform inclusive policies for tackling micronutrient deficiencies.

Tackling vitamin D deficiency at a national, regional and global level: the role of and need for data

The following is not intended in any way as a review of the means of tackling vitamin D deficiency, but rather, as aligned with the overall data-related theme of the inaugural Nutrition Society Congress, more a demonstration of the role of, and need for, data in the development of the evidence-base, guidelines and, in some cases, policies for addressing vitamin D deficiency and inadequacy of vitamin D intake. The following subsections will highlight how data in its many guises, ranging from small to very large datasets, new data as well as existing data, data from studies of human subjects, animal trials and indeed other sources and from study designs ranging from randomised controlled trials (RCT) to observational studies, all contribute to the collective evidence-base that underpins decision-making in relation to tackling vitamin D deficiency. Also, it is important to stress that the following

subsections draw on our own experience of how research in the field of vitamin D nutrition generates data that can underpin the development of such an evidence-base. In this regard, they represent exemplars and personal reflections on how data is key to decision-making in relation to guidelines and policies for addressing vitamin D deficiency.

National data on vitamin D status and intake

In 1998, the European Commission (EC) set up an expert working group to produce a report for the European Parliament on the prevention of osteoporosis⁽⁷⁾. The report contained 8 clear recommendations targeted at improving the prevention of osteoporosis. The 4th recommendation related to the need to develop, integrate and implement policies to advise the general public and health professionals about calcium and vitamin D nutrition at all stages of life⁽⁷⁾. Importantly, the report highlighted that data on dietary intake and status of vitamin D and calcium were not available in many European countries at that time. This was our first exposure to an expert body clearly identifying the priority need for such data at European member state level, which would then also inform the situation for the overall region. There was also the suggestion that there would be an audit of recommendations at member state level as a follow-on activity to the publication of the report. At around that time, there were a very limited number of publications (~4) reporting on vitamin D status in Ireland (see Fig. 1), and the studies contributing this limited data were of generally non-representative, relatively small (n in the range 29–196) samples of elderly subjects, and in some cases institutionalised individuals^(8–11). Nevertheless, these studies were hugely important in highlighting vitamin D deficiency as a potential public health problem in Ireland. It was clear to us that additional data on vitamin D intake and status for the Irish population was urgently needed, and accordingly, the research programme of our then Vitamin D Research Group (now the *Cork Centre for Vitamin D and Nutrition Research*) at University College Cork, aspects of which were funded by two EC Framework

Programme 5 projects (*OSTEODIET* and *OPTIFORD*), and by national funding from the Irish Department of Agriculture, Food and the Marine, under their Food Institutional Research Measure programme, amongst other funding sources, sought to address this key data gap.

From around 2000 onwards, a significant component of vitamin D research effort of the Centre focused on the generation of this new vitamin D status and intake data in different population subgroups in Ireland (Fig. 1); however, many of these studies were also, by necessity, relatively small, non-representative studies. Notwithstanding this, these new data, when taken with the previous data, again highlighted vitamin D status as a potential public health nutrition concern for Ireland. Importantly, the 2008–2010 nationally representative nutrition survey of adults in Ireland was resourced to include blood sampling, which allowed for the generation of much needed representative vitamin D status data for Irish adults for the first time. As part of the *National Adult Nutrition Survey*, the Centre generated this new 25-hydroxyvitamin D (25(OH)D) data based on analysis of survey serum samples by enzyme-linked immunoassay⁽¹²⁾. Fortuitously, around that time, the Centre was invited to become a member of the US National Institutes of Health-led *Vitamin D Standardization Program* (VDSP) which had as its key goal to improve the detection, evaluation and treatment of vitamin D malnutrition by making serum total 25(OH)D measurements accurate and comparable over time, location and laboratory procedure⁽¹³⁾. The VDSP, coordinated by the Office of Dietary Supplements at the National Institutes of Health, was in response to the well-recognised variability in serum 25(OH)D estimates arising from method-related differences in its measurement⁽¹⁴⁾. Accordingly, the VDSP developed protocols for standardising existing serum 25(OH)D data from national surveys⁽¹⁵⁾, and thus minimising the impact of method-related differences in their estimates. As part of the VDSP, the Centre applied these protocols to the *National Adult Nutrition Survey* serum 25(OH)D data, and based on these standardised data it was estimated that the prevalence of serum 25(OH)D < 30 nmol/l and < 50 nmol/l, reflective of risk of vitamin D deficiency and risk of inadequacy of vitamin D status, respectively⁽¹⁶⁾, was 12.3 % and 45.9 % of Irish adults, respectively⁽¹⁷⁾. Importantly, these estimates, based on standardised data, allowed for comparison with standardised data from nationally representative surveys in the US (*National Health and Nutrition Examination Survey* 2011–2014) and Canada (*Canadian Health Measures Surveys* Cycle 1 and 2) which had lower prevalence estimates (e.g. 5.0 % and 8.8 % with serum 25(OH)D < 30 nmol/l, respectively), and when limiting data to white only survey participants, making for a more valid comparison with the Irish survey data, estimates were even lower again (2.1 % and 5.9 % with serum 25(OH)D < 30 nmol/l, respectively)^(18,19). In recent times, the Centre has generated equivalent prevalence data for Irish teenagers aged 13–18 years, based on analysis of bloods from the 2019–2020 *National Teens Survey II* (NTS II) – the second national survey to have included blood sampling in its design⁽²⁰⁾. The standardised serum 25(OH)D data from this work suggests that about 21.7 % and 54.8 % of Irish teenagers, are at risk of vitamin D deficiency and inadequacy, respectively. Both the *National Adult Nutrition Survey* and NTS II also reported vitamin D intake data which allowed for benchmarking of the degree of inadequate intakes in both population subgroups. Using the US Estimated Average Requirement (EAR) value for vitamin D of 10 µg/d⁽¹⁶⁾, it was estimated that 90 % and 94 % of adults and teenagers, respectively, in Ireland had inadequate intakes of vitamin D^(20,21),

and this low intake likely contributed to the risk of vitamin D deficiency, particularly in the absence of dermal synthesis of vitamin D on exposure of skin to sunlight with sufficient ultraviolet B (UVB) radiation. Other nationally representative surveys in other population subgroups in Ireland have also reported vitamin D intake data, even where blood sampling wasn't included in their survey designs (<https://www.iuna.net/surveyreports>).

Overall, as can be seen in Fig. 1, since the 1998 EC report, there has been a noticeable increase in the number of studies of Irish population subgroups conducted by the vitamin D research community on the island of Ireland, ranging from relatively small and non-representative samples, up to regionally and, in some cases, nationally representative samples. Thus, the important data-gap around estimates of vitamin D intake and status, as highlighted by the EC report, has been largely addressed for Ireland. However, it should be stressed that as such data needs to be iterative to best capture the current situation in relation to vitamin D status and intake, this strategic data requirement needs to be addressed on an ongoing basis. It is also encouraging to see evidence of the uptake and use of such data in terms of informing national guidelines and policies. For example, three recent reports by the Scientific Committee of the *Food Safety Authority of Ireland* (FSAI) focused on vitamin D recommendations, as part of their healthy eating guidelines, used the data from the various studies since ~2000 to inform on current status of vitamin D status and/or intakes in Irish children, teenagers, adults (including pregnancy) and older adults^(22–24).

European data on vitamin D status and intake

Unlike the situation in the US and Canada, where, as mentioned already, very good estimates exist for the prevalence of vitamin D deficiency and inadequacy based on data from their representative surveys^(18,19), the equivalent estimates did not exist for Europe as a whole, only for some of its member state countries individually. In 2011, the Centre had highlighted that data on the distribution of serum 25(OH)D concentrations and vitamin D intakes, including food sources, in nationally representative populations, together with sustainable food-based strategies to bridge the gap between current and recommended intakes of vitamin D to minimise the prevalence of serum 25(OH)D concentrations < 30 nmol/l, were three critical and prioritised research requirements for Europe⁽²⁵⁾. Addressing these, and other, priority data gaps around vitamin D nutrition and health in Europe was the focus of a EC Framework Programme 7-funded *ODIN* project on vitamin D, coordinated by the *Cork Centre for Vitamin D and Nutrition Research* at University College Cork (Professor Mairead Kiely and Professor Kevin Cashman, Joint Coordinators)⁽²⁶⁾. In 2016, the *ODIN* project delivered the first internationally standardised dataset on vitamin D status and reported the prevalence of vitamin D deficiency across Europe for the first time⁽²⁷⁾. This European estimate was based on a collection of 14 nationally or regionally representative studies gathered as part of the project and their existing serum 25(OH)D data (based on a wide collection of analytical methods) were standardised as per the VDSP protocol. While the project included nationally representative nutrition surveys from Ireland, the UK and Germany, some member states in Europe did not have such nationally representative surveys. Thus, in the absence of such data, well-curated samples from regionally representative health surveys were used, as they can also achieve some degree of population coverage⁽²⁷⁾. While there was considerable variation across study populations, the prevalence of vitamin D deficiency

(standardised serum 25(OH)D concentrations < 30 nmol/l), when these representative population samples were pooled (n 55 844), was 13 %, which would broadly correspond to one in eight Europeans⁽²⁷⁾. Even a crude estimation based on the magnitude of population in Europe coupled with this population-wide prevalence estimate, suggest something in the region of 66 million individuals deficient⁽²⁸⁾. This European estimate was also considerably greater than that for the US or Canada^(18,19), as mentioned above. In terms of vitamin D inadequacy, the average yearly population prevalence of standardised serum 25(OH)D < 50 nmol/l in Europe, the US and Canada is 40.0 %, 24.0 % and 36.8 %, respectively^(18,19,27).

These are whole-population estimates and the prevalence can vary by age-grouping with a tendency for it to be lowest in childhood and possibly later life⁽²⁶⁾. It should also be stressed that the population-wide estimates, do not capture the differences by ethnicity in these regions, which can be significant. For example, the *ODIN* project showed that dark-skinned ethnic groups within Europe are worryingly at much increased risk of vitamin D deficiency compared to their white counterparts (prevalence < 30 nmol/l in the range 28–65 %, depending on the country and the ethnic group)^(27,29), which aligns with higher risk of vitamin D deficiency reported among the non-white participants in the North American surveys^(18,19). This was only a partial picture, however, as the proportion of participants from ethnic minorities was low in most studies within the *ODIN* collection. The project highlighted a lack of well-curated and characterised biobanks of ethnic minorities in Europe, representing a major research gap that should be prioritised.

Finally, while these important first of their kind estimates for Europe provided firm evidence that vitamin D deficiency is widespread across Europe, that analysis is fast approaching being 10 years old, which again underscores why this strategic data requirement needs to be addressed on an ongoing basis to maintain currency.

There is no one single underlying reason for vitamin D deficiency, but the combination of low UVB availability and/or exposure coupled with a low dietary vitamin D supply are of key importance⁽³⁰⁾. The availability of UVB of sufficient intensity to stimulate the conversion of 7-dehydrocholesterol in the skin to pre-vitamin D₃ and then vitamin D₃ is impacted by several environmental factors, such as season, latitude and prevailing weather conditions. To get a better understanding of vitamin D-effective UVB availability across Europe, the *ODIN* project modelled such availability data for nine European countries/regions⁽³¹⁾. It did this using a validated UV irradiance model which used data on atmospheric and geophysical parameters, such as local cloud, ozone and aerosol, plus topography, accessed from a number of orbiting satellites. The results showed that UVB availability decreased with increasing latitude within Europe (from 35°N to 69°N), while all locations exhibited significant seasonal variation in UVB. The number of months in which UVB availability was too low to allow for skin synthesis of vitamin D, referred to as the 'vitamin D winter', was estimated to range from being largely absent in the very south of Europe to lasting for as long as 7 or 8 months in northern Europe⁽³¹⁾. Beyond vitamin D-effective UVB availability, personal characteristics, such as skin pigmentation, age, attire, sunscreen usage, working environment, outdoor physical activity and sun exposure behaviour, can also prevent or impede vitamin D synthesis⁽³⁰⁾.

In the absence of sufficient UVB availability/exposure to enable synthesis in the skin, dietary supply of vitamin D is critical to meeting population requirements and prevention of vitamin D deficiency. However, vitamin D intake data from national surveys

of adults and children in 21 and 13 European countries, respectively, suggest habitual intakes are relatively low on average (mean intakes in the range 2.2–3.3 µg/d; depending on sex and age group)^(32,33). There was variation in the estimated mean daily vitamin D intakes across countries and most notable among adults from countries in Northern Europe compared to those in Western, Southern and Central and Eastern Europe (mean daily intakes of 7.8, 4.0, 3.5 and 1.5 µg/d, respectively, for males; and 6.1, 3.3, 2.9 and 1.1 µg/d, respectively, for females)⁽³²⁾.

Estimates of the vitamin D dietary requirement for Irish, UK, European and global populations

Since its derivation in 2011, the US Institute of Medicine's EAR for vitamin D of 10 µg/d⁽¹⁶⁾ has become a touchstone target intake with which to benchmark typical intakes of vitamin D within a population. The percentage of the population with a habitual daily nutrient intake lower than the EAR is taken as an estimate of the percentage of the population with probable inadequate intakes⁽³⁴⁾. While the prevalence of vitamin D intakes less than the EAR of 10 µg/d were not provided in many of the above mentioned European surveys^(32,33), it is clear from their mean daily intake data that many of the countries have a high majority of their respective populations with an inadequate intake of vitamin D. Roman Viñas *et al.* in 2011 projected that among European national surveys reporting vitamin D intake data, 77–100 % and 55–100 % of adults (18–64 years) and older adults (> 64 years), respectively, had intakes below 10 µg/d, which was the 2004 Nordic recommended EAR for vitamin D available at the time⁽³⁵⁾.

Estimates of dietary requirements for vitamin D as dietary targets are crucial from a public health perspective in providing a framework for prevention of vitamin D deficiency and optimising vitamin D status of individuals. The Food Standards Agency (FSA) in the UK issued a research funding call in late 2004 with the aim of adding to the evidence-base around dietary requirements for vitamin D and the significance of both dietary sources and sunlight to vitamin D status and/or functional markers. University College Cork and the University of Ulster jointly competed for and won a contract within this call to undertake research on dietary requirements for vitamin D. As a key part of this work, the two institutions conducted two winter-based, dose-related vitamin D RCT during the period 2006–2008 specifically designed to establish the distribution of dietary vitamin D required to maintain serum 25(OH)D concentrations above contemporary cut-offs (> 25, > 37.5, > 50 > 80 nmol/l) during wintertime, accounting for the impact of summer sunshine exposure and diet, in adults and in older adults, separately^(36,37). Both RCT were two-site trials (Cork; 51°N and Coleraine; 55°N) to provide the longitudinal coverage for much of the UK. While estimated dietary requirements for vitamin D at a number of selected percentiles were provided, major focus was on the vitamin D intake estimates to maintain 97.5 % of a population life-stage group above 25 nmol/l, as well as the other contemporary cut-offs, during winter, as this corresponds to a Reference Nutrition Intake (RNI) value in the UK (a Recommended Daily Allowance or Population Reference Intake in the US and EU, respectively). With the publication of the Institute of Medicine report in 2011, which highlighted an increased risk of vitamin D deficiency at serum concentrations below 30 nmol/l, we have subsequently added this threshold to the four other previously suggested ones, and derived the corresponding intake requirement estimates⁽³⁸⁾. The vitamin D intakes that would maintain serum 25(OH)D concentrations ≥ 30 nmol/l

Table 1. Estimated dietary requirements for vitamin D at the 97.5th percentile in different population subgroups to maintain wintertime serum 25(OH)D above two selected thresholds associated with increased risk of vitamin D deficiency*

Serum 25(OH)D concentration	Boys and girls aged 4–8 y (n 144)	Adolescent girls aged 11 (n 144)	Adolescent boys and girls aged 14–18 y (n 144)	Adult men and women aged 20–40 y (n 215)	Elderly men and women aged 64+ y (n 225)	Pregnant women (n 144)	White v East African women (n 68 v. 47)
	µg/d						
>25 nmol/l	6.4	8.3	10.1	8.7	8.6	11.3	NR
>30 nmol/l	8.3	10.3	13.1	13.7	12.2	13.8	8.2 v. 18.4

y, year; NR, not reported.

*Estimates from^(26–43).

during wintertime in 97.5 % of adults (20–40 years olds) and older adults (64+ years olds) were around 14 and 12 µg/d, respectively (Table 1).

In relation to the uptake and use of such data in terms of informing national guidelines and policies, in 2009, a UK FSA workshop was convened to review this new vitamin D data together with that from two other FSA-funded vitamin D projects in the 2004 call⁽⁴⁴⁾. The discussion and debate during this workshop helped inform the decision by the FSA and Department of Health in 2010 to request that the UK Scientific Advisory Committee on Nutrition (SACN) would review the existing Dietary Reference Values (DRV) for vitamin D, established in 1991 and re-iterated by a SACN update in 2007, and also make recommendations. The data from above-mentioned two vitamin D requirement RCT^(36,37), together with data from an additional analysis of vitamin D RCT data in girls (see below), were used as a basis to establish the current RNI of 10 µg/d for all those aged 4 years and older⁽⁴⁵⁾. This RNI was based on a population protective serum 25(OH)D concentration of 25 nmol/l.

Data on dietary requirements for vitamin D during childhood were extremely limited. The strategic secondary use of wintertime only data from a wider 12-month vitamin D trial in adolescent Finnish and Danish girls, which examined the impact of vitamin D supplementation on bone mass acquisition (as part of the OPTIFORD EU project⁽⁴⁶⁾), allowed us to generate equivalent estimates for this population subgroup in 2011⁽³⁹⁾. The estimates for the girls were relatively similar to those of the adults and older adults (Table 1). Since then, and as part of the ODIN project, four additional bespoke vitamin D requirement RCT were conducted in young children aged 4–8 years (Copenhagen, Denmark; 55°N), teenagers aged 14–18 years (Surrey, UK; 51°N), pregnant women (Cork, Ireland; 51°N) and in persons of ethnic minority (Helsinki, Finland; 60°N)^(40–43), as under-represented groups among the healthy population for whom data to base estimates of vitamin D requirements were scarce or absent. These vitamin D dose-related RCT estimated the dietary requirements for vitamin D to meet serum 25(OH)D thresholds of 25 and 30 nmol/l, as well as 50 nmol/l, in 97.5 % of individuals in these subgroups largely under conditions of minimal UVB exposure. The first two RCT showed that for white 4–8 year-old children and 14–18 year-old adolescents, 8 and 13 µg/d of vitamin D, respectively, prevented serum 25(OH)D concentrations falling below 30 nmol/l during wintertime (Table 1)^(40,41). Interestingly the following year, Ohlund *et al.* in a study in Northern and Southern Sweden, at 63 and 55°N, respectively⁽⁴⁷⁾ showed that white and dark-skinned children (aged 5–7 years) had varying requirements for vitamin D intakes, needing 6 and 14 µg/d, respectively, to maintain serum 25(OH)D concentrations > 30 nmol/l during winter. In relation to possible

ethnic differences in vitamin D requirements, the third ODIN vitamin D dose-response RCT, with a similar design to those mentioned above, compared vitamin D requirements of women of Finnish descent (n 69) with East African women (n 47) in Helsinki, Finland (60°N), during wintertime⁽⁴³⁾. Models adjusted for baseline differences in age, weight, vitamin D intake and serum 25(OH)D found that dietary requirements for vitamin D to maintain wintertime 25(OH)D above 30 nmol/l were significantly higher in women of East African descent (18 µg/d) than in white women of Finnish descent (8 µg/d) (Table 1)⁽⁴³⁾.

The last ODIN vitamin D dose-related RCT (conducted in Cork, Ireland; 51°N) among 144 white pregnant women estimated the vitamin D intake required to maintain maternal serum 25(OH)D in late gestation above 25, 30 and 50 nmol/l, but also at concentrations sufficient to prevent neonatal deficiency (i.e. keep concentrations in infants at or above about 25/30 nmol/l)⁽⁴²⁾. The study showed that an intake of about 14 µg/d of vitamin D prevented maternal serum 25(OH)D concentrations falling below 30 nmol/l during wintertime (Table 1). However, this would still not protect serum concentrations in the infants from falling below this threshold. The study showed that when maternal serum 25(OH)D concentration was ≥ 50 nmol/l, 95 % of cord sera had concentrations ≥ 30 nmol/l. A total vitamin D intake of 30 µg/d maintained serum 25(OH)D concentrations ≥ 50 nmol/l during pregnancy, which ensured that serum 25(OH)D concentration was ≥ 30 nmol/l in 95 % of umbilical cord sera⁽⁴²⁾.

Again in terms of uptake of these data, a number of agencies/authorities charged with the development of DRV for vitamin D have used data from these RCT either directly as is, such as the UK SACN⁽⁴⁵⁾, the German Nutrition Society⁽⁴⁸⁾ and the Irish FSAI^(22,23), or in combination with that from other relevant vitamin D RCT to provide an overall collection of aggregate data points for use in the agencies' respective meta-regression modelling to derive their dietary recommended intake values, such as the Institute of Medicine⁽¹⁶⁾, the Nordic Nutrient Recommendations⁽⁴⁹⁾ and the European Food Safety Authority (EFSA)⁽⁵⁰⁾.

In terms of establishing the vitamin D intake – serum 25(OH)D dose response, which is integral to deriving the estimates of vitamin D intake requirements, beyond the meta-regression of aggregate (or statistical summary) RCT data as used by Institute of Medicine, Nordic Nutrient Recommendations and EFSA^(16,49,50), a meta-regression based on analysis of individual participant data (IPD), in which the raw data for each RCT are used for synthesis, offers many potential advantages, is increasingly recognised as best practice and is likely to provide the most appropriate DRV estimates⁽⁵¹⁾. Again as part of the ODIN project, the Centre completed the first-ever IPD-level meta-regression from seven suitable winter-based RCT (with data from 882 participants

ranging in age from 4 to 90 years) of the vitamin D intake-serum 25(OH)D dose-response⁽⁵²⁾. Notably, vitamin D intakes and serum 25(OH)D concentrations had been analysed using the same methods in these studies, ensuring compatibility across the dose-response relationship. The IPD analysis confirmed that 10 and 13 µg/d would be required to maintain wintertime serum 25(OH)D concentrations > 25 and 30 nmol/l, respectively, in 97.5 % of individuals⁽⁵²⁾. These also represent composite estimates across the white, non-pregnant/lactating population over 5 years of age.

In the last few years, the Centre has also completed two additional vitamin D requirement IPD-level meta-analyses, one based on 11 RCT (*n* 1429) with vitamin D-fortified foods⁽⁵³⁾ and the second based on 10 RCT (*n* 677) to estimate the vitamin D dietary requirements in dark-skinned individuals (of black or South Asian descent) resident at high latitude⁽⁵⁴⁾. The IPD based on fortified food RCT in light-skinned individuals provided estimates of the vitamin D intake required to maintain 97.5 % of winter 25(OH)D concentrations ≥ 30 nmol/l as 12 µg/d⁽⁵³⁾. The IPD-derived estimate of the vitamin D intakes required by dark-skinned individuals to maintain 97.5 % of winter 25(OH)D concentrations ≥ 30 nmol/l was 27.3 µg/d and 33.2 µg/d, in South Asians and black individuals, respectively⁽⁵⁴⁾. These estimates are radically different from the equivalent estimates for white individuals and highlight the need to consider, and possibly adapt, the approach for setting vitamin D RDAs in such cases. The IPD estimates are also higher than that currently recommended internationally by several agencies, which are based predominantly on data from white individuals and derived from standard meta-regression based on aggregate data^(16,49,50). It is important to highlight and acknowledge that such extensive data-rich IPD analyses would not be possible without the generous collaboration of large numbers of willing principal investigators that see the merit of this secondary, but nonetheless strategic, use of their RCT data. The requirement intake estimates from these IPD analyses have been used as supportive evidence by the FSAI in their vitamin D recommendations⁽²²⁾.

The approach of using data from available vitamin D studies, including IPD, to derive a vitamin D intake requirement estimate for young children (0–3.9 years) has been implemented by the WHO together with the FAO in an update of their 2004 Nutrient Requirements. A large body of studies (117 in total, and including both observational studies and RCT) were used to define a serum 25(OH)D threshold of 28 nmol/l, based on its association with minimised risk of rickets in young children with adequate calcium intake⁽⁵⁵⁾. The ability to stratify on the basis of adequacy/inadequacy of calcium intake within the analysis was novel, since most exercises up to that could not distinguish, and vitamin D dietary requirements are set on the basis of assumed adequacy of calcium intake. This threshold was then used in multi-level and multivariable dose-response modelling of serum 25(OH)D to total vitamin D intake, based on RCT data from 31 trials from North America, Europe, Asia and Australasia/Oceania, with latitudes ranging from 38°S to 61°N, reflecting the global nature of the WHO-FAO recommendations. This modelling allowed for the derivation of an updated vitamin D requirement estimate of 10 µg/d for young children below 4 years of age⁽⁵⁶⁾. Interestingly, while the meta-regression was based on aggregate data (as acquisition of IPD from 31 RCT was not feasible within the timelines and resources available in the WHO-FAO update exercise), an individual variability component, as would be derived from an IPD-level meta-regression, was imputed within the modelling to allow the best estimate of the vitamin D intake requirement covering the needs of 98 % of young children possible⁽⁵⁶⁾.

Data around possible food-based strategies for improving vitamin D intakes

The Centre's work on vitamin D intake requirements over the last 2 decades suggests that intakes of between 8 to 14 µg/d are needed to prevent risk of vitamin D deficiency (reflected as serum 25(OH)D concentrations below 30 nmol/l^(16,50)) in most population subgroups and maybe higher in the case of dark-skinned ethnic subgroups. The above-mentioned representative survey data for Ireland, the UK and Europe show that habitual vitamin D intakes frequently fall well below these requirement intakes. This underscores the fact that we had previously suggested that sustainable food-based strategies to bridge the gap between current and recommended intakes of vitamin D to minimise the prevalence of serum 25(OH)D concentrations < 30 nmol/l, was one of the three critical and prioritised research requirements for Europe⁽²⁵⁾. The WHO-FAO suggest that in terms of strategies for tackling micronutrient malnutrition, while food fortification tends to have a less immediate impact compared to micronutrient supplementation, it nevertheless has a much wider and more sustained impact⁽⁵⁷⁾. Food-based strategies for improving vitamin D intakes, particularly food fortification and biofortification, have also been a focus within the Centre's wider vitamin D research programme. The Centre's collaborative studies over the last decade, funded by national agencies as well as the EC, have ranged from exploration of the potential of UV radiation of bakers' yeast and edible mushrooms in terms of enhancing their vitamin D₂ content, to addition of increased levels of vitamin D₃ and/or 25-hydroxyvitamin D₃ into the feed of aqua-cultured fish, chickens, pigs and beef cattle, so as enhance the total vitamin D content of fillets, eggs and meats, respectively, for human consumption (a process known as 'biofortification'⁽⁵⁸⁾), to more traditional addition of vitamin D to foods, such as low-fat cheese, amongst other vehicles^(59–66) (see Fig. 2). These studies have provided good evidence of the technological feasibility of the enhancement of their vitamin D content, and, in some cases where assessed, of the consumer acceptability of these fortified/biofortified food products. A number of the studies have included RCT with the vitamin D-fortified/biofortified foods which highlight the effectiveness of food fortification in terms of improving vitamin D status^(60,62,66,67), or not, as in the case of bread made from UV-irradiated yeast⁽⁵⁹⁾. Finally, the evidence from these RCT have been combined with that of other available RCT of vitamin D fortified/biofortified foods in the form of a number of systematic reviews and meta-analyses which have also highlighted their effectiveness in terms of improving vitamin D status and adding to the overall convincing evidence-base^(60,68–70).

Finally, the data on the vitamin D content that can be feasibly achieved in these fortified/biofortified foods have also been used in *in silico* dietary modelling experiments based on national nutrition survey data to predict their impact, as part of current consumption practises, on the vitamin D intake in a population subgroup(s) of interest. For example, as part of the ODIN project, such dietary modelling has been performed using data from three relevant Irish national nutrition surveys to estimate the impact of hypothetical addition of vitamin D to the food chain using some of the biofortified food strategies mentioned above, such as eggs, beef and pork, either alone or in combination with fortification of milk and cheese, on the intake of vitamin D by adults, as well as separately by children (aged 5–12 years) and adolescents in Ireland. While the impact of single biofortified foods on the distributions of vitamin D intakes were relatively small, when all three foods were combined




Figure 2. Vitamin D-fortified and vitamin D-biofortified foods investigated over the last decade as part of our vitamin D research programme.

*Vitamin D₃ and/or 25-hydroxyvitamin D₃ as the fortificant in foods or animal feeds; RCT, randomised controlled trials.

together with fortification of milk and cheese, projected mean and median intakes close to the EAR of 10 µg/d were achieved, without increasing the risk of excessive intakes^(22,26,71). This dietary modelling was also conducted across nationally representative surveys in three other EU countries and demonstrated the feasibility of achieving average intakes of ~10 µg/d vitamin D, without increasing the risk of excessive intakes⁽²⁶⁾. These modelling data complement that from other dietary modelling exercises in Ireland and the UK which were based on more traditional fortification of milk and/or wheat flour/bread, and some other foods^(21,72,73). The out-turn of this dietary modelling work has been used by the FSAI in their vitamin D recommendation reports in relation to strategies for addressing low vitamin D intakes within the population⁽²²⁾. Data on food fortification/biofortification, including the dietary modelling, together with other aspects of the Centre’s vitamin D research, such as status and intake data as well as dietary requirement estimates, have been shared at specifically organised meetings among vitamin D scientists and relevant policy decision-makers together with other stakeholders in Canada (during 2015) and Chile (during 2018), which helped inform their fortification strategies. The uptake and translation of data/knowledge around food-based approaches for tackling vitamin D deficiency to policy and action has been more variable within Europe. To further encourage the conversion of the such data/knowledge to action in Ireland, a nationally funded project called ‘Vitamin D-Deficiency Prevention IRELAND’ (acronym, VitD-DPI; <https://www.craft.do/s/o7HuSea30GNWvt>) aims to use a multi-actor approach to develop, validate and present a policy-ready national framework for vitamin D deficiency prevention in Ireland, that is endorsed by all stakeholders. The project started on the 1st of September 2022 and will be completed by the end of August 2026.

Tackling micronutrient deficiencies or hidden hunger in Europe: again, the need for data

The term ‘micronutrient deficiencies’ has been used in recent times as a synonym of ‘hidden hunger’ and/or ‘micronutrient malnutrition’⁽⁷⁴⁾, all of which, in many cases, are used interchangeably. This section of the review will again attempt to highlight how data is critically needed to address persistent data/knowledge gaps in relation to hidden hunger and, in so doing, can help to contribute to the development of strategies for their prevention. It will also point to how some of the above-mentioned data-driven

approaches in relation to tackling vitamin D deficiency, can be put to use for tackling hidden hunger. However, before that it might be useful to define what is meant by hidden hunger.

Hidden hunger, micronutrient malnutrition and micronutrient deficiencies

While not new, the terms ‘hidden hunger’ and ‘micronutrient malnutrition’ have been increasingly used in recent years, with 229 and 123 articles in PUBMED having used the respective terms in the last 5 years alone. However, over 3 decades ago, the WHO together with UNICEF convened a conference in Montreal called ‘Ending hidden hunger: a policy conference on micronutrient malnutrition’⁽⁷⁵⁾. This helped bring both terms to international prominence. The purpose of the WHO-UNICEF conference in 1991 was to reinforce the commitment and collaboration to accelerate progress towards the goals of elimination or control of micronutrient malnutrition by the year 2000, endorsed at the World Summit for Children the year before⁽⁷⁵⁾. At that time, the WHO explained that ‘micronutrient malnutrition’ is the term used when referring to the main vitamin or mineral nutritional deficiencies of public health significant, referencing iodine deficiency disorders, vitamin A deficiency and iron deficiency anaemias⁽⁷⁵⁾. The term ‘hidden hunger’, as a form of malnutrition that occurs when the intake and/or absorption of minerals and vitamins are too low to sustain good health and development as well as normal physical and mental function^(74–77), and which can exist unaccompanied by obvious clinical signs, was helpful to distinguish it from hunger and calorie deficit representing a more widely recognised form of malnutrition. A former deputy executive of UNICEF has suggested that ‘hidden hunger due to micronutrient deficiency does not produce hunger as we know it. You might not feel it in the belly, but it strikes at the core of your health and vitality’⁽⁷⁷⁾.

The prevalence of hidden hunger

Over three decades ago, a global prevalence of hidden hunger was estimated at a staggering two billion people, more than double the 805 million people who had insufficient energy intake at that point in time⁽⁷⁶⁾. This still widely quoted estimate has its origins in the WHO’s data from 1991 that 1 billion people were at risk of iodine deficiency disorders, 190 million children of pre-school age are at risk of vitamin A deficiency, and over 2 billion people are at risk of iron deficiency anaemia or are affected by some form of anaemia.

As many of the affected people are the same people in the at-risk groups, this led to the estimate that the total number of subjects is around 2 billion⁽⁷⁵⁾.

Recently, an analysis of pooled global individual-level data on micronutrient status of two key at-risk groups (pre-school-aged children and non-pregnant women of reproductive age) from nationally representative, population-based surveys of 22 countries was based on iron and zinc as well as folate or vitamin A (in the case of non-pregnant women and young children, respectively)⁽²⁾. The work showed that on a global basis, approximately half and two-thirds of the young children and women, respectively, had a deficiency of at least one of these micronutrients (representing ~1.6 billion individuals). Of note, the analysis showed that no world region, including high-income countries, was spared from the high burden of micronutrient deficiency. For Europe, prevalence estimates of 48 % and 43 % for young children and non-pregnant women, respectively, were reported for these nutrients (representing 82 million individuals)⁽²⁾. However, this is an incomplete picture of the true burden of prevalence of micronutrient deficiencies for Europe, and elsewhere. For example, the estimates for Europe were based on data from just one survey, the UK *National Diet and Nutrition Survey*. In addition, as the exercise had a global focus, it only focussed on those micronutrient deficiencies of world-wide significance and just in the two at-risk groups' population subgroups. Thus, the prevalence of hidden hunger for Europe should ideally be based on data from as many representative surveys of the various at-risk population subgroups, such as children and adolescents, women of reproductive age, pregnant women, older adults and immigrant and ethnic minority groups, as possible, and include micronutrients of public health relevance to the region.

Zero Hidden Hunger EU project will generate new data on prevalence and other aspects of hidden hunger in Europe

Thus, while addressing the public health problem of hidden hunger is a priority, it is not possible until data on the true prevalence of micronutrient deficiencies across the EU population and the causes of these deficiencies is available to predict and identify those most at risk. Without this information, discussions on how to meet dietary requirements for the priority micronutrients of public health concern, and improve their status, take place in a vacuum. This information is the key starting point towards defining food-based strategies for micronutrient deficiency prevention and health promotion throughout the life-course. In recognition of this, in 2023, the EC issued a research funding call for a large-scale (€10 M) project on eradication of micronutrient deficiencies in the EU, as part of its Horizon Europe programme. The *Cork Centre for Vitamin D and Nutrition Research* at University College Cork (Professor Kevin Cashman and Professor Mairead Kiely, Joint Coordinators), together with 18 collaborative partners across Europe, led a successful bid for this call. The four-year 'Tackling micronutrient malnutrition and hidden hunger to improve health in the EU' project, with *Zero Hidden Hunger EU* as its acronym (<http://www.zerohiddenhunger.eu>), commenced in early 2024. One of the key initial aims of *Zero Hidden Hunger EU* is to assess the true prevalence of micronutrient deficiencies, based on priority biomarker and micronutrient intake data in European populations, especially in high-risk population subgroups. It will do this by using existing high-quality data resources on population micronutrient status and intake for various European countries. As

with the *ODIN* vitamin D project, again national nutrition surveys and health surveys, as well as quality representative population cohorts, will be used in *Zero Hidden Hunger EU* to supply the needed data on micronutrient status, based on validated status markers, and intakes of the prioritised micronutrients of public health concern among the selected at risk groups within the EU population, namely, children and teenagers, women of reproductive age, pregnant women, older adults and ethnic minority groups. To complement this existing micronutrient status data, some new analysis in bio-banked blood samples will provide some key missing micronutrient status data for incorporation and use in the overall data collection. In this way, the *Zero Hidden Hunger EU* project will maximise previous EC and member state investment.

This new data on the true prevalence of micronutrient deficiencies across the EU, and information on their dietary causes, will then be utilised in the development of proposals for context-specific, foods-first, sustainable and cost-effective approaches for prevention of micronutrient deficiencies in Europe, using standardised approaches to data collection and novel dietary modelling to generate multiple predictions of the effect of food-based strategies to ensure adequate micronutrient intake, while optimising environmental outcomes. The project will also evaluate the cost effectiveness of these proposed mitigation strategies using the new prevalence data and dietary proposals. Greater detail of these planned activities will be provided elsewhere in due course.

Conclusions

There is a clear and pressing need to develop and implement inclusive policies aimed at ending hidden hunger, as well as other forms of malnutrition. In terms of tackling deficiency of vitamin D and other micronutrients – be it in relation to the quantification of the magnitude of the public health problem, establishing micronutrient intake requirement values, or proposing food-based strategies for the deficiency prevention, amongst other aspects, data is the life-blood. The present review provides additional insight into the critical importance of data, in its many forms, in the generation of the evidence-base that can inform recommendations, guidelines and other policy instruments for addressing micronutrient deficiency. Priority data can be translated into meaning and knowledge, which can inform policy decision-making and formulation, and these policies, to be effective, need to then be acted upon, if there is any chance of achieving the elimination of micronutrient deficiencies in the near future.

Acknowledgements. The oral presentation given at the inaugural Nutrition Society Congress, and this associated review article, overviewed research that has been conducted at the *Cork Centre for Vitamin D and Nutrition Research* at University College Cork together with many important collaborators nationally (including Irish Universities Nutrition Alliance; UCD; NICHE at UU, Coleraine; amongst others) and internationally (*Vitamin D Standardization Program*; our EU project collaborators, especially the *ODIN* and *Zero Hidden Hunger EU* projects; the Principal Investigators involved in our Individual Participant Data activities; as well as colleagues at the WHO and FAO in relation to vitamin D requirements; amongst others), all of whom have been key in terms of generation of data and knowledge on aspects of vitamin D and/or hidden hunger. Locally, Professor Mairead Kiely, as joint director of the Cork centre, has been at the forefront of much of the research activities which were included in the presentation/review. There is also a long list of key scientists and researchers, past and present, from the

Cork centre that have been instrumental in our research activities and the generation of the key data that was presented.

Author contributions. Kevin D. Cashman is the sole author of this review.

Financial support. No specific grant from any funding agency, commercial or not-for-profit sectors has been received for the preparation of this review article. The research referenced within the review has been on foot of many research grants and each is acknowledged in the associated citations used in the review.

Competing interests. The author does not have conflicts of interest to declare. In terms of disclosures, the author has previously served as a member of the FAO-WHO expert group on nutrient requirements for infants and young children aged 0–36 months and the UK Scientific Advisory Committee on Nutrition working group on vitamin D. He is currently a member of the Scientific Committee of the Food Safety Authority of Ireland (FSAI) and chairs its Public Health Nutrition sub-committee.

Ethical statement. None applicable.

References

- United Nations (2024) About the UN Decade of Action on Nutrition. <https://www.un.org/nutrition/about> (accessed 15 August 2024).
- Stevens GA, Beal T, Mbuya MNN, *et al.* (2022) Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: a pooled analysis of individual-level data from population-representative surveys. *Lancet Glob Health* **10**, e1590–e1599.
- Eggersdorfer M, Akobundu U, Bailey RL, *et al.* (2018) Hidden hunger: solutions for America's aging populations. *Nutrients* **10**, 1210. doi: [10.3390/nu10091210](https://doi.org/10.3390/nu10091210).
- Ngo J, Roman-Viñas B, Ribas-Barba L, *et al.* (2014) A systematic review on micronutrient intake adequacy in adult minority populations residing in Europe: the need for action. *J Immigr Minor Health* **16**, 941–950.
- United Nations (2024) SDG No. 2: Zero Hunger. <https://www.un.org/sustainabledevelopment/hunger/> (accessed 15 August 2024).
- Blackwell M (2023) 'Data is the new oil' ... so to speak. Mortgage Introducer September 2023. Page 26. https://issuu.com/keymedia/docs/mi_september-october_2023_emag?fr=xKAE9_5yMZA (accessed 31 October 2024).
- European Commission (1998) Report on Osteoporosis in the European Community – Action for Prevention Published by the European Commission in 11 languages. Manuscript completed in 1998. CE-09-97-915-EN-C ISBN 92-828-5333-0. Download PDF in English. https://www.osteoporosis.foundation/sites/infonehealth/files/2019-06/1998_EU_Report_ActionForPrevention_English.pdf (accessed 15 August 2024).
- Freaney R, McBrinn Y & McKenna MJ (1993) Secondary hyperparathyroidism in elderly people: combined effect of renal insufficiency and vitamin D deficiency. *Am J Clin Nutr* **58**, 187–191.
- Keane EM, Rochfort A, Cox J, *et al.* (1992) Vitamin-D-fortified liquid milk—a highly effective method of vitamin D administration for house-bound and institutionalised elderly. *Gerontology* **38**, 280–284.
- Meade A, Moloney M & O'Keeffe D (1986) Prevalence of vitamin D deficiency in the elderly in two rural areas in Ireland. *Ir Med J* **79**, 359.
- Vir SC & Love AH (1979) Nutritional status of institutionalized and noninstitutionalized aged in Belfast, Northern Ireland. *Am J Clin Nutr* **32**, 1934–1947.
- Cashman KD, Muldowney S, McNulty B, *et al.* (2013) Vitamin D status of Irish adults: findings from the National Adult Nutrition Survey. *Br J Nutr* **109**, 1248–1256.
- Sempos CT, Vesper HW, Phinney KW, *et al.* (2012) Vitamin D status as an international issue: national surveys and the problem of standardization. *Scand J Clin Lab Invest Suppl* **243**, 32–40.
- Binkley N, Krueger D, Cowgill CS, *et al.* (2004) Assay variation confounds the diagnosis of hypovitaminosis D: a call for standardization. *J Clin Endocrinol Metab* **89**, 3152–3157.
- Durazo-Arvizu RA, Tian L, Brooks SPJ, *et al.* (2017) The Vitamin D Standardization Program (VDSP) manual for retrospective laboratory standardization of serum 25-hydroxyvitamin D data. *J AOAC Int* **100**, 1234–1243. doi: [10.5740/jaoacint.17-0196](https://doi.org/10.5740/jaoacint.17-0196).
- Institute of Medicine (2011) *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press.
- Cashman KD, Kiely M, Kinsella M, *et al.* (2013) Evaluation of Vitamin D Standardization Program protocols for standardizing serum 25-hydroxyvitamin D data: a case study of the program's potential for national nutrition and health surveys. *Am J Clin Nutr* **97**, 1235–1242.
- Herrick KA, Storaardt RJ, Afful J, *et al.* (2019) Vitamin D status in the United States, 2011–2014. *Am J Clin Nutr* **110**, 150–157.
- Brooks SPJ, Greene-Finestone L, Whiting S, *et al.* (2017) An analysis of factors associated with 25-hydroxyvitamin D levels in white and non-white Canadians. *J AOAC Int* **100**, 1345–1354.
- Cashman KD, Kehoe L, Kearney J, *et al.* (2022) Adequacy of calcium and vitamin D nutritional status in a nationally representative sample of Irish teenagers aged 13–18 years. *Eur J Nutr* **61**, 4001–4014.
- Black LJ, Walton J, Flynn A, *et al.* (2015) Small increments in vitamin D intake by Irish adults over a decade show that strategic initiatives to fortify the food supply are needed. *J Nutr* **145**, 969–976.
- Report of the Scientific Committee of the Food Safety Authority of Ireland (2023) Vitamin D: Scientific Recommendations for 5 to 65 Year Olds Living in Ireland. <https://www.fsai.ie/publications/vitamin-d-scientific-recommendations-for-5-to-65-y> (accessed 15 August 2024).
- Report of the Scientific Committee of the Food Safety Authority of Ireland (2020) Vitamin D Scientific Recommendations for Food-Based Dietary Guidelines for Older Adults in Ireland. <https://www.fsai.ie/publications/vitamin-d-scientific-recommendations-for-food-base> (accessed 15 August 2024).
- Report of the Scientific Committee of the Food Safety Authority of Ireland (2020) Scientific Recommendations for Food-Based Dietary Guidelines for 1 to 5 Year-Olds in Ireland. <https://www.fsai.ie/publications/scientific-recommendations-for-food-based-dietary> (accessed 15 August 2024).
- Cashman KD & Kiely M (2011) Towards prevention of vitamin D deficiency and beyond: knowledge gaps and research needs in vitamin D nutrition and public health. *Br J Nutr* **106**, 1617–1627.
- Kiely M & Cashman KD (2018) Summary outcomes of the ODIN project on food fortification for vitamin D deficiency prevention. *Int J Environ Res Public Health* **15**, 2342. doi: [10.3390/ijerph15112342](https://doi.org/10.3390/ijerph15112342).
- Cashman KD, Dowling KG, Škrabáková Z, *et al.* (2016) Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr* **103**, 1033–1044.
- Cashman KD (2022) Global differences in vitamin D status and dietary intake: a review of the data. *Endocr Connect* **11**, e210282. doi: [10.1530/EC-21-0282](https://doi.org/10.1530/EC-21-0282).
- Cashman KD, Dowling KG, Škrabáková Z, *et al.* (2015) Standardizing serum 25-hydroxyvitamin D data from four Nordic population samples using the Vitamin D Standardization Program protocols: shedding new light on vitamin D status in Nordic individuals. *Scand J Clin Lab Invest* **75**, 549–561.
- Cashman KD (2020) Vitamin D deficiency: defining, prevalence, causes, and strategies of addressing. *Calcif Tissue Int* **106**, 14–29.
- O'Neill CM, Kazantzidis A, Ryan MJ, *et al.* (2016) Seasonal changes in vitamin D-effective UVB availability in Europe and associations with population serum 25-hydroxyvitamin D. *Nutrients* **8**, 533. doi: [10.3390/nu8090533](https://doi.org/10.3390/nu8090533).
- Rippin HL, Hutchinson J, Jewell J, *et al.* (2017) Adult nutrient intakes from current national dietary surveys of European populations. *Nutrients* **9**, 1288. doi: [10.3390/nu9121288](https://doi.org/10.3390/nu9121288).
- Rippin HL, Hutchinson J, Jewell J, *et al.* (2019) Child and adolescent nutrient intakes from current national dietary surveys of European populations. *Nutr Res Rev* **32**, 38–69.
- European Food Safety Authority Panel on Dietetic Products, Nutrition, and Allergies (NDA) (2010) Scientific opinion on principles for deriving and applying dietary reference values. *EFSA J* **8**, 1458. (30 pp.) www.efsa.europa.eu

35. Roman Viñas B, Ribas Barba L, Ngo J, *et al.* (2011) Projected prevalence of inadequate nutrient intakes in Europe. *Ann Nutr Metab* **59**, 84–95.
36. Cashman KD, Hill TR, Lucey AJ, *et al.* (2008) Estimation of the dietary requirement for vitamin D in healthy adults. *Am J Clin Nutr* **88**, 1535–1542.
37. Cashman KD, Wallace JM, Horigan G, *et al.* (2009) Estimation of the dietary requirement for vitamin D in free-living adults ≥ 64 years of age. *Am J Clin Nutr* **89**, 1366–1374.
38. Cashman KD (2014) A review of vitamin D status and CVD. *Proc Nutr Soc* **73**, 65–72.
39. Cashman KD, FitzGerald AP, Viljakainen HT, *et al.* (2011) Estimation of the dietary requirement for vitamin D in healthy adolescent white girls. *Am J Clin Nutr* **93**, 549–555.
40. Mortensen C, Damsgaard CT, Hauger H, *et al.* (2016) Estimation of the dietary requirement for vitamin D in white children aged 4–8 years: a randomized, controlled, dose-response trial. *Am J Clin Nutr* **104**, 1310–1317.
41. Smith TJ, Tripkovic L, Damsgaard CT, *et al.* (2016) Estimation of the dietary requirement for vitamin D in adolescents aged 14–18 years: a dose-response, double-blind, randomized placebo-controlled trial. *Am J Clin Nutr* **104**, 1301–1309.
42. O'Callaghan KM, Hennessy Á, Hull GLJ, *et al.* (2018) Estimation of the maternal vitamin D intake that maintains circulating 25-hydroxyvitamin D in late gestation at a concentration sufficient to keep umbilical cord sera ≥ 25 –30 nmol/l: a dose-response, double-blind, randomized placebo-controlled trial in pregnant women at northern latitude. *Am J Clin Nutr* **108**, 77–91.
43. Cashman KD, Ritz C, Adebayo FA, *et al.* (2019) Differences in the dietary requirement for vitamin D among Caucasian and East African women at Northern latitude. *Eur J Nutr* **58**, 2281–2291.
44. Ashwell M, Stone EM, Stolte H, *et al.* (2010) UK Food Standards Agency Workshop Report: an investigation of the relative contributions of diet and sunlight to vitamin D status. *Br J Nutr* **104**, 603–611.
45. Scientific Advisory Committee on Nutrition (2016) Report on Vitamin D and Health. http://www.sacn.gov.uk/pdfs/sacn_vitaminD_and_health_report_web.pdf (accessed 01 July 2016).
46. Mølgaard C, Larnkjaer A, Cashman KD, *et al.* (2010) Does vitamin D supplementation of healthy Danish Caucasian girls affect bone turnover and bone mineralization? *Bone* **46**, 432–439.
47. Öhlund I, Lind T, Hernell O, *et al.* (2017) Increased vitamin D intake differentiated according to skin color is needed to meet requirements in young Swedish children during winter: a double-blind randomized clinical trial. *Am J Clin Nutr* **106**, 105–112.
48. German Nutrition Society (2012) New reference values for vitamin D. *Ann Nutr Metab* **60**, 241–246.
49. NORDEN Nordic Nutrition Recommendations (2013) 5th Edition (NNR5) – Vitamin D. <http://www.slv.se/en-gb/Startpage-NNR/Public-consultation11/> (accessed August 2013).
50. EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies) (2016) Scientific Opinion on Dietary Reference Values for Vitamin D. <https://doi.org/10.2903/j.efsa.2016> (accessed February 2017).
51. Cashman KD (2018) Vitamin D requirements for the future-lessons learned and charting a path forward. *Nutrients* **10**, 533. doi: [10.3390/nu10050533](https://doi.org/10.3390/nu10050533)
52. Cashman KD, Ritz C, Kiely M, *et al.* (2017) Improved dietary guidelines for vitamin D: application of individual participant data (IPD)-level meta-regression analyses. *Nutrients* **9**, 469. doi: [10.3390/nu9050469](https://doi.org/10.3390/nu9050469).
53. Cashman KD, Kiely ME, Andersen R, *et al.* (2021) Individual participant data (IPD)-level meta-analysis of randomised controlled trials with vitamin D-fortified foods to estimate Dietary Reference Values for vitamin D. *Eur J Nutr* **60**, 939–959.
54. Cashman KD, Kiely ME, Andersen R, *et al.* (2022) Individual participant data (IPD)-level meta-analysis of randomised controlled trials to estimate the vitamin D dietary requirements in dark-skinned individuals resident at high latitude. *Eur J Nutr* **61**, 1015–1034.
55. Rios-Leyvraz M, Thacher TD, Dabas A, *et al.* (2024) Serum 25-hydroxyvitamin D threshold and risk of rickets in young children: a systematic review and individual participant data meta-analysis to inform the development of dietary requirements for vitamin D. *Eur J Nutr* **63**, 673–695.
56. Rios-Leyvraz M, Martino L & Cashman KD (2024) The relationship between vitamin D intake and serum 25-hydroxyvitamin D in young children: a meta-regression to inform WHO/FAO vitamin D intake recommendations. *J Nutr* **154**, 1827–1841.
57. Allen L, de Benoist B, Dary O, *et al.* (2006) Guidelines on Food Fortification with Micronutrients. World Health Organization and Food and Agriculture Organization of the United Nations. Geneva. http://apps.who.int/iris/bitstream/10665/43412/1/9241594012_eng.pdf (accessed 28 June 2016).
58. Hayes A & Cashman KD (2017) Food-based solutions for vitamin D deficiency: putting policy into practice and the key role for research. *Proc Nutr Soc* **76**, 54–63.
59. Itkonen ST, Skaffari E, Saaristo P, *et al.* (2016) Effects of vitamin D₂-fortified bread v. supplementation with vitamin D₂ or D₃ on serum 25-hydroxyvitamin D metabolites: an 8-week randomised-controlled trial in young adult Finnish women. *Br J Nutr* **115**, 1232–1239.
60. Cashman KD, Kiely M, Seamans KM, *et al.* (2016) Effect of ultraviolet light-exposed mushrooms on vitamin D status: liquid chromatography-tandem mass spectrometry reanalysis of biobanked sera from a randomized controlled trial and a systematic review plus meta-analysis. *J Nutr* **146**, 565–575.
61. Jakobsen J, Smith C, Bysted A, *et al.* (2019) Vitamin D in wild and farmed Atlantic salmon (*Salmo Salar*) – what do we know? *Nutrients* **11**, 982. doi: [10.3390/nu11050982](https://doi.org/10.3390/nu11050982).
62. Hayes A, Duffy S, O'Grady M, *et al.* (2016) Vitamin D-enhanced eggs are protective of wintertime serum 25-hydroxyvitamin D in a randomized controlled trial of adults. *Am J Clin Nutr* **104**, 629–637.
63. Duffy SK, Kelly AK, Rajauria G, *et al.* (2018) The use of synthetic and natural vitamin D sources in pig diets to improve meat quality and vitamin D content. *Meat Sci* **143**, 60–68.
64. Duffy SK, O'Doherty JV, Rajauria G, *et al.* (2017) Cholecalciferol supplementation of heifer diets increases beef vitamin D concentration and improves beef tenderness. *Meat Sci* **134**, 103–110.
65. Duffy SK, O'Doherty JV, Rajauria G, *et al.* (2018) Vitamin D-biofortified beef: a comparison of cholecalciferol with synthetic versus UVB-mushroom-derived ergosterol as feed source. *Food Chem* **256**, 18–24.
66. Manios Y, Moschonis G, Mavrogianni C, *et al.* (2017) Reduced-fat Gouda-type cheese enriched with vitamin D₃ effectively prevents vitamin D deficiency during winter months in postmenopausal women in Greece. *Eur J Nutr* **56**, 2367–2377.
67. Grønberg IM, Tetens I, Christensen T, *et al.* (2020) Vitamin D-fortified foods improve wintertime vitamin D status in women of Danish and Pakistani origin living in Denmark: a randomized controlled trial. *Eur J Nutr* **59**, 741–753.
68. Dunlop E, Kiely ME, James AP, *et al.* (2021) Vitamin D food fortification and biofortification increases serum 25-hydroxyvitamin D concentrations in adults and children: an updated and extended systematic review and meta-analysis of randomized controlled trials. *J Nutr* **151**, 2622–2635.
69. Cashman KD & O'Neill CM (2024) Strategic food vehicles for vitamin D fortification and effects on vitamin D status: a systematic review and meta-analysis of randomised controlled trials. *J Steroid Biochem Mol Biol* **238**, 106448. doi: [10.1016/j.jsbmb.2023.106448](https://doi.org/10.1016/j.jsbmb.2023.106448)
70. Nyakundi PN, Némethné Kontár Z, Kovács A, *et al.* (2023) Fortification of staple foods for household use with vitamin D: an overview of systematic reviews. *Nutrients* **15**, 3742. doi: [10.3390/nu15173742](https://doi.org/10.3390/nu15173742).
71. Buttriss JL, Lanham-New SA, Steenson S, *et al.* (2022) Implementation strategies for improving vitamin D status and increasing vitamin D intake in the UK: current controversies and future perspectives: proceedings of the 2nd Rank Prize Funds Forum on vitamin D. *Br J Nutr* **127**, 1567–1587.
72. Allen RE, Dangour AD, Tedstone AE, *et al.* (2015) Does fortification of staple foods improve vitamin D intakes and status of groups at risk of deficiency? A United Kingdom modeling study. *Am J Clin Nutr* **102**, 338–344.
73. McCourt A, McNulty BA, Walton J, *et al.* (2020) Efficacy and safety of food fortification to improve vitamin D intakes of older adults. *Nutrition*

- 75–76, 110767. doi: [10.1016/j.nut.2020.110767](https://doi.org/10.1016/j.nut.2020.110767). Epub 2020 Feb 14. Erratum in: *Nutrition*. 2020 Jul - Aug;75–76:110837.
74. Das JK & Padhani ZA (2022) Alleviating hidden hunger: an infallible bridge to improved health and nutrition. *Lancet Glob Health* **10**, e1539–e1540. doi: [10.1016/S2214-109X\(22\)00421-1](https://doi.org/10.1016/S2214-109X(22)00421-1).
75. World Health Organization (WHO) (1991) *National Strategies for Overcoming Micronutrient Malnutrition*. Geneva: World Health Organization.
76. Food and Agriculture Organisation (FAO) (2013) The State of Food and Agriculture. <https://fao.org/4/i3300e/i3300e.pdf> (accessed 27 May 2024).
77. International Food Policy Research Institute (IFPRI) (2014) Chapter 3 Addressing the Challenge of Hidden Hunger. In ‘2014 Global Hunger Index – The Challenge of Hidden Hunger’ pp. 20–27. <https://www.ifpri.org/sites/default/files/ghi/2014/contents.html> (accessed 27 May 2024).