

Abundance and genetic variation of the coral-killing cyanobacteriosponge *Terpios hoshinota* in the Spermonde Archipelago, SW Sulawesi, Indonesia

ESTHER VAN DER ENT^{1,2}, BERT W. HOEKSEMA¹ AND NICOLE J. DE VOOGD^{1,3}

¹Department of Marine Zoology, Naturalis Biodiversity Center, P.O. Box 9517, 2300RA Leiden, the Netherlands, ²Department of Biomarine Sciences, Utrecht University, the Netherlands, ³Institute for Ecosystem Dynamics, University of Amsterdam, 1090 GE Amsterdam, the Netherlands

The cyanobacteriosponge Terpios hoshinota is expanding its range across the Indo-Pacific. This species can have massive outbreaks on coral reefs, actively kill corals it overgrows and affect the entire benthic community. Although it has received much attention on the post-outbreak follow-up, little is known about its ecology, habitat preferences, and the possible environmental triggers that cause its outbreaks. We present a baseline study in the Spermonde Archipelago, Indonesia, where T. hoshinota was first observed in 2012. We surveyed 27 reefs and recorded patches between 14 and 217 cm², at four reef sites (~15% cover). The sponge was found on both mid-shelf and outer shelf reefs but not close to the coast and the city of Makassar. Differences in benthic community structure, as well as spatial variables relating to the on-to-offshore gradient in the Spermonde archipelago, neither constrained nor promoted its expansion. Patches of the sponge were mostly overgrowing branching corals, belonging to Acroporidae species. Genetic variation within T. hoshinota was studied by sequencing partitions of the mitochondrial CO1 and nuclear ribosomal 28S gene. Two haplotypes were found within the Spermonde archipelago, which differed from the CO1 sequence in GenBank. The present study provides an indication of habitat preferences of T. hoshinota in non-outbreak conditions, although it is still unclear which environmental conditions may lead to the onset of its outbreaks.

Keywords: Competitive sponge-coral interaction, outbreak, benthic community, CO1, 28S

Submitted 21 November 2014; accepted 4 March 2015; first published online 11 May 2015

INTRODUCTION

Coral reefs worldwide have been in decline for decades under increasing anthropogenic disturbances, such as overfishing, pollution and coral bleaching resulting from global climate change (Pandolfi *et al.*, 2000). It is crucial to gain more knowledge on the multitude of biotic and abiotic stressors that threaten coral reefs in order to successfully implement conservation plans for their protection.

The encrusting cyanobacteriosponge *Terpios hoshinota* Rützler & Muzik (1993) has come to the attention as another potential threat to Indo-Pacific coral reefs. Although sponges are an important component of coral reef biodiversity, and together with stony and soft corals form complex three-dimensional structures that provide shelter to numerous marine organisms, they are also very successful competitors for space on reefs, at the expense of corals (Aerts & van Soest, 1997; de Voogd *et al.*, 2004; Wulff, 2012). Generally, sponges have a competitive advantage over corals because they can successfully resist coral defences and persevere in

environments that are unfavourable to corals (Chadwick & Morrow, 2011). Additionally, the sponge *T. hoshinota* can actively kill and overgrow live coral tissue. This sponge occasionally exhibits outbreaks that can cover huge areas, resulting in the mass mortality of corals and other sessile organisms (Rützler & Muzik, 1993; Liao *et al.*, 2007). During outbreaks, *T. hoshinota* can rapidly decimate corals due to growth rates of approximately 1–2 mm per day, and its capacity to occupy reef substrata for up to several years (Plucer-Rosario, 1987; Schils, 2012). This may prevent the settlement of coral larvae, hereby hindering reef recovery (Bryan, 1973; Plucer-Rosario, 1987). It is clear that this sponge is very persistent on coral reefs, and once it has invaded a new area, it appears to remain there permanently (Reimer *et al.*, 2011a, b). However, the outbreaks do not always lead to irreversible reef degradation because coral recovery has also been observed with the near disappearance of *T. hoshinota* (Plucer-Rosario, 1987; Reimer *et al.*, 2011b).

Terpios hoshinota has been described from the Ryukyu Archipelago, Japan and Guam (Rützler & Muzik, 1993), but outbreaks of the sponge have been reported and studied earlier in Guam, the Caroline Islands, the Philippines, Taiwan and American Samoa (Bryan, 1973; Plucer-Rosario, 1987). More recently, the sponge has been reported in the Great Barrier Reef (Fujii *et al.*, 2011), Indonesia (de Voogd

Corresponding author:
N.J. de Voogd
Email: Nicole.deVoogd@naturalis.nl

et al., 2013), Malaysia (Hoeksema et al., 2014b) and the Maldives (Montano et al., 2015). Most information about *T. hoshinota* is obtained through studies during and after outbreaks. However, pre-outbreak data on *T. hoshinota* are still lacking, and baseline data concerning its ecology, habitat preferences and outbreak triggers are therefore urgently needed.

Recently, small patches of *T. hoshinota* were observed on an offshore reef in the Spermonde Archipelago, Makassar Strait, Indonesia (de Voogd et al., 2013). Its identity was confirmed by morphological analyses and cytochrome oxidase I mitochondrial DNA sequences. The reefs of the Spermonde Archipelago have been extensively studied during sponge surveys since 1997 (de Voogd et al., 2004, 2005, 2006; Cleary & de Voogd, 2007; Hoeksema et al., 2014a, b), but *T. hoshinota* has not been recorded before 2012. Hence, this area is highly suitable for a baseline study. Future surveys in this area may reveal whether the occurrence of *T. hoshinota* is stable, or whether its presence will inevitably always lead to an outbreak event.

The objectives for the present baseline study were (1) to map the current distribution of *T. hoshinota* in the Spermonde Archipelago; (2) to study its distribution and abundance in relation to cross-shelf variation in benthic community structure; (3) to record its coral substrate preference under regular (non-outbreak) conditions; and (4) to analyse the genetic diversity within a *T. hoshinota* population by comparing sequences obtained by use of the Folmer partition and the I3-M11 partition of the mitochondrial CO1 and the

D3–D5 region of the nuclear ribosomal 28S gene with sequences from the marine region of China available in GenBank.

MATERIALS AND METHODS

Study area

The Spermonde Archipelago in south-west Sulawesi is situated within the Coral Triangle, the centre of maximum marine benthic biodiversity (Hoeksema, 2007). The shelf ranges 32–56 km in width and approximately 128 km in length and is covered by over ~100 coral cay islands and submerged reefs (Figure 1). Its depth ranges towards a maximum depth of > 50 m near the offshore shelf edge in the Makassar Strait. Several near-shore reefs are located close to the port of Makassar, which houses over a million people and exerts environmental stress on the nearby reefs by sewage seepage and industrial pollution. The mouth of the Jene Berang River, south of Makassar, discharges terrigenous sediments and inland-pollution into the Spermonde Archipelago. The near-shore reefs off Makassar, up to 4 km from the coastline, are directly influenced by this river through high sedimentation and freshwater discharge causing these reefs to have shallower euphotic zones, higher nutrient levels, higher silt and sand content and lower salinity (Edinger et al., 1998; Renema & Troelstra, 2001; Renema et al., 2001; Cleary




Fig. 1. Bathymetric map of the Spermonde Archipelago showing ○ = surveyed sites. ⊙ *T. hoshinota* observation. Inset showing the location of the Spermonde Archipelago in relation to the island of Sulawesi.

et al., 2005), but they are more sheltered from wave energy and cold-water upwelling (Hoeksema, 2012a). Most of the coral cay islands are inhabited by fishing communities, who may use destructive fishing methods such as blast fishing (Pet-Soede *et al.*, 1999; Hoeksema, 2004). Furthermore, corals in this area are being collected for the aquarium trade (Knittweis & Wolff, 2010). Although the reefs around the most offshore island on the shelf, Kapoposang, have been designated a Marine Protected Area (MPA) category V by IUCN, there is little surveillance in the area.

The Spermonde Archipelago thus presents a diverse array of environmental conditions and cross-shelf diversity, making it a very suitable area for research. It is one of the best-explored marine regions in Indonesia and has been well-documented in literature, as extensive marine biological and physical geographic studies have been conducted during the last 35 years (Moll, 1983; Verheij & Erftemeijer, 1993; Verheij & Prud'homme van Reine, 1993; Erftemeijer, 1994; Sterrenburg *et al.*, 1995; de Voogd *et al.*, 2004, 2005, 2006; Cleary *et al.*, 2005; Becking *et al.*, 2006; Cornils *et al.*, 2011; Hoeksema & Crowther, 2011; Sawall *et al.*, 2011; Hoeksema, 2012a, b; Hoeksema & de Voogd, 2012).

Benthic community of reef sites

A total of 27 reef sites, spread over 20 islands in the Spermonde Archipelago, were thoroughly surveyed by scuba diving for the presence of *T. hoshinota* from 15 August to 2 September 2013. The sessile benthic community of the surveyed reefs was assessed by carrying out line intercept transects (LIT). Rather than determining the benthic community at species level, the LIT method is based on classification of sessile life forms in broader, environmentally significant categories, in order to give an impression of the general state of coral reefs in the Spermonde Archipelago (English *et al.*, 1997). At each site two 30-m parallel transects were laid down, at 3 and 6 m. The transects were recorded using a digital underwater camera. The LIT recordings were analysed and percentages in which the different benthic life form categories occurred over the transect line were calculated, which were categorized as 'live coral', 'dead coral (with algae)', 'algae', 'sponges', 'rubble', 'rock', 'sand' and 'others'. The category 'live coral' was comprised of the categories 'branching', 'encrusting', 'foliose', 'massive' and 'submassive' corals. The reef areas surrounding the transects were surveyed for patches of *T. hoshinota* by using the Roving Diver Technique (RDT) (Munro, 2005; Hoeksema & Koh, 2009).

Sampling

Sponge identification *in situ* was based on the morphological characteristics given by Rützler & Muzik (1993) and by comparison of sponge specimens with those presented in pictures of *T. hoshinota* in other localities (Reimer *et al.*, 2011a, b; de Voogd *et al.*, 2013). All sponges were photographed using a digital camera, along with a ruler for scale. Additionally, the surrounding coral substrates were photographed for coral host identification. Fragments of approximately 2 cm² of sponge tissue were collected for DNA and morphological analyses. Fragments were immediately stored in ethanol (96%) and kept at -20°C. Vouchers are housed in the sponge collection of Naturalis Biodiversity Center, Leiden, the Netherlands.

Molecular analyses

DNA was extracted from the sponge tissue samples using the DNeasy Blood & Tissue kit (Qiagen) following the instructions of the manufacturer. Partitions of a mitochondrial and nuclear marker were amplified: two regions of the cytochrome oxidase subunit 1 (CO1), the standard DNA-barcoding partition called the 'Folmer partition', as well as the I3-M11 partition, and the D3-D5 region of the nuclear ribosomal 28S gene (28S).

The Folmer partition of CO1 was amplified by using a specific forward primer designed for the family *Suberitidae*, SUB-CO1F (5'-GGAATGATCGGGACAGCTTTTAGCATG-3') designed by Becking *et al.* (2013) and the degenerated reverse primer from Folmer *et al.* (1994) designed by Meyer *et al.* (2005), dgHCO2198 (5'-TAAACTTCAGGGTGACCAARAAYCA-3'). These primers amplified a fragment of 600 bp. Amplifications were carried out in 25 µL reaction volumes containing 15.75 µL of MilliQ water, 2.5 µL 10× reaction buffer, 1 µL MgCl₂ (25 mM), 2 µL dNTPs (2.5 mM), 1 µL of each primer (10 µM), 0.25 µL Taq polymerase, and 1.5 µL of DNA. The PCR profile was: initial denaturing step of 94°C for 4 min, 38 cycles of denaturing at 94°C for 30 s, annealing at 54°C for 30 s, extension at 72°C for 40 s, and a final extension step at 72°C for 4 min. The I3-M11 partition of CO1 was amplified by a nested approach using primers designed by Misof *et al.* (2000) and Erpenbeck *et al.* (2004). The first primers used were C1J2165 (5'-GAAGTTTATATTTTAAATTTTACCDGG-3') and C1Npor2760 (5'-TCTAGGTAATCCAGCTAAACC-3'), which amplified a fragment of 544 bp. Amplifications were carried out in 25 µL reaction volumes containing 15.75 µL of MilliQ water, 2.5 µL 10× reaction buffer, 1 µL MgCl₂ (25 mM), 2 µL dNTPs (2.5 mM), 1 µL of each primer (10 µM), 0.25 µL Taq polymerase and 1.5 µL of DNA. The PCR profile was: initial denaturing step of 94°C for 4 min, 38 cycles of denaturing at 94°C for 30 s, annealing at 50°C for 30 s, extension at 72°C for 45 s, and a final extension step at 72°C for 4 min. Then, 1 µL of the obtained PCR product was used as a DNA template in a consecutive amplification using primers CO1porF1 (5'-CCNCANTTNKCNMNAAAAAACA-3') and CO1porR1 (5'-AANTGNTGNGGRAARAANG-3'), which amplified a fragment of 473 bp. Amplifications were carried out in 25 µL reaction volumes containing 15.75 µL of MilliQ water, 2.5 µL 10× reaction buffer, 1 µL MgCl₂ (25 mM), 2 µL dNTPs (2.5 mM), 1 µL of each primer (10 µM), 0.25 µL Taq polymerase and 1.5 µL of DNA. The PCR profile was: initial denaturing step of 94°C for 4 min, 35 cycles of denaturing at 94°C for 30 s, annealing at 45°C for 30 s, extension at 72°C for 45 s and a final extension step at 72°C for 4 min. The D3-D5 region of 28S was amplified using primers McCormack & Kelly (2002), RD3A (5'-GACCCGTCTTGAAACACGA-3') and RD5B2 (5'-ACACACTCCTTAGCGGA-3'), which amplified a fragment of 596 bp. Amplifications were carried out in 25 µL reaction volumes containing 8.9 µL of MilliQ water 5 µL 5× Phire[®] reaction buffer, 5 µL Q-solution, 2 µL dNTPs (2.5 mM), 1.3 µL of each primer (10 µM), 0.5 µL Phire[®] Hotstart II Taq polymerase and 1 µL DNA. The PCR profile was: initial denaturing step of 98°C for 30 s, 35 cycles of denaturing at 98°C for 5 s, annealing at 50°C for 5 s, extension at 72°C for 15 s and a final extension step at 72°C for 5 min.

Following PCR, amplification success was checked on a 1.5% agarose gel. PCR products were then sent to Macrogen Europe for sequencing. The obtained sequences were checked, trimmed and aligned using Geneious version 4.8.3. The poriferan origin of the sequences was confirmed through BLAST searches in GenBank, which revealed best matches of the CO1 and 28S data with sequences of *T. hoshinota* (KJ008098; Yongxing Island, South China Sea) and other species of the family Suberitidae, respectively. Haplotype and nucleotide diversities were calculated for both the Folmer partition as I3-M11 partition using DnaSP (version 5.0, Librado & Rozas, 2009).

Additionally, one sequence of the Folmer partition of CO1 of a *T. hoshinota* specimen that was sampled during a previous survey in the Spermonde Archipelago (2012) was used for comparison, as well as one sequence of sponge tissue sampled during fieldwork in the Thousand Islands, Java (de Voogd et al., 2013).

RESULTS

Presence of *Terpios hoshinota* in the Spermonde Archipelago

A total of 18 patches of *T. hoshinota* were found at four of the 27 surveyed sites (Figures 1 & 2; Table 1). Three patches of *T. hoshinota* were found at Samalona Reef (~7.5 km from Makassar), varying in size between 86 and 217 cm² and a total cover of ~431 cm². *Terpios hoshinota* was most abundant on the reef of Pulau Badi (distance from Makassar ~23.5 km) with 10 individuals. Patch size varied between 14 and 305 cm² and the total covered reef area was ~667 cm². Small patches of *T. hoshinota* were also observed at the outer reef of Lankai (~37 km from Makassar), where it was first recorded during the survey in 2012. These patches varied between 50 and 194 cm² in size, covering a total reef area of 620 cm². Finally a single patch of ~172 cm² was found within the MPA of Kapoposang (~71 km from Makassar). The sponge was not encountered on surveyed reefs that were most near shore, i.e. Lae-Lae, Bone Lola and Bone Baku (Figure 1).

Distribution in relation to the benthic community

Some variables of the benthic community composition showed a distinct distribution with respect to the distance to Makassar (Figure 3). The total live coral cover at the inner-shelf reefs, ≤10 km away from Makassar, was low (ranging from 9 to 15%). Over the mid-shelf it varied greatly, ranging between 6% at Lumulumu (3 m depth) and 68% at Barrang Lompo (3 m depth). Although Barrang Lompo is one of the largest and most densely populated islands in the Spermonde Archipelago, the reef was largely covered by large foliose corals (35%; Figure 3), spanning the entire northern reef area. The outer-shelf reefs had a lower live coral cover (7.5–30%). The total live coral cover encompassed all major coral growth forms (Figure 4). Branching and massive corals were the dominant coral growth forms, although most were small (<20 cm). Furthermore, the total algae cover showed high variation, in the range of 1–52%, although it was

generally high throughout the Spermonde Archipelago. The dead coral rock cover (also bare hard substrate) varied throughout the archipelago (range 0–20%). The benthic community composition was not significantly different between the two depths for any category except massive coral cover, which showed a higher cover at 3 m depth (two-tailed *t*-test; *P* = 0.02). The benthic environment of the four sites where the sponge *T. hoshinota* was found is also depicted (Figures 3 & 4).

Substrate preferences

The coral substrate of each observed *T. hoshinota* sponge was identified (Table 1) following Veron (2000). All *T. hoshinota* patches were overgrowing live corals, most of which belonged to species of the genera *Acropora*, *Isopora* and *Montipora* (Acroporidae), *Seriatopora* and *Pocillopora* (Pocilloporidae) and the species *Pavona venosa* (Agariciidae) (Figure 2).

Genetic diversity

All sequences were submitted to NCBI GenBank under accession numbers KP764919 (28S), KP764915–KP764916 (Folmer partition of CO1) and KP764917–KP764918 (I3-M11 partition of CO1).

Final alignments for the sponge *T. hoshinota* were obtained for the Folmer partition of CO1 of 600 bp (Tables 2 & 3). These alignments revealed five variable sites, which all exhibited C-T transitions and all resulting in silent mutations. This sequence variation revealed two haplotypes in the sequences obtained from samples from the Spermonde Archipelago, separating haplotype C2, comprising three specimens from the mid-shelf island of Pulau Badi and one specimen from the outer-shelf island of Kapoposang, from haplotype C3, comprising one specimen from the inner-shelf island of Samalona and three specimens from the outer-shelf island of Lankai. The sequence obtained from Lankai during the 2012 survey, was identical to haplotype C3. The sequence from the Thousand Islands, Java, however was similar to haplotype C2. The sequence from GenBank of the Folmer partition of CO1 of *T. hoshinota* (KJ008098) is 1 bp different from C2 and 5 bp different from C3, separating it in another haplotype, C1. Mitochondrial diversity based on the sequences were 0.636 (SD = 0.089) (haplotype diversity) and 0.00394 (SD = 0.00052) (nucleotide diversity).

Final alignments for the I3-M11 partition of CO1 of 473 bp were obtained from all eight samples from the Spermonde Archipelago (Tables 2 & 3). This region exhibited almost no sequence variation; only one variable site was observed, resulting in two haplotypes. These two haplotypes, I1 and I2, resulted in the same separation of specimens as haplotype C2 and C3, respectively. The two haplotypes resulted from a C-T transition at position 61 of the fragment. This missense mutation causes the encoding of different amino acids, creating cysteine in I1, and arginine in I2. Mitochondrial diversity based on the sequences were 0.571 (SD = 0.094) (haplotype diversity) and 0.00121 (SD = 0.00020) (nucleotide diversity).

The sequences of the D3–D5 region of 28S showed no nuclear variation within the specimens from the Spermonde Archipelago.

Since the Folmer partition and the I3-M11 partition are both located on the same mitochondrial marker (CO1), the haplotypes found for these partitions could be combined,




Fig. 2. (A) Patch of *Terpios hoshinota* with directional growth to the left, toward a live coral colony of *Acropora retusa*. A colony of *Seriatopora caliendrum* is already overgrown (although a few living branches can still be seen). Filamentous algae start to cover the dead coral substrate. (B) A thick tissue thread growing away from the substrate (encircled). The encircled tissue was observed to grow only on the site of the coral that was exposed to sunlight, and not underneath it. (C) The sponge shows discolouration at the growth front with the coral *Acropora muricata*. (D) Hairy tips can be seen at the growth front in an interaction with the coral.

creating two haplotypes in total, separating haplotype C2I1, comprising specimens from the mid-shelf island of Pulau Badi and the outer-shelf island of Kapoposang, from haplotype C3I2, comprising specimens from the inner-shelf island of Samalona and the outer-shelf island of Lankai.

DISCUSSION

Since its first sighting in 2012 in the Spermonde Archipelago on the reef of Lankai (de Voogd *et al.*, 2013), *T. hoshinota* has been advancing throughout the archipelago. During the survey in 2012 it was found at only one out of 14 surveyed reefs in the Spermonde Archipelago (~7%), on the off-shore reef of Lankai. The present study confirms the occurrence of *T. hoshinota* in the Spermonde Archipelago at four of the 27 surveyed reef sites (~15%).

The sites where *T. hoshinota* has been observed during the present survey are all environmentally and spatially very different from each other (Edinger *et al.*, 1998; Renema *et al.*,

2001; Cleary *et al.*, 2005; Cleary & de Voogd, 2007; Hoeksema, 2012a). The sponge was found on the mid-shelf reef at Samalona and Pulau Badi, as well as at the outer edge of the Spermonde Archipelago, at Lankai and Kapoposang, indicating that it can also occur in relatively pristine reefs. These findings are supported by a survey of the Ryukyu Archipelago by Reimer *et al.* (2011a), where *T. hoshinota* was also found in more pristine reefs, as well as reefs that suffered under anthropogenic stressors. Schils (2012) recorded an outbreak after volcanic eruptions, indicating that *T. hoshinota* and its symbiotic cyanobacteria benefit from the induced nutrient enrichment by volcanic ash. A massive *T. hoshinota* outbreak in an area with high turbidity and 'murky' water was also recorded by Rützler & Muzik (1993). However, the present study showed no large variation in patch size between the four sites where *T. hoshinota* was observed, indicating that no clear relation can be found between the size of the sponges and the distance offshore.

In the present study, the sponge was observed across the whole shelf width except for the inner-shelf reefs, near the

Table 1. Specifications of locations where *Terpios hoshinota* was found in the Spermonde Archipelago, Indonesia, including the coral substrates that the individual patches were covering. (◆) Indicates the patches that were sampled for DNA analyses.

Locality	Coordinates	Distance from Makassar (km)	Visibility during dive	Depth of patches (m)	Size of patch (cm ²)	Voucher RMNH	Scleractinian substrate
Samalona	5°7'21.17"S	7.3	17 m	3	216.8		<i>Isopora brueggemanni</i> (Brook, 1893)
	119°20'21.22"E				85.6		<i>Acropora muricata</i> (Linnaeus, 1758)
Pulau Badi	4°58'10.53"S 119°16'55.04"E	23.5	19 m	6	128.6	POR8729	<i>Acropora muricata</i> ◆
					22.7	POR8730	<i>Acropora muricata</i> ◆
					47.3		<i>Acropora muricata</i> /dead coral
					58.6		<i>Montipora foliosa</i> (Pallas, 1766)/ dead coral
					32.8		<i>Seriatopora hystrix</i> (Dana, 1846)/ dead <i>Acropora</i>
					14.3		<i>Montipora stellate</i> (Bernard, 1897)
					18.3		<i>Montipora capitata</i> (Dana, 1846)
					38.0		<i>Montipora stellate</i>
105.6		<i>Seriatopora hystrix</i>					
305.4	POR8731	Dead <i>Acropora</i> ◆					
23.5	POR8732	<i>Seriatopora hystrix</i> / <i>Montipora foliosa</i> ◆					
Lankai	5°1'44.58"S 119°5'8.79"E	37.2	17 m	3	187.1	POR8733	Dead <i>Acropora</i> ◆
					189.5	POR8734	<i>Pocillopora verrucosa</i> (Ellis & Solander, 1786)◆
					49.6		<i>Acropora tenuis</i> (Dana, 1846)
					193.5	POR8735	<i>Seriatopora caliendrum</i> / <i>Acropora retusa</i> (Dana 1846)◆
Kapoposang	4°41'28.77"S 118°56'43.95"E	71.3	30+ m	3	172.2	POR8736	<i>Pavona venosa</i> (Ehrenberg, 1834)◆

city of Makassar, where the photic zone was much shallower and visibility during dives was less than 5 m. A previous study in the Spermonde Archipelago showed that water velocity and salinity and the variation in water velocity and temperature increased from the inner- to the outer-shelf due to the decreasing influence of the Jene Berang river and to increasing oceanic influences, which had a distinct influence on the cross-shelf sponge community (Cleary & de Voogd, 2007). While *T. hoshinota* in the present study was found at various locations across the shelf, indicating that it can survive under different abiotic conditions, it is not clear whether these abiotic variables constrain or promote the expansion of the sponge.

The benthic community did not vary between the two surveyed depths with exception of the cover of massive corals. *Terpios hoshinota* was found both at shallow depth (3 m) and deeper on the reef slope (6 m). Reimer *et al.* (2011a) also recorded *T. hoshinota* mostly on shallow reefs (<5 m; 87%) although it was also recorded as deep as 17 m. The largest patches in the present study were observed at Pulau Badi at 6 m depth, which is arguably still relatively shallow and confirms *T. hoshinota* as a shallow reef sponge. Its symbiotic relationship with intercellular bacteria (5×10^5 cells cm⁻²), mainly cyanobacteria (61–98%) also makes it essential for *T. hoshinota* to grow in the light (Rützler & Muzik, 1993; Tang *et al.*, 2011; Hoeksema *et al.*, 2014b). *Terpios hoshinota* was not observed beyond a depth of 6 m in the present study.

The benthic community shows much cross-shelf variation, although both live coral, algae and rubble were dominant on all reefs. The cover of rubble, caused by storms, anchor damage and blast fishing, was generally high, up to 83%, throughout the Spermonde Archipelago. Reefs on the inner-shelf, less than 10 km away from Makassar, showed a very low cover of live coral, while over the mid-shelf area the cover varied extremely between sites and between all benthic variables, and the most off-shore reef had a relatively low coral cover again. A similar unimodal pattern was observed by Cleary *et al.* (2008) in the Thousand Islands, Jakarta, where *T. hoshinota* has also been recorded. It is clear that the Spermonde Archipelago suffers from human impact. Although branching and massive corals were the dominant coral growth forms, most branching corals that occurred mid-shelf were very small colonies (<20 cm), growing on top of large piles of coral rubble. Thus the high cover of branching corals does not imply that healthy reefs were present, which could sustain large fish populations, but rather that reefs in the Spermonde Archipelago are in the process of recovery.

The four sites where patches of *T. hoshinota* were discovered also varied distinctly from each other in benthic community structure; the sponges were found at sites with both high, intermediate and low cover of live corals, algae, rubble and sand. This shows that the sponge can survive within different benthic community structures, at least prior to an outbreak, but it does not explain why it has not expanded to other reefs in the Spermonde. Since an outbreak has not occurred




Fig. 3. Variation in the cover of live coral, total algae, dead coral (with algae), coral rock, rubble and sand, as a function of distance from Makassar.

in either of these structurally variable sites, *Terpios hoshinota* outbreaks are probably not related to (variation in) benthic community structure. It is noteworthy that at all four sites, *T. hoshinota* was almost exclusively found on branching corals, yet the benthic community of the reefs where they were found had a generally low cover of branching corals; with the exception of Pulau Badi (16%), all sponges were found at sites with a branching coral cover of less than 2%. During massive outbreaks *T. hoshinota* is known to overgrow any coral form and other benthic structures (Rützler & Muzik, 1993; Reimer *et al.*, 2011a).

Terpios hoshinota was mostly covering and interacting with living corals. Bryan (1973) suggested that the killing of live coral tissue may provide the sponge with additional nutrients, which would explain why the sponge prefers expanding over on live corals. However, Plucer-Rosario (1987) demonstrated that cleaned coral substrate resulted in even higher growth rates (1–2 mm/day) and concluded that *T. hoshinota* may be killing corals in competition for space on a reef.

In the present study all live coral substrates were identified to species level. Sponge patches were mostly found on corals of the family Acroporidae, and also to a lesser extent on corals of the family Pocilloporidae. This study is the first to record such a preference in regular conditions outside the context of an outbreak, and although *T. hoshinota* is also known to grow over any benthic structure during massive outbreaks (Rützler & Muzik, 1993; Reimer *et al.*, 2011a), this information can be crucial to understanding colonization preference

before an outbreak of *T. hoshinota*. In the Thousand Islands, Jakarta, where no outbreak event had occurred yet, the sampled *T. hoshinota* patches were also overgrowing corals of the family Acroporidae, specifically the foliose coral *Montipora* sp. (de Voogd *et al.*, 2013).

When expanding over live coral tissue, *T. hoshinota* shows directional growth and has an active growth front, which features structures like thin tissue threads, hairy tips, compact edges and disintegrated tissues. The hairy tips tissue threads contain cyanobacteria, sponge tissue and spicules, and play an active role in killing adjacent corals (Soong *et al.*, 2009; Wang *et al.*, 2012). In a patch more cryptically hidden in the reef, tissue threads were observed to grow towards the light. This growth strategy further highlights the potential of *T. hoshinota* to rapidly invade reefs.

Finally a study was performed on the genetic diversity in the sponge *T. hoshinota*. The D₃–D₅ region of the nuclear ribosomal 28S gene has been used to distinguish genera and species of a wide range of demosponge taxa, but it has also been found to be too conserved to discriminate between closely related species (Reveillaud *et al.*, 2010; Becking *et al.*, 2013). Indeed, no genetic variation in sequences was observed in the D₃–D₅ region of the 28S gene. However, two distinct haplotypes of *T. hoshinota* could be distinguished by sequencing the Folmer partition and the I₃-M₁₁ partition of the mitochondrial CO₁ in the present samples. These two haplotypes both occur in the Spermonde in the same frequency; haplotype C₂I₁ represents specimens from the mid-shelf




Fig. 4. Variation in the cover of the individual growth forms of live corals, as a function of distance from Makassar.

Table 2. Polymorphic positions of *Terpios hoshinota* for two regions of the mitochondrial marker COI, the standard DNA-barcoding partition (Folmer partition) and the I₃-M₁₁ partition. Three haplotypes (C₁–3) are found for the Folmer partition with a total of five variable sites. Two haplotypes (I₁–I₂) are found for the I₃M₁₁ partition, with one variable site. When these two regions of the mitochondrial marker CO₁ are combined, three haplotypes are found (C₁, C₂I₁, C₃I₂).

	Polymorphic positions of Folmer partition					Polymorphic positions of I ₃ -M ₁₁ partition		CO ₁ partitions combined	Sampling location
Folmer (N = 11)	81	189	261	561	594	I ₃ -M ₁₁ (N = 8)	61	(Folmer + I ₃ -M ₁₁ (N = 8))	
C ₁	C	C	C	C	T		n/a	C ₁	South China Sea
C ₂	C	T	C	C	T	I ₁	T	C ₂ I ₁	Thousand islands (N = 1) Kapoposang (N = 1) Pulau Badi (N = 3)
C ₃	T	T	T	T	C	I ₂	C	C ₃ I ₂	Lankai 2012 (N = 1) Samalona (N = 1) Lankai (N = 3)

Table 3. Genetic composition and mitochondrial diversity (based on the Folmer partition and I₃-M₁₁ partition of the mitochondrial marker CO₁) of *Terpios hoshinota*. N = number of samples, nH = number of haplotypes, H = haplotype (gene) diversity, π = nucleotide diversity (SD in parentheses). C₁–C₃ and I₁–I₂ refer to the relative frequencies of haplotypes.

	n	nH	H	π	Frequency of haplotype		
					C ₁	C ₂ /I ₁	C ₃ /I ₂
Folmer partition	11	3	0.636 (\pm 0.089)	0.00394 (\pm 0.00052)	0.091	0.455	0.455
I ₃ -M ₁₁	8	2	0.571 (\pm 0.094)	0.00121 (\pm 0.00020)		0.500	0.500

island of Pulau Badi and the outer-shelf island of Kapoposang, and also includes the specimen from the Thousand islands, Jakarta, and haplotype C3I2 represents specimens from the inner-shelf island of Samalona and the outer-shelf island of Lankai. The CO1 sequence of *T. hoshinota* obtained from GenBank, originates from Yongxing Island, South China Sea, where the sponge has had outbreak events in 2008 and 2010 (Shi *et al.*, 2012). This sequence varied slightly from those found in the Spermonde Archipelago, and was separated in another haplotype, C1.

The sponge *T. hoshinota* is recognized as an aggressive coral-killing sponge and has drawn the attention for extensive studies. There are a multitude of encrusting sponge species that are able to kill corals, though usually by simply overgrowing and smothering them to death (Aerts & van Soest, 1997). Some of the most aggressive coral-killing sponges belonging to the family *Clionidae* are encrusting and excavating sponges that are overgrowing and killing already stressed corals in the Caribbean at a disturbingly fast rate (Rützler, 2002; Chaves-Fonnegra & Zea, 2011). Another coral-killing cyanobacteriosponge, *Chalinula milnei*, exhibits similar aggressive growth strategies as *T. hoshinota* (Hoeksema *et al.*, 2014a) and was more abundantly observed than *T. hoshinota* throughout the entire Spermonde Archipelago during surveys in 2012 and 2013 (F. Dekker, unpublished data). However, it has received much less attention than *T. hoshinota*, while it may also impose a threat to coral reefs.

Recently, *T. hoshinota* was not observed at Yonama, Southern Japan where a massive outbreak of the 'black disease' occurred a quarter of a century ago (Reimer *et al.*, 2011b). In addition, this outbreak did not cause irreversible damage to the reefs. However, in the present study we observed dead coral next to *T. hoshinota* covered by filamentous algae. It is known that a phase-shift from a coral to more algae-dominated reef may in fact be irreversible (Bellwood *et al.*, 2004), and this way an outbreak of *T. hoshinota* can indirectly cause irreversible degradation of coral reefs.

Currently, there has been no major outbreak event in the Spermonde Archipelago yet, making this site and the *T. hoshinota* patches recorded during this study of particular interest for further research. The present results suggest that the sponge tolerates large variations in benthic community structure and other environmental and spatial conditions, indicating there are still other unknown mechanisms or stressors, either biotic or abiotic, that may play a role in the onset of *T. hoshinota* outbreaks. Liao *et al.* (2007) recorded the sponge in Taiwan, suggesting that *T. hoshinota* is widening its distribution in the north-western Pacific. Indeed, *T. hoshinota* has now been recorded as far west as the Maldives in the Indian Ocean (Montano *et al.*, 2015) and as far south-east as the Great Barrier Reef (Fujii *et al.*, 2011). It is unclear whether additional records are a consequence of range expansion, more outbreaks, or easier recognition as the species is more widely known than before.

ACKNOWLEDGEMENTS

Research permits were issued by the Indonesian State Ministry for Research and Technology (RISTEK) and the Indonesian Institute of Sciences (PPO-LIPI). Hasanuddin University at Makassar provided local support. We thank F. Dekker and S. Sradaputta for their help during fieldwork. We thank

K. Beentjes, M. Eurlings, B. Reijnen and C. de Leeuw for advice and help with lab techniques and molecular analyses.

FINANCIAL SUPPORT

This research was funded by the Treub-Maatschappij, Stichting Jan Joost ter Pelkwijk Fonds and Mej. A.M. Buitendijk Fonds.

REFERENCES

- Aerts L.A.M. and van Soest R.W.M. (1997) Quantification of sponge/coral interactions in a physically stressed reef community, NE Colombia. *Marine Ecology Progress Series* 148, 125–134.
- Becking L.E., Cleary D.F.R., de Voogd N.J., Renema W., de Beer M., van Soest R.W.M. and Hoeksema B.W. (2006) Betadiversity of tropical marine assemblages in the Spermonde Archipelago, Indonesia. *Marine Ecology* 27, 76–88.
- Becking L.E., Erpenbeck D., Peijnenburg K.T. and de Voogd N.J. (2013) Phylogeography of the sponge *Suberites diversicolor* in Indonesia: insights into the evolution of marine lake populations. *PLoS ONE* 8, e75996.
- Bellwood D.R., Hughes T.P., Folke C. and Nyström M. (2004) Confronting the coral reef crisis. *Nature* 429, 827–833.
- Bryan P.G. (1973) Growth rate, toxicity, and distribution of the encrusting sponge *Terpios* sp. (Hadromerida: Suberitidae) in Guam, Mariana Islands. *Micronesica* 9, 237–242.
- Chadwick N.E. and Morrow K.M. (2011) Competition among sessile organisms on coral reefs. In Dubinsky Z. and Stambler N. (eds) *Coral reefs: an ecosystem in transition*. Dordrecht: Springer, pp. 347–371. http://dx.doi.org/10.1007/978-94-007-0114-4_20.
- Chaves-Fonnegra A. and Zea S. (2011) Coral colonization by the encrusting excavating Caribbean sponge *Cliona delitrix*. *Marine Ecology* 32, 162–173.
- Cleary D.F.R., Becking L.E., de Voogd N.J., Renema W., de Beer M., van Soest R.W.M. and Hoeksema B.W. (2005) Variation in the diversity and composition of benthic taxa as a function of distance offshore, depth and exposure in the Spermonde Archipelago, Indonesia. *Estuarine, Coastal and Shelf Science* 65, 557–570.
- Cleary D.F.R., De Vantier L., Vail L., Manto P., de Voogd N.J., Ravello-Dolmen P.G. and Hoeksema B.W. (2008) Relating variation in species composition to environmental variables: a multi-taxon study in an Indonesian coral reef complex. *Aquatic Sciences* 70, 419–431.
- Cleary D.F.R. and de Voogd N.J. (2007) Environmental associations of sponges in the Spermonde Archipelago, Indonesia. *Journal of the Marine Biological Association of the United Kingdom* 87, 1669–1676.
- Cornils A., Schulz J., Schmitt P., Lanuru M., Richter C. and Schnack-Schiel S. (2011) Mesozooplankton distribution in the Spermonde Archipelago (Indonesia, Sulawesi) with special reference to the Calanoida (Copepoda). *Deep-Sea Research II* 57, 2076–2088.
- de Voogd N.J., Becking L.E., Noor A., Hoeksema B.W. and van Soest R.W.M. (2004) Sponge interactions with spatial competitors in SW Sulawesi. *Bollettino dei Musei e degli Istituti Biologici dell'Università di Genova* 68, 253–261.
- de Voogd N.J., Cleary D.F.R., Hoeksema B.W., Noor A. and van Soest R.W.M. (2006) Sponge betadiversity in the Spermonde Archipelago, Indonesia. *Marine Ecology Progress Series* 309, 131–142.

- de Voogd N.J., Cleary D.F.R. and Dekker F. (2013) The coral-killing sponge *Terpios hoshinota* invades Indonesia. *Coral Reefs* 32, 755.
- de Voogd N.J., Haftka J.J.H. and Hoeksema B.W. (2005) Evaluation of the ecological function of amphitoxin in the reef-dwelling sponge *Callyspongia (Euplaccella) biru* (Haplosclerida: Callyspongiidae) at southwest Sulawesi, Indonesia. *Contributions to Zoology* 74, 53–61.
- Edinger E., Jompa J. and Limmon G. (1998) Reef degradation and coral biodiversity in Indonesia: effects of land-based pollution, destructive fishing practices and changes over time. *Marine Pollution Bulletin* 36, 617–630.
- English S., Wilkinson C. and Baker V. (1997) *Survey manual for tropical marine resources*, 2nd edition. Townsville: Australian Institute of Marine Science.
- Erftemeijer P.L.A. (1994) Differences in nutrient concentrations and resources between seagrass communities on carbonate and terrigenous sediments in South Sulawesi, Indonesia. *Bulletin of Marine Science* 54, 403–419.
- Erpenbeck D., Knowlton A.L., Talbot S.L., Highsmith R.C. and Van Soest R.W.M. (2004) A molecular comparison of Alaskan and North East Atlantic *Halichondria panicea* (Pallas 1766) (Porifera: Demospongiae) populations. *Bollettino dei Musei e degli Istituti Biologici dell'Università di Genova* 68, 319–325.
- Fujii T., Keshavmurthy S., Zhou W., Hirose E., Chen C.A. and Reimer J.D. (2011) Coral-killing cyanobacteriosponge (*Terpios hoshinota*) on the Great Barrier Reef. *Coral Reefs* 30, 483.
- Folmer O., Black M., Hoeh W., Lutz R. and Vrijenhoek R. (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3, 294–299.
- Hoeksema B.W. (2004) Biodiversity and the natural resource management of coral reefs in Southeast Asia. In Visser L.E. (ed.) *Challenging coasts. Transdisciplinary excursions into integrated coastal zone development*. Amsterdam: Amsterdam University Press, pp. 49–71.
- Hoeksema B.W. (2007) Delineation of the Indo-Malayan centre of maximum marine biodiversity: the Coral Triangle. In Renema W. (ed.) *Biogeography, time and place: distributions, barriers and islands*. Dordrecht: Springer, pp. 117–178.
- Hoeksema B.W. (2012a) Distribution patterns of mushroom corals (Scleractinia: Fungiidae) across the Spermonde Shelf, Indonesia. *Raffles Bulletin of Zoology* 60, 183–212.
- Hoeksema B.W. (2012b) Evolutionary trends in onshore-offshore distribution patterns of mushroom coral species (Scleractinia: Fungiidae). *Contributions to Zoology* 81, 199–221.
- Hoeksema B.W. and Crowther A.L. (2011) Masquerade, mimicry and crypsis of the polymorphic sea anemone *Phyllo-discus semoni* Kwietniewski, 1897, and its aggregations in South Sulawesi. *Contributions to Zoology* 80, 251–268.
- Hoeksema B.W., Dekker F. and de Voogd N.J. (2014a) Free-living mushroom corals strike back by overtopping a coral-killing sponge. *Marine Biodiversity* 44, 3–4.
- Hoeksema B.W. and de Voogd N.J. (2012) On the run: free-living mushroom corals avoiding interaction with sponges. *Coral Reefs* 31, 455–459.
- Hoeksema B.W. and Koh E.G.L. (2009) Depauperation of the mushroom coral fauna (Fungiidae) of Singapore (1860s–2006) in changing reef conditions. *Raffles Bulletin of Zoology Supplement* 22, 91–101.
- Hoeksema B.W., Waheed Z. and de Voogd N.J. (2014b) Partial mortality in corals overgrown by the sponge *Terpios hoshinota* at Tioman Island, Peninsular Malaysia (South China Sea). *Bulletin of Marine Science* 90, 989–990.
- Knittweis L. and Wolff M. (2010) Live coral trade impacts on the mushroom coral *Heliofungia actiniformis* (Scleractinia: Fungiidae) populations in the Indo-Malay Archipelago: implications for live coral trade management efforts. *Conservation Genetics* 10, 241–249.
- Liao M.H., Tang S.L., Hsu C.M., Wen K.C., Wu H., Chen W.M. and Chen C.A. (2007) The “Black Disease” of reef-building corals at Green Island, Taiwan – outbreak of a cyanobacteriosponge, *Terpios hoshinota* (Suberitidae; Hadromerida). *Zoological Studies* 46, 520.
- Librado P. and Rozas J. (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452.
- McCormack G.P. and Kelly M. (2002) New indications of the phylogenetic affinity of *Spongosorites suberitoides* Diaz et al. 1993 (Porifera Demospongiae) as revealed by 28 s ribosomal DNA. *Journal of Natural History* 36, 1009–1021.
- Meyer C.P., Geller J.B. and Paulay G. (2005) Fine scale endemism on coral reefs: archipelagic differentiation in turbinid gastropods. *Evolution* 59, 113–125.
- Misof B., Erpenbeck D. and Sauer K.P. (2000) Mitochondrial gene fragments suggest paraphyly of the genus *Panorpa* (Mecoptera, Panorpidae). *Molecular Phylogenetics and Evolution* 17, 76–84.
- Moll H. (1983) Zonation and diversity of scleractinia on reefs off S. W. Sulawesi, Indonesia. PhD dissertation, Leiden University, the Netherlands.
- Montano S., Chou W.H., Chen C.A., Galli P. and Reimer J.D. (2015) First record of the coral-killing sponge *Terpios hoshinota* in the Maldives and Indian Ocean. *Bulletin of Marine Science* 91, 97–98.
- Munro C. (2005) Diving systems. In Eleftheriou A. and McIntyre A. (eds) *Methods for the study of marine benthos*, 3rd edition. Oxford: Blackwell Science, pp. 112–159.
- Pandolfi J.M., Bradbury R.H., Sala E., Hughes T.P., Bjorndal K.A., Cooke R.G., McArdle D., McClenachan L., Newman M.J., Paredes G., Warner R.R. and Jackson J.B. (2003) Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301, 955–958.
- Pet-Soede C., Cesar H.S.J. and Pet J.S. (1999) An economic analysis of blast fishing on Indonesian coral reefs. *Environmental Conservation* 26, 83–93.
- Plucer-Rosario G. (1987) The effect of substratum on the growth of *Terpios*, an encrusting sponge which kills corals. *Coral Reefs* 5, 197–200.
- Reimer J.D., Mizuyama M., Nakano M., Fujii T. and Hirose E. (2011a) Current status of the distribution of the coral-encrusting cyanobacteriosponge *Terpios hoshinota* in southern Japan. *Galaxea, Journal of Coral Reef Studies* 13, 35–44.
- Reimer J.D., Nozawa Y. and Hirose E. (2011b) Domination and disappearance of the black sponge: a quarter century after the initial *Terpios* outbreak in southern Japan. *Zoological Studies* 50, 394.
- Renema W., Hoeksema B.W. and Van Hinte J.E. (2001) Larger benthic foraminifera and their distribution patterns on the Spermonde shelf, South Sulawesi. *Zoologische Verhandlungen* 344, 115–150.
- Renema W. and Troelstra S.R. (2001). Larger foraminifera distribution on a mesotrophic carbonate shelf in SW Sulawesi (Indonesia). *Palaeogeography, Palaeoclimatology, Palaeoecology* 175, 125–146.
- Reveillard J., Remerie T., van Soest R., Erpenbeck D., Cárdenas P., Derycke S. and Vanreusel A. (2010) Species boundaries and phylogenetic relationships between Atlanto-Mediterranean shallow-water and deep-sea coral associated *Hexadella* species (Porifera, Ianthellidae). *Molecular Phylogenetics and Evolution* 56, 104–114.
- Rützler K. (2002) Impact of crustose cionid sponges on Caribbean reef corals. *Acta Geologica Hispanica* 37, 61–72.

- Rützler K. and Muzik K.** (1993) *Terpios hoshinota*, a new cyanobacteriosponge threatening Pacific reefs. *Scientia Marina* 57, 395–403.
- Sawall Y., Teichberg M., Seemann J., Litaay M., Jompa J. and Richter C.** (2011) Nutritional status and metabolism of the coral *Stylophora subseriata* along a eutrophication gradient in Spermonde Archipelago (Indonesia). *Coral Reefs* 30, 841–853.
- Schils T.** (2012) Episodic eruptions of volcanic ash trigger a reversible cascade of nuisance species outbreaks in pristine coral habitats. *PLoS ONE* 7, e46639.
- Shi Q., Liu G.H., Yan H.Q. and Zhang H.L.** (2012) Black disease (*Terpios hoshinota*): a probable cause for the rapid coral mortality at the northern reef of Yongxing Island in the South China Sea. *Ambio* 41, 446–455.
- Soong K., Yang S. and Chen C.A.** (2009) A novel dispersal mechanism of a coral-threatening sponge, *Terpios hoshinota* (Suberitidae, Porifera). *Zoological Studies* 48, 596–596.
- Sterrenburg F.A.S., Erfteimeijer P.L.A. and Nienhuis P.H.** (1995) Diatoms as epiphytes on seagrasses in South Sulawesi (Indonesia) comparison with growth on inert substrata. *Botanica Marina* 38, 1–8.
- Tang S.L., Hong M.J., Liao M.H., Jane W.N., Chiang P.W., Chen C.B. and Chen C.A.** (2011) Bacteria associated with an encrusting sponge (*Terpios hoshinota*) and the corals partially covered by the sponge. *Environmental Microbiology* 13, 1179–1191.
- Verheij E. and Erfteimeijer P.L.A.** (1993) Distribution of seagrasses and associated macroalgae in South Sulawesi, Indonesia. *Blumea* 38, 45–64.
- Verheij E. and Prud'homme van Reine W.F.** (1993) Seaweeds of the Spermonde Archipelago, SW Sulawesi, Indonesia. *Blumea* 37, 385–510.
- Veron J.E.N.** (2000) *Corals of the World*, Volumes 1–3. Townsville: Australian Institute of Marine Science.
- Wang J.T., Chen Y.Y., Meng P.J., Sune Y.H., Hsu C.M., Wei K.Y. and Chen C.A.** (2012) Diverse interactions between corals and the coral-killing sponge, *Terpios hoshinota* (Suberitidae: Hadromerida). *Zoological Studies* 51, 150–159.
- and
- Wulff J.** (2012) Ecological interactions and the distribution, abundance, and diversity of sponges. *Advances in Marine Biology* 61, 146–166.

Correspondence should be addressed to:

N.J. de Voogd
 Department of Marine Zoology, Naturalis Biodiversity Center,
 P.O. Box 9517, 2300RA Leiden, the Netherlands
 email: Nicole.deVoogd@naturalis.nl