


cambridge.org/ags

S. S. Souza, J. A. Flôres, A. P. Coelho , J. M. Deienno, L. B. Lemos
and G. S. Rolim

Crops and Soils Research Paper

Cite this article: Souza SS, Flôres JA, Coelho AP, Deienno JM, Lemos LB, Rolim GS (2022). Intercropping and cover crop effects on maize nitrogen requirement for optimal growth. *The Journal of Agricultural Science* **160**, 325–337. <https://doi.org/10.1017/S0021859622000478>

Received: 18 November 2021
Revised: 9 July 2022
Accepted: 16 July 2022
First published online: 1 August 2022

Key words:

Crotalaria spectabilis; grain yield; *Urochloa ruziziensis*; *Zea mays* L

Author for correspondence:

A. P. Coelho,
E-mail: anderson_100ssp@hotmail.com

School of Agricultural and Veterinarian Sciences, São Paulo State University (Unesp), Via de Acesso Prof. Paulo Donatto Castellane, km 5, s/n. 14884-900, Jaboticabal, São Paulo, Brazil

Abstract

Intercropping maize (*Zea mays* L.) with cover crops (*Crotalaria spectabilis* and *Urochloa ruziziensis*) is a sustainable cultivation strategy that can generate ecological benefits combined with grain yield (GY). However, cover crops may require nitrogen (N) fertilization management to be adjusted to obtain a high GY in intercropping systems. This study was carried out over 2 years in southeastern Brazil using randomized complete block design in a split-plot scheme. The plots were composed of three cropping systems: sole maize, maize + *U. ruziziensis* and maize + *C. spectabilis*. The subplots consisted of four N levels: 0, 70, 140 and 210 kg/ha. The GY of the maize + *U. ruziziensis* may be reduced by 13% compared to the sole maize. The GY of the sole maize increased up to 11.3 Mg/ha for 95 kg/ha of N in the first year and 6.7 Mg/ha for 169 kg/ha of N in the second year. The GY of the maize + *U. ruziziensis* intercropping system showed linear increments of 120 kg/ha for every 10 kg/ha of N applied. In the maize + *C. spectabilis* intercropping, GY increased up to 9.8 Mg/ha for 201 kg/ha of N in the first year, without any variation during the second year. The land equivalent ratio (LER) of intercropping was increased by more than 35% compared to that of sole maize, and N fertilization increased the LER of the sole maize and maize + *U. ruziziensis* intercropping. Maize intercropping had higher LER values with greater demand for nitrogen to obtain similar yields compared to sole maize cropping.

Introduction

Food production will remain viable over time if sustainable production methods are practised. Hence, many countries have adopted systems for improving agricultural yields that are associated with ecosystem conservation, soil improvements and adaptation to climate change (United Nations, 2021). Satisfactory results have been achieved with the use of cover crops as they contribute to improved soil quality, due to high organic matter (Schipanski *et al.*, 2014) and nutrient cycling (Pacheco *et al.*, 2013). Consequently, easing the over-reliance on fertilizers and pesticides, which in turn are environmentally unsustainable. Moreover, cover crops could control pests (Schipanski *et al.*, 2014) and weeds (Martins *et al.*, 2016), by covering the space or area needed by weeds.

Intercropping is the best management to introduce cover crops into cropping systems. This system results in minimal changes to the routine of farmers and combines economic returns with ecosystem services for the environment and soil conservation. Overall, intercropping systems combine growing several species with each other during a season as an alternative to monoculture. Intercropping has been attributed to the fourth green revolution since it makes agriculture ecologically promising by producing more biomass in the same area than sole cropping (Martin-Guay *et al.*, 2018).

Among cereal crops, maize is prioritized for intercropping with cover crops. This is because, in addition to its economic and social importance (USDA, 2021), maize has vigorous vegetative growth stages. These traits are advantageous in intercropped systems, mitigating the competition between cover crops and grain yield (GY) of the cereal crop (de Oliveira *et al.*, 2010; Kappes and Zancanaro, 2015). Intercropping systems of maize with grasses (Kluthcouski *et al.*, 2000) and legumes (de Oliveira *et al.*, 2010) are successful examples of maize intercropping systems that have generated economic, social and environmental benefits.

Using forage grasses, such as *Urochloa ruziziensis*, in the cropping system contributes to the chemical and physical aspects of the soil (Silveira *et al.*, 2011; Mingotte *et al.*, 2020; Tan, 2020). Additionally, they stand out for their high straw yield, further contributing to sustainability (Mingotte *et al.*, 2014). Legume species, such as crotalarias, are called green manure crops which are associated with microorganisms that are capable of fixing nitrogen biologically (BNF), receive nutrients and release them into the soil through root exudates (Murrell *et al.*, 2017) or decompose the crop remains (Marcelo *et al.*, 2012). *Crotalaria spectabilis*

© The Author(s), 2022. Published by
Cambridge University Press



Table 1. Soil chemical attributes of the 0.00–0.20 m layer analysed prior to the start of the experiment

Year	pH ^a	OM (g/dm ³)	P ^b (mg/ dm ³)	K ⁺ (mmol _c / dm ³)	Ca ²⁺ (mmol _c / dm ³)	Mg ²⁺ (mmol _c / dm ³)	H + Al (mmol _c / dm ³)	SB (mmol _c / dm ³)	CEC (mmol _c / dm ³)	V (%)
2016/17	5.1	23	47	3.5	34	14	37	51	88	58
2017/18	5.4	27	72	5.8	42	21	34	69	103	67

OM, organic matter; SB, sum of bases; CEC, cation exchange capacity; V, base saturation.
^apH CaCl₂; ^bP resin.

can supply up to 160 kg/ha of nitrogen (N) to the soil, reducing N fertilizer usage in the production system (Calegari and Carlos, 2014).

High N levels are usually recommended for crop fertilization management because N is the most extracted nutrient by plants and has complex dynamics in the soil–plant–atmosphere system. However, several recommendations do not consider the cropping systems, such as intercropped systems, used by crop producers. Hence, more specific N fertilization recommendations are needed in this context to optimize its use in agriculture.

Maize crops require 20–25 kg/ha of N to produce 1 Mg/ha of grains (Fan et al., 2020). In the context of sole maize cropping, the N requirement for obtaining high maize yields reaches 30 kg/ha at sowing and 140 kg/ha with top-dressing fertilization (Raij and Cantarella, 1997). However, the intercropping of cover crops with maize may alter the N requirement for obtaining high maize yields because these cover crops may absorb the applied N, resulting in a lower N availability for maize. The species used for intercropping (grass or legumes) may also interfere with the dynamics of N in the soil, directly affecting the management of N fertilization. Therefore, further studies are needed to determine the amount of N required by maize in intercropping systems.

The present study thus aimed to verify whether the management of top-dressing N fertilization is changed by intercropping maize with grass (*U. ruziziensis*) and legume (*C. spectabilis*) in comparison with that by sole cropping of maize.

Materials and methods

Experimental area location and characterization

Experiments were conducted during the first cropping season (summer) of 2016/17 and 2017/18, respectively. Both experiments were conducted in the same area, located at the São Paulo State University (Unesp), School of Agricultural and Veterinarian Sciences, in Jaboticabal, São Paulo, Brazil (21°14'59"S, 48°17'13"W and 565 m of altitude).

The climate of the site is 'Aw', characterized as humid tropical with rainy season in summer and dry season in winter, according to Köppen's classification (Rolim et al., 2007). The soil is an Oxisol (Soil Survey Staff, 2014), with 540 g/kg of clay, was previously cultivated with maize in the conventional tillage system, and was kept in fallow until installation of the experiments.

Experimental design and treatments

The experimental design used was randomized complete block design in a split-plot structure, with four replicates. The plots consisted of three cropping systems (sole maize, maize intercropped with *U. ruziziensis* and maize intercropped with *C. spectabilis*)

and the subplots were four N levels (0, 70, 140 and 210 kg/ha), applied broadcast as top-dressing at V₆ stage of maize. The fertilizer used as the source of the nutrient was Kimcoat® polymer-coated urea, with 44% of N. Each subplot consisted of five maize rows with length of 10 m and spaced at 0.45 m, in a total area of 22.5 m². The three central rows were considered usable area, disregarding 0.5 m at the ends.

Experiment installation and conduction

Pretreatment

Before the installation of the experiment in 2 years, the soil attributes were determined in the 0.00–0.20 m layer (Table 1).

Conventional tillage was performed in both years, with plowing as well as operations with disc harrow and levelling harrow. In addition, 2 t/ha of dolomite limestone with 90% relative total neutralizing power were applied 90 days before experiment in 2017/18. This management was necessary because 6 months after the harvest during the first year of the experiment, we verified that the soil base saturation was below 50%. Thus, the application of lime was necessary to increase the soil base saturation level and promote ideal conditions for plant growth in the second year of the experiment. After liming, the soil base saturation increased to 67% (Table 1).

Sowing

Sowing in the cropping systems was performed at the same location in the two experiments. Cover crops were distributed broadcast, using 20 kg/ha of *C. spectabilis* seeds and 15 kg/ha of *U. ruziziensis* seeds. Then, a seeder-fertilizer was used to sow the maize hybrids at the site, which led to the deposition of soil on the seeds of the cover crops.

The maize sowing in the first year was carried out on 11/22/2016, using the hybrid DOW 2B710 PW, in the interrow spacing of 0.45 m and with an adjusted population of 60 000 plants per ha. The crop management used for this maize hybrid was carried out as recommended by the manufacturer. The sowing furrow fertilization was calculated as a function of the soil attributes determined before the experiment installation and according to the recommendations of Raij and Cantarella (1997). At seed furrow, 20 kg/ha of N, 70 kg/ha of P₂O₅ and 40 kg/ha of K₂O were applied using the formulated fertilizer 04-14-08 (urea + single superphosphate + potassium chloride).

The maize sowing in the second year was carried out on 11/10/2017, using the hybrid DOW 2B810 PW, in the interrow spacing of 0.45 m and with an adjusted population of 88 000 plants per ha. The crop management used for this maize hybrid was carried out as recommended by the manufacturer. The seed furrow fertilization was calculated as a function of the soil attributes determined before the experiment started and according to the




Fig. 1. Maximum and minimum air temperature ($^{\circ}\text{C}$) and precipitation (mm) during the experimental period. Dashed arrows indicate the times of sowing of maize and cover crops (1), top-dressing nitrogen fertilization (2) and harvest (3) of sole maize and intercropped systems, in the 2016/17 (a) and 2017/18 (b) cropping seasons.

recommendations of Raji and Cantarella (1997). At seed furrow, 25 kg/ha of N, 89 kg/ha of P_2O_5 and 51 kg/ha of K_2O were applied using the formulated fertilizer 08-28-16 (urea + triple superphosphate + potassium chloride).

Fertilization input

In two years, top-dressing fertilization was performed at the V_6 stage of the crop, on 12/24/2016 and 11/28/2017, with N levels defined by the treatments. In addition to N, 45 kg/ha of K_2O was also applied on the same day in both years, using potassium chloride as a source.

Climate characterization in the period

The meteorological data recorded during the experiments are presented in Fig. 1. In 2016/17, the average maximum and minimum temperatures were 31.1 and 19.4 $^{\circ}\text{C}$, respectively, and the accumulated precipitation was 761.6 mm. In 2017/18, the average maximum and minimum temperatures were 30.5 and 19.5 $^{\circ}\text{C}$, respectively, and the accumulated precipitation was 760.4 mm.

The plants were grown under rainfed conditions for two years; however, on the first and second day of maize sowing for 2016/17, conventional sprinkler irrigation with 10 mm water depths was

performed, for the establishment of the plants. When necessary, the crop practices related to the control of pests and diseases were carried out based on monitoring. Chemical management of weeds was performed in the system with sole maize, at 25 days after emergence, using the herbicide potassium glyphosate at a rate of 2500 g/ha of a.i. in both years. Maize plants were mechanically harvested on 04/16/2017 and 04/06/2018, respectively.

Evaluations

The evaluated variables analysed in the two years were the same. Leaf nitrogen content (g/kg) was evaluated during the flowering (R_1 stage) of maize, with the collection of the middle third without the midrib of ten leaves, located below and on the opposite side of the main ear, in ten plants per subplot (Cantarella and Raji, 1997). The samples were washed with running water and detergent at 1 ml/l, dried in an oven with forced air circulation at 65–70 $^{\circ}\text{C}$, and then processed in a Wiley-type mill to determine the N content, according to semi-micro Kjeldahl method (Carmo *et al.*, 2000).

Stem diameter (mm) was obtained using a digital caliper, for five plants per subplot, in the second internode above ground, at the maize physiological maturity. Plant height (cm) was

Table 2. Leaf nitrogen content (LNC), stem diameter (SD), plant height (PH) and main ear height (EH) of maize in different cropping systems and subjected to top-dressing nitrogen levels in 2016/17 (year 1) and 2017/18 (year 2) cropping seasons

Treatments	LNC (g/kg)		SD (mm)		PH (cm)		EH (cm)	
	Years							
	16/17	17/18	16/17	17/18	16/17	17/18	16/17	17/18
Cropping systems (CS)	Mean							
M + U ^a	27.20	30.97	20.55	20.04	229.8	234.4	126.2 b ^b	127.7 ab
M + C ^c	27.59	32.27	20.81	20.58	231.4	238.6	130.2 a	130.8 a
M ^d	28.04	36.55	21.58	20.62	215.9	233.6	124.1 b	126.6 b
LSD (5%)	1.92	6.41	1.28	1.15	16.77	6.34	3.20	4.14
F test	F value							
CS	0.92 ^{ns}	3.89 ^{ns}	3.36 ^{ns}	1.49 ^{ns}	4.81 ^{ns}	3.28 ^{ns}	17.26**	5.16*
N levels (D)	8.58**	11.36**	12.05**	3.44*	0.01 ^{ns}	0.18 ^{ns}	0.29 ^{ns}	0.23 ^{ns}
CS × D	0.76 ^{ns}	2.87*	0.58 ^{ns}	0.08 ^{ns}	0.05 ^{ns}	0.04 ^{ns}	0.67 ^{ns}	0.39 ^{ns}
CV (%) – CS	6.43	17.78	5.62	5.19	6.85	2.48	2.32	2.97
CV (%) – D	7.64	11.17	4.19	5.00	7.69	3.22	4.45	4.54
Overall Mean	27.61	33.26	20.98	20.41	225.71	235.54	126.84	128.33

^aMaize intercropped with *U. ruziziensis*.

^bDifferent letters correspond to the significant difference at 5% probability level ($P \leq 0.05$) by Tukey test.

^cMaize intercropped with *C. spectabilis*.

^dSole maize.

**Differs by F test at 1% probability level; *differs by F test at 5% probability level; ^{ns}not significant.

measured at the time of physiological maturity of the maize crop, for five plants per subplot, as the distance from the soil surface to the flag leaf, with a graduated ruler. Main ear height (cm) was determined as the distance from the soil surface to the main ear, with a graduated ruler, for five plants per subplot at the physiological maturity of the maize crop.

At the maize harvest time, ten ears were randomly collected to determine the production components of maize: number of rows per ear (n°), number of grains per row (n°), number of grains per ear (n°), thousand-grain weight (g) and grain mass per ear. For the determination of thousand-grain weight and grain mass per ear, the grain moisture was corrected to 0.13 g/g.

Maize GY was obtained by manually harvesting the ears of the usable area in each subplot, which were mechanically threshed. The obtained grains were weighed and the value was corrected to a moisture content of 0.13 g/g.

To verify the effect of cropping systems and top-dressing N fertilization on grain production, the land equivalent ratio (LER) index was calculated according to Mead and Willey (1980) (Eqn (1)). This index allows assessing the area required by the sole cropping system to equal the total grain production of an intercropped system. Here, we considered the sole maize cropping system at a level of N equal to 0 as the control treatment. The average data of the two cropping seasons were used to calculate the LER:

$$LER = \frac{Y_{MM0}}{Y_{CS}} + \frac{Y_{CCM}}{Y_{CS}} \quad (1)$$

where LER = land equivalent ratio; Y_{MM0} = sole maize GY at level 0 (kg/ha); Y_{CS} = maize GY of the cropping system of interest (kg/ha) and Y_{CCM} = sole cover crop dry mass yield (kg/ha).

The average production of the total dry mass by the sole cropping of *C. spectabilis* and *U. ruziziensis* for the two cropping seasons was 6.50 and 6.00 t/ha, respectively. These values were used to calculate the LER for each treatment.

Statistical analyses

For the study of the cropping systems, the data were subjected to analysis of variance by the F test, and the means were compared by the Tukey test at the 5% probability level. We use mixed models for statistical analysis, adopting treatments as fixed effects and blocks as random effects. We used Shapiro–Wilk test to assess the data normality (Royston, 1995) and Levene's test to assess the homoscedasticity of variances (Gastwirth et al., 2009). Regarding the effect of N levels and their interaction with the cropping systems, polynomial regression analyses were performed. The statistical program Sisvar was used (Ferreira, 2011).

In addition, due to the structure of dependence between the variables, multivariate statistics by principal components (PCs) was used to analyse the agronomic variables of maize, as a function of the factors cropping systems and N levels. All variables were standardized before analysis. The number of PCs was determined based on the Kaiser criterion, using those with eigenvalues greater than 1 (Kaiser, 1958). For multivariate analysis, the statistical program used was Statistica v.7.

Results

Growth variables

Among the aspects that characterize nutritional diagnosis and growth of maize (Table 2), only main ear height was influenced by the cropping systems (first year: $P < 0.001$; second year:




Fig. 2. Leaf N content (a) and stem diameter (b) for maize as a function of nitrogen levels applied as top-dressing in the 2016/17 cropping season. ▲ Sole maize; ■ maize intercropped with *C. spectabilis*; ● maize intercropped with *U. ruziziensis*.

$P < 0.05$). In both years, the main ears of maize intercropped with *C. spectabilis* were located in the upper region of the stem. In the first year, leaf N content was affected ($P < 0.01$) differently between the cropping systems as a function of nitrogen fertilization (Fig. 2(a)). This variable increased quadratically up to the N level of 144 kg/ha in sole maize and linearly in the other cropping systems. Thus, the maximum value of leaf N content was not reached with the studied levels for the intercropping with *C. spectabilis* and *U. ruziziensis*, respectively increasing by 0.18 and 0.22 g/kg with the addition of 10 kg/ha of N. Stem diameter increased with the increment in N levels, and every 10 kg/ha of the nutrient added led to increment of approximately 0.09 mm in stem diameter (Fig. 2(b)).

In the second year, for leaf N content, based on the interaction between the factors cropping systems and nitrogen fertilization ($P < 0.001$), there was a quadratic increase up to the N levels of 144, 125 and 198 kg/ha in sole maize, intercropped with *C. spectabilis* and intercropped with *U. ruziziensis*, respectively (Fig. 3(a)). In addition, when there was no supply of N as top-dressing (level 0), sole maize had a higher value of leaf N content (35 g/kg) compared to maize intercropped with *C. spectabilis* (27 g/kg) and *U. ruziziensis* (23 g/kg).




Fig. 3. Leaf N content (a), stem diameter (b) and main ear height (c) for maize crop as a function of nitrogen levels applied as top-dressing in the 2017/18 cropping season. ▲ Sole maize; ■ maize intercropped with *C. spectabilis*; ● maize intercropped with *U. ruziziensis*.

Stem diameter and main ear height increased linearly as a function of nitrogen fertilization, at rates of 0.05 mm and 0.08 cm, respectively, for every 10 kg/ha of the nutrient added (Figs 3(b) and (c)).

Yield components and GY

Regarding the variables related to ear and GY, the cropping systems affected ($P < 0.01$) only in GY in the first year (Table 3).

Table 3. Number of rows per ear (NRE), number of grains per row (NGR), number of grains per ear (NGE), thousand-grain weight (1000GW), grain mass per ear (GME) and GY of maize in different cropping systems and subjected to top-dressing nitrogen levels in 2016/17 (year 1) and 2017/18 (year 2) cropping seasons

Treatments	NRE (n°)		NGR (n°)		NGE (n°)		1000GW (g)		GME (g)		GY (t/ha)	
	Years											
	16/17	17/18	16/17	17/18	16/17	17/18	16/17	17/18	16/17	17/18	16/17	17/18
Cropping systems (CS)	Mean											
M + U ^a	20	19	20	35	558	665	269.6	228.5	151.1	121.7	8.11 b ^b	6.75
M + C ^c	19	19	19	35	560	671	265.2	230.3	146.3	128.6	8.80 ab	6.83
M ^d	19	19	19	36	573	677	270.8	235.8	156.6	128.1	9.32 a	6.37
LSD (5%)	0.88	0.89	2.49	1.45	86.9	47.76	18.14	16.19	23.84	1.95	0.71	1.04
F test	F value											
CS	2.64 ^{ns}	0.09 ^{ns}	0.35 ^{ns}	0.61 ^{ns}	0.16 ^{ns}	0.31 ^{ns}	0.50 ^{ns}	1.04 ^{ns}	0.87 ^{ns}	3.80 ^{ns}	13.75 ^{**}	1.09 ^{ns}
N levels (D)	0.50 ^{ns}	1.03 ^{ns}	3.05 [*]	2.43 ^{ns}	0.39 ^{ns}	3.09 [*]	1.84 ^{ns}	4.26 ^{ns}	0.81 ^{**}	4.32 [*]	35.06 ^{**}	6.24 ^{**}
CS × D	0.90 ^{ns}	0.80 ^{ns}	0.55 ^{ns}	0.68 ^{ns}	0.24 ^{ns}	0.38 ^{ns}	0.47 ^{ns}	0.69 ^{ns}	0.6 ^{ns}	0.28 ^{ns}	14.11 ^{**}	1.42 ^{**}
CV (%) – CS	4.18	4.35	7.86	3.76	14.21	6.57	6.23	6.45	14.53	6.19	7.5	14.43
CV (%) – D	4.68	3.59	6.89	4.65	9.33	5.39	4.97	6.09	9.38	9.51	7.87	10.15
Overall Mean	19	19	29	35	564	671	268.5	231.5	151.33	126.1	8.75	6.66

^aMaize intercropped with *U. ruziziensis*.

^bDifferent letters correspond to significant difference at 5% probability level ($P \leq 0.05$) by Tukey test.

^cMaize intercropped with *C. spectabilis*.

^dSole maize.

^{**}Differs by F test at 1% probability level; ^{*}differs by F test at 5% probability level; ^{ns}not significant.




Fig. 4. Number of grains per row (a), number of grains per ear (b), thousand-grain weight (c) and GY (d) for maize crop as a function of nitrogen levels applied as top-dressing in the 2016/17 cropping season. ▲ Sole maize; ■ maize intercropped with *C. spectabilis*; ● maize intercropped with *U. ruziziensis*.

Thus, sole maize produced approximately 1200 kg/ha (15%) more than maize intercropped with *U. ruziziensis* (Table 3).

Regarding the increment of N in the top-dressing fertilization for the first year, there was a quadratic increase in the number of grains per row (Fig. 4(a)) and thousand-grain weight (Fig. 4(c)) up to the N levels of 204 and 169 kg/ha, respectively. As for the number of grains per ear, the increase was linear, and every 10 kg/ha of N led to an increment of 1.7 grains on the ear (Fig. 4(b)).

For GY, different responses were observed between the cropping systems and N levels (first year: $P < 0.001$; second year: $P < 0.01$) (Fig. 4(d)). There was a quadratic increase up to 95 and 201 kg/ha of N for sole maize and intercropped with *C. spectabilis*, respectively. For maize intercropped with *U. ruziziensis*, the increment was linear, equal to 150 kg/ha for every 10 kg/ha of N added.

In the second year, the treatments did not affect the grain and ear variables (Table 3). As for nitrogen fertilization, every 10 kg/ha of N added led to an increase of 0.07 grains per row (Fig. 5(a)), 0.8 g in thousand-grain weight (Fig. 5(c)) and 0.7 g in the grain mass per ear (Fig. 5(d)). The number of grains per ear (Fig. 5(b)), however, increased quadratically up to the N level of 164 kg/ha. Based on the interaction between the factors cropping systems and N levels, GY increased up to the N level of 169 kg/ha for sole maize (Fig. 5(e)). On the other hand, in the intercropping with *C. spectabilis* there was no increase in GY as a function of N levels. For maize intercropped with *U. ruziziensis*, every 10 kg/ha

of N applied led to an increment of approximately 90 kg/ha of grains, not reaching the maximum of the variable with the tested levels. In addition, without nitrogen fertilization, the yields were equal to 5.58, 6.83 and 5.84 t/ha, for sole maize, maize intercropped with *C. spectabilis* and maize intercropped with *U. ruziziensis*, respectively.

PCs analysis

Multivariate analysis of PCs for the agronomic variables of maize in the first year showed that the two first PCs were responsible for 68.84% of the total variability (Fig. 6(a)). The variables that most discriminated PC1 and PC2 were related to the production performance and vegetative growth of maize, respectively. Grain mass per ear (−0.94), stem diameter (−0.91), 1000-grain weight (−0.80), number of grains per ear (−0.85), GY (−0.65), number of grains per row (−0.86), leaf N content (−0.79) were correlated with PC1, called agronomic performance. On the other hand, PC2, called plant growth, was characterized by ear height (−0.92) and plant height (−0.92). The number of rows per ear was not part of any component this year.

Analysis of the biplot showed that the cropping systems with sole maize at N levels 70, 140 and 210 kg/ha, maize intercropped with *C. spectabilis* at the N level of 210 kg/ha and maize intercropped with *U. ruziziensis* at N levels of 140 and 210 kg/ha were the ones with the highest performance. This fact confirms




Fig. 5. Number of grains per row (a), number of grains per ear (b), thousand-grain weight (c), grain mass per ear (d) and GY (e) for maize crop as a function of nitrogen levels applied as top-dressing in the 2017/18 cropping season. ▲ Sole maize; ■ maize intercropped with *C. spectabilis*; ● maize intercropped with *U. ruziziensis*.

that the requirement for nitrogen fertilization is lower for sole maize.

In the second year, principal components analysis (PCA) showed for the agronomic variables of maize (Fig. 6(b)), that the first two PCs were responsible for 73.14% of the total variability and that the variables that most discriminated PC1 and PC2 were related, respectively, to the agronomic performance and vegetative growth of maize, as in the first year. Thus, the variables that most discriminated PC1 were stem diameter (−0.93), number of grains per row (−0.83), number of grains per ear (−0.86),

thousand-grain weight (−0.90), grain mass per ear (−0.94) and GY (−0.73). The variables most correlated with PC2 were leaf N content (0.60), plant height (0.87) and main ear (0.70). The number of rows per ear was not part of any component this year.

Based on the direct correlations between the variables and PC1, it was observed that the higher the agronomic performance of maize, the higher the GY, stem diameter, grain mass per ear, thousand-grain weight, number of grains per ear, number of grains per row and leaf N content. Furthermore, the biplot analysis showed that the highest agronomic performance occurred




Fig. 6. Biplot of the PCs for the distribution of cropping systems under nitrogen levels as a function of the agronomic variables of maize in experiments I (a) and II (b) in Jaboticabal, SP, Brazil. The cropping systems are indicated by M: sole maize, M + C: maize intercropped with *C. spectabilis* and M + U: maize intercropped with *U. ruziziensis*. The numbers accompanying the cropping system indicate the N level (kg/ha). The agronomic variables are: NRE = number of rows per ear; GME = grain mass per ear; SD = stem diameter; 1000GW = 1000-grain weight; NGE = number of grains per ear; GY = grain yield; NGR = number of grains per row; LNC = leaf N content; EH = ear height; PH = plant height.

for sole maize at the N level of 210 kg/ha, maize intercropped with *C. spectabilis* at the N levels of 70, 140 and 210 kg/ha and maize intercropped with *U. ruziziensis* at N levels of 70, 140 and 210 kg/ha.

Land equivalent ratio

The intercropping systems, either with *C. spectabilis* or *U. ruziziensis*, showed a higher LER than the sole maize cropping system (Fig. 7). Between the two intercropping systems, the maize

+ *U. ruziziensis* intercropping system showed a higher LER with increasing N levels in the top-dressing. Top-dressing N fertilization increased the LERs of the sole maize cropping and maize + *U. ruziziensis* intercropping systems. The maximum LER value for the former system (1.14) was obtained at the 128 kg/ha N level, whereas linear increments of 0.012 for every 10 kg/ha of N added to the top-dressing fertilization were observed in the latter. In the intercropping system with *C. spectabilis*, top-dressing N fertilization did not affect the LER, which showed an average value of 1.36.




Fig. 7. LER of intercropped systems as a function of nitrogen levels applied as top-dressing in the average of the two years. M: sole maize; M+Cs: maize intercropped with *C. spectabilis*; M+Ur: maize intercropped with *U. ruziziensis*. * $P < 0.05$; ** $P < 0.01$.

Discussion

Cover crops enhance the sustainability of cropping systems owing to the various ecological benefits generated by their cultivation, including producing straw for soil cover thus mitigate evaporation from the soil and run-off, improve nutrient cycling and weed control (Pacheco et al., 2013; Schipanski et al., 2014). Although sustainability in cropping systems is one of the main agricultural goals in many countries (United Nations, 2021), these benefits alone do not ensure the implementation of intercropped systems by farmers. This is because cover crops do not provide direct economic benefits. Thus, intercropping systems involving cover crops are a better alternative to the sole cultivation of cover crops since they combine the ecosystem benefits of cover crops and the economic benefits of grain production in the same area. However, an increased acceptance of cover crop cultivation should not affect the GY.

Our results showed that, although the cover crops do not interfere with the growth of maize, in the presence of *C. spectabilis* the ears were located higher up on the stem, in the two years. Thus, the fact that maize intercropped with *C. spectabilis* has ears in a higher region of the stem without necessarily having excessive growth of the plant is beneficial, because it avoids lodging and facilitates the mechanized harvesting operation, especially in intercropping systems, allowing the harvest of only the ear and not of parts of cover plants, which could lead to problems in the harvesters.

The increase in the variables related to plant growth as a function of nitrogen fertilization is justified by the N soil solution availability increase, favouring the uptake by plants, which use this element in their metabolism. Thus, the increase in leaf N content in the first year in an intercropping system agrees with that found in the literature (Souza et al., 2011; dos Santos et al., 2013; Pivetta et al., 2019).

In the second year, the results of leaf N content showed higher requirement for nitrogen fertilization by maize intercropped with *U. ruziziensis*, and in the absence of the nutrient as top-dressing, the only value below the range considered adequate in the diagnostic leaf of maize, from 27 to 35 g/kg (Cantarella and Rajj, 1997), occurred in this intercropping. Therefore, for the leaf N

content of maize intercropped with *U. ruziziensis* to reach the recommended level, N fertilization of at least 36 kg/ha needs to be applied.

The increase in stem diameter as a function of N levels in both years is consistent with that observed by Gonçalves et al. (2016) and Marini et al. (2015). For being an organ with structural and reserve function, a larger stem diameter is advantageous to avoid lodging of plants and contribute with translocated photoassimilates to grain filling (da Silva Brum et al. 2016).

The results of PCA confirmed the direct relationship between stem diameter and the variables of grains and ear, confirming the capacity of the stem to translocate reserves for grain filling. In addition, especially in the first year, it was possible to observe the greater dependence of nitrogen fertilization for maize intercropped with *U. ruziziensis* in comparison with other cropping systems.

The negative loadings of all variables in PC1 indicate direct correlations between the variables, showing a relationship between stem diameter and the variables of ear and grains, since thicker stems indicate greater reserve of photoassimilates and, consequently, provide better development of grains and ear. Therefore, the higher the agronomic performance of maize, the higher the GY, stem diameter, grain mass per ear, thousand-grain weight, number of grains per ear, number of grains per row and leaf N content.

A similar result regarding the lower GY for maize intercropped with *U. ruziziensis* compared to sole maize was verified by Arf et al. (2018). Souza et al. (2019), on the other hand, observed no differences between sole maize and intercropped with *U. ruziziensis* and *C. spectabilis* for GY, as occurred in the second year of our study.

The differences reported in the literature for GY are related to crop management, intercropping modalities used and climate and soil factors. Thus, the favourable conditions of soil fertility possibly met the nutritional needs of maize in the second year of our study, avoiding losses in GY when the cover plant was present, especially *U. ruziziensis*. The study plots were allocated to the same location during the two years of experimentation. Hence, while the maize crops did not benefit from the greater amounts of N from the cover crop residues during the first year, they did so during the second year. Deienno et al. (2021) reported that the intercropping systems of maize with *C. spectabilis* and *U. ruziziensis* left up to 66 and 94% more N, respectively, in the soil than the N left by the sole maize cropping system. This suggests that the higher amounts of N left in the soil after the first year of the experiment helped mitigate the lower availability of N for maize intercropping systems during the second year. Duete et al. (2008) found that, although the use of 39% of nitrogen fertilizer in the form of N¹⁵ urea by maize, in an Oxisol, the main source of the nutrient for the crop was the soil.

The quadratic or linear increase as a function of the increase of N of the variables number of grains per row, number of grains per ear, thousand-grain weight and GY in the first year, as well as these variables plus the grain mass per ear in the second year, occurred due to the greater availability for absorption of the nutrient by the plants. This is because N acts on the synthesis of proteins, hormones, chlorophylls and other compounds important for plant development; therefore, vegetative and reproductive aspects are favoured by the increase in the nutrient (Costa et al., 2014).

Regarding to GY in the first year, the top-dressing nitrogen fertilization in sole maize in the present study is consistent with

the value recommended by Cantarella and Rajj (1997), close to 100 kg/ha of N. On the other hand, the intercropped systems showed greater demand for nitrogen to obtain similar yields compared to sole maize. This occurs because in intercropped systems, cover crops take advantage of the applied N for maize top-dressing fertilization. This is even more severe in intercropping systems where cover crops are sown simultaneously with maize. In this situation, at the time of top-dressing fertilization, cover crops have a high development and are able to compete with maize for the nutrient.

In addition, among the intercropping systems, the presence of *U. ruziziensis* requires N levels even higher than that of *C. spectabilis* to obtain maximum yields. This is because, in addition to the requirement for N of the grass being higher than that of the legume, it does not have a contribution of BNF for N supply. Thus, the intercropping of maize with *U. ruziziensis*, when sown simultaneously, must receive top-dressing nitrogen fertilization with higher N levels compared to sole maize, or even the splitting of N, increasing fertilization efficiency (Du *et al.*, 2019). Almeida *et al.* (2017) observed in the intercropping of maize with *U. ruziziensis*, with simultaneous sowing, competition of the grass with maize up to the N level of 100 kg/ha. However, when the intercropping occurs simultaneously with top-dressing nitrogen fertilization in maize, there is no competition of the forage species with maize for abiotic factors, including N (Borghetti *et al.*, 2014).

Several studies have shown that there is a reduction in maize yield when it is intercropped (Pariz *et al.*, 2011; Crusciol *et al.*, 2013; Cambaúva *et al.*, 2019). Cambaúva *et al.* (2019) observed that the intercropping of maize + *Crotalaria juncea* reduces maize GY. However, for *C. spectabilis*, the same used in the present study, the GY of maize was equal to that obtained with sole maize. According to the authors, the reduction in the yield of maize intercropped with *C. juncea* is due to the greater competition of this plant with maize compared to *C. spectabilis*, due to its greater plant height (Cambaúva *et al.*, 2022).

Intercropping systems can also promote competition for water because the leaf area index in intercropped systems is higher than that in the sole crop systems, increasing transpiration (Martin-Guay *et al.* 2018; Cambaúva *et al.*, 2019). While assessing the temporal variability of soil moisture in the sole maize and maize + *C. spectabilis* intercropping systems, Trevisan *et al.* (2021) observed lower soil moisture during some periods in the intercropped system, reducing water availability for maize. Silva *et al.* (2020) found that soil moisture can be up to 10% lower in the maize + *U. ruziziensis* intercropping system than in the sole maize cropping system, especially during the grain-filling stage. It is noteworthy that in regions with high-intensity rainfall, a higher leaf area index may mitigate the impact of rain on the soil, reducing surface runoff and the potential for soil erosion, especially at the beginning of maize development.

Another benefit of intercropping is the higher total yield compared to sole cropping of maize and cover crops. This was verified by the LERs obtained in the present study, in which the intercropping systems presented LER values that were 35% higher than those of the sole cropping system, thereby indicating that for a total production similar to that of the intercropped systems, the sole cropping area should be increased by more than 35%. This value was similar to that observed by Martin-Guay *et al.* (2018), who found that the LER of intercropping was 1.30, increasing the gross energy produced per area by 38%.

Depending on the cropping system, fertilization management is essential for increasing land yield. The LER of the sole maize increased by up to 14% and the LER of the maize + *U. ruziziensis* intercropping system increased at a rate of 0.023 for every 10 kg/ha of N. These results were similar to those observed by Salama *et al.* (2022) who evaluated the effect of intercropping maize with cowpea and N fertilization management on LER. They found that some intercropping configurations increased the LER by more than 30% and that better N fertilization management increased the LER by more than 25%. In the present study, the highest LER of the sole maize was obtained under 128 kg/ha N level, a value close to that recommended for maize in this region, which is 140 kg/ha of N (Cantarella and Rajj, 1997).

The results obtained here suggest that the maize intercropping, associated with an adequate management of top-dressing nitrogen fertilization, may be a great strategy for farmers and technicians to increase sustainability in their crops. In addition to the ecosystem benefits of intercropping, our findings show that it is possible to associate the same land area with more production and more than one economic activity. This was evidenced by the 35% increase in the LER using intercropped systems and top-dressing fertilization, generating scientific information to prevent the opening of new cultivation areas in regions considered marginal, either due to climate or environmental issues.

Conclusion

Maize intercropping with *U. ruziziensis* and *C. spectabilis* increases the requirement for N fertilization to obtain an agronomic performance similar to that of the sole maize cropping system. This is especially true for intercropping with *U. ruziziensis*, where the increase in GY as a function of N levels is linear, with an average increase of 120 kg/ha for every 10 kg/ha of N applied. For other intercropping, N fertilization can either increase the GY quadratically or not interfere with this attribute. Compared to a sole maize crop, intercropping and N fertilization help increase land yield, with an increase in the LER by more than 35%. We determine that intercropping of maize with *U. ruziziensis* or *C. spectabilis* and with an adequate top-dressing N management is a potential strategy to assist farmers in using a more conservation-based cropping system. This will increase yields and promote a more efficient land use.

Acknowledgements. The present study builds on the PhD thesis of Stefany Silva de Souza published on 9 July 2020 (available from the São Paulo State University repository).

Author contributions. Conceptualization: SSS, JAF and LBL; data curation: SSS, JAF and APC; formal analysis: SSS, JAF and JMD; funding acquisition: SSS, LBL and GSR; investigation: SSS, LBL and GSR; methodology: SSS, JAF and LBL; project administration: LBL; resources: LBL; supervision: LBL and GSR; validation: APC and GSR; visualization: JMD and LBL; writing – original draft: SSS and APC; writing – review and editing: LBL and GSR

Financial support. This study was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), finance code 001.

Conflict of interest. The authors declare there are no conflicts of interest.

References

- Almeida REM, Favarin JL, Otto R, Pierozan C, Oliveira SM, Tezotto T and Lago BC (2017) Effects of nitrogen fertilization on yield components in a corn-palisadegrass intercropping system. *Australian Journal of Crop Science* 11, 352–359.

- Arf O, Meirelles FC, Portugal JR, Buzetti S, Sá ME and Rodrigues RAF (2018) Benefícios do milho consorciado com gramínea e leguminosas e seus efeitos na produtividade em sistema plantio direto. *Revista Brasileira de Milho e Sorgo* 17, 431–444.
- Borghesi E, Crusciol CAC, Trivelin PCO, Nascente AS, Costa C and Mateus GP (2014) Nitrogen fertilization ($15\text{NH}_4\text{NO}_3$) of palisadegrass and residual effect on subsequent no-tillage corn. *Revista Brasileira de Ciência do Solo* 38, 1457–1468.
- Calegari A and Carlos JAD (2014) Recomendações de plantio e. Informações gerais sobre o uso de espécies para adubação verde no Brasil. In de Lima Filho OF, Ambrosano EJ, Rossi F and Carlos JAD (eds). *Adubação verde e plantas de cobertura no Brasil: Fundamentos e prática*. Brasília: Embrapa, pp. 451–477.
- Cambaiva V, Leal FT and Lemos LB (2019) Crescimento, produtividade e palhada de milho exclusivo e consorciado com crotalárias em diferentes espaçamentos. *Revista Brasileira de Milho e Sorgo* 18, 99–111.
- Cambaiva V, Leal FT, Coelho AP and Lemos LB (2022) Do *Crotalaria* plant height and maize inter-row spacing affect intercropped maize yield? *Revista Brasileira de Milho e Sorgo* 21, e1260.
- Cantarella H and Raij BV (1997) Cereais. In Raij BV, Cantarella H, Quaggio JÁ and Furlani AMC (eds), *Recomendação de adubação e calagem para o Estado de São Paulo*, 2nd edn. Campinas: Instituto Agronômico de Campinas, pp. 48–49.
- Carmo CAFS, de Araújo WS, Bernardi ACC and Saldanha MFC (2000) *Métodos de análise de tecidos vegetais utilizados na Embrapa Solos*. Rio de Janeiro: Embrapa Solos, 41 p. (Embrapa Solos. Circular Técnica, 6).
- Costa NR, Andreotti M, Lopes KSM, Santos FG and Pariz CM (2014) Adubação nitrogenada em capins do gênero *Urochloa* implantados em consórcio com a cultura do milho. *Revista Brasileira de Ciências Agrárias* 9, 376–383.
- Crusciol CAC, Nascente AS, Mateus GP, Borghi E, Leles EP and Santos ND (2013) Effect of intercropping on yields of corn with different relative maturities and palisadegrass. *Agronomy Journal* 105, 599–606.
- da Silva Brum M, dos Cunha VS, Stecca JDL, Grando LFT and Martin TN (2016) Components of corn crop yield under inoculation with *Azospirillum brasilense* using integrated crop-livestock system. *Acta Scientiarum Agronomy* 38, 485–492.
- Deianno J, Souza SS, Coelho AP and Lemos LB (2021) Maize intercropping and nitrogen fertilization aiming grain yield and implement a no-till system. *Brazilian Journal of Maize and Sorghum* 20, e1225.
- de Oliveira P, Kluthcouski J, Favarin JL and de Santos DC (2010) *Sistema Santa Brígida – Tecnologia Embrapa: Consorciação de Milho com Leguminosas*. Santo Antonio de Goiás: Embrapa Arroz e Feijão, 16 p. (Embrapa Arroz e Feijão. Circular Técnica, 88).
- dos Santos LPD, Aquino LA, Nunes PHMP and Xavier FO (2013) Doses de nitrogênio na cultura do milho para altas produtividades de grãos. *Revista Brasileira de Milho e Sorgo* 12, 270–279.
- Du X, Xi M and Kong L (2019) Split application of reduced nitrogen rate improves nitrogen uptake and use efficiency in sweetpotato. *Scientific Reports* 9, 1–11.
- Duete RRC, Muraoka T, da Silva EC, Trivelin PCO and Ambrosano EJ (2008) Manejo da adubação nitrogenada e utilização do nitrogênio (^{15}N) pelo milho em Latossolo Vermelho. *Revista Brasileira de Ciência do Solo* 32, 161–171.
- Fan Y, Wang Z, Liao D, Ali Raza M, Wang B, Zhang J, Chen J, Feng L, Wu X, Liu C, Yang W and Yang F (2020) Uptake and utilization of nitrogen, phosphorus and potassium as related to yield advantage in maize soybean intercropping under different row configurations. *Scientific Reports* 10, 1–10.
- Ferreira DF (2011) Sisvar: a computer statistical analysis system. *Ciência e Agrotecnologia* 35, 1039–1042.
- Gastwirth JL, Gel YR and Miao W (2009) The impact of Levene's test of equality of variances on statistical theory and practice. *Statistical Science* 24, 343–360.
- Gonçalves AKA, Silva TRB and Brandão AG (2016) Manejo de adubação nitrogenada em milho solteiro e consorciado com *Brachiaria ruziziensis*. *Revista Brasileira de Milho e Sorgo* 15, 318–327.
- Kaiser HF (1958) The varimax criterion for analytic rotation in factor analysis. *Psychometrika* 23, 187–200.
- Kappes C and Zancanaro L (2015) Sistemas de consórcios de braquiária e de crotalárias com a cultura do milho. *Revista Brasileira de Milho e Sorgo* 14, 219–234.
- Kluthcouski J, Cobucci T, Aidar H, Yokoyama LP, Oliveira IP, Costa JLS, Silva JG, Vilela L, Barcellos AO and Magnabosco CU (2000) *Sistema Santa Fé – Tecnologia Embrapa: integração lavoura-pecuária pelo consórcio de culturas anuais com forrageiras, em áreas de lavoura, nos sistemas de plantio direto e convencional*. Santo Antônio de Goiás: Embrapa Arroz e Feijão, 28 p. (Embrapa Arroz e Feijão. Circular Técnica, 38).
- Marcelo AV, Corá JE and Fernandes C (2012) Sequências de culturas em sistema de semeadura direta II – decomposição e liberação de nutrientes na entressafra. *Revista Brasileira de Ciência do Solo* 36, 1568–1582.
- Marini D, Guimarães VF and Dartora J, do Lana MC and Pinto Júnior AS (2015) Growth and yield of corn hybrids in response to association with *Azospirillum brasilense* and nitrogen fertilization. *Revista Ceres* 62, 117–123.
- Martin-Guay M-O, Paquette A, Dupras J and Rivest D (2018) The new Green Revolution: sustainable intensification of agriculture by intercropping. *Science of the Total Environment* 615, 767–772.
- Martins D, Gonçalves CG and Silva Junior AC (2016) Coberturas mortas de inverno e controle químico sobre plantas daninhas na cultura do milho. *Revista Ciência Agronômica* 47, 649–657.
- Mead R and Willey R (1980) The concept of a 'land equivalent ratio' and advantages in yields from intercropping. *Experimental Agriculture* 16, 217–228.
- Mingotte FLC, Yada MM, Jardim CA, Fiorentin CF, Lemos LB and Fornasieri Filho D (2014) Sistemas de cultivo antecessores e doses de nitrogênio em cobertura no feijoeiro em plantio direto. *Bioscience Journal* 30, 696–706.
- Mingotte FLC, Jardim CA, Yada MM, Amaral CB, Chiamolera TPLC, Coelho AP, Lemos LB and Fornasieri Filho D (2020) Impact of crop management and no-tillage system on grain and straw yield of maize crop. *Cereal Research Communications* 48, 399–407.
- Murrell EG, Schipanski ME, Finney DM, Hunter MC, Burgess M, LaChance JC, Baraibar B, White CM, Mortensen DA and Kaye JP (2017) Achieving diverse cover crop mixtures: effects of planting date and seeding rate. *Agronomy Journal* 109, 259–271.
- Pacheco LP, Barbosa JM, Leandro WM, de Machado PLOA, de Assis RL, Madari BE and Petter FA (2013) Ciclagem de nutrientes por plantas de cobertura e produtividade de soja e arroz em plantio direto. *Pesquisa Agropecuária Brasileira* 48, 1228–1236.
- Pariz CM, Andreotti M, Azenha MV, Bergamaschine AF, Mello LMM and Lima RC (2011) Corn grain yield and dry mass of *Brachiaria* intercrops in the crop-livestock integration system. *Ciência Rural* 41, 875–882.
- Pivetta LA, Jordão LM, Larini WF, Luchese AV, Gasparin CE, Debuss AL, Carvalho MG and Silva WO (2019) Doses e épocas de aplicação de nitrogênio em milho safrinha. *Journal of Agronomic Sciences* 8, 1–13.
- Raij BV and Cantarella H (1997) Milho para grão e silagem. In Raij BV, Cantarella H, Quaggio JA and Furlani AMC (eds). *Recomendação de adubação e calagem para o Estado de São Paulo*. Campinas: Instituto Agronômico de Campinas (IAC), pp. 56–59 (Boletim Técnico, 100).
- Rolim GS, Camargo MBP, Lania DG and Moraes JFL (2007) Classificação climática de Köppen e de Thornthwaite e sua aplicabilidade na determinação de zonas agroclimáticas para o estado de São Paulo. *Bragantia* 66, 711–720.
- Royston P (1995) Remark AS R94: a remark on algorithm AS 181: the W-test for normality. *Journal of the Royal Statistical Society* 44, 547–551.
- Salama HAS, Nawar AI and Khalil HE (2022) Intercropping pattern and n fertilizer schedule affect the performance of additively intercropped maize and forage cowpea in the Mediterranean region. *Agronomy* 12, 107.
- Schipanski ME, Barbercheck M, Douglas MR, Finney DM, Haider K, Kaye JP, Kemanian AR, Mortensen DA, Ryan MR, Tooker J and White C (2014) A framework for evaluating ecosystem services provided by cover crops in agroecosystems. *Agricultural Systems* 125, 12–22.
- Silva GSF, Andrade Júnior ASD, Cardoso MJ and Araújo Neto RBD (2020) Soil water dynamics and yield in maize and *Brachiaria ruziziensis* intercropping. *Pesquisa Agropecuária Tropical* 50, e59809.
- Silveira PM, Silva JH, Lobo Junior M and Cunha PCR (2011) Atributos do solo e produtividade do milho e do feijoeiro irrigado sob sistema integração lavoura-pecuária. *Pesquisa Agropecuária Brasileira* 46, 1170–1175.

- Soil Survey Staff** (2014) *Keys to Soil Taxonomy*, 12th ed, Washington, DC: Department of Agriculture. Natural Resources Conservation Service, 372 p.
- Souza JA, Buzetti S, Teixeira Filho MCM, Andreotti M, Sá ME and Arf O** (2011) Adubação nitrogenada na cultura do milho safrinha irrigado em plantio direto. *Bragantia* **70**, 447–454.
- Souza SS, Couto Júnior PA, de Flôres JA, Mingotte FLC and Lemos LB** (2019) Maize cropping systems and response of common bean in succession subjected to nitrogen fertilization. *Pesquisa Agropecuária Tropical* **49**, 1–11.
- Tan M** (2020) Macro- and micromineral contents of different quinoa (*Chenopodium quinoa* Willd.) varieties used as forage by cattle. *Turkish Journal of Agriculture and Forestry* **44**, 46–53.
- Trevisan M, Silva LFS, Fontanetti A, Souza Gallo A and França Guimarães N** (2021) Soil temperature and moisture in the intercropping of maize with *Crotalaria spectabilis* and *Cajanus cajan* in an organic system. *Research, Society and Development* **10**, e539101422443.
- United Nations** (2021) *Transforming our World: The 2030 Agenda for Sustainable Development*. Available in: <https://sdgs.un.org/2030agenda> (Accessed 31 May 2021).
- USDA - United States Department of Agriculture** (2021) World Agricultural Production. Foreign Agricultural Service/USDA Global Market Analysis, Circular Series WAP 2-20. Available in: <https://apps.fas.usda.gov/psdonline/circulars/production.pdf> (Accessed 3 January 2021).