

The mitochondrial genome of *Gyrodactylus salaris* (Platyhelminthes: Monogenea), a pathogen of Atlantic salmon (*Salmo salar*)

T. HUYSE^{1,*}, L. PLAISANCE^{2,3}, B. L. WEBSTER¹, T. A. MO⁴, T. A. BAKKE², L. BACHMANN² and D. T. J. LITTLEWOOD¹

¹ Parasitic Worms Group, Department of Zoology, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

² The Natural History Museum, Department of Zoology, University of Oslo, P.O. Box 1172, Blindern, NO-0318 Oslo, Norway

³ Marine Biology Research Division 0202, Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA 92093-0202, USA

⁴ National Veterinary Institute, P.O. Box 8156 Dep., NO-0033 Oslo 1, Norway

(Received 12 September 2006; revised 25 October 2006; accepted 26 October 2006; first published online 11 December 2006)

SUMMARY

In the present study, we describe the complete mitochondrial (mt) genome of the Atlantic salmon parasite *Gyrodactylus salaris*, the first for any monogenean species. The circular genome is 14 790 bp in size. All of the 35 genes recognized from other flatworm mitochondrial genomes were identified, and they are transcribed from the same strand. The protein-coding and ribosomal RNA (rRNA) genes share the same gene arrangement as those published previously for neodermatan mt genomes (representing cestodes and digeneans only), and the genome has an overall A + T content of 65%. Three transfer RNA (tRNA) genes overlap with other genes, whereas the secondary structure of 3 tRNA genes lack the DHU arm and 1 tRNA gene lacks the TΨC arm. Eighteen regions of non-coding DNA ranging from 4 to 112 bp in length, totalling 278 bp, were identified as well as 2 large non-coding regions (799 bp and 768 bp) that were almost identical to each other. The completion of the mt genome offers the opportunity of defining new molecular markers for studying evolutionary relationships within and among gyrodactylid species.

Key words: mitochondrial genome, *Gyrodactylus salaris*, Monopisthocotylea, *Salmo salar*, gyrodactylosis.

INTRODUCTION

Gyrodactylus salaris Malmberg, 1957 is a monogenean flatworm that is among the most serious threats to wild and farmed Atlantic salmon (*Salmo salar*) today. Ectoparasitic, viviparous and with a direct life-cycle, this species can occur in such high numbers on its hosts, and throughout entire river systems, that it can cause the collapse of fish stocks. Losses are estimated at more than 15% of the total wild salmon catch in Norway alone, with concomitant costs from the loss of fisheries, tourism and the need to survey and eradicate infected stocks, amounting to more than USD 50 million per year (Mo *et al.* 2004). Salmon stocks in 46 watercourses in Norway are threatened or have been lost (Hansen *et al.* 2003; Mo and Norheim, 2005), and other fish species such as rainbow trout (*Oncorhynchus mykiss*), brook trout (*Salvelinus fontinalis*) and Arctic charr (*S. alpinus*) readily act as suitable hosts (Bakke *et al.*

2002; Robertsen *et al.* 2006). In Europe, particularly in the UK, economically important salmon populations are at severe risk from the introduction of *G. salaris* with live fish, especially rainbow trout (Peeler and Thrush, 2004). No convincing, parasite-specific control measures are yet available. Presently, the only methods to eradicate *G. salaris* include killing all fish by the use of the biocide rotenone or by aqueous aluminium to selectively kill the parasites (Soleng *et al.* 1999, 2005; Poléo *et al.* 2004). Despite all eradication attempts, the parasite is still increasing its geographical distribution and in approximately 30% of the treated rivers, the parasite has reappeared (Mo and Norheim, 2005). Many EU states have instigated codes of practice for the movement of *G. salaris* (see Peeler *et al.* 2006), but there are few tools available to accurately track and understand the origins, spread and epidemiology of infections (Bakke *et al.* 2006).

The accurate diagnosis of infection is central to the control of *G. salaris*. Traditional methods for identification of gyrodactylids are based on morphometry of the attachment sclerites. However, more than 400 nominal species have been described within the genus *Gyrodactylus* (Harris *et al.* 2004), and additional species are continuously discovered as

* Corresponding author: Parasitic Worms Group, Department of Zoology, The Natural History Museum, Cromwell Road, London SW7 5BD, UK. Tel: +442 079426115. Fax: +442079425151. E-mail: T.Huyse@nhm.ac.uk

Table 1. List of PCR primer combinations for all overlapping regions

Primers	5' to 3'
Universal*	
<i>U12SR</i>	TAACCGCRRMTGCTGGCACTG
<i>U12SF</i>	CAGTGCCAGCAKYYGCGGTTA
<i>CytbF</i>	GGWTAYGTWYTWCCWTGRGGWCARAT
<i>CytbR</i>	GCRTAWGCRAAWARRAARTAYCAYTCWGG
<i>UNAD5F</i>	TTRGARGCNATGCGBGCHCC
<i>UNAD5R</i>	GGWGC MCGCATNGCYTCYAA
<i>UND1F</i>	CGHAAGGNCCNAAHAAGGT
Long PCR*	
<i>Gsal_16SF</i>	CCGGTGTAAGCCAGGTTGGTTTC
<i>Gsal_CytbR</i>	GGTTAGTACCGTGGCWCGCCA
<i>Gsal_COI_544F</i>	T TACTACGGATGGTGTTCGCC
<i>Gsal_COI_204R</i>	GAAATACCAGACAGGTGAAGCG
<i>Gsal_CytbF</i>	TGGGCWGCCACGGTACTAACC
<i>Gsal-MIT3</i>	TGGCATCAATAGCCAAGCCCTTAAAGC
Sequencing primers	
<i>16Snew</i>	AAGTCAACATCGAGGTAGC
<i>16S_γ2</i>	CGGTCTTAACTCAAGAGCTTCA
<i>16STL-3'</i>	TYACRCCGGTCTKAACTC
<i>16STL_5'</i>	KTRCCTTTTGYATCATG
<i>COI_mono5</i>	TAATWGGTGGKTTTGGTAA
<i>COI_mono3</i>	TAATGCATMGGAAAAAACA
<i>Platymt12S_F1</i>	GTGCCAGDCYCGGTTA
<i>Gsal_COI_F1</i>	ATCGGAGGAGTGACAGGGATAGTG
<i>Gsal_COI_F2</i>	CGTGTATGCTTGTATGACGACTC
<i>Gsal_16S_R1</i>	GCTCTTAGGGTCTTTCCGTC
<i>Gsal_COI_F3</i>	GAGTAGTAGGGTGGTAAACGG
<i>Gsal_COI_204R</i>	GAAATACCAGACAGGTGAAGCG
<i>Gsal_ND5_seqR</i>	CAGGAACAGAGCATTATTAGGC
<i>Gsal_NC2_seqR</i>	GCCTCATCTGCCTACTTATTTG
<i>Gsal_cox2_segF</i>	GGAACCAGAGGCTTGTAATAATGTC
<i>Gsal_cox2_segF2</i>	GGAGTTTTCGTCCGATACTG

* General and long PCR primers have also been used as sequencing primers.

new suitable host species are examined. Also, within *G. salaris*, there is significant variation in morphology. Its hook morphology closely resembles those of other related species. Discriminating among them requires high quality scanning electron micrographs and the application of sophisticated statistical tools (Shinn *et al.* 2001). However, the discrimination between *G. salaris* and *G. thymalli* Zitnan, 1960 has proved difficult, since at least 1 Norwegian *G. thymalli* population fell within the morphological variation of *G. salaris* (Shinn *et al.* 2004; unpublished observations). Thus, morphology alone does not easily lend itself to routine use in a diagnostic laboratory. To date, molecular identification methods have mainly used the internal transcribed spacers (ITS) of the nuclear ribosomal DNA (rDNA), since they display a low degree of variation within species (Zietara *et al.* 2002; Huysse *et al.* 2003; Matejusová *et al.* 2003). However, although *G. thymalli* has been described as a distinct species based on morphological, ecological and pathological grounds, no variation was detected in the ITS region when compared with *G. salaris* (see Cunningham, 1997). Limited success has been achieved with the sequencing of the intergenic spacer (IGS) of rDNA

(Sterud *et al.* 2002; Cunningham *et al.* 2003; Hansen *et al.* 2006), but a suite of alternative markers is still needed for the detection of population variation, to further understand the taxonomy and biology of the parasite, and to study its transmission and dispersion in space and time.

Mitochondrial (mt) genomes offer a wealth of informative characters, useful in phylogenetic and population genetic studies (Boore *et al.* 2005). Many commercially and economically important parasites are now the focus of mitogenomic studies, as their mt genomes are now relatively easy to characterize and can be used as the basis for the design of molecular markers (Place *et al.* 2005). Thus far, partial sequences of the cytochrome *c* oxidase subunit 1 gene (*cox1*) have been used in studying the phylogeographic structure of *G. salaris* and *G. thymalli* (see Hansen *et al.* 2003, 2006; Meinila *et al.* 2004). These studies have revealed a high level of intra- and interspecific variation and could not confirm monophyly of either species. However, more variable regions are likely to be present in the other parts of the mt genome, offering higher resolution for studies on more recent evolutionary processes in *G. salaris*, such as host switching and speciation processes.




Fig. 1. Mitochondrial genome map of *Gyrodactylus salaris* from Atlantic salmon in the River Signaldalselva, North Norway. Protein-coding genes are shown as open arrows, ribosomal RNA genes as shaded arrows, and tRNAs as arrowed lines. The non-coding regions NC1 and NC2 share a 722 bp repeated element (see text for further details).

Comparing the mt genome sequences of different species will identify regions of high sequence divergence. In the present study, we sequenced and characterized the mt genome of *G. salaris*, the first for any monogenean. We describe the gene order, codon usage, tRNA features and gene boundaries, and compare these features with those found in other parasitic flatworms. This study provides the first step to defining new molecular mt markers, which will assist in studying the evolution, ecology and epidemiology of this important fish pathogen and its control.

MATERIALS AND METHODS

Sample collection and preparation

Parasite samples of *G. salaris* from wild Atlantic salmon parr were collected in the river Signaldalselva, North Norway in September 2001. This

river is infected *via* migrating infected parr from the river Skibotnelva, which, in turn, has been infected by an accidental introduction of Swedish salmon parr. Hansen *et al.* (2003) sequenced the partial *cox1* of this strain (haplotype B), and found this haplotype in other Swedish rivers which drain into the Baltic Sea and are close to the Hölle hatchery, Sweden. The authors proposed that River Signaldalselva is most likely infected with *G. salaris* specimens originating from the Hölle hatchery, which is the type locality for this species. The samples were kept in 96% (v/v) ethanol at 5 °C. For genomic DNA extraction, a total of 5–10 parasites were briefly air dried to remove the ethanol, pooled in 5 µl of H₂O and digested by the addition of 5 µl of lysis solution, 1× PCR buffer (Eurogentec; 16 mM (NH₄)₂SO₄, 67 mM Tris-HCl (pH 8.8 at 25 °C), 0.01% Tween-20), 0.45% (v/v) Tween 20, 0.45% (v/v) NP 40 and 60 µg/ml of proteinase K (Sigma). The

Table 2. Start and stop positions for individual protein-coding genes in the mitochondrial genome of *Gyrodactylus salaris* from River Signaldalselva, North Norway

(Inferred start and stop codons are indicated (see also Fig. 1).)

Genes	Length		Codon		Position 5' to 3'	Genes	Length bp	Position 5' to 3'
	bp	aa	Start	Stop				
Protein						RNAs		
<i>cox3</i>	639	213	ATG	TAA	1–639	<i>rrnL</i>	956	8933–9889
<i>cytb</i>	1074	358	ATG	TAG	705–1778	<i>rrnS</i>	712	9950–10661
<i>nad4L</i>	249	83	ATG	TAA	1784–2032	<i>trnH</i>	63	639–701
<i>nad4</i>	1275	425	ATG	TAA	2005–3213	<i>trnF</i>	66	3216–3281
<i>atp6</i>	513	171	ATG	TAA	4081–4593	<i>trnV</i>	64	5457–5521
<i>nad2</i>	897	299	ATG	TAA	4600–5496	<i>trnA</i>	70	5520–5589
<i>nad1</i>	888	296	ATG	TAA	5657–6544	<i>trnD</i>	66	5590–5656
<i>nad3</i>	351	117	ATG	TAG	6828–7178	<i>trnN</i>	67	6555–6621
<i>cox1</i>	1548	516	ATG	TAA	7311–8858	<i>trnP</i>	66	6627–6692
<i>cox2</i>	582	194	ATG	TAA	10662–11243	<i>trnI</i>	72	6686–6757
<i>nad6</i>	483	161	ATG	TAA	11430–11912	<i>trnK</i>	65	6759–6824
<i>nad5</i>	1551	517	ATG	TAG	13156–14706	<i>trnS1</i>	72	7177–7238
						<i>trnW</i>	64	7243–7306
Non-coding						<i>trnT</i>	66	8868–8933
NC 1	799				3282–4080	<i>trnC</i>	60	9890–9949
NC 2	768				12184–12951	<i>trnE</i>	72	11355–11426
						<i>trnY</i>	67	11916–11982
						<i>trnL1</i>	68	11983–12050
						<i>trnQ</i>	63	12051–12113
						<i>trnM</i>	65	12120–12184
						<i>trnS2</i>	73	12952–13024
						<i>trnL2</i>	67	13020–13086
						<i>trnR</i>	67	13089–13155
						<i>trnG</i>	68	14720–14787

sample was incubated at 65 °C for 25 min, followed by 10 min at 95 °C to inactivate the proteinase.

Conventional and long polymerase chain reaction (PCR)

Part of the cytochrome *b* gene (*cytb*) was amplified (in 25 µl reaction volumes) using universal primers *CytbF*–*CytbR* (see Table 1) (Boore and Brown, 2000) and puRe *Taq* Ready-to-Go PCR beads (Amersham Biosciences). The solution consisted of 20 ng of gDNA and 40 pM of each PCR primer; beads contained 1.5 U *Taq* polymerase, 10 mM Tris-HCl (pH 9.0), 50 mM KCl, 1.5 mM MgCl₂, 200 µM each dNTP and stabilisers including BSA. Cycling conditions were as follows: initial denaturation for 3 min at 95 °C, followed by 40 cycles of 30 sec at 95 °C, 30 sec at 45 °C, 2 min at 72 °C, and a final extension for 10 min at 72 °C. Specific reverse and forward primers (see Table 1) were designed from the sequence of this fragment and from partial *cox1* from *G. salaris* available from GenBank (Accession no. AY258372). The mt genome was amplified by long PCR using the GeneAmp XL PCR kit (Applied Biosystems) or the Expand Long Template PCR System (Roche Applied Science) in 3 fragments employing primer pairs Gsal_CO1_544F–U12SR (2015 bp) Gsal_16SF–Gsal_CytbR (6109 bp) and

Gsal_CytbF–Gsal_MIT3R (6537 bp). Additional overlapping fragments were amplified, using, for example, primer pairs U12SF–UNAD5R, UND1F–Gsal_CO1_204R). Cycling conditions were: an initial denaturation for 30 sec at 94 °C, followed by 40 cycles of 20 sec at 94 °C, 30 sec at 58–65 °C, 8 min at 64–68 °C, and final extension of 10 min at 68 °C.

Cloning and sequencing

PCR products, covering the whole genome, were individually cloned using the TOPO[®]XL PCR Cloning Kit (Invitrogen), following manufacturer's instructions. Ten clones from each of the cloning reactions were grown for 15 h in 3 ml volumes of Luria-Bertani (LB) medium, shaking (220 rpm) at 37 °C. Plasmid DNA was extracted using QIAprep Spin Miniprep Kit (Qiagen). Clones were examined for inserts by digestion with the restriction endonuclease *EcoRI* (Invitrogen). Two positive clones from each of the reactions were selected for sequencing, carried out using Big Dye Chemistry (version 1.1) in a 3730 DNA Analyser (Applied Biosystems). The flanking regions of the inserts were sequenced with forward and reverse M13 primers, and sequence identity was verified using the Basic Local Alignment Search Tool (BLAST) (available at www.ncbi.nih.



Fig. 2. Alignment of the shared portions of non-coding regions NC1 and NC2 (see Fig. 1), illustrating sequence conservation, differences (shaded bases) and putative start codon (white on black bases) although no convincing open reading frames were detected.

gov/BLAST/). The remaining fragments were sequenced by primer walking (Table 1).

Annotation of sequences

Contiguous sequence fragments were assembled using Sequencher™ 4.5 (GeneCodes Corporation). Genome annotation was performed using MacVector® 7.2.3 (Accelrys). The sequence identity of open reading frames (ORFs) was verified using BLAST, and individual genes were aligned with published mt genomes of other flatworms to identify start and stop codons. The flatworms included: Digenea – *Schistosoma japonicum* (Accession no. NC_002544), *S. mekongi* (NC_002529), *S. mansoni* (NC_002545), *S. haematobium* (NC_008074), *S. spindale* (NC_008067), *Paragonimus westermani* (NC_002354) and *Fasciola hepatica* (NC_002546); Cestoda – *Echinococcus granulosus* (NC_008075), *E. multilocularis* (NC_000928), *Hymenolepis diminuta* (NC_002767), *Taenia crassiceps* (NC_002547), *T. asiatica* (NC_004826) and *T. solium* (NC_004022). The program tRNAscan-SE 1.21 (available at

Table 3. Base composition

	A %	C %	G %	T %	AT %
Protein genes					
1st codon position	30.8	16.4	22.9	29.9	60.7
2nd codon position	19.1	14.4	20.1	46.4	65.5
3rd codon position	32.1	24.0	20.7	23.1	55.2
All positions (total)	27.3	18.3	21.2	33.1	60.4
rRNA genes	33.9	14.9	17.5	33.7	67.6
tRNA genes	32.6	13.1	18.7	34.3	66.9
All coding sites	29.1	17.2	20.5	33.2	62.3
Non-coding NC1	35.8	16.5	19.5	28.2	63.0
Non-coding NC2	36.1	16.0	19.3	28.6	64.7

www.genetics.wustl.edu/eddy/tRNAscan-SE/) was used to identify tRNAs and their secondary structures (Lowe and Eddy, 1997). The tRNAs, which were not detected by tRNA scan-SE 1.21, were identified by searching for the conserved motif YUxxxR, where xxx denotes the anticodon, and by detecting stem and loop regions by eye. The 2




Fig. 3. Comparison of mean length of mitochondrial protein coding genes from parasitic flatworms; \pm S.D. Monogenea – *G. salaris* (this study); Digenea – *Schistosoma japonicum*, *S. mekongi*, *S. mansoni*, *S. haematobium*, *S. spindale*, *Paragonimus westermani* and *Fasciola hepatica*; Cestoda – *Echinococcus granulosus*, *E. multilocularis*, *Hymenolepis diminuta*, *Taenia crassiceps*, *T. asiatica* and *T. solium* (see text for GenBank Accession numbers).

Start and stop codons

All initiation codons predicted are ATG, whereas TAG (for *cytb*, *nad3* and *nad5*) and TAA (for *cox3*, *nad4L*, *nad4*, *atp6*, *nad2*, *nad1*, *cox1*, *cox2* and *nad6*) are used as stop codons. Other invertebrate initiation codons, such as ATC and GTG, were not identified. Protein-coding genes were translated using the flatworm (rhabditophoran) mitochondrial code (see Table 9 in GenBank; Telford *et al.* 2000).

tRNA secondary structures, and overlap between adjacent genes

All of the 22 tRNA genes were identified. All but 8 were identified using the program tRNAscan; the remaining tRNA genes were identified by examining for putative anti-codon sequences and determining characteristic flanking stem and loop features. The 2 leucine tRNAs code for CTA and TTA (*trnL1* and *trnL2*, respectively), and the 2 serine tRNAs code for AGC and TCA (*trnS1* and *trnS2*, respectively). Three tRNA genes overlap with other genes, namely *trnV* (39 bp with *nad2*), *trnS1* (5 bp with *trnP*), *trnS2* (2 bp with *nad3*). Of all of the predicted secondary structures for the tRNA genes, *trnS1*, *trnS2* and *trnC* lack the DHU arm, and *trnP* appears to lack the T Ψ C arm.

Nucleotide composition and codon usage

The overall nucleotide composition of all coding sites is: A (29.1%), C (17.2%), G (20.5%), and T (33.2%) (see Table 3). The genome has an overall A+T content of 62.3%. The protein-coding genes display a relatively low A+T content, particularly at the third

codon positions (55.2%). The highest A+T content is in the rRNA and tRNA genes (67.6 and 66.9%). These findings fall within the ranges reported for other mt genomes of flatworms (Le *et al.* 2002; Johnston 2006). The codons predicted to be most frequently used are AUA (214), CUA (187), UUC (169), UUA (150), and the least frequently used (G+C-rich) codons are CCG, CGC, CGT (11), CAG and CGG (9), not including the stop codons UAA (9) and UAG (3) (Table 4).

Non-coding regions

There are 18 non-coding regions ranging from 4 bp to 112 bp. Typical control regions are not readily identifiable within the mt genomes of flatworms. However, the origin of replication may be located in the non-coding region between *cox2* and *trnE* (positions 11244–11354), as this region folds with hairpins but without a T-rich loop (features of the control region as described by Wolstenholme (1992); nevertheless, this designation remains putative). Control regions are often targeted as a source of genetic markers, since they tend to vary considerably between species. Therefore, this region warrants characterization for additional strains and species of gyrodactylids. The 2 large non-coding regions, NC1 and NC2 (see Fig. 1) are 798 bp and 767 bp long, with an A+T content of 63 and 64.7%, respectively. NC1 is situated between *trnP* and *atp6* and NC2 within a cluster of tRNA genes (*trnW*, *trnL1*, *trnQ*, *trnM*, NC2, *trnS2*, *trnL2* and *trnR*). No repetitive regions were found in either of these non-coding regions. An alignment of NC1 and NC2 shows that almost the entire region (722 bp) is conserved (see Fig. 3), with only 6 bp (<0.1%) difference (Fig. 2).

The conserved fragment begins in frame on a recognized initiation codon, ATG. However, in spite of extensive searches, no significant ORFs (for putative genes) were detected. The stop codon TAA occurred frequently in the 3 reading frames within NC1 and NC2. Repeat regions, particularly tandemly repeated regions, can yield variable length polymorphisms through slippage events, but NC1 and NC2 are separated by several kilobases from one another, leading us to speculate as to their possible origin and function within the mt genome. One may hypothesize that the NC1 and NC2 are not of mitochondrial but of nuclear origin. However, these non-coding regions have been identified in different PCR products amplified by long PCR. None of the coding genes detected in these long PCR products displayed degeneration which is typical for nuclear copies of mt DNA. Sequences obtained from PCR products using different primer pairs were consistently the same. Furthermore, independent PCR and sequencing conducted at the Natural History Museums in London and in Oslo confirmed the results. Therefore, we conclude that a nuclear origin of the non-coding NC1 and NC2 regions can be ruled out.

In summary, the mt genome of *G. salaris*, from a pathogenic population infecting Atlantic salmon parr in northern Norway, has been sequenced and annotated. Short non-coding regions, a putative control region and 2 longer, almost identical but separated non-coding regions, offer the opportunity of discovering rapidly evolving regions of the genome suitable for the definition of strain-specific markers. The variability in these and other regions may now be explored by comparing *G. salaris* from various geographical locations with reference strains and with other species of *Gyrodactylus*. These markers can then be used to 'genotype' *G. salaris* occurring on other salmonid hosts (e.g. *Oncorhynchus mykiss*, *Salvelinus fontinalis* and *S. alpinus*) to recognize host-associated haplotypes, and to genetically characterize *G. salaris* strains with varying degrees of virulence and host-specificity to salmon (Mo, 2006; Olstad *et al.* 2006). Having available mt markers will enable tools to be developed for studying the transmission dynamics of various species of *Gyrodactylus* between/among different host species and river systems, providing crucial information for an improved understanding of the spread and epidemiology of this pathogen. Additional mt genomes from a diversity of flatworm taxa, including polyopisthocotylean monogeneans and turbellarians, will allow an assessment of the phylogenetic utility of mt genomes at deeper evolutionary levels within the phylum.

We are grateful to Dr Catherine Collins and Dr Carey Cunningham for providing starting material for this project. Dr Le Thanh Hoa provided useful primer sequences prior to their publication. We thank Julia Llewellyn-Hughes and Claire Griffin of the NHM

Sequencing Unit for expert technical assistance. T. H. was funded by a Marie Curie Fellowship (Meif-ct-2004-501684). D. T. J. L. and B. L. W. were funded by the Wellcome Trust, through a Senior Research Fellowship (043965) to D. T. J. L. L. P., T. A. B. and L. B. were supported by the National Centre for Biosystematics (currently NCB 146515/420; co-funded by NHM, UiO, and the Norwegian Research Council, NRC) and the 'Wild Salmon Program' (currently NRC 145861/720). Note on authorship: T. H. and D. T. J. L. conceived the project. T. A. M. and T. B. provided parasite samples. T. H. sequenced and annotated the genome with assistance from B. L. W. and D. T. J. L. L. P. assisted with primer design. T. H. and D. T. J. L. wrote the manuscript.

REFERENCES

- Bakke, T. A., Cable, J. and Harris, P. D.** (2006). The biology of gyrodactylid monogeneans: the "Russian Doll-killers". *Advances in Parasitology* (in the Press).
- Bakke, T. A., Harris, P. D. and Cable, J.** (2002). Host specificity dynamics: observations on gyrodactylid monogeneans. *International Journal for Parasitology* **32**, 281–308.
- Benson, G.** (1999). Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research* **27**, 573–580.
- Boore, J. L. and Brown, W. M.** (2000). Mitochondrial genomes of *Galathealimum*, *Helobdella*, and *Platynereis*: sequence and gene arrangement comparisons indicate that Pogonophora is not a phylum and Annelida and Arthropoda are not sister taxa. *Molecular Biology and Evolution* **17**, 87–106.
- Boore, J. L., Macey, J. R. and Medina, M.** (2005). Sequencing and comparing whole mitochondrial genomes of animals. *Molecular Evolution: Producing the Biochemical Data, Part B* **395**, 311–348.
- Cunningham, C. O.** (1997). Species variation within the internal transcribed spacer (ITS) region of *Gyrodactylus* (Monogenea: Gyrodactylidae) ribosomal RNA genes. *Journal of Parasitology* **83**, 215–219.
- Cunningham, C. O., Collins, C. M., Malmberg, G. and Mo, T. A.** (2003). Analysis of ribosomal RNA intergenic spacer (IGS) sequences in species and populations of *Gyrodactylus* (Platyhelminthes: Monogenea) from salmonid fish in northern Europe. *Diseases of Aquatic Organisms* **57**, 237–246.
- Després, L., Imbert-Establet, D., Combes, C., Bonhomme, F. and Monnerot, M.** (1991). Isolation and polymorphism in mitochondrial DNA from *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* **47**, 139–142.
- Hansen, H., Bachmann, L. and Bakke, T. A.** (2003). Mitochondrial DNA variation of *Gyrodactylus* spp. (Monogenea, Gyrodactylidae) populations infecting Atlantic salmon, grayling, and rainbow trout in Norway and Sweden. *International Journal for Parasitology* **33**, 1471–1478.
- Hansen, H., Martinsen, L., Bakke, T. A. and Bachmann, L.** (2006). The incongruence of nuclear and mitochondrial DNA variation supports conspecificity of the monogenean parasites *Gyrodactylus salaris* and *G. thymalli*. *Parasitology* **133**, 639–650.
- Harris, P. D., Shinn, A. P., Cable, J. and Bakke, T. A.** (2004). Nominal species of the genus *Gyrodactylus* von

- Nordmann 1832 (Monogenea: Gyrodactylidae), with a list of principal host species. *Systematic Parasitology* **59**, 1–27.
- Huyse, T., Audenaert, V. and Volckaert, F. A. M.** (2003). Speciation and host-parasite relationships in the parasite genus *Gyrodactylus* (Monogenea, Platyhelminthes) infecting gobies of the genus *Pomatoschistus* (Gobiidae, Teleostei). *International Journal for Parasitology* **33**, 1679–1689.
- Johnsen, B. O., Møkkelgjerd, P. I. and Jensen, A. J.** (1999). The parasite *Gyrodactylus salaris* on salmon parr in Norwegian rivers, status report at the beginning of year 2000. (In Norwegian, English summary). *NINA Oppdagsmelding* **617**, 1–129.
- Johnston, D. A.** (2006). Genomes and genomics of parasitic flatworms. In *Parasitic Flatworms: Molecular Biology, Biochemistry, Immunology and Physiology* (ed. Maule, A. G. and Marks, N. J.), pp. 37–80. CABI, Wallingford.
- Le, T. H., Blair, D. and McManus, D. P.** (2002). Mitochondrial genomes of parasitic flatworms. *Trends in Parasitology* **18**, 206–213.
- Le, T. H., Humair, P. F., Blair, D., Agatsuma, T., Littlewood, D. T. J. and McManus, D. P.** (2001). Mitochondrial gene content, arrangement and composition compared in African and Asian schistosomes. *Molecular and Biochemical Parasitology* **117**, 61–71.
- Littlewood, D. T. J., Lockyer, A. E., Webster, B. L., Johnston, D. A. and Le, T. H.** (2006). The complete mitochondrial genomes of *Schistosoma haematobium* and *Schistosoma spindale* and the evolutionary history of mitochondrial genome changes among parasitic flatworms. *Molecular Phylogenetics and Evolution* **39**, 452–467.
- Lowe, T. M. and Eddy, S. R.** (1997). tRNAscan-SE: a program for improved transfer RNA detection in genomic sequence. *Nucleic Acids Research* **25**, 955–964.
- Matejusová, I., Gelnar, M., Verneau, O., Cunningham, C. O. and Littlewood, D. T. J.** (2003). Molecular phylogenetic analysis of the genus *Gyrodactylus* (Platyhelminthes: Monogenea) inferred from rDNA ITS region: subgenera versus species groups. *Parasitology* **127**, 603–611.
- Meinila, M., Kuusela, J., Zietara, M. S. and Lumme, J.** (2004). Initial steps of speciation by geographic isolation and host switch in salmonid pathogen *Gyrodactylus salaris* (Monogenea: Gyrodactylidae). *International Journal for Parasitology* **34**, 515–526.
- Mo, T. A.** (2006). Chapter 2.1.14. Gyrodactylosis (*Gyrodactylus salaris*) [online manual]. World Organization for Animal Health. http://www.oie.int/eng/normes/fmanual/A_00031.htm
- Mo, T. A. and Norheim, K.** (2005). The surveillance and control programme for *Gyrodactylus salaris* in Atlantic salmon and rainbow trout in Norway. In *Annual Report 2004*, pp. 137–139. National Veterinary Institute.
- Mo, T. A., Norheim, K. and Hellesnes, I.** (2004). The surveillance and control programme for *Gyrodactylus salaris* in Atlantic salmon and rainbow trout in Norway. (In Norwegian, English summary.) *Norsk Veterinærtidsskrift* **3**, 157–163.
- Olstad, K., Robertsen, G., Bachmann, L. and Bakke, T. A.** (2006). Intraspecific differences in host preference among *Gyrodactylus salaris* (Monogenea) strains: an experimental approach. *Parasitology* (in the Press).
- Peeler, E. J. and Thrush, M. A.** (2004). Qualitative analysis of the risk of introducing *Gyrodactylus salaris* into the United Kingdom. *Diseases of Aquatic Organisms* **62**, 103–113.
- Peeler, E. J., Thrush, M. A., Paisley, L. and Rodgers, C.** (2006). An assessment of the risk of spreading the fish parasite *Gyrodactylus salaris* to uninfected territories in the European Union with the movement of live Atlantic salmon (*Salmo salar*) from coastal waters. *Aquaculture* **258**, 187–197.
- Place, A. R., Feng, X. J., Steven, C. R., Fourcade, H. M. and Boore, J. L.** (2005). Genetic markers in blue crabs (*Callinectes sapidus*) II. Complete mitochondrial genome sequence and characterization of genetic variation. *Journal of Experimental Marine Biology and Ecology* **319**, 15–27.
- Poléo, A. B. S., Schjolden, J., Hansen, H., Bakke, T. A., Mo, T. A., Rosseland, B. O. and Lydersen, E.** (2004). The effect of various metals on *Gyrodactylus salaris* (Platyhelminthes, Monogenea) infections in Atlantic salmon (*Salmo salar*). *Parasitology* **128**, 1–9.
- Robertsen, G., Hansen, H., Bachmann, L. and Bakke, T. A.** (2006). Arctic charr (*Salvelinus alpinus*) is a suitable host for *Gyrodactylus salaris* (Monogenea, Gyrodactylidae) in Norway. *Parasitology* (in the Press).
- Shinn, A. P., Gibson, D. I. and Sommerville, C.** (2001). Morphometric discrimination of *Gyrodactylus salaris* Malmberg (Monogenea) from species of *Gyrodactylus* parasitising British salmonids using novel parameters. *Journal of Fish Diseases* **24**, 83–97.
- Shinn, A. P., Hansen, H., Olstad, K., Bachmann, L. and Bakke, T. A.** (2004). The use of morphometric characters to discriminate specimens of laboratory-reared and wild populations of *Gyrodactylus salaris* and *G. thymalli* (Monogenea). *Folia Parasitologica* **51**, 239–252.
- Soleng, A., Poleo, A. B. S., Alstad, N. E. W. and Bakke, T. A.** (1999). Aqueous aluminium eliminates *Gyrodactylus salaris* (Platyhelminthes, Monogenea) infections in Atlantic salmon. *Parasitology* **119**, 19–25.
- Soleng, A., Poleo, A. B. S. and Bakke, T. A.** (2005). Toxicity of aqueous aluminium to the ectoparasitic monogenean *Gyrodactylus salaris*. *Aquaculture* **250**, 616–620.
- Sterud, E., Mo, T. A., Collins, C. M. and Cunningham, C. O.** (2002). The use of host specificity, pathogenicity, and molecular markers to differentiate between *Gyrodactylus salaris* Malmberg, 1957 and *G. thymalli* Zitnan, 1960 (Monogenea: Gyrodactylidae). *Parasitology* **124**, 203–213.
- Telford, M. J., Herniou, E. A., Russell, R. B. and Littlewood, D. T. J.** (2000). Changes in mitochondrial genetic codes as phylogenetic characters: two examples from the flatworms. *Proceedings of the National Academy of Sciences, USA* **97**, 11359–11364.
- Wolstenholme, D. R.** (1992). Animal mitochondrial DNA: structure and evolution. *International Reviews in Cytology* **141**, 173–216.
- Zietara, M. S., Huyse, T., Lumme, J. and Volckaert, F. A.** (2002). Deep divergence among subgenera of *Gyrodactylus* inferred from rDNA ITS region. *Parasitology* **124**, 39–52.