

PROBLEMS AND SOLUTIONS

PROBLEMS

04.2.1. A Range Equality for Block Matrices with Orthogonal Projectors

Yongge Tian

Queen's University, Kingston, Canada

When considering the general Gauss–Markov model $Y = X\beta + e$, where $E(e) = 0$, $\text{Cov}(e) = V$, one often needs to know whether the range inclusion $\text{range}(X) \subseteq \text{range}(V)$ holds. Show that if V is the following tridiagonal block matrix:

$$V = \begin{bmatrix} P + Q & PQ & & & \\ QP & P + Q & \ddots & & \\ & \ddots & \ddots & PQ & \\ & & QP & P + Q & \end{bmatrix}_{n \times n},$$

where P and Q are two orthogonal projectors of the same size, then

$$\text{range}(V) = \text{range} \begin{bmatrix} P + Q & & & \\ & \ddots & & \\ & & & P + Q \end{bmatrix}_{n \times n}.$$

04.2.2. Characterizations of Hermitian Projectors

Geert Dhaene and Luc Lauwers

Center for Economic Studies, K.U. Leuven, Belgium

Let P in $\mathbb{C}^{n \times n}$ be a projector with P^+ its Moore–Penrose inverse and P^H its conjugate transpose. Werner (2002) provides a list of equivalent conditions for P to be Hermitian: (i) $P = PP^H P$, (ii) $P^+ P^+ = P^+$, (iii) $P^+ = P$, and (iv) $P^+ = P^H$. Extend this list and show that also condition (a) (resp. (b)) is sufficient and necessary for a projector P to be Hermitian:

- (a) the composition $P^H P$ is a projector,
- (b) the composition PP^H is a projector.

REFERENCE

Werner, H.J. (2002) Partial isometry and idempotent matrices. Solution 28-7.5. *IMAGE, The Bulletin of the International Linear Algebra Society* 29, 31–32.

SOLUTION

04.2.1. Fixed Effects Estimation of the Population-Averaged Slopes in a Panel Data Random Coefficient Model—Solution

Jeffrey M. Wooldridge (the poser of the problem)
Michigan State University

a. Just substitute $\mathbf{b}_i = \boldsymbol{\beta} + \mathbf{d}_i$ into (1) and rearrange:

$$y_{it} = a_i + \mathbf{x}_{it}(\boldsymbol{\beta} + \mathbf{d}_i) + u_{it} = a_i + \mathbf{x}_{it}\boldsymbol{\beta} + (\mathbf{x}_{it}\mathbf{d}_i + u_{it})$$

$$\equiv a_i + \mathbf{x}_{it}\boldsymbol{\beta} + v_{it}, \quad t = 1, \dots, T,$$

where $v_{it} \equiv \mathbf{x}_{it}\mathbf{d}_i + u_{it}$.

b. The time-demeaned equation is simply

$$\ddot{y}_{it} = \ddot{\mathbf{x}}_{it}\boldsymbol{\beta} + (\ddot{\mathbf{x}}_{it}\mathbf{d}_i + \ddot{u}_{it}) = \ddot{\mathbf{x}}_{it}\boldsymbol{\beta} + \ddot{v}_{it}, \quad t = 1, \dots, T, \tag{5}$$

where $\ddot{v}_{it} \equiv \ddot{\mathbf{x}}_{it}\mathbf{d}_i + \ddot{u}_{it}, t = 1, \dots, T$.

c. The fixed effects estimator is simply the pooled ordinary least squares estimator applied to the time-demeaned equation (5). So we can write the fixed effects estimator as

$$\hat{\boldsymbol{\beta}} = \left(N^{-1} \sum_{i=1}^N \sum_{t=1}^T \ddot{\mathbf{x}}_{it}'\ddot{\mathbf{x}}_{it} \right)^{-1} \left(N^{-1} \sum_{i=1}^N \sum_{t=1}^T \ddot{\mathbf{x}}_{it}'\ddot{y}_{it} \right)$$

$$= \boldsymbol{\beta} + \left(N^{-1} \sum_{i=1}^N \sum_{t=1}^T \ddot{\mathbf{x}}_{it}'\ddot{\mathbf{x}}_{it} \right)^{-1} \left(N^{-1} \sum_{i=1}^N \sum_{t=1}^T \ddot{\mathbf{x}}_{it}'\ddot{v}_{it} \right)$$

$$= \boldsymbol{\beta} + \left(N^{-1} \sum_{i=1}^N \ddot{\mathbf{X}}_i'\ddot{\mathbf{X}}_i \right)^{-1} \left(N^{-1} \sum_{i=1}^N \sum_{t=1}^T \ddot{\mathbf{x}}_{it}'\ddot{v}_{it} \right).$$

By the rank condition (3) and the weak law of large numbers (WLLN), $(N^{-1} \sum_{i=1}^N \ddot{\mathbf{X}}_i'\ddot{\mathbf{X}}_i)^{-1} \xrightarrow{p} \mathbf{A}^{-1}$, where $\mathbf{A} \equiv E(\ddot{\mathbf{X}}_i'\ddot{\mathbf{X}}_i)$. Consistency follows, again by the WLLN, if

$$E(\ddot{\mathbf{x}}_{it}'\ddot{v}_{it}) = \mathbf{0}, \quad t = 1, \dots, T.$$

Sufficient is $E(\ddot{\mathbf{x}}_{it}'\ddot{\mathbf{x}}_{it}\mathbf{d}_i) = \mathbf{0}$ and $E(\ddot{\mathbf{x}}_{it}'\ddot{u}_{it}) = \mathbf{0}, t = 1, \dots, T$. But $E(\ddot{u}_{it}|\ddot{\mathbf{x}}_{it}) = 0, t = 1, \dots, T$, under (2), which implies $E(\ddot{\mathbf{x}}_{it}'\ddot{u}_{it}) = \mathbf{0}, t = 1, \dots, T$. Further, (4) is equivalent to $E(\mathbf{d}_i|\ddot{\mathbf{x}}_{it}) = \mathbf{0}, t = 1, \dots, T$; the law of iterated expectations then implies that $E(\ddot{\mathbf{x}}_{it}'\ddot{\mathbf{x}}_{it}\mathbf{d}_i) = \mathbf{0}$.

Condition (4) only restricts the mean dependence between \mathbf{b}_i , and the time-demeaned variables, $\ddot{\mathbf{x}}_{it}$; it places no direct restriction on the dependence between \mathbf{b}_i , and $\bar{\mathbf{x}}_i$. Often, it is reasonable to think that although the time averages of some covariates would be correlated with unobserved heterogeneity—in particular, with \mathbf{b}_i —deviations about the time average might not be. So the fixed

effects estimator, although not robust to arbitrary dependence between \mathbf{b}_i and $(\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{iT})$, does have some useful robustness properties in the general random coefficient panel data model.

d. Zero correlation does not suffice because we need \mathbf{b}_i to be uncorrelated with $\ddot{\mathbf{x}}_{it}'\ddot{\mathbf{x}}_{it}$, $t = 1, 2, \dots, T$.