

Research Article

Cite this article: Beets JP, Haug EJ, Sperry BP, Thum RA, and Richardson RJ (2024) Response of four *Vallisneria* taxa to aquatic herbicides. *Invasive Plant Sci. Manag* **17**: 297–304. doi: [10.1017/inp.2024.33](https://doi.org/10.1017/inp.2024.33)






Received: 31 July 2024
Revised: 13 September 2024
Accepted: 16 September 2024
First published online: 29 October 2024

Associate Editor:
Ryan M. Wersal, Minnesota State University

Keywords:
Aquatic; herbicide; invasive species; native species; submersed vegetation

Corresponding author: Jens P. Beets; Email: jens.beets@usda.gov

Response of four *Vallisneria* taxa to aquatic herbicides

Jens P. Beets¹ , Erika J. Haug² , Benjamin P. Sperry³ , Ryan A. Thum⁴  and Robert J. Richardson⁵ 

¹Research Ecologist, U.S. Department of Agriculture, Agricultural Research Service, Invasive Species and Pollinator Health Unit, Davis, CA, USA; ²Environmental Specialist, North Carolina Division of Water Resources, Intensive Surveys Branch, Raleigh, NC, USA; ³Research Biologist, U.S. Army Engineer Research & Development Center, Gainesville, FL, USA; ⁴Professor, Department of Plant Sciences and Plant Pathology, Montana State University, Bozeman, MT, USA and ⁵Professor, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA

Abstract

Native aquatic macrophytes such as American eelgrass (*Vallisneria americana* Michx.) are often desirable in aquatic ecosystems due to the ecological benefits they provide but are threatened by competition from invasive taxa including non-native *Vallisneria* taxa and hydrilla [*Hydrilla verticillata* (L. f.) Royle]. Identifying potential selective herbicide management options can provide tools to minimize impacts to native taxa in restoration and aquatic invasive plant management programs. Greenhouse mesocosm experiments were conducted in 2023 to investigate herbicide efficacy on two native eelgrass species (*V. americana* and *Vallisneria neotropicalis* Vict.), two non-native taxa (*Vallisneria australis* S.W.L. Jacobs & Les and *Vallisneria spiralis* L. × *Vallisneria denseserrulata* Makino), and *H. verticillata*. Herbicide applications included endothall, diquat, florypyrauxifen-benzyl, fluridone, and flumioxazin and select combinations of these herbicides used in *H. verticillata* management. Endothall alone provided 90% to 100% aboveground biomass reduction at 3,000 µg L⁻¹ with at least 24 h of continuous or intermittent exposure to all native and invasive species at 6 wk after exposure, whereas florypyrauxifen-benzyl applied alone resulted in minimal aboveground biomass reduction. A 45-d of exposure of fluridone (10 µg L⁻¹) resulted in 95% biomass reduction of *V. americana* and 7% to 48% of other tested taxa. The combination of flumioxazin and florypyrauxifen-benzyl resulted in 90% to 100% aboveground biomass reduction, and endothall combined with florypyrauxifen-benzyl resulted in 93% to 100% aboveground biomass reduction across taxa. Reductions in belowground biomass mirrored trends observed in aboveground biomass. No treatments selectively controlled invasive *Vallisneria* without injury to native *Vallisneria*, although efficacy was observed on *H. verticillata*. These insights provide an understanding for differences between these *Vallisneria* for researchers moving forward with selectively targeting *H. verticillata* in the presence of native *Vallisneria* species and two new aquatic invasive plants. Future research should expand treatment scenarios, increase the study period, and identify potential integrated plant management strategies for field scenarios.

Introduction

Diverse, native submersed aquatic vegetation (SAV) assemblages are essential for aquatic ecosystems as they provide fish and wildlife habitat and sediment stabilization, reduce wave action, and improve water quality (Dodd et al. 2021; Gettys and Haller 2013; Korschgen et al. 1987; Moore et al. 2010; Owens et al. 2008). However, native aquatic plant communities are sensitive to invasion by non-native plants that reduce biodiversity and habitat quality as well as negatively impact human uses of aquatic systems such as hydropower, flood control, irrigation, and recreation (Langeland 1996; Madsen and Sand-Jensen 1991; Smart et al. 1994; Zhang and Boyle 2010). Aquatic ecosystem restoration efforts often include re-vegetation plantings of native plants (Canfield and Hoyer 1992; Gettys and Haller 2013). In the United States, American eelgrass (*Vallisneria americana* Michx.) is arguably the most common and sought after submersed plant species used in restoration projects due to ease of propagation; resistance to invasive species pressure; tolerance of wave action; growth in a variety of soil and water characteristics; and ecosystem services provided, including stabilizing water and soil quality and serving as a food source or host to food sources for invertebrate and vertebrate species (Gettys and Haller 2013; Henry 2017; Korschgen and Green 1988; Moore et al. 2010).

Vallisneria is a genus with global distribution; however, *V. americana* is native to North America, primarily found in eastern North America, with some distribution in Europe and Asia as well (Korschgen and Green 1988; Les et al. 2008; Martin and Mort 2023; Mesterházy et al. 2021). *Vallisneria* taxa are herbaceous and monocotyledonous and form rosettes at the

Management Implications

Diverse communities of submersed aquatic vegetation are ecologically beneficial to aquatic ecosystems; however, non-native plants commonly disrupt these communities and their associated benefits. *Vallisneria* species are recognized as desirable plants for aquatic resource managers, but recent documentation of multiple non-native *Vallisneria* taxa in the United States has prompted concerns in management of aquatic systems. The primary objective was to evaluate the response of two native eelgrass taxa (*Vallisneria americana* and *Vallisneria neotropicalis*) and two non-native taxa (*Vallisneria australis* and hybrid *Vallisneria spiralis* × *Vallisneria denseserrulata*) to common aquatic herbicide treatments in a small-scale mesocosm study. This research also provided efficacy data for new aquatic invasive taxa. Managers may control native and invasive *Vallisneria* taxa with endothall or combinations of endothall and diquat, endothall and florpyrauxifen-benzyl, and florpyrauxifen-benzyl and flumioxazin. Florpyrauxifen-benzyl applied alone does not provide nominal above- or belowground biomass control compared with some of the other treatments tested. Fluridone reduced native *V. americana* above- and belowground biomass but is less likely to suppress *V. australis*, *V. neotropicalis*, or *V. spiralis* × *V. denseserrulata*. Control of invasive *Hydrilla verticillata* (hydrilla) with minimal damage to native *Vallisneria* was also observed with florpyrauxifen-benzyl treatments. These observations may be of particular interest to managers looking to control *H. verticillata* growing intermingled with native *Vallisneria* taxa, especially in restoration sites.

hydrosol surface (Godfrey and Wooten 1979). They are dioecious, with female flowers in long sessile stalks at the water surface and male flowers on short stalks near the basal rosette that detach and float free to the water surface (Les et al. 2008; Martin and Mort 2023). *Vallisneria* also reproduces asexually via stolons, which produce clonal daughter plants and some species, such as *V. americana*, also reproduce via subterranean turions. Unlike most aquatic macrophytes, which are canopy forming, *Vallisneria* is a meadow-forming plant with biomass more evenly distributed in the water column (Best et al. 2004; Haller and Sutton 1975; Hauxwell et al. 2007).

Recently, genetic divergence between U.S. native populations of *V. americana* was recognized, which suggested the resurrection of *Vallisneria neotropicalis* Vict., previously consolidated with *V. americana* (Gorham et al. 2021; Les et al. 2008; Martin and Mort 2023). These taxa were formerly considered separate ecotypes of *V. americana*, despite considerable phenological differences, including annual versus perennial habits of overwintering and the lack of turion production in southern ecotypes, thought to be a function of geography or environmental factors (Godfrey and Wooten 1979; Korschgen and Green 1988; McFarland and Schafer 2008; Smart et al. 2005). Anecdotal evidence indicates there is a geographic delineation separating *V. americana* and *V. neotropicalis*, but the exact distribution is not currently understood and requires further investigation.

The presence of non-native *Vallisneria* species, cultivars, and hybrids has also been recently documented in the United States, primarily *Vallisneria spiralis* L. and *Vallisneria spiralis* L. × *Vallisneria denseserrulata* Makino in the southeastern United States and *Vallisneria australis* S.W.L. Jacobs & Les in California (CDFA 2021; Gorham et al. 2021; Martin and Mort 2023). Given the prevalence of different *Vallisneria* species in the aquarium and

water garden communities, it is possible some of these new infestations have arisen from accidental releases. *Vallisneria australis* and *V. neotropicalis* have been identified as new invasions in Europe, and the hybrid *V. spiralis* L. × *V. denseserrulata* has been documented in Japan (Mesterházy et al. 2021; Wasekura et al. 2016). Hybrids between *V. australis* and one of the U.S. native species (*V. americana* or *V. neotropicalis*) are commercially available in the aquarium trade mistakenly labeled as *V. americana* (Martin and Mort 2023). Unfortunately, some restoration efforts have unknowingly planted non-native *Vallisneria* taxa, as plants collected from Crystal River, FL, were identified as *V. spiralis* L. × *V. denseserrulata* following a restoration project (Martin and Mort 2023). However, we cannot yet determine the contribution of spread that restoration efforts may have played compared with aquarium-based releases.

Differentiation between cryptic species and hybrids via plant morphology alone is often difficult with closely related taxa, especially *Vallisneria*. Hybridization between macrophytes is likely more common than currently recognized and has been observed in *Myriophyllum* and *Nymphoides* (Harms et al. 2021; Parks et al. 2016; Tavalire et al. 2012). Multiple *Vallisneria* taxa have been documented to coexist in waterbodies, so selective management actions would be desirable to suppress invasive populations while mitigating adverse effects to native populations (Gorham et al. 2021).

To combat invasions by other species and restore SAV communities, aquatic resource managers have been motivated to identify and cultivate *Vallisneria* populations with growth characteristics favorable for restoration efforts. This includes establishing or reestablishing self-sustaining populations that are fast growing; tolerant to chemical and nonchemical management activity; and resistant to invasion, competition, and environmental disturbance. Unfortunately, these characteristics are also common in many weedy species. *Vallisneria* may also slow or prevent invasions by introduced plant species, especially in low-nutrient sediments, where it has been shown to outcompete hydrilla [*Hydrilla verticillata* (L. f.) Royle] in mesocosm trials (Owens et al. 2008; Van et al. 1999). Chadwell and Engelhardt (2008) observed reduced *H. verticillata* establishment via fragmentation alone in field trays with *V. americana*, but in plots where *H. verticillata* subterranean turions were not removed, *V. americana* did not prevent colonization. The competitive pressure posed by exotic macrophytes such as *H. verticillata* and Eurasian watermilfoil (*Myriophyllum spicatum* L.) has bolstered the need for selective management options of ecologically important species such as *Vallisneria* (Beets et al. 2019; Mudge 2013; Netherland et al. 1997).

Effective and selective control of aquatic invasive plants is commonly achieved using registered herbicides and is dependent on a concentration and exposure time (CET) that is plant and herbicide specific. CET requirements have been widely studied primarily in *H. verticillata* and *M. spicatum*. The required exposure times are not always feasible in water bodies where increased water exchange occurs. In these systems, increasing application rate is not always desired, as this can reduce selectivity and increase costs due to required herbicide volumes (Nault et al. 2014; Netherland 2015; Netherland and Getsinger 1995; Netherland et al. 1991, 1997). Netherland (2015) observed equivalent efficacy on monoecious *H. verticillata* between intermittent and continuous exposure to fluridone in mesocosm experiments. These intermittent exposures represent the CET requirements separated by nontreated periods, and these separation periods may need to be adjusted based on mode of action (Darnell 2022). The concept of intermittent exposures builds on the practice of injecting fluridone

over long periods of time. Results have indicated that pausing application during a dilution event would not negatively impact efficacy, and subsequent applications may build on the initial exposure period (Darnell 2022; Netherland 2015).

Vallisneria, presumed to be *V. americana* has shown sensitivity to fluridone at concentrations of 1.5 to 20 $\mu\text{g L}^{-1}$ in mesocosms; however, field studies indicate recovery from initial phytotoxicity often occurs and increased *Vallisneria* frequency can occur (Mudge 2013; Netherland et al. 1997; Smith and Pullman 1997; Valley et al. 2006). Fluridone concentrations below 12 $\mu\text{g L}^{-1}$ with 35 to 60 d of exposure are generally employed to control *H. verticillata* and *M. spicatum* while providing selectivity (Gettys and Leon 2021; Netherland and Getsinger 1995; Netherland et al. 1993). Endothall applications of 0.5 to 3.4 mg L^{-1} with 24 h to 3 wk of exposure have provided control of various *Vallisneria* taxa in mesocosm and field studies (Dugdale et al. 2022; Mudge 2013; Skogerboe and Getsinger 2002). Mudge (2013) reported suppression of suspected *V. americana* with diquat at 100 $\mu\text{g L}^{-1}$ and excellent control with combinations of endothall and flumioxazin at 500 $\mu\text{g L}^{-1}$ + 50 $\mu\text{g L}^{-1}$. Combinations of endothall and diquat have proven effective on *H. verticillata*, with indication of additive effects when the two herbicides are combined (Chiconela and Haller 2013; Pennington et al. 2001; Skogerboe et al. 2004). *Vallisneria* has shown limited sensitivity to florpyrauxifen-benzyl at 3 to 48 $\mu\text{g L}^{-1}$ for 6 h to 15 d, although this may be dependent on plant age and size (Beets et al. 2019; Dodd et al. 2022; Sperry et al. 2021). There is a lack of knowledge concerning efficacy of combinations of florpyrauxifen-benzyl and other aquatic registered herbicides.

Previous studies have largely focused on selectivity for what were thought to be one or two native ecotypes of *V. americana*, but there is a lack information concerning differential herbicide sensitivity in *Vallisneria* taxa (Beets et al. 2019; Mudge 2013; Netherland and Glomski 2014; Skogerboe and Getsinger 2002). Additionally, these studies only differentiated northern and southern *Vallisneria*, or narrowleaf and broadleaf, assuming these plants to be *V. americana*. This is likely no longer a valid assumption in future studies without confirmation through genetic testing. *Hydrilla verticillata* has historically been identified as a major threat to native *Vallisneria* populations and has a large body of research for comparison of efficacious management options. As such, *H. verticillata* was included for comparison of *Vallisneria* control with a well-researched species. The objective of this research was to evaluate the response of *Vallisneria* taxa to a variety of aquatic herbicide CETs used in *H. verticillata* management and determine their utility in management of invasive *Vallisneria* taxa.

Materials and Methods

Mesocosm experiments were conducted and repeated under greenhouse conditions in the summer of 2023 (Trial 1 in June and Trial 2 in July) at North Carolina State University in Raleigh, NC (35.810278°N, 78.721714°W) to evaluate the response of *Vallisneria* taxa to common aquatic herbicide treatments. Plant material for the four *Vallisneria* taxa was sourced from the following sites: *V. americana* from the Connecticut River (41.48333°N, 72.50656°W), *V. neotropicalis* from Lake Gaston, NC (36.49792°N, 77.84371°W), *V. australis* from Lake Mattamuskeet, NC (35.52013°N, 76.1088°W), and *V. spiralis* × *V. denseserrulata* from Wheeler Lake, AL (34.62456°N, 86.98359°W). Monoecious *H. verticillata* was originally sourced from an impoundment in Granville County, NC (36.136993°N, 78.794979°W). Genetic confirmation of *Vallisneria* was performed using ITS sequencing (Les et al. 2008) at Montana State University.

Table 1. Herbicides evaluated in a greenhouse trial examining control of four *Vallisneria* taxa and *Hydrilla verticillata* in Raleigh, NC, in 2023.

Common name	Trade name	Manufacturer
Diquat	Reward®	Syngenta, Greensboro, NC, USA
Endothall dipotassium salt	Aquathol® K	UPL, King of Prussia, PA, USA
Florpyrauxifen-benzyl	Procellacor® SC	SePRO, Carmel, IN, USA
Flumioxazin	Clipper® SC	NuFarm, Alsip, IL, USA
Fluridone	Sonar Genesis®	SePRO, Carmel, IN, USA

Single rosettes of each *Vallisneria* population were planted in 0.52-L pots containing topsoil (Timberline Soil Top Soil, Oldcastle® Lawn & Garden, Atlanta, GA) with slow-release fertilizer (Osmocote® Smart Release 15-9-12, Scotts, Marysville, OH) at a rate of 3 g L^{-1} soil and covered with a sand cap to reduce nutrient leaching into the water column. Water used in this study was conditioned tap water (API Tap Water Conditioner®, Mars Fishcare North America, Chalfont, PA) with a pH of 7.8 to 8.2. A single sprouted apical stem of *H. verticillata* was planted in a 0.09-L cup with the same amended topsoil to serve as a bioindicator for treatment efficacy. Plants were established for four wk before study initiation in 16-L mesocosms.

Following the establishment period, one pot of each taxon was placed in a 16-L mesocosm containing 12 L of conditioned tap water. Each mesocosm was considered an experimental unit and was treated with one of 11 applications containing diquat, endothall, florpyrauxifen-benzyl, flumioxazin, fluridone, or a combination of these herbicides (Table 1). Nontreated controls and a pretreatment biomass harvest control were included in the experiment as references. Following predetermined herbicide exposure periods, plants were placed in new mesocosms containing nontreated water for a 6-wk recovery period. Treatments were applied via syringe using herbicide stock solutions. The experiment was set up as a randomized complete block design with four replications. Mesocosms were maintained under 50% shade to reduce light stress, minimize algal growth, and regulate water temperature. Mean water temperature was 26.7 °C and pH was 8.1 at time of treatment. Mesocosms were maintained at a consistent water level via addition of conditioned tap water.

Six weeks after treatment (WAT), above- and belowground biomass was harvested, placed in a forced-air dryer for 72 h at 65 °C, and weighed. Fluridone treatments were harvested 6 wk after the end of their exposure period due to long exposure requirements of fluridone. Likewise, an additional set of nontreated controls was included for each fluridone harvest time. Biomass data were normalized using their respective nontreated controls to calculate percent biomass reduction using the following equation:

$$\text{Biomass reduction (\%)} = \frac{\text{Mean control biomass} - \text{Sample biomass}}{\text{Mean control biomass}} \times 100 \quad [1]$$

Biomass reduction was subjected to mixed-model ANOVA for each herbicide treatment, with trial and block as random effects and taxa as a fixed effect. Where appropriate, means were separated using a Tukey's honest significant difference (HSD) in JMP Pro 17 (JMP Pro v. 17, SAS Institute, Cary, NC).

Table 2. Aboveground biomass reduction of four *Vallisneria* taxa 6 wk after in-water herbicide application (6 wk after end of exposure period for fluridone).

	Herbicide							
	Diquat	Endothall	Endothall	Endothall ^a	FB ^b	FB ^{b,c}	Fluridone	Fluridone ^d
Concentration ($\mu\text{g L}^{-1}$)	370	2,000	3,000	3,000	30	30	10	10
Exposure time (h)	12	12	24	24	72	72	45 d	45 d
	% Biomass reduction ^e							
Taxa								
<i>V. americana</i>	84 a	99 a	100 a	99 a	-44 b	-40 b	95 a	64 a
<i>V. australis</i>	59 a	70 bc	94 b	98 a	20 a	20 a	7 b	21 b
<i>V. neotropicalis</i>	11 b	82 c	99 a	100 a	-1 ab	6 ab	6 b	36 ab
<i>V. spiralis</i> × <i>V. denseserrulata</i>	63 a	97 ab	100 a	100 a	23 a	38 a	-48 c	60 a

^aIntermittent exposure (8 h + 40-h rest [3×]).

^bFlorpyrauxifen-benzyl.

^cIntermittent exposure (24 h + 6-d rest [3×]).

^dIntermittent exposure (15 d + 6-d rest [3×]).

^eMean responses within a column followed by the same letter do not differ according to Fisher's protected LSD ($P \leq 0.05$). Negative values indicate plant growth.

Table 3. Belowground biomass reduction of four *Vallisneria* taxa 6 wk after in-water herbicide application (6 wk after end of exposure period for fluridone).

	Herbicide							
	Diquat	Endothall	Endothall	Endothall ^a	FB ^b	FB ^{b,c}	Fluridone	Fluridone ^d
Concentration ($\mu\text{g L}^{-1}$)	370	2,000	3,000	3,000	30	30	10	10
Exposure time (h)	12	12	24	24 ^b	72	72	45 d	45 d
	% Biomass reduction ^e							
Taxa								
<i>V. americana</i>	64 a	91 a	95 a	96 a	24 a	17 a	83 a	76 a
<i>V. australis</i>	-1 a	25 b	86 a	95 a	-11 a	-14 a	3 b	10 c
<i>V. neotropicalis</i>	6 a	86 a	99 a	99 a	-16 a	-1 a	15 b	35 bc
<i>V. spiralis</i> × <i>V. denseserrulata</i>	43 a	72 a	91 a	91 a	33 a	24 a	-74 c	41 b

^aIntermittent exposure (8 h + 40-h rest [3×]).

^bFlorpyrauxifen-benzyl.

^cIntermittent exposure (24 h + 6-d rest [3×]).

^dIntermittent exposure (15 d + 6-d rest [3×]).

^eMean responses within a column followed by the same letter do not differ according to Fisher's protected LSD ($P \leq 0.05$). Negative values indicate plant growth.

Results and Discussion

Trial and block factors were not significant when included in separate mixed-model ANOVAs for each herbicide, so results between trials were pooled. *Vallisneria americana* was highly sensitive to most herbicide treatments, resulting in increased control compared with the other *Vallisneria* taxa tested. Diquat alone for 12 h at $370 \mu\text{g L}^{-1}$ resulted in 84% biomass reduction of *V. americana* at 6 WAT (Table 2). Biomass reduction from this treatment was similar in *V. australis* and *V. spiralis* × *V. denseserrulata* (59% and 63% respectively) but was significantly lower in *V. neotropicalis* (11%). Visual injury was observed at 1 to 2 WAT in sensitive taxa. Reductions in belowground biomass did not significantly differ between the four tested taxa when exposed to diquat alone despite a wide range of biomass reductions (Table 3).

Endothall alone for 12 h at $2,000 \mu\text{g L}^{-1}$ resulted in 99% aboveground biomass reduction of *Vallisneria americana* and 97% reduction of *V. spiralis* × *V. denseserrulata* biomass at 6 WAT (Table 2). Aboveground biomass reduction was significantly lower for *V. australis* (70%) and *V. neotropicalis* (82%) than for the other two taxa. Visual injury was observed in all taxa within 1 WAT. Belowground biomass reduction was significantly lower in *V. australis* (25%) than the other tested taxa (Table 3). One *V. americana* replicate in this treatment produced a winter bud/turion despite a substantial amount of injury from this contact herbicide. Endothall alone ($3,000 \mu\text{g L}^{-1}$) under a 24-h continuous exposure resulted in 99% to 100%

aboveground biomass reduction of *V. americana*, *V. spiralis* × *V. denseserrulata*, and *V. neotropicalis*, but biomass reduction was slightly lower in *V. australis* (94%). A similar trend was observed in belowground biomass, but there was no significant difference in the reduction among the tested taxa. When the 24-h exposure of endothall was split into three 8-h exposures and a 40-h rest between applications, there was no significant difference in above- or belowground biomass reduction between the *Vallisneria* taxa, and biomass reduction was comparable to continuous endothall exposure at the same CET.

Florpyrauxifen-benzyl alone for 72 h at $30 \mu\text{g L}^{-1}$ resulted in 20% and 23% aboveground biomass reduction for *V. australis* and *V. spiralis* × *V. denseserrulata*, respectively, while growth was observed in *V. americana* (-44%) and no impact to *V. neotropicalis* was observed at 6 WAT (Table 2). This CET resulted in minimal (-11% to 33%) impact to belowground biomass, and no significant differences between taxa were observed (Table 3). Similar trends were observed when the 72-h exposure period was broken up into three 24-h exposure periods. Aboveground biomass of *V. australis* and *V. spiralis* × *V. denseserrulata* was reduced 20% and 38%, respectively, with growth (-40%) of *V. americana* and no effect on *V. neotropicalis*. Minimal impacts on belowground biomass (-14% to 24%) were observed in the intermittent applications of florpyrauxifen-benzyl. No injury to *Vallisneria* was observed when florpyrauxifen-benzyl was applied alone. The observed growth in aboveground biomass may be a result

of hormesis, or the augmented growth due to sublethal auxin-mimic exposure, which has been observed in Canadian waterweed (*Elodea canadensis* Michx.) and Brazilian waterweed (*Egeria densa* Planch.) when exposed to florpyrauxifen-benzyl (Cedergreen et al. 2007; Howell et al. 2022; Mudge et al. 2021).

Fluridone applied at 10 µg L⁻¹ with a 45-d exposure reduced aboveground biomass of *Vallisneria americana* by 95%, *V. australis* by 7%, and *V. neotropicalis* by 6% (Table 2). Growth and a lack of visual injury were observed in *V. spiralis* × *V. denseserrulata* with this treatment (-48% biomass reduction). This trend was also observed for belowground biomass of all taxa (Table 3). Interestingly, when fluridone was split into three 15-d applications with a 6-d rest period, aboveground biomass reduction on *V. spiralis* × *V. denseserrulata* was 60%, which was similar to the aboveground biomass reduction on *V. americana* and significantly higher than the efficacy on *V. australis*. Intermittent applications of fluridone resulted in a 10% reduction in *V. australis* belowground biomass and 35% for *V. neotropicalis* belowground biomass. Chlorosis, a common symptom of fluridone applications (Netherland and Getsinger 1995; Netherland et al. 1997; Sprecher et al. 1998) was observed in *V. australis* leaves treated with fluridone; however, this symptom did not persist, which is reflected in the lack of biomass reduction at 6 wk after the end of the exposure period.

A 12-h exposure of endothall (2,000 µg L⁻¹) plus diquat (370 µg L⁻¹) reduced *V. americana* aboveground biomass by 98% and *V. spiralis* × *V. denseserrulata* by 100% at 6 WAT (Table 4). Aboveground biomass reductions from this herbicide combination were lower in *V. australis* (11%) and *V. neotropicalis* (62%). A similar trend was observed in belowground biomass, with reductions in belowground biomass of 41% and 100% for *V. australis* and *V. americana*, respectively (Table 5). The combination of flumioxazin at 300 µg L⁻¹ and florpyrauxifen-benzyl at 30 µg L⁻¹ for a 48-h exposure resulted in 97%, 98%, and 99% aboveground biomass reduction of *V. spiralis* × *V. denseserrulata*, *V. australis*, and *V. americana*, respectively. Efficacy of this combination treatment against aboveground biomass remained high for *V. neotropicalis* (91%). There were no significant differences in belowground biomass reduction between taxa when treated with this herbicide combination (79% to 99%). Combinations of 2,000 µg L⁻¹ endothall and 30 µg L⁻¹ florpyrauxifen-benzyl were highly efficacious against all *Vallisneria*, with 100% aboveground biomass reduction of *V. americana* and *V. spiralis* × *V. denseserrulata* and 99.9% in *V. neotropicalis*. Aboveground biomass reduction was significantly lower in *V. australis* (94%), but still considered efficacious. Similar to the previous herbicide combination, there were no differences in belowground biomass reduction across plant species (77% to 100%).

Hydrilla verticillata was included as a bioindicator in each treatment, as these treatments were selected from mesocosm trials targeting monoecious and dioecious *H. verticillata* (Darnell 2022; JPB, unpublished data). Rapid symptomology was observed in *H. verticillata* treated with all herbicides except fluridone and the 12-h exposure of endothall. At 6 WAT, *H. verticillata* aboveground biomass was reduced 80% to 100% except for the 12-h exposure of endothall (Table 6). This treatment only resulted in a 7% reduction in aboveground biomass. While an 83% reduction in belowground biomass was observed, this was likely due to several replicates in this treatment detaching from the sediment and persisting as fragments. Reductions in belowground biomass did not significantly differ between treatments, with several treatments resulting in 100% belowground biomass reduction.

Table 4. Aboveground biomass reduction of four *Vallisneria* taxa 6 wk after in-water herbicide applications.

	Herbicide ^a		
	Endothall + diquat	Flumioxazin + FB ^b	Endothall + FB ^b
Concentration (µg L ⁻¹)	2,000 + 370	300 + 30	2,000 + 30
Exposure time (h)	12	48	48
	----- % Biomass reduction ^c -----		
Taxa			
<i>V. americana</i>	98 a	99 a	100 a
<i>V. australis</i>	11 b	98 a	94 b
<i>V. neotropicalis</i>	62 b	91 b	99 a
<i>V. spiralis</i> × <i>V. denseserrulata</i>	100 a	97 a	100 a

^aHerbicides were applied simultaneously at indicated rates.

^bFlorpyrauxifen-benzyl.

^cMean responses within a column followed by the same letter do not differ according to Fisher's protected LSD (P ≤ 0.05).

Table 5. Belowground biomass reduction of four *Vallisneria* taxa 6 wk after in-water herbicide applications.

	Herbicide ^a		
	Endothall + diquat	Flumioxazin + FB ^b	Endothall + FB ^b
Concentration (µg L ⁻¹)	2,000 + 370	300 + 30	2,000 + 30
Exposure time (h)	12	48	48
	----- % Biomass reduction ^c -----		
Taxa			
<i>V. americana</i>	100 a	99 a	100 a
<i>V. australis</i>	41 c	95 a	83 a
<i>V. neotropicalis</i>	58 bc	80 a	77 a
<i>V. spiralis</i> × <i>V. denseserrulata</i>	85 ab	79 a	90 a

^aHerbicides were applied simultaneously at indicated rates.

^bFlorpyrauxifen-benzyl.

^cMean responses within a column followed by the same letter do not differ according to Fisher's protected LSD (P ≤ 0.05).

These results are promising for long-term *H. verticillata* control, as reductions in belowground biomass can be indicative of low regrowth potential via root crown or subterranean turions (Haller et al. 1976; Nawrocki et al. 2016; Netherland et al. 1997; Steward 1980). The high level of efficacy demonstrated by the intermittent applications of endothall, florpyrauxifen-benzyl, and fluridone also further validates preliminary studies for intermittent applications. Monoecious *H. verticillata* overwinters as subterranean turions, so early-season management is often desirable before biomass has peaked and has proven highly efficacious, and small plants such as those utilized in this study can be representative of plants at this growth stage in the field (Langeland and Pesacreta 1986; Nawrocki et al. 2016). While not the main objective of the study, we find it important to highlight that *H. verticillata* bioindicator plants were well managed utilizing florpyrauxifen-benzyl treatments, with biomass reductions of 96% while native *V. americana* remained relatively unharmed (Table 6). Fluridone applications provided similar selectivity between *H. verticillata* and *V. neotropicalis* (Tables 2 and 3). *Hydrilla verticillata* can outcompete native species such as *V. americana* and *V. neotropicalis*. Therefore, identification of selective management

Table 6. Biomass reduction of *Hydrilla verticillata* 6 wk after in-water herbicide application (6 wk after end of exposure period for fluridone).

Herbicide	Rate μg L ⁻¹	Exposure time	% Biomass reduction ^a	
			Aboveground	Belowground
Diquat	370	12 h	96 a	100 a
Endothall	2,000	12 h	7 b	83 a
Endothall	3,000	24 h	91 a	86 a
Endothall	3,000	24 h	99 a	100 a
Florpyrauxifen-benzyl	30	72 h	96 a	100 a
Florpyrauxifen-benzyl	30	72 h	97 a	100 a
Fluridone	10	45 d	80 a	75 a
Fluridone	10	45 d	99 a	100 a
Endothall + diquat	2,000 + 370	48 h	98 a	100 a
Flumioxazin + Florpyrauxifen-benzyl	300 + 30	48 h	99 a	100 a
Endothall + florpyrauxifen-benzyl	3,000 + 30	48 h	100 a	100 a

^aMean responses within a column followed by the same letter do not differ according to Fisher's protected LSD ($P \leq 0.05$).

options is important, even if they are not effective against the exotic *Vallisneria* taxa (Beets et al. 2019; Chadwell and Engelhardt 2008; Haller and Sutton 1975).

There is a currently a lack of knowledge about management strategies for the newly documented invasive hybrid species *V. spiralis* × *V. denseserrulata*, and the populations of *Vallisneria australis* that have been identified in the United States. The results of this research can provide an initial understanding of potentially efficacious management actions. Additionally, comparisons with historical datasets are complicated and may not be accurate, because previous research may not have accurately identified the *Vallisneria* species being studied without genetic confirmation of the tested species (Gorham et al. 2021; Martin and Mort 2023).

Native *Vallisneria* of various previously classified ecotypes have shown sensitivity to endothall at concentrations of 0.5 mg L⁻¹ and higher, thus validating the findings of this research (Mudge 2013; Skogerboe and Getsinger 2001, 2002). These previous studies also indicated recovery from viable root crowns and turions, so treatment timing may play an important role in recovery, and further investigation into direct effects on *Vallisneria* turions is warranted to improve strategies for effective selective management of these native species. Similarly, Turnage and Madsen (2015) observed expansion of *Vallisneria* in Minnesota lakes following diquat applications, and Mudge (2013) observed minimal biomass reduction in *Vallisneria* exposed to diquat.

The observation of minimal impact on native *Vallisneria* with florpyrauxifen-benzyl is corroborated by previous mesocosm and field studies, indicating that florpyrauxifen-benzyl alone is not an effective control method of *Vallisneria* but does provide a selective management option for *H. verticillata* when *Vallisneria* is a non-target species (Beets et al. 2019; Dodd et al. 2022; Sperry et al. 2021). *Vallisneria* sensitivity to fluridone has been variable in previous mesocosm and field studies, which may be partially explained by differentiation between *V. americana* and *V. neotropicalis* or other *Vallisneria* taxa, as seen in this study (Getsinger et al. 2001, 2002; Nelson et al. 1998; Netherland et al. 1997; Poovey et al. 2004). Several divergences in efficacy compared with previous studies were observed. Dugdale et al. (2012) observed a 90% reduction in *V. australis* treated with diquat, while diquat efficacy was minimal in this study (Tables 2 and 3). Previous studies have indicated low efficacy of flumioxazin when applied alone and in combination with endothall or diquat (Mudge 2013; Mudge et al. 2010). However, combinations with flumioxazin have not been fully investigated at higher doses, especially in combination with florpyrauxifen-benzyl.

With the resurrection of the distinction between *V. americana* and *V. neotropicalis*, future studies must confirm which species are being utilized, and new evaluations are warranted to properly correlate selectivity with the species present. This study presents new evidence that previous differences in efficacy between *V. americana* populations may in fact have been differences in response between these two species, especially when comparing northern and southern populations in the United States. While the observed differences in herbicide response are indicative of variation between species, it is possible that the observed differences are limited to the tested populations. Further investigation and replication are needed to see whether there is a noticeable within-taxa variation to herbicide response in *Vallisneria*.

Previous field and mesocosm studies have indicated that *Vallisneria australis* is sensitive to but not completely controlled by similar concentrations and exposure times that were effective in reducing above- and belowground biomass in this study (Clements et al. 2018; Dugdale et al. 2019, 2022). Dugdale et al. (2019) also observed improved endothall efficacy on *Vallisneria australis* in flowing mesocosms compared with quiescent mesocosms. There is a general lack of information concerning the efficacy of other herbicides on *V. australis* to further corroborate our findings. However, the findings of this research with *V. australis*, especially those surrounding florpyrauxifen-benzyl, flumioxazin, and fluridone, are similar to the results observed with other species. Further investigation into the effects of water flow on herbicide efficacy can greatly improve management strategies, especially in lotic systems such as the San Joaquin River delta in California, where invasions of *V. australis* have been identified (CDFA 2021).

Female flowers were observed in all *Vallisneria* taxa included in this study. However, no male flowers were observed in the hybrid *V. spiralis* × *V. denseserrulata* or *V. australis*. Male flowers have not been observed in the hybrid *Vallisneria* in other studies, indicating propagation may primarily be occurring via asexual reproduction (Gorham et al. 2021; Wasekura et al. 2016). Considerable genetic plasticity has been observed in *V. australis*, and its documented hybridization with native *Vallisneria* indicates these female flowers are not sterile or male plants are also present in the United States (Martin and Mort 2023). The possibility of hybridization between these invasive and native taxa should be investigated further, and additional field monitoring is needed to confirm the lack of male plants in the United States, considering the issues that have arisen with hybridization between native and invasive species such as *Myriophyllum spicatum* L. × *Myriophyllum sibiricum* Kom. and

Nymphoides cristata (Roxb.) Kuntze × *Nymphoides aquatica* (J.F. Gmel.) Kuntze (Harms et al. 2021; Parks et al. 2016).

In conclusion, these small-scale mesocosm experiments provide an initial understanding of herbicide response in two new invasive aquatic species, as well as two native species that were previously understood to be the same species. No herbicide treatments tested definitively provided selectivity between invasive and native *Vallisneria* taxa. However, *H. verticillata* was selectively managed while minimal damage to native *Vallisneria* was observed for several well-established treatments, and these results further document the efficacy of intermittent exposures. Future research should scale up the size and study length in experimental design (i.e., larger mesocosms or small field plots and evaluation periods longer than 6 wk) to provide a more operational understanding of selective management practices. Given the observed resilience of *V. australis* and the hybrid *V. spiralis* × *V. denseserrulata*, rapid endeavors should be made to improve understanding of efficacious management options as well as phenological and growth patterns to improve timing of management efforts. Turion production was not consistent throughout the trial for statistical analysis and was only observed in *V. americana*, thus future studies are needed to determine what factors lead to turion formation, including temperature, season, and herbicide application.

There are several key concepts future research endeavors should seek to investigate to improve management of a spreading problematic invasive plant. (1) Identify potential hybridization of non-native *Vallisneria* taxa with native *Vallisneria* taxa, as well as other *Vallisneria* taxa that may be present in the United States, given the genus's prevalence in the aquarium and water garden trade. (2) Implement improvements in genetic confirmation of *Vallisneria* populations before treatment to inform herbicide treatment decisions and improve potential selectivity, as plants in this genus demonstrate similar morphology. (3) Improve the understanding of the ecological and anthropological impacts these non-native *Vallisneria* taxa have on aquatic ecosystems. (4) Identify phenological and growth differences between *Vallisneria* taxa to better inform management actions and the potential spread of these plants. (5) Identify new herbicide use patterns and potential new herbicide chemistries and use integrated pest management strategies to provide selective management options for *Vallisneria* taxa.

Acknowledgments. The authors would like to thank Kara Foley, Andrew Howell, Michael Punt, Delaney Davenport, and Logan Wilson for their assistance with plant propagation, treatment, and harvesting. The manuscript was reviewed in accordance with U.S. Army Engineer Research and Development Center policy and approved for publication. Citation of trade names does not constitute endorsement or approval of the use of such commercial products. The content of this work does not necessarily reflect the position or policy of the U.S. government and no official endorsement should be inferred. The U.S. Department of Agriculture is an equal opportunity employer and provider.

Funding statement. Funding for this research was provided by the U.S. Army Corps of Engineers Environmental Research and Development Center Aquatic Plant Control Research Program and the Michael D. Netherland Graduate Student Scholarship through the Aquatic Plant Management Society.

Competing interests. The authors declare no conflicts of interest.

References

- Beets J, Heilman M, Netherland MD (2019) Large-scale mesocosm evaluation of floryprauxifen-benzyl, a novel arylpicolinate herbicide, on Eurasian and hybrid watermilfoil and seven native submersed plants. *J Aquat Plant Manag* 57:49–55
- Best EPH, Kiker GA, Boyd WA (2004) A Simulation Model on the Competition for Light of Meadow-forming and Canopy-forming Aquatic Macrophytes at High and Low Nutrient Availability. Technical Report TR-04-14. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 79 p
- [CDFA] California Department of Food & Agriculture (2021) California Pest Rating Proposal for *Vallisneria australis* S.W.L. Jacobs and Les: Australian Eelgrass, Ribbonweed. <https://blogs.cdfa.ca.gov/Section3162/wp-content/uploads/2021/10/Vallisneria-australis-Australian-eelgrass.pdf>. Accessed: October 30, 2023
- Canfield DE, Hoyer MV (1992) Aquatic Macrophytes and Their Relation to the Limnology of Florida Lakes. Final Report. Tallahassee: Bureau of Aquatic Plant Management, Florida Department of Natural Resources. 176 p
- Cedergreen N, Streibig JC, Kudsk P, Mathiassen SK, Duke SO (2007) The occurrence of hormesis in plants and algae. *Dose Response* 5:150–162
- Chadwell TB, Engelhardt KAM (2008) Effects of pre-existing submersed vegetation and propagule pressure on the invasion success of *Hydrilla verticillata*. *J Appl Ecol* 45:515–523
- Chiconela TF, Haller WT (2013) Herbicide combinations for the enhancement of diquat phytotoxicity for hydrilla control. *ARNP J Agric Biol Sci* 8:555–562
- Clements D, Butler KL, Hunt TD, Liu Z, Dugdale TM (2018) Efficacy of endothall dimethyl alkylamine salt applied to static irrigation channels during winter to control aquatic weeds in temperate Australia. *J Aquat Plant Manag* 56:84–92
- Darnell TL (2022) Reproductive Biology and Management of Dioecious *Hydrilla verticillata* (L.f.) Royle. Master's thesis. [Gainesville]: University of Florida. 132 p
- Dodd LL, Harms NE, Schad AN (2021) Reciprocal competitive effects of congeneric invaders, *Trapa natans* L. and *Trapa bispinosa* Roxb. var. *iinumai* Nakano, in established freshwater plant cultures. *Aquat Bot* 174:103419
- Dodd LL, Mudge CR, Schad AN (2022) Comparative trials of herbicides for control of *Trapa natans* and *T. bispinosa* var. *iinumai* in the presence of *Heteranthera dubia* and *Vallisneria americana*. *J Aquat Plant Manag* 60:66–74
- Dugdale TM, Hunt TD, Clements D, Butler K (2012) Potential new herbicides for submersed aquatic weeds in Victoria. Pages 41–44 in *Proceedings of the 18th Australasian Weeds Conference*. Palmerston North, New Zealand: Council of Australasian Weed Societies
- Dugdale TM, Islam MS, Hunt TD, Liu Z, Butler KL, Clements D (2022) Operational-scale validation of a winter-use pattern for endothall to control submersed aquatic weeds in ponded Australian irrigation canals. *J Aquat Plant Manag* 60:39–47
- Dugdale TM, Islam MS, Hunt TD, Liu Z, Butler KL, Clements D, Netherland MD (2019) Hydrodynamic exposure and time since application influence endothall amine potency against submersed aquatic plants. *Aquat Bot* 155:18–24
- Getsinger KD, Madsen JD, Koschnick TJ, Netherland MD, Stewart RM, Honnell DR, Staddon AG, Owens CS (2001) Whole Lake Applications of Sonar for Selective Control of Eurasian Watermilfoil. ERDC/EL TR-01-7. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 59 p
- Getsinger KD, Poovey AG, James WF, Stewart RM, Grodowitz MJ, Maceina MJ, Newman RM (2002) Management of Eurasian Watermilfoil in Houghton Lake, Michigan: Workshop Summary. ERDC/EL TR02-24. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 88 p
- Gettys LA, Haller WT (2013) Effect of ecotype, sediment composition, and fertility level on productivity of eight Florida ecotypes of American eelgrass (*Vallisneria americana*). *J Aquat Plant Manag* 51:127–131
- Gettys LA, Leon RG (2021) A population genetics approach for the study of fluridone resistance in hydrilla. *Aquat Invasions* 16:28–42
- Godfrey RK, Wooten JW (1979) *Aquatic and Wetland Plants of Southeastern United States: Monocotyledons*. Athens, GA: University of Georgia Press. 712 p
- Gorham SB, Seyoum S, Furman BT, Darnell KM, Reynolds LK, Tringali MD (2021) Molecular detection of a non-native hybrid eelgrass, *Vallisneria spiralis* Linnaeus (1753) × *V. denseserrulata* Makino (1921), in the southeastern United States. *Aquat Bot* 175:103445

- Haller WT, Miller JL, Garrad LA (1976) Seasonal production and germination of hydrilla vegetative propagules. *J Aquat Plant Manag* 14:26–29
- Haller WT, Sutton DL (1975) Community structure and competition between hydrilla and vallisneria. *Hyacinth Control J* 13:48–50
- Harms NE, Thum RA, Gettys LA, Markovich IJ, French A, Simantel L, Richardson R (2021) Hybridization between native and invasive *Nymphoides* species in the United States. *Biol Invasions* 10:3003–3011
- Hauxwell J, Frazer TK, Osenberg CW (2007) An annual cycle of biomass and productivity of *Vallisneria americana* in a subtropical spring-fed estuary. *Aquat Bot* 87:61–68
- Henry AL (2017) Monoecious Hydrilla (*Hydrilla verticillata* (L.f) Royle) Growth and Phenology in Two Dissimilar Climates. MS thesis. Raleigh, NC: North Carolina State University. 127 p
- Howell AW, Hofstra DE, Heilman MA, Richardson RJ (2022) Susceptibility of native and invasive submersed plants in New Zealand to florypyrauxifen-benzyl in growth chamber exposure studies. *Invasive Plant Sci Manag* 15:133–140
- Korschgen, CE, George LS, Green WL (1987) Feeding ecology of canvasback staging on Pool 7 of the Upper Mississippi River. Pages 237–249 in Weller ME, ed. *Waterfowl in Winter*. Minneapolis: University of Minnesota Press
- Korschgen CE, Green WL (1988) American wildcelery (*Vallisneria americana*): ecological considerations for restoration. U.S. Fish and Wildlife Service, Fish and Wildlife Technical Report 19. Northern Prairie Wildlife Research Center Online, Jamestown, ND. 24 p
- Langeland KA (1996) *Hydrilla verticillata* (LF) Royle (Hydrocharitaceae), “the perfect aquatic weed.” *Castanea* 61:293–304
- Langeland KA, Pesacrete GJ (1986) Management Program for Hydrilla (a Monoecious Strain) in North Carolina. UNC-WRRI-86-225. Raleigh: Water Resources Research Institute of the University of North Carolina. 33 p
- Les DH, Jacobs SW, Tippery NP, Chen L, Moody ML, Wilstermann HM (2008) Systematics of *Vallisneria* (Hydrocharitaceae). *Syst Bot* 33:49–65
- Madsen JD, Sand-Jensen KAJ (1991) Photosynthetic carbon assimilation in aquatic macrophytes. *Aquat Bot* 41:5–40
- Martin AP, Mort ME (2023) *Vallisneria* (Hydrocharitaceae): novel species, taxonomic revisions, and hybridization. *Aquat Bot* 188:103669
- McFarland DG, Schafer DJ (2008) Factors influencing reproduction in American wild celery: a synthesis. *J Aquat Plant Manag* 46:129–144
- Mesterházy A, Somogyi G, Efremov A, Verloove F (2021) Assessing the genuine identity of alien *Vallisneria* (Hydrocharitaceae) species in Europe. *Aquat Bot* 174:103431
- Moore KA, Shields EC, Jarvis JC (2010) The role of habitat and herbivory on the restoration of tidal freshwater submersed aquatic vegetation populations. *Restor Ecol* 18:596–604
- Mudge CR (2013) Impact of aquatic herbicide combinations on nontarget submersed plants. *J Aquat Plant Manag* 51:39–44
- Mudge CR, Haller WT, Netherland MD, Kowalsky JK (2010) Evaluating the influence of pH-dependent hydrolysis on the efficacy of flumioxazin for hydrilla control. *J Aquat Plant Manag* 48:25–30
- Mudge CR, Sartain, BT, Sperry BP, Getsinger KD (2021) Efficacy of Florpyrauxifen-Benzyl for Eurasian Watermilfoil Control and Nontarget Illinois Pondweed, Elodea, and Coontail Response. ERDC/TN APCRP CC-24. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 7 p
- Nault ME, Netherland MD, Mikulyuk A, Skogerboe JG, Asplund T, Hauxwell J, Toshner P (2014) Efficacy, selectivity, and herbicide concentrations following a whole-lake 2,4-D application targeting Eurasian watermilfoil in two adjacent northern Wisconsin lakes. *Lake Reserv Manag* 30:1–10
- Nawrocki JJ, Richardson RJ, Hoyle ST (2016) Monoecious hydrilla tuber dynamics following various management regimes on four North Carolina reservoirs. *J Aquat Plant Manag* 54:12–19
- Nelson LS, Shearer JF, Netherland MD (1998) Mesocosm evaluation of integrated fluridone-fungal pathogen treatment on four submersed plants. *J Aquat Plant Manag* 36:73–77
- Netherland MD (2015) Laboratory and greenhouse response of monoecious hydrilla to fluridone. *J Aquat Plant Manag* 53:178–184
- Netherland MD, Getsinger KF (1995) Potential control of hydrilla and Eurasian watermilfoil under various fluridone half-life scenarios. *J Aquat Plant Manag* 33:36–42
- Netherland MD, Getsinger KF, Skogerboe JD (1997) Mesocosm evaluation of the species-selective potential of fluridone. *J Aquat Plant Manag* 35:41–50
- Netherland MD, Getsinger KF, Turner EG (1993) Fluridone concentration and exposure time requirements for control of Eurasian watermilfoil and hydrilla. *J Aquat Plant Manag* 31:189–194
- Netherland MD, Glomski LM (2014) Mesocosm evaluation of triclopyr on Eurasian watermilfoil and three native submersed species: the role of treatment timing and herbicide exposure. *J Aquat Plant Manag* 52:57–64
- Netherland MD, Green WR, Getsinger KD (1991) Endothall concentration and exposure time relationships for the control of Eurasian watermilfoil and hydrilla. *J Aquat Plant Manag* 29:61–67
- Owens CS, Smart RM, Dick GO (2008) Resistance of *Vallisneria* to invasion from hydrilla fragments. *J Aquat Plant Manag* 46:113–116
- Parks S R, McNair JN, Hausler P, Tynning P, Thum RA (2016) Divergent responses of cryptic invasive watermilfoil to treatment with auxinic herbicides in a large Michigan lake. *Lake Reserv Manag* 32:366–372
- Pennington TG, Skogerboe JG, Getsinger KD (2001) Herbicide/copper combinations for improved control of *Hydrilla verticillata*. *J Aquat Plant Manag* 39:56–58
- Poovey AG, Skogerboe JG, Getsinger KD (2004) Efficacy of AVAST! Fluridone Formulation against Eurasian Watermilfoil and Nontarget Submersed Plants. ERDC/EL TR-04-9. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 29 p
- Skogerboe JG, Getsinger KD (2001) Endothall species selectivity evaluation: southern latitude aquatic plant community. *J Aquat Plant Manag* 39:129–135
- Skogerboe JG, Getsinger KD (2002) Endothall species selectivity evaluation: northern latitude aquatic plant community. *J Aquat Plant Manag* 40:1–5
- Skogerboe JG, Pennington T, Hyde J, Aguillard (2004) Combining Endothall with Other Herbicides for Improved Control of Hydrilla—A Field Demonstration. ERDC/TN APCRP-CC-04. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 7 p
- Smart RM, Barko JW, McFarland DG (1994) Competition between *Hydrilla verticillata* and *Vallisneria americana* under Different Environmental Conditions. APCRP Technical Reports Collection. ERDC/TR A-94-1. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 85 p
- Smart RM, Dick GO, Snow JR (2005) Update to the Propagation and Establishment of Aquatic Plants Handbook. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 44 p
- Smith CS, Pullman GD (1997) Experiences using Sonar® A.S. Aquatic Herbicide in Michigan. *Lake Reserv Manag* 12:338–346
- Sperry BP, Leary JK, Jones KD, Ferrell JA (2021) Observations of a submersed field application of florypyrauxifen-benzyl suppressing hydrilla in a small lake in central Florida. *J Aquat Plant Manag* 59:20–26
- Sprecher SL, Netherland MD, Stewart AB (1998) Phytoene and carotene response of aquatic plants to fluridone under laboratory conditions. *J Aquat Plant Manag* 36:111–120
- Steward KK (1980) Retardation of hydrilla (*Hydrilla verticillata*) regrowth through chemical control of vegetative reproduction. *Weed Sci* 28:245–251
- Tavalire HF, Bugbee GE, LaRue EA, Thum RA (2012) Hybridization, cryptic diversity, and invasiveness in introduced variable-leaf watermilfoil. *Evol Appl* 5:892–900
- Turnage G, Madsen JD (2015) Management of Flowering Rush Using the Contact Herbicide Diquat in Detroit Lakes, Minnesota 2014. Geosystems Research Institute Report 5065. Mississippi State, MS: Geosystems Research Institute, Mississippi State University. 48 p
- Valley RD, Crowell W, Welling CH, Proulx N (2006) Effects of a low-dose fluridone treatment on submersed aquatic vegetation in a eutrophic Minnesota lake dominated by Eurasian watermilfoil and coontail. *J Aquat Plant Manag* 44:19–25
- Van TK, Wheeler GS, Center TD (1999) Competition between *Hydrilla verticillata* and *Vallisneria americana* as influenced by soil fertility. *Aquat Bot* 62:225–233
- Wasekura H, Horie S, Fujii S, Maki M (2016) Molecular identification of alien species of *Vallisneria* (Hydrocharitaceae) species in Japan with a special emphasis on the commercially traded accessions and the discovery of hybrid between nonindigenous *V. spiralis* and native *V. denseserrulata*. *Aquat Bot* 128:1–6
- Zhang C, Boyle KJ (2010) The effect of an aquatic invasive species (Eurasian watermilfoil) on lakefront property values. *Ecol Econ* 70:394–404