

## Commentary to Special Article

**Cite this article:** Purgato M, Jayaram G, Surkan PJ, Bass J, Bolton P (2019). Encompassing a global mental health perspective into psychotherapy research: a critique of approaches to measuring the efficacy of psychotherapy for depression. *Epidemiology and Psychiatric Sciences* **28**, 275–277. <https://doi.org/10.1017/S2045796018000781>

Received: 26 November 2018  
Accepted: 2 December 2018  
First published online: 15 January 2019

### Key words:

Depression; global mental health; meta-analysis; psychotherapy; systematic review

### Author for correspondence:

Marianna Purgato, E-mail: [marianna.purgato@univr.it](mailto:marianna.purgato@univr.it)

# Encompassing a global mental health perspective into psychotherapy research: a critique of approaches to measuring the efficacy of psychotherapy for depression

Marianna Purgato<sup>1,2</sup>, Geetha Jayaram<sup>3,4</sup>, Pamela J. Surkan<sup>5</sup>, Judith Bass<sup>6</sup>  
and Paul Bolton<sup>5,6</sup>

<sup>1</sup>Department of Neuroscience, Biomedicine and Movement Sciences, Section of Psychiatry, WHO Collaborating Centre for Research and Training in Mental Health and Service Evaluation, University of Verona, Verona, Italy; <sup>2</sup>Cochrane Global Mental Health, University of Verona, Verona, Italy; <sup>3</sup>Department of Medicine, Johns Hopkins University, Baltimore, MD, USA; <sup>4</sup>Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA; <sup>5</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA and <sup>6</sup>Department of Mental Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Commentary paper on: Munder T, Flückiger C, Leichsenring F, Abbas A, Hilsenroth M, Luyten P, Rabung S, Steinert C and Wampold B. Is psychotherapy effective? A re-analysis of treatments for depression. *Epidemiol Psychiatr Sci.* 2018 Jul 30:1–7. doi: 10.1017/S2045796018000355

At the beginning of 2018, Cuijpers *et al.* published a paper on the role of bias in influencing the estimation of the effects of psychotherapy for depression (Cuijpers *et al.*, 2018a). Prior to undertaking the analyses presented in that paper, the authors made a series of decisions and established criteria regarding the parameters of their analyses including: (1) excluding studies of waiting list controls and non-Western studies; (2) including only studies with low risk of bias and (3) using a methodology to estimate the number of unpublished studies as a measure of publication bias and the effect on effect size when data from missing studies were imputed. Cuijpers and colleagues found that the mean effect size for psychotherapies for depression was 0.31. This is below previous estimates, which included all the studies comparing psychotherapy with any control group, and corresponds to a small effect according to Cohen (Cohen, 1992; Higgins, 2011). As a follow-up to this paper, and given their interest in meta-analyses for psychological interventions (Flückiger *et al.*, 2018; Munder and Barth, 2018), Munder and colleagues re-analysed Cuijpers' data (Munder *et al.*, 2018). Starting from different theoretical assumptions – mainly related to the choice of waiting list condition as the most appropriate control, the decision to not consider the study's risk of bias, and the inclusion of non-Western studies – this new analysis produced an estimate close to 0.70. This estimate was consistent with the moderate effect size identified in the Cuijpers' analysis that included all the studies.

These papers by Munder and Cuijpers show how widely findings of meta-analyses may differ when performed based on different assumptions, with important implications in terms of results and interpretation. In this commentary we describe some methodological issues when considering the work of Munder and Cuijpers from a global mental health perspective.

First, the choice of control group is particularly relevant to research in low- and middle-income countries. In the Cuijpers' paper, the choice of excluding the waiting list as a control condition was motivated by the issue of limiting participants seeking care for their mental condition elsewhere because they are waiting for the therapy (Mohr *et al.*, 2014; Cuijpers and Cristea, 2016). This is not a concern in many low- and middle-income countries where alternative treatments are not available. One way to mitigate concern over this problem in the future might be to limit data to people with long-lasting conditions. In many low- and middle-income countries, participants suffer from long-lasting and even chronic conditions because they lack the possibility of receiving evidence-based treatments. Although Cuijpers favours the choice of treatment as usual (TAU) as a control condition, he also recognises that TAU may vary according to populations and contexts, to the point that being in the TAU condition sometimes corresponds to not getting treatments at all (Cuijpers *et al.*, 2018b), and differentiating TAU from no treatment or from waiting list control might become difficult. This is true in many low- and middle-income countries where there are no available treatment options outside the study context.

Second, we stress the importance of including studies in which psychotherapy is provided by para-professionals. In many parts of the world there are limited resources to deliver

standard psychotherapeutic interventions defined in Munder's paper (Munder *et al.*, 2018). Evidence is mounting for the effectiveness of psychotherapeutic interventions delivered by para-professionals in low- and middle-income countries without extensive training or experience in mental health, but trained for the delivery of a specific intervention (Morina *et al.*, 2017; Patel *et al.*, 2018; Purgato *et al.*, 2018). While Cuijpers *et al.* included in his analysis several types of psychotherapy, like self-help or web-based interventions, Munder *et al.* applied stricter criteria regarding what constitutes psychotherapy: face-to-face meetings and having special experience or training in the handling of human relationships. For example, Munder criticised the choice of Cuijpers of including a study in which a short cognitive behavioural therapy (CBT)-based intervention was delivered by nurses with no specific mental health expertise except a 4-day training on how to detect depression and how to deliver techniques based upon self-management and CBT (i.e. reattribution of negative cognitions and problem solving) (Lamers *et al.*, 2010). While including interventions provided by para-professionals may increase heterogeneity of the interventions (Fletcher, 2007; Purgato and Adams, 2012), however it may allow for inclusion of more studies conducted in countries with limited financial and human resources (Singla *et al.*, 2017, 2018; Cuijpers *et al.*, 2018b; Purgato *et al.*, 2018). If heterogeneity is a concern, an option for including these studies in systematic reviews could consist of conducting separate meta-analyses within the same review, or planning subgroup analyses, for example accounting for the type of region in which studies were conducted, the type of interventionist, or the presence/type of humanitarian crisis. Exclusion of these studies would result in missing an important group of rigorously tested treatments that actually share many characteristics with standard psychotherapies (Barbui *et al.*, 2017).

Moreover, a similar reasoning on heterogeneity applies to other characteristics of psychotherapeutic interventions, as these interventions are complex and many variables contribute to increased heterogeneity. For example pooling interventions with different number of sessions (i.e. one session *v.* 60 sessions), or targeting different populations (i.e. young *v.* elderly) in the same meta-analysis is problematic. At present, psychotherapy literature has been mainly focused on the comparison between (different types of) psychotherapies together against 'inactive' controls as on waiting lists, no treatment, or TAU. However, the papers by Cuijpers and Munder highlighted that it is difficult to draw firm clinical conclusions from meta-analyses. First because each 'inactive' control presents specific characteristics that may contribute to influence the effect size. Differences across control groups have been detected also in a subgroup analysis accounting for the type of control group (waiting list, no treatment, ETAU) on the outcome depressive symptoms at endpoint ( $p = 0.02$ ) and at 1–4 month follow-up ( $p = 0.0001$ ) in a recent systematic review on psychotherapies in low- and middle-income countries (Purgato *et al.*, 2018). The second reason is that even when we follow strict criteria in defining psychotherapy, meta-analysing different interventions even in the same category often results in clinically meaningful statistical heterogeneity (i.e.  $I^2 \geq 75\%$ ). In this case the risk of merging 'apples and oranges' (Eysenck, 1978) applies within interventions belonging to the same category.

Finally, many types of psychological and psychosocial interventions have been introduced and tested in low- and middle-income countries. Research in global mental health might benefit from head-to-head comparisons considering

separately each type of psychotherapeutic interventions *v.* each other, *v.* 'inactive' controls, but also *v.* other intervention option including less resource-intensive psychosocial interventions in homogeneous population groups. Moreover, given that efficacy has been proven for many types of psychological interventions (Tol *et al.*, 2011; Morina *et al.*, 2017), it would be worth investigating the mechanism of action of these interventions (i.e. mediators and moderators). This could help inform the best matches between population groups and evidence-based interventions. This would enable the generation of more informative research for clinicians and for policy makers involved in the management of health systems that incorporate mental health interventions across contexts.

## References

- Barbui C, Purgato M, Churchill R, Adams CE, Amato L, Macdonald G, McCleery J, Minozzi S and Syed Sheriff R (2017) Cochrane For global mental health. *The Lancet Psychiatry* 4, e6.
- Cohen J (1992) A power primer. *Psychological Bulletin* 112, 155–159.
- Cuijpers P and Cristea IA (2016) How to prove that your therapy is effective, even when it is not: a guideline. *Epidemiology and Psychiatric Sciences* 25, 428–435.
- Cuijpers P, Karyotaki E, Reijnders M and Ebert DD (2018a) Was Eysenck right after all? A reassessment of the effects of psychotherapy for adult depression. *Epidemiology and Psychiatric Sciences* 1–10. <https://doi.org/10.1017/S2045796018000057>
- Cuijpers P, Karyotaki E, Reijnders M, Purgato M and Barbui C (2018b) Psychotherapies for depression in low- and middle-income countries: a meta-analysis. *World Psychiatry* 17, 90–101.
- Eysenck H (1978) An exercise in mega-silliness. *American Psychologist* 33, 517.
- Fletcher J (2007) What is heterogeneity and is it important? *British Medical Journal* 334, 94–96.
- Fluckiger C, Del Re AC, Barth J, Hoyt WT, Levitt H, Munder T, Spielmanns GI, Swift JK, Visla A and Wampold BE (2018) Considerations of how to conduct meta-analyses in psychological interventions. *Psychotherapy Research* 28, 329–332.
- Higgins JPT and Green SE (2011) Cochrane Handbook for systematic reviews of interventions. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration. Available at <http://handbook.cochrane.org> (Accessed 26 November 2018).
- Lamers F, Jonkers CC, Bosma H, Kempen GI, Meijer JA, Penninx BW, Knottnerus JA and Van Eijk JT (2010) A minimal psychological intervention in chronically ill elderly patients with depression: a randomized trial. *Psychotherapy and Psychosomatics* 79, 217–226.
- Mohr DC, Ho J, Hart TL, Baron KG, Berendsen M, Beckner V, Cai X, Cuijpers P, Spring B, Kinsinger SW, Schroder KE and Duffecy J (2014) Control condition design and implementation features in controlled trials: a meta-analysis of trials evaluating psychotherapy for depression. *Translational Behavioural Medicine* 4, 407–423.
- Morina N, Malek M, Nickerson A and Bryant RA (2017) Psychological interventions for post-traumatic stress disorder and depression in young survivors of mass violence in low- and middle-income countries: meta-analysis. *British Journal of Psychiatry* 210, 247–254.
- Munder T and Barth J (2018) Cochrane's Risk of Bias tool in the context of psychotherapy outcome research. *Psychotherapy Research* 28, 347–355.
- Munder T, Fluckiger C, Leichsenring F, Abbas AA, Hilsenroth MJ, Luyten P, Rabung S, Steinert C and Wampold BE (2018) Is psychotherapy effective? A re-analysis of treatments for depression. *Epidemiology and Psychiatric Sciences*, 1–7. <https://doi.org/10.1017/S2045796018000355>
- Patel V, Saxena S, Lund C, Thornicroft G, Baingana F, Bolton P, Chisholm D, Collins PY, Cooper JL, Eaton J, Herrman H, Herzallah MM, Huang Y, Jordans MJD, Kleinman A, Medina-Mora ME, Morgan E, Niaz U, Omigbodun O, Prince M, Rahman A, Saraceno B, Sarkar BK, De Silva M, Singh I, Stein DJ, Sunkel C and Unutzer J (2018) The Lancet

- Commission on global mental health and sustainable development. *Lancet* **392**, 1553–1598.
- Purgato M and Adams CE** (2012) Heterogeneity: the issue of apples, oranges and fruit pie. *Epidemiology and Psychiatric Sciences* **21**, 27–29.
- Purgato M, Gastaldon C, Papola D, Van Ommeren M, Barbui C and Tol WA** (2018) Psychological therapies for the treatment of mental disorders in low- and middle-income countries affected by humanitarian crises. *Cochrane Database of Systematic Reviews* **7**, Cd011849.
- Singla DR, Kohrt BA, Murray LK, Anand A, Chorpita BF and Patel V** (2017) Psychological treatments for the world: lessons from low- and middle-income countries. *Annual Review of Clinical Psychology* **13**, 149–181.
- Singla DR, Raviola G and Patel V** (2018) Scaling up psychological treatments for common mental disorders: a call to action. *World Psychiatry* **17**, 226–227.
- Tol WA, Barbui C, Galappatti A, Silove D, Betancourt TS, Souza R, Golaz A and Van Ommeren M** (2011) Mental health and psychosocial support in humanitarian settings: linking practice and research. *Lancet* **378**, 1581–1591.