

# *Babesia bovis*: lipids from virulent S2P and attenuated R1A strains trigger differential signalling and inflammatory responses in bovine macrophages

G. GIMENEZ<sup>1\*</sup>, M. L. BELAUNZARÁN<sup>1</sup>, C. V. PONCINI<sup>1</sup>, F. C. BLANCO<sup>2</sup>, I. ECHAIDE<sup>3</sup>, P. I. ZAMORANO<sup>4</sup>, E. M. LAMMEL<sup>1</sup>, S. M. GONZÁLEZ CAPPA<sup>1</sup> and E. L. D. ISOLA<sup>1</sup>

<sup>1</sup>*Instituto de Microbiología y Parasitología Médica, Universidad de Buenos Aires-Consejo Nacional de Investigaciones Científicas y Tecnológicas (IMPaM, UBA-CONICET). Facultad de Medicina, Paraguay 2155 piso 13, C1121ABG Buenos Aires, Argentina*

<sup>2</sup>*Instituto de Biotecnología, Centro de Investigaciones en Ciencias Veterinarias y Agronómicas, INTA-Castelar, Nicolas Repetto y de los Reseros s/n, 1686 Hurlingham, Buenos Aires, Argentina*

<sup>3</sup>*Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria Rafaela, Ruta 34 km 227, 2300 Santa Fe, Argentina*

<sup>4</sup>*Instituto de Virología, Centro de Investigaciones en Ciencias Veterinarias y Agronómicas, INTA-Castelar, Nicolas Repetto y de los Reseros s/n, 1686 Hurlingham, Buenos Aires, Argentina*

(Received 7 August 2012; revised 21 September and 9 October 2012; accepted 12 October 2012; first published online 3 January 2013)

## SUMMARY

The intra-erythrocytic protozoan *Babesia bovis* is an economically important pathogen that causes an acute and often fatal infection in adult cattle. Babesiosis limitation depends on the early activation of macrophages, essential cells of the host innate immunity, which can generate an inflammatory response mediated by cytokines and nitric oxide (NO). Herein, we demonstrate in bovine macrophages that lipids from *B. bovis* attenuated R1A strain (L<sub>A</sub>) produced a stronger NO release, an early TNF $\alpha$  mRNA induction and 2-fold higher IL-12p35 mRNA levels compared to the lipids of virulent S2P strain (L<sub>V</sub>). Neither L<sub>A</sub> nor L<sub>V</sub> induced anti-inflammatory IL-10. Regarding signalling pathways, we here report that L<sub>A</sub> induced a significant phosphorylation of p38 and extracellular signal-regulated kinases 1 and 2 (ERK1/2) whereas L<sub>V</sub> only induced a reduced activation of ERK1/2. Besides, NF- $\kappa$ B was activated by L<sub>A</sub> and L<sub>V</sub>, but L<sub>A</sub> produced an early degradation of the inhibitor I $\kappa$ B. Interestingly, L<sub>V</sub> and the majority of its lipid fractions, exerted a significant inhibition of concanavalin A-induced peripheral blood mononuclear cell proliferation with respect to L<sub>A</sub> and its corresponding lipid fractions. In addition, we determined that animals infected with R1A developed a higher increase in IgM anti-phosphatidylcholine than those inoculated with S2P. Collectively, S2P lipids generated a decreased inflammatory response contributing to the evasion of innate immunity. Moreover, since R1A lipids induced a pro-inflammatory profile, we propose these molecules as good candidates for immunoprophylactic strategies against babesiosis.

Key words: Lipids, *Babesia bovis*, bovine macrophages, signalling, innate immunity, anti-phospholipid antibodies.

## INTRODUCTION

*Babesia bovis* is an intraerythrocytic apicomplexan parasite responsible for babesiosis in bovines, ruminants that are the natural hosts of this parasitosis. This tick-borne disease, prevalent in tropical and subtropical areas worldwide, is poorly controlled by the existing chemotherapy and immunoprophylaxis, remaining as an important constraint for the development of cattle industries (Bock *et al.* 2004). *Babesia bovis*-infected erythrocytes undergo sequestration by attachment to capillary endothelium, resulting in organ damage, cerebral dysfunction and pulmonary

oedema, similar to the most severe form of human malaria caused by another apicomplexa, *Plasmodium falciparum* (Wright *et al.* 1988). In *B. bovis* infection, innate immune mechanisms have been recognized to have an important role in the limitation of acute infection, whereas acquired immunity is likely to be more important for resistance to parasite challenge (Fell and Smith, 1998; Brown and Palmer, 1999). In this sense, it has been demonstrated that innate immune molecules derived from *B. bovis*-activated macrophages, such as NO and inflammatory cytokines, contribute to that limitation (Shoda *et al.* 2000). Although pro-inflammatory mediators function as elements of early protective immunity against haemoprotozoan parasites, as infection progresses in severity, overproduction of these molecules has been implicated in the pathological sequelae of disease (Krause *et al.* 2007).

As concerns lipids from microorganisms, these compounds have an important modulatory role in the

\* Corresponding author: Instituto de Microbiología y Parasitología Médica, Universidad de Buenos Aires-Consejo Nacional de Investigaciones Científicas y Tecnológicas (IMPaM, UBA-CONICET). Facultad de Medicina, Paraguay 2155 piso 13, C1121ABG Buenos Aires, Argentina. Tel: +54 11 5950 9500 ext. 2191. Fax: +54 11 5950 9577. E-mail: paradife@fmed.uba.ar

immune system (Schofield and Hackett, 1993; Roach and Schorey, 2002; Quesniaux *et al.* 2004). In *B. bovis*, it has been reported that this parasite possesses a high rate of phosphatidylcholine (PC) and phosphatidic acid (PA) biosynthesis (Florin-Christensen *et al.* 2000). Furthermore, we have demonstrated that the lipid composition of 2 *B. bovis* strains with polar behaviour, attenuated R1A and virulent S2P, present quantitative differences that contribute to the distinct inflammatory responses observed in a murine experimental model (peritoneal macrophages) (Gimenez *et al.* 2010). It is well known that macrophage stimulation leads to the activation of transcription factors related to inflammatory cytokines and NO production, and that the mitogen activated-protein kinase (MAPK) and the nuclear factor- $\kappa$ B (NF- $\kappa$ B) pathways are crucially involved in this process (Roach and Shorey, 2002; Hayden and Ghosh, 2008). In this regard, we have previously determined in murine macrophages that lipids of *B. bovis* R1A strain activate extracellular signal-regulated kinases 1/2 (ERK1/2) as well as the involvement of this pathway in pro-inflammatory cytokine release (Gimenez *et al.* 2010).

In the present work the immunomodulatory effect of lipids from these *B. bovis* strains in cells from its natural host was studied. With this aim, we analysed in bovine macrophages the induction of NO and the cytokines TNF $\alpha$ , IL-12 and IL-10, the activation of MAPKs and NF- $\kappa$ B signalling pathways and investigated the effect of these lipids in the proliferation of bovine peripheral blood mononuclear cells. Moreover, as we previously demonstrated in non-infected bovines the presence of natural IgM anti-PC and anti-phosphatidylethanolamine (PE) (Gimenez *et al.* 2007), we also here analysed the anti-phospholipid antibody response in bovines experimentally infected with each strain.

## MATERIALS AND METHODS

### Reagents and antibodies

Silica gel 60 thin layer chromatography plates and organic solvents were purchased from Merck (Darmstadt, Germany). Egg L- $\alpha$ -phosphatidylcholine (PC), egg sphingomyelin (SM) and egg L- $\alpha$ -phosphatidylethanolamine (PE) were purchased from Avanti Polar Lipids (Alabaster, Alabama, USA). Lipid standards, concanavalin A (Con A), protease inhibitors cocktail, Histopaque-1077, LPS from *Escherichia coli* (serotype O127:b8), sulphanilamide and naphthylethyldiamine dihydrochloride were acquired from Sigma Chemical Co. (St Louis, MO, USA). The Roche blocking reagent<sup>®</sup> was purchased from Roche (Mannheim, Germany). Peroxidase-labelled (HRP) anti-bovine IgM and anti-bovine IgG antibodies were obtained from KPL (Gaithersburg, Maryland, USA). O-phenyldiamine was acquired from

DakoCytomation (Denmark). Recombinant bovine IFN $\gamma$  (Ciba-Geigy) was kindly provided by Dr Carlos Suárez (WSU, Pullman, USA). Rabbit polyclonal anti-mouse  $\beta$ -actin, goat anti-rabbit IgG-HRP, anti-mouse IgG-HRP, anti-phospho p38 and anti- $\kappa$ B $\alpha$  antibodies were purchased from Santa Cruz Biotechnology Inc (Santa Cruz, CA, USA). Rabbit anti-phospho ERK and anti-total ERK antibodies were obtained from Cell Signaling Technology, Inc (Danvers, MA, USA). Nitrocellulose membrane and autoradiographic films were from GE Healthcare (Piscataway, NJ, USA). Supersignal<sup>®</sup> West Pico Chemiluminescent substrate was purchased from Pierce (Rockford, IL, USA). Protein and DNA molecular weight markers, random primers, SuperScript II Reverse Transcriptase, Taq Polymerase, DNase and oligonucleotides were obtained from Invitrogen Corporation (CA, USA). Maxisorp 96-well ELISA plates, 6- and 96-well culture plates and 100-mm polystyrene Petri dishes were purchased from Nunc (Roskilde, Denmark). Hanks balanced salt solution, TRIzol, RPMI 1640 medium were obtained from GIBCO BRL (Grand Island, New York, USA). The RNA extraction kit and PCR SYBR green QuantiTec were purchased from Qiagen Sciences Inc. (Venlo, Holland).

### Parasites

Merozoites from 2 strains of *B. bovis* with polar behaviour were employed: R1A, attenuated and S2P, virulent. R1A is a vaccinal strain derived from a virulent isolate responsible for a babesiosis outbreak in Santa Fe, Argentina, and attenuated by a series of 22 passages of parasitized blood in splenectomized calves (Anziani *et al.* 1993). The virulent S2P strain was isolated from a bovine during a babesiosis outbreak in Salta, Argentina and has been adapted to and maintained in continuous *in vitro* culture (Mangold *et al.* 1993; Baravalle *et al.* 2012). Continuous *in vitro* multiplication does not modify the attenuated or virulent phenotypes of *B. bovis* strains (Baravalle *et al.* 2012). Parasites from both strains were cultured in bovine erythrocytes under micro-aerophilic conditions (Levy and Ristic, 1980). To isolate R1A- and S2P-free merozoites, cultures were left overnight at 4 °C to favour parasite release from bovine erythrocytes to the extracellular media. Cultures were then centrifuged at 400 **g** for 10 min at 4 °C and parasites contained in the resulting supernatants were further separated by centrifugation at 12800 **g** for 20 min at 4 °C. Pellets were stored at -20 °C until used for lipid extraction.

### Experimental cattle

Six Holstein calves (6 months old), from INTA Experimental Station, Rafaela, Santa Fe, Argentina,

were infected with R1A or S2P *B. bovis*-parasitized erythrocytes. Holstein calves, located at INTA-Castelar, Argentina, serologically negative to *B. bovis* were used to obtain cells from whole blood samples for *in vitro* studies. First,  $10^7$  (R1A) or  $10^5$  (S2P) *B. bovis*-parasitized erythrocytes suspended in 2 ml of PBS, were packed into 2.5 ml individual sterile syringes and inoculated into 3 calves each via the jugular vein within 30 min after harvest from the tissue culture. The concentration of parasitized erythrocytes was determined in a Neubauer haemocytometer after staining the DNA with fluorochrome acridine orange. The calves were provided with water *ad libitum*, standard hay and commercial cattle concentrate. Calves were observed daily throughout the experimental period. All animal usage was according to protocols from the Animal Ethics Committee at INTA, Argentina, under the guidelines of the European Union (EU directive 86/609/EEC).

#### Serum samples

Blood samples were collected from *B. bovis*-inoculated calves following experimental infection protocols. Calves were bled weekly from the jugular vein, starting 1 week before the experimental infection (day 0) and ending during week 10 (day 66). Serum samples were stored at  $-20^{\circ}\text{C}$  until used for anti-phospholipid antibodies analysis.

#### Detection of anti-phospholipid antibodies in the sera of *B. bovis* experimentally infected bovines

The serum samples previously obtained were analysed by ELISA (Gimenez *et al.* 2007). Briefly, 100  $\mu\text{l}$  aliquots of PC, PE or SM in ethanol at 50  $\mu\text{g}/\text{ml}$  were loaded into test wells of microtitre plates. The solvent was evaporated and blocking was performed with a commercial blocking system (Roche Blocking Reagent<sup>®</sup>) for 2 h at  $37^{\circ}\text{C}$ . The plates were then washed with Tris-buffered saline (TBS), incubated with 60  $\mu\text{l}$  of the different serum samples (1:100, v/v) for 2 h at  $37^{\circ}\text{C}$ , washed again and incubated with 60  $\mu\text{l}$  of the conjugates anti-bovine IgM-HRP or anti-bovine IgG-HRP (1:500, v/v) for 2 h at  $37^{\circ}\text{C}$ . Plates were then washed and incubated with  $\text{H}_2\text{O}_2$  + O-phenyldiamine (OPD). After 4 min the absorbance was measured in a microplate reader Model 450 (Bio-Rad) at 490 nm. Results are expressed as the ratio of A490 of each day post-inoculation (p.i.)/0 days p.i. (basal level).

#### Preparation of total lipid extracts from R1A ( $L_A$ ) and S2P ( $L_V$ ) *B. bovis* merozoites

The method of Bligh and Dyer (1959) was used for lipid extraction. The solvents were evaporated under nitrogen to a constant weight and lipids from

different batches of R1A ( $L_A$ ) and S2P ( $L_V$ ) *B. bovis* merozoites were re-suspended individually in ethanol for macrophage stimulation assays or in chloroform for preparation of lipid fractions. Lipid extracts were stored at  $-20^{\circ}\text{C}$  until used.

#### Preparation of lipid fractions from $L_A$ and $L_V$

Each chloroform lipid extract previously obtained was separated by thin-layer chromatography (TLC) in silica gel plates, as previously described (Gimenez *et al.* 2010). Briefly, a double solvent system was employed: chloroform/methanol/acetic acid/water (65:35:6:2, v/v) up to the middle of the plate to separate phospholipids and hexane/diethylether/acetic acid (90:10:1, v/v) up to the top to separate neutral lipids (Florin-Christensen *et al.* 2000). For mass determination, plates were sprayed with 10%  $\text{CuSO}_4$  in 8%  $\text{H}_3\text{PO}_4$  and charred by exposure to  $150^{\circ}\text{C}$  for 13 min (Baron *et al.* 1984). Lipids were identified by comparison with authentic standards and the following fractions: sphingomyelin + lysophosphatidylcholine (SM+LPC), PC, phosphatidylserine + phosphatidylinositol (PS+PI), PE, PA, free fatty acids (FFA) and cholesterol + diacylglycerol (CHO+DG) were eluted from the TLC plates with  $2 \times 2$  ml portions of chloroform/methanol (1:1, v/v). The solvents were evaporated under nitrogen to a constant weight and lipids were re-suspended in ethanol for macrophage stimulation assays.

#### Culture of bovine monocyte-derived macrophages

Monocyte-derived macrophages were isolated from peripheral blood mononuclear cells (PBMC) by a modification of a method previously described by Stich *et al.* (1998). Briefly, bovine peripheral blood was collected from the jugular vein into heparinized 60-ml syringes and centrifuged at 1200 *g* for 30 min. Buffy coats were collected and diluted approximately to 30 ml with 2 volumes of PBS, under layered with 15 ml of Histopaque-1077, and centrifuged at 1500 *g* for 20 min. The PBMC were then removed from the interface between the plasma and Ficoll solution, pooled, diluted at least 1:3 with RPMI medium, and centrifuged at 500 *g* for 15 min. The pellets containing PBMC were washed repeatedly at 250 *g* until the supernatants were clear, re-suspended in complete RPMI medium (with 10% bovine autologous plasma), placed in 100-mm polystyrene Petri dishes at approximately  $1-2 \times 10^8$  PBMC/ml (10 ml/dish) and incubated overnight at  $37^{\circ}\text{C}$  in 5%  $\text{CO}_2$ . Non-adherent cells were removed by washing the plates 3 times with complete RPMI 1640 medium, and adherent cells were cultured in 10 ml of complete RPMI at  $37^{\circ}\text{C}$  for an additional 6–8 days to allow maturation into macrophages. Cell morphology was controlled by optic microscopy and the culture

medium was changed every 2–3 days. After approximately 1 week, adherent macrophages were washed once with complete RPMI and once with  $Mg^{2+}$ - and  $Ca^{2+}$ -containing Hanks balanced salt solution (HBSS), and removed by vortexing and incubated at 37 °C for 1 h in 0.5 mM EDTA in  $Mg^{2+}$ - and  $Ca^{2+}$ -free HBSS. Macrophages were harvested and used for nitric oxide (NO) and cytokine induction assays, as well as for the study of p38, ERK1/2 and NF- $\kappa$ B signalling pathways.

#### Nitric oxide detection

NO production was determined by measurement of nitrite ( $NO_2^-$ ) levels, a stable derivative of NO by the Griess assay, according to Shoda *et al.* (2000). Briefly, macrophages were cultured for 48 h at a concentration of  $10^5$  cells/well (100  $\mu$ l/well) in 96-well flat-bottom plates with complete RPMI medium, vehicle (0.5% ethanol) +50 U/ml bovine IFN $\gamma$  (control), 10  $\mu$ g/ml LPS +50 U/ml bovine IFN- $\gamma$  (positive control) or the different batches of  $L_A$  or  $L_V$  (50  $\mu$ g/ml) +50 U/ml bovine IFN- $\gamma$ . In all the assays vehicle was added in the equivalent amount as present in each of the lipid mixtures at each concentration. Culture supernatants were transferred (50  $\mu$ l/well) to new 96-well flat-bottom plates, and 100  $\mu$ l of 1% sulfanilamide in 2.5%  $H_3PO_4$ /well followed by 100  $\mu$ l of 0.1% naphthylethylenediamine dihydrochloride in 2.5%  $H_3PO_4$ /well were added to the supernatants. Absorbance at 540 nm was compared to a  $NaNO_2$  standard curve.

#### Cytokine mRNA analysis

Macrophages ( $1 \times 10^7$  cells/well) were plated into 6-well plates and left to adhere for 2 h at 37 °C in 5%  $CO_2$ . Cells were incubated overnight with 50 U/ml bovine IFN $\gamma$  and then stimulated with the different batches of  $L_A$  or  $L_V$  (50  $\mu$ g/ml) or vehicle (0.5% ethanol) for 0, 3, 6 and 16 h, at 37 °C in 5%  $CO_2$ . Total cellular RNA was isolated from macrophage cultures using the TRIzol reagent RNA isolation method as specified by the manufacturer (GIBCO) and the sediment was cleaned up following the protocol of a commercial kit for RNA extraction (Qiagen). Quality and quantity of total RNA were estimated by UV spectrophotometry (Nanodrop, Wilmington, DE, USA) and electrophoresis was performed on 0.8% agarose gels. DNA-free RNA (1  $\mu$ g) was mixed with 50 ng of random primers (Invitrogen) in 20  $\mu$ l of final volume and reverse transcribed to total cDNA with SuperScript II reverse transcriptase (Invitrogen) according to the manufacturer's instructions. One  $\mu$ l of the cDNA was used as template for each real-time quantitative PCR (qPCR) reaction. The primers for bovine: TNF $\alpha$  (F: CCCCAGAGGGAAGAGTCC, R: GGGCTACCGGCTTGTTACTTG, melting temperature (Tm)

84 °C), IL-12p35 (F: TAGCCACGAATGAGAGT-TGCC, R: TTTCCAGAAGCCAGACAATGC, Tm 78 °C), IL-10 (F: GGAAGAGGTGATGCC-ACAGG, R: AGGGCAGAAAGCGATGACAG, Tm 84 °C) and pol II genes (F: GGACCCGTGT-GGACAAGAAT, R: ACAAGCCCCAGGTAA-TCATCC, Tm 76 °C) were those previously described (Blanco *et al.* 2009). qPCR reactions were performed as previously described (Coussens *et al.* 2004). Briefly, SYBR green QuantiTec Mastermix (Qiagen) was employed and reactions were made on Applied Biosystem 7000 SDS using standard cycling conditions. All reactions were performed in duplicate and qPCR data were analysed using the 2- $\Delta\Delta$ CT with efficiency correction according to Pfaffl *et al.* (2002). REST beta 9 software (<http://www.gene-quantification.info/>, [rest-mcs-beta-9august2006]) was used for final calculations and statistical analysis. The value for each treatment was compared with the value of the control by using a Pair Wise Fixed Reallocation Randomization test attached to software REST beta 9. Results were presented as the relative expression ratio of each target gene (TNF $\alpha$ , IL-12p35 or IL-10), which considers the response of the target gene of the sample versus control (cells incubated overnight with IFN $\gamma$ +vehicle), and also compared to a reference non-regulated gene (Pol II).

#### Signalling pathway analysis by immunoblot

Macrophages ( $4 \times 10^6$  cells/well) were plated into 6-well plates and left to adhere 2 h at 37 °C in 5%  $CO_2$ . Cells were incubated overnight with 50 U/ml IFN- $\gamma$  and then stimulated with the different batches of  $L_A$  or  $L_V$  (50  $\mu$ g/ml) or vehicle (0.5% ethanol) for 5, 15 and 30 min at 37 °C in 5%  $CO_2$ . Cells were washed thrice with PBS and treated for 30 min with cold lysis buffer (50 mM Tris, pH 7, 1 mM EDTA, 1% Triton X-100, NaF 0.1 mM, protease inhibitor cocktail). Proteins from cells adhered to the 6-well plates were re-suspended in Laemmli's sample buffer, boiled and stored at -20 °C until used for phospho-ERK1/2, phospho-p38 and I $\kappa$ B immunoblot analysis. Samples were separated in a 10% SDS-PAGE, transferred to nitrocellulose membranes using a Trans-blot semi-dry transfer cell (Bio-Rad, California, USA) and blocked with TBS +0.05% Tween 20 (TBST) +5% skimmed milk for 2 h at room temperature. After 3 washes with TBST, the membranes were incubated with the following antibodies: anti-phospho-ERK1/2 (1:1000, v/v), anti-phospho-p38 (1:1000, v/v) or anti-I $\kappa$ B $\alpha$  (1:1000, v/v) in TBST overnight at 4 °C. Membranes were then washed thrice with TBST, incubated with goat polyclonal anti-rabbit IgG-HRP (1:2000, v/v) or anti-mouse IgG-HRP (1:2000, v/v) for 1 h at room temperature and washed thrice before detection. HRP-labelled conjugates were detected using Supersignal<sup>®</sup> West Pico chemiluminescent

substrate; membranes were exposed to autoradiographic films and scanned using an HP Scanjet 2400 apparatus.

For the loading control, detection of total ERK1/2 or  $\beta$ -actin was performed. The same membranes were blocked as described above and incubated with rabbit anti-total ERK1/2 antibody (1:1000, v/v) or rabbit polyclonal anti-mouse  $\beta$ -actin (1:5000, v/v) overnight at 4 °C. After 3 washes membranes were incubated with goat polyclonal anti-rabbit - HRP (1:2000, v/v) for 1 h at room temperature and washed thrice; HRP - labelled conjugates were detected as described above. The intensity of the signal of phospho-ERK1/2, phospho-p38 and  $I\kappa B\alpha$  bands was quantified by densitometry using Gel-Pro<sup>®</sup> Analyzer 4.0 software and normalized to the intensity of the corresponding ERK or  $\beta$ -actin bands.

#### Bovine PBMC proliferation assays

Cells ( $2 \times 10^5$  cells/well) were incubated for 48 h at 37 °C in 5% CO<sub>2</sub>, in 96-well flat-bottom plates with culture medium, vehicle (0.5% ethanol) or 2  $\mu$ g/ml concanavalin A (Con A) alone or together with: the different batches of L<sub>A</sub> or L<sub>V</sub> (0.5, 5 and 50  $\mu$ g/ml) and the lipid fractions SM+LPC, PC, PS+PI, PE, PA, FFA or CHO+DG of each *B. bovis* strain (0.5 or 5  $\mu$ g/ml). The assays were conducted in triplicate and the mixtures were radio-isotope labelled with 1  $\mu$ Ci/well [<sup>3</sup>H]-thymidine during the final 18 h of culture, harvested and counted in a liquid scintillation counter (Rackbeta, Liquid Scintillation counter, Amersham-Pharmacia).

#### Statistical analysis

Results were expressed as mean  $\pm$  S.E.M. or S.D. Unpaired Student's *t*-test was used to compare between groups, using GraphPad Prism 4 software for Windows. One- or two- way analysis of variance (ANOVA) was used for multi-group analysis depending on the assay.

## RESULTS

#### L<sub>A</sub> induced a significant release of NO in bovine macrophages

Considering that *B. bovis* lipids are relevant antigens (Goodger *et al.* 1990; Shoda *et al.* 2000) that can be recognized by innate immune receptors such as TLR2 (Gimenez *et al.* 2010), we here analysed the production of NO, a macrophage secretory product with inhibitory effect in *in vitro* growth of intracellular *B. bovis* (Montealegre *et al.* 1985; Shoda *et al.* 2000). First, we assessed the induction of NO in bovine macrophages stimulated with the lipid extracts of *B. bovis* merozoites from attenuated R1A (L<sub>A</sub>) or virulent S2P (L<sub>V</sub>) strains, in the presence of IFN $\gamma$ . Even though the lipids from both strains




Fig. 1. L<sub>A</sub> induced a higher release of NO than L<sub>V</sub>. Results are presented as the mean  $\pm$  S.E.M. of triplicate determinations and are representative of 3 independent experiments performed with different L<sub>A</sub> or L<sub>V</sub> batches and macrophages from different cattle. \*Statistically significant for macrophages stimulated with L<sub>A</sub> compared to L<sub>V</sub> ( $P < 0.05$ ). Lipids from both strains induced significant levels of nitrite compared to control cells ( $P < 0.05$ ). L<sub>A</sub>: total lipids of R1A *Babesia bovis* attenuated strain; L<sub>V</sub>: total lipids of S2P *B. bovis* virulent strain.

induced significant levels of NO compared to control cells, L<sub>A</sub> induced 4-fold higher levels than L<sub>V</sub>, indicating that the lipids of the attenuated strain generated a stronger inflammatory response (Fig. 1).

#### L<sub>A</sub> produced an earlier/stronger induction of inflammatory cytokine mRNA in bovine macrophages compared to L<sub>V</sub>

Cytokines IL-12 and TNF $\alpha$ , produced by macrophages and other antigen-presenting cells, are critical for generating and regulating innate and acquired immune responses against many pathogens (Trinchieri, 1995). On the other hand, IL-10 has been shown to suppress a broad range of inflammatory responses (Sabat *et al.* 2010). Herein, we analysed the induction of TNF $\alpha$ , IL-12 and IL-10 by real-time qPCR in bovine macrophages stimulated with L<sub>A</sub> or L<sub>V</sub> at different times. Control cells did not display variations in cytokine mRNA levels in the times assayed (data not shown) and all results were presented as the relative expression ratio of each target gene which considers the response of the sample versus control. Figure 2A shows that lipids from both strains induced a TNF $\alpha$  mRNA peak of similar intensity but L<sub>V</sub>-stimulated macrophages reached the peak 3 h later than those treated with L<sub>A</sub>. As regards IL-12p35, L<sub>A</sub> and L<sub>V</sub> induced significant levels of this cytokine mRNA at 3 h post-stimulation, reaching their maximum levels at 6 h post-stimulation (Fig. 2B). Noteworthy, at this time the expression ratio of IL-12p35 mRNA was 2-fold higher in macrophages stimulated with L<sub>A</sub> with respect to those stimulated with L<sub>V</sub> (Fig. 2B).




Fig. 2. L<sub>A</sub> produced an earlier/stronger induction of inflammatory cytokine mRNA compared to L<sub>V</sub>. Results are expressed as the relative expression ratio of each cytokine of each sample versus control and are representative of 3 independent experiments performed with different L<sub>A</sub> or L<sub>V</sub> batches and macrophages from different cattle. \*Statistically significant for IL-12p35 mRNA expressed in macrophages at 6 h post-L<sub>A</sub> stimulation compared to L<sub>V</sub> ( $P < 0.005$ ). L<sub>A</sub>: total lipids of R1A *Babesia bovis* attenuated strain; L<sub>V</sub>: total lipids of S2P *B. bovis* virulent strain.

As concerns IL-10, neither L<sub>A</sub> nor L<sub>V</sub> induced significant mRNA levels in the times assayed (Fig. 2C). Collectively, these results indicate that lipids of the virulent strain induced a reduced/retarded inflammatory cytokine response with respect to that induced by lipids of the attenuated strain.

#### *L<sub>A</sub> and L<sub>V</sub> induced a differential activation of p38 and ERK1/2 MAPKs and NF- $\kappa$ B signalling pathways in bovine macrophages*

Immunoblot analysis showed that L<sub>A</sub> induced a transient phosphorylation of p38 (pp38) with a

maximum at 5 min, decreasing at 30 min post-stimulation compared with control cells; in contrast, cells stimulated with L<sub>V</sub> did not display phosphorylation signals (Fig. 3A). No changes were observed at the different times assayed in control cells stimulated with IFN $\gamma$  + vehicle (basal phosphorylation pattern) (data not shown). The densitometry analysis of these results is shown in Fig. 3B.

In the case of ERK1/2, L<sub>A</sub> induced an increasing and sustained phosphorylation with the maximum levels at 30 min post-stimulation (Fig. 3C). At variance, macrophages stimulated with L<sub>V</sub> rapidly reached (5 min) the maximum levels of pERK but with lower values than those obtained with L<sub>A</sub>, recovering the basal levels at 30 min (Fig. 3C). No changes were observed in control cells (basal phosphorylation pattern) at the different times assayed (data not shown). The densitometry analysis of these results is shown in Fig. 3D.

NF- $\kappa$ B is an inducible transcription factor whose activation is regulated by the cytoplasmic inhibitor I $\kappa$ B; herein we evaluated the degradation of I $\kappa$ B as a hallmark of the activation of the NF- $\kappa$ B pathway (Pahlevan *et al.* 1999; Chen *et al.* 2000; Eligini *et al.* 2002). Macrophages stimulated with L<sub>A</sub> showed at 5 min an important decrease of I $\kappa$ B compared to control cells, recovering the basal levels at 30 min post-stimulation (Fig. 3E); whereas, in cells stimulated with L<sub>V</sub> a significant degradation of I $\kappa$ B was observed but at 30 min post-stimulation (Fig. 3E). No changes were observed in I $\kappa$ B levels at the different times assayed in control cells (data not shown). The densitometry analysis of these results is shown in Fig. 3F. Taken together, these results indicate that L<sub>A</sub> lipids promote a rapid/strong activation of p38, ERK1/2 and NF- $\kappa$ B pathways in contrast to L<sub>V</sub> that generated a delayed/reduced stimulation of these signalling cascades.

#### *L<sub>V</sub> produced a strong inhibition of bovine PBMC proliferation*

Having demonstrated that L<sub>A</sub> and L<sub>V</sub> were able to induce different bovine macrophage activation patterns, we studied the effect of both lipid extracts on PBMC proliferation as an early indicator of lymphocyte activation. As shown in Fig. 4A, both L<sub>A</sub> and L<sub>V</sub> inhibited ConA-induced PBMC proliferation in a dose-dependent manner. Interestingly, this effect was significantly higher in cells stimulated with L<sub>V</sub> with respect to those stimulated with L<sub>A</sub>, indicating that the former might have a stronger inhibitory effect over lymphocyte response. Next, we analysed the effect of the lipid fractions: SM+LPC, PC, PS+PI, PE, PA, CHO+DG and FFA from R1A and S2P strains, at 2 concentrations (0.5 or 5  $\mu$ g/ml). With respect to R1A lipid




Fig. 3. L<sub>A</sub> and L<sub>V</sub> induced a differential activation of p38 and ERK1/2 MAPKs and NF- $\kappa$ B signalling pathways. Immunoblot analysis of (A) p38 phosphorylation, (B) ERK1/2 phosphorylation, (C) I $\kappa$ B degradation, was performed. Blots were re-probed for  $\beta$ -actin or total ERK detection, as the loading control. Cells incubated overnight with IFN $\gamma$  + vehicle (IFN $\gamma$ ) show the basal phosphorylation profile (control). The blots are representative of 3 independent assays with different cattle. The intensity of the signals of (D) pp38, (E) ERK and (F) I $\kappa$ B were quantified by densitometry using Gel-Pro<sup>®</sup> analyser 4.0 software, normalized to the intensity of the corresponding  $\beta$ -actin or total ERK bands and expressed as arbitrary units (AU) with respect to control cells. The values are the mean  $\pm$  s.d. of duplicate determinations and are representative of 3 independent experiments performed with different L<sub>A</sub> or L<sub>V</sub> batches and macrophages from different cattle. \*Statistically significant with respect to control cells ( $P < 0.05$ ). L<sub>A</sub>: total lipids of R1A *Babesia bovis* attenuated strain; L<sub>V</sub>: total lipids of S2P *B. bovis* virulent strain; pp38: phospho-p38; pERK: phospho-ERK.

fractions, only FFA at the higher concentration produced a significant inhibitory effect in Con A-induced PBMC proliferation (Fig. 4B). In the case of S2P lipid fractions, most of them exerted an inhibitory effect, except for SM+LPC and PA in both concentrations and PC in the lower concentration (Fig. 4C). These results are in accordance with those previously obtained with the whole lipid extracts (L<sub>A</sub> and L<sub>V</sub>), corroborating that the marked inhibitory effect of L<sub>V</sub> in the lymphocyte proliferative response is a consequence of the individual inhibitory activities that the majority of the lipid fractions exert.

#### *Anti-phospholipid antibodies increased in bovines infected with R1A or S2P strains*

Previously, we determined in non-infected bovines the presence of natural IgM anti-PE and anti-PC antibodies (approximately 5-fold higher levels against PE than PC) and no detectable activity of anti-SM antibodies (Gimenez *et al.* 2007). Herein, we analysed anti-phospholipid antibody levels during the course of R1A or S2P *B. bovis* infection. For R1A the parasitaemia peak (approx. 0.02%) appeared at 7–8 days p.i. and lasted 1–2 days and for S2P, appeared at 8–10 days p.i. (approx. 2%) and




Fig. 4. S2P lipids produced a higher inhibitory effect in ConA-induced PBMC proliferation compared R1A lipids. Bovine PBMC were stimulated for 48 h with ConA alone and in the presence of: (A)  $L_A$  or  $L_V$  (0.5, 5 or 50  $\mu\text{g/ml}$ ), (B) lipid fractions of R1A (0.5 or 5  $\mu\text{g/ml}$ ) or (C) lipid fractions of S2P (0.5 or 5  $\mu\text{g/ml}$ ). Control cells were incubated with culture medium (C) or vehicle. The values are the mean  $\pm$  S.E.M. of triplicate determinations and are representative of 3 independent experiments performed with different  $L_A$  or  $L_V$  batches and PBMC from different cattle. The results obtained with  $L_A$ ,  $L_V$  or the different lipid fractions were statistically significant with respect to those obtained with ConA alone ( $*P < 0.05$ ,  $**P < 0.001$ ).  $L_A$ : total lipids of R1A *B. bovis* attenuated strain;  $L_V$ : total lipids of S2P *Babesia bovis* virulent strain; ConA: Concanavalin A, SM: sphingomyelin, LPC: lysophosphatidylcholine, PC: phosphatidylcholine, PS: phosphatidylserine, PI: phosphatidylinositol, PE: phosphatidylethanolamine, PA: phosphatidic acid, CHO: cholesterol, DG: diacylglycerol, FFA: free fatty acids.

lasted for more than 3 days (data not shown). Figure 5 shows that the increase in anti-phospholipids antibodies against PC and PE (IgM isotype) was




Fig. 5. Anti-phospholipid antibodies in *Babesia bovis*-infected bovines. IgM anti-PC (A) or anti-PE (B) antibodies in bovines infected with R1A-attenuated (black square) or S2P virulent (grey circle) *B. bovis* strains were detected by ELISA. The graphics show the ratio between absorbance at 490 nm (A490) of each day post-inoculation (days p.i.) and 0 days p.i. Results represent the mean  $\pm$  S.E.M. of triplicate measurements of 3 independent animals for each strain. At 21 days p.i., the values for the IgM anti-PC increase in animals inoculated with R1A were significantly higher with respect to those obtained in animals inoculated with S2P ( $P < 0.03$ ). On the same day, the values for the IgM anti-PC increase were significantly higher than IgM anti-PE in animals inoculated with R1A ( $P < 0.02$ ). PC: phosphatidylcholine, PE: phosphatidylethanolamine.

significantly higher in animals infected with each *B. bovis* strain with respect to their corresponding basal levels (0 days p.i.), reaching a peak at 21 days p.i. At 21 days p.i., the R1A strain produced a significantly higher increase in IgM anti-PC levels than that of IgM anti-PE, whereas S2P generated a similar increase for both antibodies. In addition, bovines infected with the R1A strain developed at 21 days p.i. a significantly higher increase in IgM anti-PC levels than those inoculated with S2P strain (Fig. 5A), whereas IgM anti-PE did not show significant differences between animals inoculated with each strain (Fig. 5B). No IgM anti-SM antibodies were detected, as expected (data not shown). Regarding the IgG isotype, no reactivity was demonstrated



against the phospholipids here studied (data not shown).

#### DISCUSSION

Macrophages, crucial cells of innate immunity, can inhibit early *Babesia bovis* replication in acute infection through the generation of soluble factors such as cytokines and inflammatory mediators (Brown *et al.* 2006). This is the first study that demonstrates in bovine macrophages the induction of NO by lipids purified from merozoites of 2 *B. bovis* strains with polar behaviour: attenuated R1A or virulent S2P ( $L_A$  or  $L_V$ ). It has been reported that NO has a relevant role in *B. bovis* growth inhibition (Johnson *et al.* 1996; Shoda *et al.* 2000), therefore the present finding that, in the early stage of parasite lipids-host cell interaction  $L_A$  induced the highest levels of NO, could be related to the attenuated virulence of R1A (vaccinal strain). At variance, the low NO induction by  $L_V$  could contribute to the establishment of this virulent strain in host cells. However, in long-term infections, the increase in the concentration of inflammatory mediators may correlate with the increase in the number and severity of clinical manifestations such as anaemia, coagulopathy, respiratory distress syndrome, encephalopathy and even death (Krause *et al.* 2007).

Regarding cytokine induction,  $L_V$  generated in bovine macrophages lower levels of IL-12p35 (Th1 polarizing cytokine) and a delayed induction of the inflammatory cytokine TNF $\alpha$  compared to  $L_A$ . This could correlate with the virulent behaviour of S2P through the generation of a lower inflammatory response and consequent evasion of host immune mechanisms. These results are in line with those previously described in murine peritoneal macrophages, another experimental model (Gimenez *et al.* 2010). Other authors have reported that lipids from erythrocytes infected with *B. bovis* Mexico strain did not induce TNF $\alpha$  or IL-12p35 or p40 by bovine macrophages (Shoda *et al.* 2000). These differences could be mainly attributed to the fact that they used a lipid mixture of erythrocytes plus *B. bovis* merozoites, a different strain and lipid quantities, as well as to variations in the kinetics of cytokine induction and determination of cytokine mRNA. On the other hand, the lipid structure and the relative quantity of each lipid in the total extracts of the *B. bovis* strains used, might account for the differential bovine macrophage activation here observed. In this respect, it has been demonstrated that lipoarabinomannans isolated from non-pathogenic mycobacterias stimulated a stronger cytokine response in macrophages, compared to those from pathogenic mycobacterias, demonstrating that slight variations in these molecules have a relevant impact in immunogenicity (Roach *et al.* 1993; Riedel and Kaufmann, 2000). Concerning IL-10, neither  $L_A$  nor  $L_V$  induced this

anti-inflammatory cytokine in bovine macrophages. In the case of  $L_A$ , this was in accordance with the pro-inflammatory profile here described for this lipid extract, whereas the results obtained with  $L_V$  were unexpected. In this sense, similar results have been reported for a lipid fraction of a pathogenic strain of *M. ulcerans* that inhibits the release of anti-inflammatory IL-10 as well as pro-inflammatory TNF $\alpha$  in human monocytes (Pahlevan *et al.* 1999).

Mitogen activated-protein kinases (MAPK) including ERK1/2 and p38, and the NF- $\kappa$ B pathways are important cellular targets of diverse pathogens (Ruckdeschel *et al.* 1997; Feng *et al.* 1999; Reiling *et al.* 2001; Roach and Schorey, 2002; Surewicz *et al.* 2004; Yadav *et al.* 2004; Souza *et al.* 2007; Hayden and Ghosh, 2008). The inhibition of the p38 pathway and the transient and lower activation of ERK 1/2 cascades by  $L_V$  compared to  $L_A$  here observed, are in agreement with the results described above. In addition, other studies have reported decreased MAPK activity only for pathogenic mycobacteria-infected cells (Roach and Schorey, 2002). Moreover, the high ERK1/2 activation induced by  $L_A$  is in accordance with the high release of TNF $\alpha$  since it has been reported that this signalling pathway is involved in the expression of this cytokine in human and murine macrophages stimulated with different pathogens or derived molecules (Ruckdeschel *et al.* 1997; Surewicz *et al.* 2004; Gimenez *et al.* 2010). As TNF $\alpha$  is one of the target genes regulated by NF- $\kappa$ B and also an activator of this signalling pathway (Verma *et al.* 1995), the reduced and delayed activation of NF- $\kappa$ B cascade produced by  $L_V$  with respect to  $L_A$ , is congruent with the retarded induction of TNF $\alpha$  detected in bovine macrophages stimulated by  $L_V$ . Altogether, the distinct activation patterns of these cascades induced by the lipids of each *B. bovis* strain could account for the contrasting effects observed in the inflammatory response and consequently with the outcome of infection.

As regards bovine PBMC proliferation assays, our results indicated that *B. bovis* lipids could modulate the LT response in a dose-dependent manner, being the lipids of the virulent S2P strain ( $L_V$ ) able to exert a higher inhibitory effect in the adaptive immune response with respect to  $L_A$ . Accordingly, it has been reported that a lipid fraction from a pathogenic strain of *M. ulcerans*, but not from a non-pathogenic strain, had an inhibitory effect in the proliferation of human PBMC/LT (Pahlevan *et al.* 1999). In addition, *M. leprae* and *M. tuberculosis* lipoarabinomannans suppress *in vitro* proliferation of LT from lepromatous patients in response to mitogens and antigens (Kaplan *et al.* 1987). In fact, we also observed different responses between the lipid fractions from  $L_A$  and  $L_V$  that could be attributed to qualitative differences in the structure of bioactive molecules, since they were tested in the same concentrations.

Finally, the significant increase in IgM anti-PC antibodies detected in bovines infected with R1A or S2P *B. bovis* strains with respect to non-infected animals could have resulted from the high rate of PC synthesis in *B. bovis* merozoites (Florin-Christensen *et al.* 2000) and the recognition of this antigenic molecule by the immune system. The finding that R1A generated a higher increase in anti-PC antibody response than S2P, despite having the former lower and short-term parasitaemias than the latter, might contribute in the better control of infection that R1A establishes with less or null clinical manifestations. In this sense, it has been described that the pre-immunization of bovines with a lipid extract of *B. bovis*-infected erythrocytes, developed lower and delayed parasitaemias after the challenge with a virulent strain of the same parasite (Goodger *et al.* 1990). As we only detected IgM anti-phospholipid antibodies that decreased with time, we propose that they could partially restrict the early phase of *B. bovis* infection. On the other hand, despite the fact that animals inoculated with S2P displayed higher parasitaemias than R1A-infected animals, the lower humoral response here observed could be attributed to the higher antigen offer, thus interfering with the efficiency of this immune response, as reported for malaria (Hisaeda *et al.* 2005; Wykes *et al.* 2007).

In summary, the strong inflammatory response generated by lipids from attenuated R1A strain could contribute to limit *B. bovis* acute infection and promote host survival, whereas the reduced inflammatory response induced by lipids from virulent S2P might collaborate in an early stage to evade the innate immune system, thus allowing *B. bovis* establishment in the host. Further studies should be performed using a larger set of strains of polar phenotypes in order to confirm and generalize the conclusions of this study. Despite the discovery of *B. bovis* over a century ago, there are still no safe and effective vaccines that protect cattle against this pathogen, therefore the ability of R1A lipids to stimulate protective innate immune responses here described, opens up the possibility that they might be a key factor in designing vaccine adjuvants for bovine babesiosis prophylaxis.

#### ACKNOWLEDGEMENTS

We thank Drs Silvina Wilkowsky for helpful discussions and Fabiana Bigi for allowing the performance of qPCR assays in her laboratory (INTA-Castelar, Argentina).

#### FINANCIAL SUPPORT

This work was supported by a grant from Agencia Nacional de Promoción Científica y Tecnológica (FONCYT) (PICT 08-12074).

#### REFERENCES

- Anziani, O. S., Guglielmone, A. A., Abdala, A. A., Aguirre, D. H. and Mangold, A. J. (1993). Protección conferida por *Babesia bovis* vacunal en novillos Holando Argentino. *Revista Medicina Veterinaria* **74**, 47–49.
- Baravalle, M. E., Thompson, C., Valentini, B., Ferreira, M., Torioni de Echaide, S., Florin-Christensen, M. and Echaide, I. (2012). *Babesia bovis* biological clones and the inter-strain allelic diversity of the Bv80 gene support subpopulation selection as a mechanism involved in the attenuation of two virulent isolates. *Veterinary Parasitology* <http://dx.doi.org/10.1016/j.vetpar.2012.06.037>.
- Baron, C. B., Cunningham, M., Strauss, J. F., 3rd and Coburn, R. F. (1984). Pharmacomechanical coupling in smooth muscle may involve phosphatidylinositol metabolism. *Proceedings of the National Academy of Sciences, USA* **81**, 6899–6903.
- Blanco, F. C., Schierloh, P., Bianco, M. V., Caimi, K., Meikle, V., Alito, A. E., Cataldi, A. A., Sasiain M del, C. and Bigi, F. (2009). Study of the immunological profile towards *Mycobacterium bovis* antigens in naturally infected cattle. *Microbiology and Immunology* **53**, 460–467. doi:10.1111/j.1348-0421.2009.00141.x.
- Bligh, E. G. and Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemical Physiology* **37**, 911–917.
- Bock, R., Jackson, L., de Vos, A. and Jorgensen, W. (2004). Babesiosis of cattle. *Parasitology* **129**, S247–269. doi:10.1017/S0031182004005190.
- Brown, W. C., Norimine, J., Knowles, D. P. and Goff, W. L. (2006). Immune control of *Babesia bovis* infection. *Veterinary Parasitology* **138**, 75–87. doi:10.1016/j.vetpar.2006.01.041.
- Brown, W. C. and Palmer, G. H. (1999). Designing blood-stage vaccines against *Babesia bovis* and *B. bigemina*. *Parasitology Today* **15**, 275–281.
- Coussens, P. M., Verman, N., Coussens, M. A., Elfman, M. D. and McNulty, A. M. (2004). Cytokine gene expression in peripheral blood mononuclear cells and tissues of cattle infected with *Mycobacterium avium* subsp. paratuberculosis: evidence for an inherent proinflammatory gene expression pattern. *Infection and Immunity* **72**, 1409–1422.
- Chen, C.-C., Sun, Y.-T., Chen, J.-J. and Chiu, K.-T. (2000). TNF- $\alpha$ -induced cyclooxygenase-2 expression in human lung epithelial cells: involvement of the phospholipase C- $\gamma$ 2, protein kinase C- $\alpha$ , tyrosine kinase, NF- $\kappa$ B-inducing kinase, and I- $\kappa$ B kinase 1/2 pathway. *Journal of Immunology* **165**, 2719–2728.
- Eligini, S., Brambilla, M., Banfi, C., Camera, M., Sironi, L., Barbieri, S. S., Auwerx, J., Tremoli, E. and Colli, S. (2002). Oxidized phospholipids inhibit cyclooxygenase-2 in human macrophages via nuclear factor- $\kappa$ B/I- $\kappa$ B- and ERK2-dependent mechanisms. *Cardiovascular Research* **55**, 406–415.
- Fell, A. H. and Smith, N. C. (1998). Immunity to asexual blood stages of Plasmodium: is resistance to acute malaria adaptive or innate? *Parasitology Today* **14**, 364–369.
- Feng, G. J., Goodridge, H. S., Harnett, M. M., Wei, X. Q., Nikolaev, A. V., Higson, A. P. and Liew, F. Y. (1999). Extracellular signal-related kinase (ERK) and p38 mitogen-activated protein (MAP) kinases differentially regulate the lipopolysaccharide-mediated induction of inducible nitric oxide synthase and IL-12 in macrophages: Leishmania phosphoglycans subvert macrophage IL-12 production by targeting ERK MAP kinase. *Journal of Immunology* **163**, 6403–6412.
- Florin-Christensen, J., Suarez, C. E., Florin-Christensen, M., Hines, S. A., McElwain, T. F. and Palmer, G. H. (2000). Phosphatidylcholine formation is the predominant lipid biosynthetic event in the hemoparasite *Babesia bovis*. *Molecular and Biochemical Parasitology* **106**, 147–156.
- Gimenez, G., Florin-Christensen, M., Belaunzarán, M. L., Isola, E. L., Suarez, C. E. and Florin-Christensen, J. (2007). Evidence for a relationship between bovine erythrocyte lipid membrane peculiarities and immune pressure from ruminal ciliates. *Veterinary Immunology and Immunopathology* **119**, 171–179. doi:10.1016/j.vetimm.2007.05.012.
- Gimenez, G., Magalhaes, K. G., Belaunzarán, M. L., Poncini, C. V., Lammel, E. M., González Cappa, S. M., Bozza, P. T. and Isola, E. L. (2010). Lipids from attenuated and virulent *Babesia bovis* strains induce differential TLR2-mediated macrophage activation. *Molecular Immunology* **47**, 747–755. doi:10.1016/j.molimm.2009.10.014.
- Goodger, B. V., Commins, M. A., Waltisbuhl, D. J., Wright, I. G. and Rode-Bramanis, K. (1990). *Babesia bovis*: immunity induced by vaccination with a lipid enriched fraction. *International Journal for Parasitology* **20**, 685–687.
- Hayden, M. S. and Ghosh, S. (2008). Shared principles in NF- $\kappa$ B signaling. *Cell* **132**, 344–362. doi:10.1016/j.cell.2008.01.020.
- Hisaeda, H., Yasutomo, K. and Himeno, K. (2005). Malaria: immune evasion by parasites. *The International Journal of Biochemistry & Cell Biology* **37**, 700–706.

- Johnson, W. C., Cluff, C. W., Goff, W. L. and Wyatt, C. R. (1996). Reactive oxygen and nitrogen intermediates and products from polyamine degradation are Babesiicidal in vitro. *Annals of the New York Academy of Sciences* **791**, 136–147.
- Kaplan, G., Gandhi, R. R., Weinstein, D. E., Levis, W. R., Patarroyo, M. E., Brennan, P. J. and Cohn, Z. A. (1987). *Mycobacterium leprae* antigen-induced suppression of T cell proliferation in vitro. *Journal of Immunology* **138**, 3028–3034.
- Krause, P. J., Daily, J., Telford, S. R., Vannier, E., Lantos, P. and Spielman, A. (2007). Shared features in the pathobiology of babesiosis and malaria. *Trends in Parasitology* **23**, 605–610.
- Levy, M. G. and Ristic, M. (1980). *Babesia bovis*: continuous cultivation in a microaerophilous stationary phase culture. *Science* **207**, 1218–1220.
- Mangold, A. J., Aguirre, D. H., Cafrune, M. M., de Echaide, S. T. and Guglielmone, A. A. (1993). Evaluation of the infectivity of a vaccinal and a pathogenic *Babesia bovis* strain from Argentina to *Boophilus microplus*. *Veterinary Parasitology* **51**, 143–148.
- Montealegre, F., Levy, M. G., Ristic, M. and James, M. A. (1985). Growth inhibition of *Babesia bovis* in culture by secretions from bovine mononuclear phagocytes. *Infection and Immunity* **50**, 523–526.
- Pahlevan, A. A., Wright, D. J., Andrews, C., George, K. M., Small, P. L. and Foxwell, B. M. (1999). The inhibitory action of *Mycobacterium ulcerans* soluble factor on monocyte/T cell cytokine production and NF-kappa B function. *Journal of Immunology* **163**, 3928–3935.
- Pfaffl, M. W., Horgan, G. W. and Dempfle, L. (2002). Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Research* **30**, e36.
- Quesniaux, V. J., Nicolle, D. M., Torres, D., Kremer, L., Guerardel, Y., Nigou, J., Puzo, G., Erard, F. and Ryffel, B. (2004). Toll-like receptor 2 (TLR2)-dependent-positive and TLR2-independent-negative regulation of proinflammatory cytokines by mycobacterial lipomannans. *Journal of Immunology* **172**, 4425–4434.
- Reiling, N., Blumenthal, A., Flad, H. D., Ernst, M. and Ehlers, S. (2001). Mycobacteria-induced TNF-alpha and IL-10 formation by human macrophages is differentially regulated at the level of mitogen-activated protein kinase activity. *Journal of Immunology* **167**, 3339–3345.
- Riedel, D. D. and Kaufmann, S. H. (2000). Differential tolerance induction by lipoarabinomannan and lipopolysaccharide in human macrophages. *Microbes and Infection* **2**, 463–471.
- Roach, S. K. and Schorey, J. S. (2002). Differential regulation of the mitogen-activated protein kinases by pathogenic and nonpathogenic mycobacteria. *Infection and Immunity* **70**, 3040–3052. doi:10.1128/IAI.70.6.3040-3052.2002.
- Roach, T. I., Barton, C. H., Chatterjee, D. and Blackwell, J. M. (1993). Macrophage activation: lipoarabinomannan from avirulent and virulent strains of *Mycobacterium tuberculosis* differentially induces the early genes c-fos, KC, JE, and tumor necrosis factor-alpha. *Journal of Immunology* **150**, 1886–1896.
- Ruckdeschel, K., Machold, J., Roggenkamp, A., Schubert, S., Pierre, J., Zumbihl, R., Liautard, J. P., Heesemann, J. and Rouot, B. (1997). *Yersinia enterocolitica* promotes deactivation of macrophage mitogen-activated protein kinases extracellular signal-regulated kinase-1/2, p38, and c-Jun NH2-terminal kinase. Correlation with its inhibitory effect on tumor necrosis factor-alpha production. *Journal of Biological Chemistry* **272**, 15920–15927.
- Sabat, R., Grütz, G., Warszawska, K., Kirsch, S., Witte, E., Wolk, K. and Geginat, J. (2010). Biology of interleukin-10. *Cytokine Growth Factor Reviews* **21**, 331–344.
- Schofield, L. and Hackett, F. (1993). Signal transduction in host cells by a glycosylphosphatidylinositol toxin of malaria parasites. *Journal of Experimental Medicine* **177**, 145–153.
- Shoda, L. K., Palmer, G. H., Florin-Christensen, J., Florin-Christensen, M., Godson, D. L. and Brown, W. C. (2000). *Babesia bovis*-stimulated macrophages express interleukin-1beta, interleukin-12, tumor necrosis factor alpha, and nitric oxide and inhibit parasite replication in vitro. *Infection and Immunity* **68**, 5139–5145.
- Souza, C. D., Evanson, O. A. and Weiss, D. J. (2007). Role of the mitogen-activated protein kinase pathway in the differential response of bovine monocytes to *Mycobacterium avium* subsp. paratuberculosis and *Mycobacterium avium* subsp. avium. *Microbes and Infection* **9**, 1545–1552.
- Stich, R. W., Shoda, L. K., Dreewes, M., Adler, B., Jungi, T. W. and Brown, W. C. (1998). Stimulation of nitric oxide production in macrophages by *Babesia bovis*. *Infection and Immunity* **66**, 4130–4136.
- Surewicz, K., Aung, H., Kanost, R. A., Jones, L., Hejal, R. and Toossi, Z. (2004). The differential interaction of p38 MAP kinase and tumor necrosis factor-alpha in human alveolar macrophages and monocytes induced by *Mycobacterium tuberculosis*. *Cellular Immunology* **228**, 34–41. doi:10.1016/j.cellimm.2004.03.007.
- Trinchieri, G. (1995). Interleukin 12: a proinflammatory cytokine with immunoregulatory functions that bridge innate resistance and antigen-specific adaptive immunity. *Annual Review of Immunology* **13**, 251–276. doi:10.1146/annurev.iy.13.040195.001343.
- Verma, I. M., Stevenson, J. K., Schwarz, E. M., Van Antwerp, D. and Miyamoto, S. (1995). Rel/NF-kappa B/I kappa B family: intimate tales of association and dissociation. *Genes and Development* **9**, 2723–2735. doi:10.1101/gad.9.22.2723.
- Wright, I. G., Goodger, B. V. and Clark, I. A. (1988). Immunopathophysiology of *Babesia bovis* and *Plasmodium falciparum* infections. *Parasitology Today* **4**, 214–218.
- Wykes, M., Keighley, C., Pinzon-Charry, A. and Good, M. F. (2007). Dendritic cell biology during malaria. *Cellular Microbiology* **9**, 300–305.
- Yadav, M., Roach, S. K. and Schorey, J. S. (2004). Increased mitogen-activated protein kinase activity and TNF-alpha production associated with *Mycobacterium smegmatis*- but not *Mycobacterium avium*-infected macrophages requires prolonged stimulation of the calmodulin/calmodulin kinase and cyclic AMP/protein kinase A pathways. *Journal of Immunology* **172**, 5588–5597.