






ORIGINAL ARTICLE

Audio-visual entrainment, cranio-electro stimulation, and sensory involvement: rival effects on attention and L2 vocabulary retention

Reza Pishghadam^{1,2} , Nasim Boustani¹ , Johannes Gerwien³ , Shaghayegh Shayesteh¹  and Taqi Al Abdwani² 

¹Ferdowsi University of Mashhad, Mashhad, Iran, ²Gulf College, Al Khuwair, Oman and ³Universität Heidelberg, Heidelberg, Germany

Corresponding author: Reza Pishghadam; Emails: pishghadam@um.ac.ir; reza@gulfcollege.edu.om

(Received 14 July 2024; revised 15 October 2024; accepted 4 November 2024; first published online 23 December 2024)

Abstract

It is believed that the attentional engagement of language learners may reinforce deeper neuronal processing and promote later retrieval. To address language learners' needs and facilitate language learning, we used audio-visual entertainment (AVE) and cranio-electro stimulation (CES), in addition to multisensory-based instruction, to modify attention and retention processes. Thus, we taught a set of words with the common procedure of audio-visual instruction to 32 English language learners in the control group, CES, and AVE sessions. However, they received five sensory involvements (i.e., auditory, visual, tactile, olfactory, and gustatory) for the target words in the multisensory session. Following each instruction, a pragmatic-Stroop task and a true/false test were conducted to examine the subjects' attention and retention processes, respectively. Analyzing the response times acquired from the pragmatic-Stroop task, it was found that multisensory-based instruction led to quicker responses in comparison to the audio-visual method preceded by AVE and CES stimulations. The response accuracy results from the retention test also revealed that the subjects provided more accurate responses to the words taught during the multisensory session. The implication is that the enriched multisensory inputs can improve L2 learners' mental agility and facilitate successful retention and retrieval of information after a short interval period.

Keywords: Attention; audio-visual entertainment (AVE); cranio-electro stimulation (CES); multisensory input; retention

Introduction

Discovering the functions of the brain is connected to deciphering neuronal codes apropos of the synchronizations and desynchronizations of neuronal activities.

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

On this subject, information on the excitability of neurons and neural oscillations is crucial in understanding its functions. Synchronous neural oscillations pertain to the periodic co-occurrences of a population of neurons arising from neuronal spiking (Jensen *et al.*, 2007). That is, the rhythmic neuronal synchronization is dynamically driven by the effectiveness of communications between higher and lower cortical regions; on the other hand, the synchrony influences a myriad of cognitive processes, like attention, retention, and perception (Engel *et al.*, 2001; Jensen *et al.*, 2007), which is generally known as brainwave entrainment. In other words, brainwave entrainment is “the use of external stimuli to produce a frequency following response of brainwaves to match the frequency of the external stimuli” (Basu & Banerjee, 2020, p. 269).

Grounded in the synchronization of the brainwave entrainment mechanism, multiple neurocognitive products have been introduced (e.g., audio-visual entertainment [AVE] and cranio-electro stimulation [CES]). These noninvasive cognitive therapies convert energies from sensory receptors into neural codes (as in AVE) or modify sympathetic and parasympathetic relations via neuromodulatory impacts (as in CES). Generally, the clinical and non-clinical applications of these devices are in line with the remedy of various experiences (e.g., attention deficit, insomnia, pain, anxiety, etc.) via constant stimulations at specific frequencies (i.e., gamma, beta, alpha, theta, and delta) over the cortex (Collura & Siever, 2009; Frederick *et al.*, 1999; Siever, 2007, 2012). For instance, Wuchrer (2009) enhanced sustained attention as a particular function of the brain by exciting the nerve pathways through the constant flashes of light (photic stimulation) and pulses of tones (auditory stimulation) at SMR/Beta frequency in a 20-minute AVE session. In addition, Lee *et al.* (2019) and Schroeder and Barr (2001) underlined the transmission of electrical potentials through the vagus nerve into the thalamus and examined the enhancement of attention in response to a 100 Hz mode via CES. They reported that the 100 Hz (vs. 0.5 Hz) mode activates the alpha and beta waves, leading to the modifications of thinking activities, in general, and mental load during attention and concentration processes, in particular. The outcomes of these studies provided some initial evidence for the assumption that the use of these safe and cost-effective treatments could give immediate improvements in learners’ attention. It is indeed believed that the attentional engagement of learners may change the initial encoding, reinforce deeper neuronal processing, and enhance learning. This engagement may protect against the loss of information and promote later retrieval.

In addition to the entrainment techniques, recent sensory processing models argue how, in the presence of continuous visual, auditory, tactile, and gustatory stimuli, sensory information can travel through the thalamus (i.e., the thalamo-cortical loop) and entrain the brain areas (Collura & Siever, 2009). Sensory transductions create efficacious perceptual information, and their contributions (i.e., known as multisensory, intersensory, or crossmodal) establish arch-perception of the world (Ernst, 2008; Rosenberg, 2015; Stein *et al.*, 2009). Recent results from behavioral (e.g., Pishghadam & Shayesteh, 2016; Pishghadam *et al.*, 2018) and electrophysiological studies (Boustani *et al.*, 2021; Pishghadam, Daneshvarfard, & Shayesteh, 2021; Pishghadam, Jajarmi, & Shayesteh, 2021; Shayesteh *et al.*, 2020) verify that the multisensory input processing is in alignment with cortical and

neocortical modulations. Accentuating the interplay between multisensory inputs and cognitive processes, studies (e.g., Battich et al., 2020; Boustani et al., 2021; Shayesteh, 2019; Talsma et al., 2010) have clarified the power of multisensory inputs on attention mechanism. In their electrophysiological study, Boustani et al. (2021) argued how different combinations of the senses in instructional practices can yield different results in sentence comprehension and retention mechanisms. However, it is not clear whether these sensory practices can compete with the common noninvasive cognitive therapies of AVE and CES devices.

To compare the rival effects of AVE and CES therapies on attention and retention manipulations, the experiments were designed using vocabulary instructions in L2. Similar investigations about the attention and retention mechanisms were arranged in subjects' non-native languages (Boustani & Al Abdwani, 2023; Boustani et al., 2021; Pishghadam, Daneshvarfard, & Shayesteh, 2021; Pishghadam, Jajarmi, & Shayesteh, 2021). Believing that the neuroelectric oscillations at SMR/Beta frequency serve as crucial protocols in improving attention (designed based on Collura & Siever, 2009; Wuchrer, 2009), for the purpose of this exploratory study, we applied DAVID Delight Pro and conducted an AVE session with the Brain Booster protocol. Furthermore, in considering the propositions about the excitability states created by the CES and the linkages that have been emphasized between neurostimulations and brain alterations, we used a 100 Hz mode (by DAVID Delight Pro for CES) with the intention of increasing attention (designed based on Lee et al., 2019; Schroeder & Barr, 2001). On the other hand, taking the role of enriched sensory inputs into consideration in attention modifications, we drew upon Pishghadam's (2015) emotioncy model and conducted a multisensory-based language instruction by simply enforcing the involvement of five senses during learning.

Hence, to highlight the efficacy of incorporating AVE and CES treatments into conventional classroom instructions (i.e., audio-visual) and compare it with that of the multisensory-based instruction for attention and retention mechanisms, we conducted four sessions: (1) control, (2) multisensory, (3) AVE, and (4) CES for each subject. In the control session, they learned a set of target words through the audio-visual method of teaching, but in the AVE and CES sessions, the teaching method was preceded by a round of AVE and CES. However, in the multisensory session, a list of words was taught through the auditory, visual, tactile, olfactory, and gustatory senses. At the end of each session, a pragmatic-Stroop test and a series of true/false statements were arranged to assess the learners' attention and retention processes, respectively. Following the results of the AVE- (Wuchrer, 2009), CES- (Lee et al., 2019; Schroeder & Barr, 2001), and multisensory-based (Boustani et al., 2021) studies, we hypothesized that the experiences in the experimental sessions (vs. control session) might manifest differences in the learners' attention and L2 vocabulary retention processes.

Methodology

Subjects

Among the 63 volunteers who participated in the exploratory study, 18 were excluded since they did not comply with the inclusion criteria (Table 1), and 12 of

Table 1. Inclusion criteria and acceptable range

Criteria	Test	Acceptable range
English Language Proficiency	Oxford Quick Placement Test (Allan, 1992)	30–40
Working Memory Span	Wechsler Adult Intelligence Scale III (Wechsler, 1981)	10–13
Attention Span	Sustained Attention to Response Task (Robertson et al., 1997)	≥ 10%
Emotioncy Level	The Emotioncy Scale (Borsipour, 2016)	0
Handedness	The Edinburgh Inventory of Handedness (Oldfield, 1971)	10–12

them did not take part in the four-session instruction, leaving 33 native Persian speakers (15 males and 18 females). After handling outliers, one subject was excluded from the final analysis, resulting in a total of 32 participants. This sample size is consistent with Lakens' (2022) justification and is influenced by statistical as well as practical considerations, including the availability of participants as well as time and logistical constraints. In addition, the high-power values ($1-\beta$) show that our sample size was sufficient for detecting the effects. They were right-handed adults (checked through Oldfield's (1971) Edinburgh Inventory of Handedness test) and ranged from 18 to 30 years of age. Based on a self-report questionnaire, we found that none of them reported any history of epilepsy, seizure, brain injury, stroke, tumor, ADHD, and motion sickness. The subjects had no heart pacer, neurological impairment, or psychiatric disorder and were with normal or corrected-to-normal vision. They also did not take drugs or use alcohol.

Moreover, they were intermediate English language learners (Allan, 1992) with normal working memory (Wechsler, 1981). It is worth noting that the motivation for using the English proficiency measures was to ascertain the participants' general level of English comprehension used in both the instructions and test sessions and control it as a variable that may impact EFL learners' attention (Segalowitz & Frenkiel-Fishman, 2005) and retention (Li et al., 2023) processes. In addition, to ensure the subjects' homogeneity assumption of attention, the Sustained Attention to Response Task (Robertson et al., 1997) was used to withhold their sensitivity to rare (vs. frequent) stimuli. Drawing on previous findings (e.g., Alloway & Alloway, 2012; Ralph et al., 2015; Robertson et al., 1997), we accepted those volunteers whose scores were less than 10%. Subsequently, we conducted Borsipour's (2016) emotioncy scale to accept those volunteers who did not have any prior familiarity with the target words. More specifically, it was scaled from not familiar (0 point); heard (1 point), heard and seen (2 points); heard, seen, and touched (3 points); heard, seen, touched, and used (tasted) (4 points); and heard, seen, touched, used, and done research on (5 points).




Figure 1. The task design.

Target words

To determine the target words for instruction, a 30-item list (designed based on Borsipour, 2016) including plants, fruits, and vegetables with their Persian equivalent was initially distributed among 130 respondents who were not the subjects of the experiment. Finally, 20 English words (some of them were low-frequency loanwords), with which 93% of the respondents had no familiarity, were chosen as the target words. To control the interfering effect of variables, we selected nouns with a defined number of letters (4–10) and syllables (2–3) (see Figure 1). Remarkably, to make the learning experience more realistic and applicable to real-world situations, the subjects were engaged with real (vs. unreal and pseudo) words. This approach enhanced the ecological validity of our findings and yielded reliable and generalizable results. Moreover, studies (e.g., Binder et al., 2020; Chen et al., 2020) in the field of language learning and cognitive psychology support the use of real words in experimental settings to investigate learning processes.

Procedure

Figure 1 illustrates the procedure. The details have been explained in the instruction (section 2.3.1), brain wave entrainment devices (section 2.3.2), behavioral measurements (section 2.3.3), and data analysis sections (section 2.3.4). Notably, each subject participated in all four conditions—AVE, control, multisensory, and CES—at different times. This within-subjects design controlled for individual variations in attention and working memory, ensuring that any differences in the results can be attributed to the experimental conditions rather than pre-existing differences between groups.

Instructions

Five words were taught to the subjects in each session (i.e., control, multisensory, CES, and AVE). The reason behind choosing five words came from the Stroop-

based studies (e.g., MacLeod, 1991; Parris *et al.*, 2021) pointing to the set-size effects of stimuli. Considering the number of items in this color-word test, we selected five words to meet the stimulus set size limitation.

The explanations in the instructions characterized the physical properties of the target words, like their size, color, and taste. To minimize the bias effect, and maintain the integrity and reliability of the results, the instructions were randomized. Additionally, it was crucial to ensure that there were at least two days between the AVE and CES sessions to prevent any potential carry-over effects. It is remarkable that the order of the words was counterbalanced across the participants. However, to maintain consistency within each learning condition and control the potential effect of confounding variables, the words were not counterbalanced across the sessions. This allowed us to ensure any differences in learning outcomes could be attributed to the specific intervention.

Moreover, the amount of information transferred and the instruction time devoted to the subjects and for each target item were much the same, and the instructions were done by one of the researchers. Importantly, to control the interfering effect of repetition, there were no repetitions of the target words, and the subjects were exposed to them only once. Each session was devoted to one participant, and the instruction time was preferably organized between 8 AM and 1 PM to be consistent with all the subjects. Each subject experienced AVE, control, multisensory, and CES methods on four separate days, with each session lasting for approximately 60 to 100 minutes. This time comprised signing the informed consent, receiving the instruction, preparing the AVE and CES devices, running a sample task, and then doing the attention and retention tasks.


Audio-visual instruction. Following the procedure employed commonly in academic settings, the subjects in the control group, CES, and AVE sessions merely received auditory and visual inputs. More specifically, they underwent verbal explanations while receiving visual input via a PowerPoint presentation. The colored photos were presented at the center of a white background, and the name of each item was typed in black font. The font size was Times New Roman 60 pt bold on the top of a 55-inch television screen.

Multisensory-based instruction. To conduct the multisensory-based language instruction, we relied upon previous studies (Boustani *et al.*, 2021; Pishghadam, Daneshvarfard, & Shayesteh, 2021; Shayesteh *et al.*, 2020). The subjects received full sensory involvement for the five target words in the multisensory session. In fact, they were exposed to tactile, olfactory, and gustatory inputs besides auditory and visual inputs. In particular, to teach the target words, real plants and fruits, besides, an oral presentation and a PowerPoint were used. Table 2 delineates a sample instruction for the fruit kiwano, in which the subjects received auditory, visual, tactile, olfactory, and gustatory inputs.

Brain wave entrainment devices

Audio-visual entrainment device (AVE). The Digital Audio-Visual (David Delight Pro by Mind Alive, Inc., Edmonton, Canada) was applied. The photic and auditory stimulations were administered with the Brain Booster (Left stim 14 Hz, Right stim

Table 2. A sample instruction

Kiwano	
<p>Now, look at this fruit. This is called kiwano. As you see, kiwano is oval with orange skin... Touch it. How does it feel? Right... It's rough. And can you eat the skin? Of course, not... The skin is inedible. Let's cut the fruit. Oh, look... Kiwano flesh is green. How is the flesh? Aha right... Jellylike. In fact, kiwano flesh is green and jellylike. Do you see the white seeds? Touch them... Are they soft or hard? They're soft... Aha, so they're edible. Let's smell it... , Kiwano smells fruity. Do you like to taste this fruit? Yeah... How does it taste? Aha right... It tastes sweet and somewhat like a cucumber.</p>	

20 Hz) session to help the subjects' focus and concertation processes (Wuchrer, 2009). Following the handbook of "Operator's Manual" (<https://mindalive.com/manuals/>) and experiments (e.g., Collura & Siever, 2009; Wuchrer, 2009), we specifically chose the 24-minute SMR/Beta protocol (Left: 13.5 Hz/Right: 18 Hz) to create a gentle and entraining session that helps to focus mental functioning and memory.

The Tru-Vu Omniscreen Multi-Color Eyeset used yellow (to improve cognition and focus [14–20 Hz]) and cyan (to enhance entrainment while still being calming and relaxing) flickering lights via LEDs implanted on the inner side of the darkened eyeglasses. Adjusting the brightness of the LEDs, the subjects were instructed to close their eyes, be relaxed but not fall asleep, and attend to the lights during the photic training period. Running a sound sync session, we selected the binaural-beat entrainment mode to improve the subjects' attention (Aparecido-Kanzler et al., 2021; Lane et al., 1998; Park et al., 2018).

To achieve the best results, we conducted the AVE session in the morning (Collura & Siever, 2009) in a sound-attenuated and dimly lit room. The subjects were in a sedentary position in a comfortable chair, and they were asked to drink a glass of water 15 minutes before its commencement (Collura & Siever, 2009). They were also foretold not to consume any drugs and not to drink any caffeine products before the session (On et al., 2013). All the subjects signed an informed consent form approved by the Ferdowsi University of Mashhad Ethics Committee.

Cranio-electro stimulation device (CES). The CES ear clip stimulus cable was connected to the CES output jack of the Delight Pro device (Mind Alive, Inc., Edmonton, Canada) powered by a 9-volt alkaline battery. The 20-minute CES session was run at 100 Hz mode to create a positive impact on attention (Lee et al., 2019; Schroeder & Barr, 2001). Entraining the CES pulsed electrical current, two sponges were placed on the left and right earlobes while the subjects remained seated in a comfortable chair. For better conductance, we dampened the earlobes with a small amount of water. Each subject was informed about the device with a neutral expectation prime and neutral language (Sansevere et al., 2022) before signing the informed consent. Due to the enhancements in the production of the neurotransmitters, we had a 20-minute time break between the CES

Table 3. Examples of the congruent and incongruent target word

Target word	Condition
Salak	Congruent
Salak	Incongruent
Salak	Incongruent
Salak	Incongruent
Salak	Incongruent

implementation and the audio-visual instruction (Liss & Liss, 1996; Southworth, 1999). Furthermore, the AVE and CES sessions were separated by at least two days each to decrease any carry-over effects.

Behavioral measurements

Immediately following each instruction, a Stroop and a true/false test were arranged to examine the subjects' attention and retention processes, respectively.

Attention test. To measure the subjects' attention to the target words, we used SuperLab (Version 4.0) and designed the neuropsychological test of Stroop (Stroop, 1935). Customizing this color-word test, we designed a pragmatic-Stroop test consisting of the target words (five words for each group), in addition to equal stimuli set size as fillers (4–10 letters and 2–3 syllables) in green, orange, brown, pink, and purple ink colors. The subjects had to determine the congruence (10 items) and incongruence (40 items) of each word in harmony with the appeared color in a 2.5-minute block with 50 trials by pressing the defined keyboard keys. To judge the congruence or incongruence state of the words, the subjects were informed to decide based on the color of the plants, fruits, and vegetables taught (Table 3). For instance, salak skin is brown, so the word salak, which was written in brown, was congruent; however, those words in pink, green, orange, and purple ink colors were incongruent.

The stimuli appeared randomly in a word-by-word format in the center of a 17-inch computer screen in Times New Roman 58 pt bold font size at the center of a white background. The stimuli were presented for 2000 milliseconds on the computer screen (designed based on <https://www.pytoolkit.org>), and the inter-stimulus interval between the experimental trials was 1000 milliseconds (Ben-Haim *et al.*, 2016) (see Figure 2).

Retention test. The computerized version (by SuperLab 4.0) of the retention test was designed with 50 (25 correct and 25 pragmatically incorrect) sentences with the target words for each group (see Table 4). The subjects were required to judge the acceptability of the sentences, which appeared randomly in the center of a 17-inch computer screen in black font color and Times New Roman 48 pt bold font size at the center of a white background by pressing “x” or “m” key for correct and incorrect sentences, respectively. To improve the reliability of measures and to maintain the complexity features, the sentences were in a specific range of word




Figure 2. Stroop trial samples (Rambutan and Longan).

counts (i.e., 3–4) and complexity (simple present tense, singular type, and active voice).

Data analysis

To analyze the subjects' response time (RT) acquired from the pragmatic-Stroop test and response accuracy (RA) from the retention test across different sessions (i.e., control, multisensory, AVE, and CES), IBM SPSS Statistics software (version 25) was run. Additionally, Microsoft Power BI Desktop (version 2.81.5831.1181 64-bit) was utilized for data visualization. An outlier was identified by examining the distribution of the RT and RA data. The data from a subject, which included inconsistent scores with the general pattern of responses, were excluded from both the RA and RT analyses to prevent distortion of the overall results. The normality of data was examined by the classical Kolmogorov–Smirnov test and an alpha level of 0.05 was applied in the post hoc analysis with Bonferroni correction.

Results

Response time


The distributions of data points in the RT dataset across AVE, CES, control, and multisensory groups are depicted in the violin plot (Figure 3). The width of each curve represents the density of RTs in milliseconds. The data for the AVE and control groups suggest multiple peaks, indicating a multimodal distribution, in contrast to the CES and multisensory groups. The central box plots illustrate the median scores and interquartile ranges, summarizing the central tendency and variability within each group. The figure shows that the multisensory group has the lowest median and mean values. This means that the subjects had the fastest reaction time in judging the match or mismatch between colors and words in the multisensory group. The following section provides a detailed statistical analysis of the data.

To systematically analyze subject-level variability in a within-subjects design, a linear mixed-effects model was conducted for each dependent variable. This model involved both fixed effects (i.e., the groups) and random effects (i.e., the subjects), allowing us to examine the subjects' reaction times in judging color-word congruency or incongruency across the control, multisensory, CES, and AVE

Table 4. Examples of the correct and incorrect sentences

Sentence	Condition
Kumquat is oval.	True
Kumquat is round.	False

Note: The target words are boldfaced.

**Figure 3.** Violin plot for response time.

groups. Consistent with the conventional Stroop effect analysis, which involves calculating the difference in reaction times between incongruent and congruent trials (MacLeod, 1991), the dependent variable for the RT analysis included both congruent and incongruent trials. The Type III tests of fixed effects confirmed that the group effect was significant [$F(3, 93) = 11.186, p < 0.01$] and the subjects surpassed in multisensory group ($M = 1196.385$ ms) compared to the control ($M = 1242.638$ ms), CES ($M = 1338.472$ ms), and AVE ($M = 1325.953$ ms) groups. Precisely, the pairwise post hoc results (Table 5) presented that the multisensory group performed significantly faster than the CES ($MD = -142.088$ ms, $p < 0.001$) and AVE ($MD = -129.568$ ms, $p < 0.001$) groups. The mean differences were also significant between the control and CES group ($MD = -95.834$, $p < 0.01$) and the control and AVE group ($MD = -83.315$ ms, $p < 0.01$). As a result, the subjects' attention process and RT meaningfully varied across different groups (i.e., Control/ Multisensory < CES/ AVE).

Table 5. Pairwise comparisons of the groups for response time

Group comparison	Mean difference
Control vs. Multisensory	46.253
Control vs. CES	-95.834**
Control vs. AVE	-83.315*
Multisensory vs. CES	-142.088***
Multisensory vs. AVE	-129.568***
CES vs. AVE	12.520

Based on estimated marginal means.
p value significance level: *** < .001, ** < .01, * < .05.
 Note: The standard error is 28.73 and the degree of freedom is 93.




Figure 4. Violin plot for response accuracy.

Response accuracy

The violin plot (Figure 4) below visualizes the distributions of RA across AVE, CES, control, and multisensory groups. The width of each violin corresponds to the density of accurate responses. Accordingly, the AVE data distribution indicates a multimodal distribution compared to the other groups. The central box plots present the median scores and interquartile ranges, illustrating the central tendency and variability within each group. Based on the figure, the multisensory group has the highest median and mean values, while the control group has the lowest. This conveys that the subjects outperformed in the multisensory group. To delineate the subjects’ performances across the four groups, a mixed-effects statistical analysis was conducted, as explained below.

A linear mixed-effects model was run to evaluate the average subjects’ assessment of sentence correctness across the four groups (i.e., control, multisensory, CES, and

Table 6. Pairwise comparisons of the groups for response accuracy

Group comparison	Mean difference
Control vs. Multisensory	-8.000***
Control vs. CES	-4.688***
Control vs. AVE	-5.156***
Multisensory vs. CES	3.313**
Multisensory vs. AVE	2.844*
CES vs. AVE	-0.469

Based on estimated marginal means.

p value significance level: *** < .001, ** < .01, * < .05

Note: The standard error is 0.939 and the degree of freedom is 93.

AVE) and inter-subject variability. The model included both fixed effects (i.e., the groups) and random effects (i.e., the subjects). The Type III tests of fixed effects analysis presented the significant impact of the group [$F(3, 93) = 24.92, p < 0.01$] with the outstanding performance of the multisensory group ($M = 43.625$ correct sentences) over control ($M = 35.625$ correct sentences), CES ($M = 40.312$ correct sentences), and AVE ($M = 40.781$ correct sentences) groups. Specifically, the pairwise comparisons (Table 6) revealed significant variations between multisensory and control ($MD = 8, p < 0.001$), between multisensory and CES ($MD = 3.313, p < 0.01$), and between multisensory and AVE ($MD = 2.844, p < 0.05$) groups. In addition, the post hoc pairwise comparisons revealed significant differences between the control and CES ($MD = -4.688, p < 0.001$) and between the control and AVE ($MD = -5.156, p < 0.001$) groups. Thus, the various treatments had noteworthy impacts on the subjects' retention and judgment of the correct and pragmatically incorrect sentences (i.e., Control < CES/ AVE < Multisensory).

Discussion

To discover modifications in attention and retention mechanisms, we used two noninvasive cognitive therapies in competition with multisensory practices. Thus, for the purpose of this experiment, we compared and contrasted the efficacy of DAVID Delight Pro (for AVE and CES) and that of multisensory-based instruction for L2 vocabulary learning and retention. Finally, the learners' attention and retention processes were assessed by researchers-made pragmatic-Stroop tasks and true/false acceptability judgment tests, respectively. In what follows, we discussed the behavioral results elicited by multiple exposures across the sessions.

Response time

The results revealed that the multisensory input had a meaningful impact on RT. The accumulating evidence has progressively revealed the facilitatory effect of optimal combinations of senses on performances (e.g., Bolognini *et al.*, 2005; Boustani *et al.*, 2021; Frassinetti *et al.*, 2002; Lovelace *et al.*, 2003; Pishghadam *et al.*,

2020; Shayesteh et al., 2020). In fact, the presence of significant differences in the agility of attending to the congruence and incongruence of the color-word pragmatic-Stroop test adheres to the implication that the involvement of the senses is key to fostering reaction time. The effect of the enriched sensory inputs on RT is specifically consistent with Boustani et al.'s (2021) study, which pronounced the predominance of responses to the sentences with the involvement of the five sensory inputs. However, in their study, the weight of senses drove results in a sentence comprehension task, whereas in the current experiment, the multimodality of senses affected the comprehension of words in connection with their color. The impact is specifically accentuated when the sensory inputs are congruent (Li & Deng, 2023). Examining the presentation of information through multiple sensory channels and congruency in inputs, Li and Deng propose that multisensory integration allows for the concurrent processing of various information. However, the process can lead to faster recognition and RT in conjunction with inputs that complement each other (i.e., congruent vs. incongruent inputs). Applying Li and Deng's idea to the current study, the instructor's explanations (i.e., auditory inputs) were consistent with the presented stimuli features (i.e., visual, tactile, gustatory, and olfactory). This is also in line with Pahor et al.'s (2021) findings, which discuss that engaging multiple senses can reduce cognitive load and maintain attention, leading to enhanced processing speed. In effect, the multisensory presentation of information enables the simultaneous processing of various inputs, ameliorating the processing speed. Such accounts provide the implication that multisensory inputs enhance agility by directing the learners' attention toward the instruction.

Moreover, the results revealed that the learners did not surpass significantly after the AVE session in comparison to the control and multisensory sessions. A justification could be the deep relaxation effect that the device creates by the dissociation mechanism disconnecting the subjects from their "thoughts and somatic awareness" (Siever, 2007). The tool may induce a restabilization effect as a result of the cyan flickering light via LEDs, leading to the relaxation of muscles (Siever, 2007), hence, a slow RT. The other potential factor that slows down the RT may be the alterations in the neurotransmitters and cerebral blood flow (CBF). Siever (2007) believes that during cognitive tasks, the brain's requirement for CBF increases. On the other hand, they are accompanied by the secretion of norepinephrine and adrenaline because of anxiety (Aston-Jones et al., 1991; Bremner, 2002), degrading performance on tasks. Accordingly, adjusting the flash of photic stimulation regarding the wide array of frequencies is a way to boost CBF (Fox & Raichle, 1985; Sappey-Marinier et al., 1992; Siever, 2007). However, it seems that the contemporaneous impact of facilitative (vs. debilitating) anxiety (Alpert & Haber, 1960) may be required for the promotion of learners' mental agility, while the stimulation apparatus may suppress both facilitative and debilitating anxieties. To minimize the potential confounding effect of the AVE apparatus, incorporating other photic stimulation colors and comparing their impacts may help identify any variable associated with this device.

Another probable justification for the poor performance of the subjects may reside in the influences of binaural beats. Following several studies (Aparecido-Kanzler et al., 2021; Lane et al., 1998; Park et al., 2018), we ran a binaural-beat-based stimulation to entrain the attention-related brain waves, though the exploratory

findings did not prove a positive modification in RT. Finding the reason, Robison and colleagues (2022) claim that the binaural-beat stimulation “exhibited a shallower vigilance decrement,” evidencing that this stimulation does not significantly modulate the reaction time and sustained attention. In effect, losses in vigilance and alertness undermine operator effectiveness and performance (Matthews et al., 2010). In other words, “measures of attentional resource availability may predict the operator’s subsequent vigilance performance” and CBF velocity as an index of “energization predicted sensory and cognitive vigilance, consistent with resource theory” (Matthews et al., 2010, p. 187). Similarly, electroencephalography recordings after a 20-minute binaural-beat stimulation revealed no electrophysiological improvement in focus and concentration mechanisms (Crespo et al., 2013). Siever (2007) claims that the AVE mechanisms reassure us that the excess experience of relaxation harnesses the speed of mental processes. A suggestion is to replicate the study with the employment of monoaural beats and alterations in the duration of exposure.

Moreover, the absence of a significant performance in attention after a 20-minute session of CES 100 Hz mode runs counter to the results of some studies (e.g., Lee et al., 2019; Schroeder & Barr, 2001). The changes were consistent with an evaluation of the brain’s current density that reported a decline in beta activity with concomitant alleviations of anxiety (Kennerly, 2006). In effect, the beta brainwave is commonly associated with being conscious, vigilant, and awake (Addante et al., 2021; Siever, 2008), whereby the suppression of this wave imposes a sleep state. Another reason may be the secretion of serotonin (as cited in Operator’s Manual, 2021 [<https://mindalive.com/manuals/>]), creating a calm state of mind. The combination of these findings provides a fair understanding of impairments in mental alacrity after a CES round. It is recommended to apply other voltages across different time spans and engage subjects with tasks designed to intentionally inhibit sleep.

Response accuracy

The results obtained from the retention task confirmed the hypothesis, remarking that the meaningful effects of the experimental sessions orchestrated the successful retention of information in competition with the control session. The significant performance after multisensory inputs was also in accordance with previous studies (e.g., Boustani et al., 2021; Frassinetti et al., 2002; Lovelace et al., 2003; Shayesteh, 2019; Shayesteh et al., 2020) evidencing the major contribution of optimal sensory inputs in learners’ performances. Enriched sensory inputs exert specific impacts on retention, which are apparent not only in immediate recall (Shayesteh, 2019) but also in the long-term retention of information (Boustani et al., 2021). In fact, the meaningful differences in the number of correct answers highlighted that the multisensory-based instruction maximized comprehension and retention processes. Engaging multiple sensory pathways can stimulate deeper encoding of information, thus enhancing subsequent retention and retrieval of information (Dunn & Dunn, 1993). Delving into the neural pathways of multisensory learning, Jensen (2009) provides insights into the involvement of various brain regions contributing to holistic cognitive development. The process fosters neural connections and memory

retention (Bahri Roudposhti & Al Abdwani, 2024; Dunn & Dunn, 1993; Pishghadam et al., 2024). Aligning with the findings, we find that multisensory learning can be a promising solution to retaining and retrieving information effectively. Among the myriad of benefits, we found that this approach can support immersive learning experiences and optimize retention processes.

The accurate performances of learners in the CES and AVE groups, compared to the control group, demonstrate the brain's need to reduce overarousal for effective information retention. Practically, the use of AVE and CES devices in current learning situations, which are merely based on visual and auditory inputs, can be effective in the consolidation of information. In common educational settings, learners receive an enormous amount of information without being involved in or completely understanding it, which is a source of anxiety (Siever, 2012). Hence, to degrade the possible effects, the AVE and CES control variations in levels of neurotransmitters and CBF (Siever, 2008). Such a result may be embodied in a host of alterations occurring in neuronal excitability states. In sum, taking the mechanisms of photic and audio stimulations as well as the 100 Hz CES mode into account, we can implicate that an appropriate level of altered CBF is the prerequisite of the retention process.

Last but not least, there are numerous open questions about the effect of the AVE sessions and CES modes on learners' performances. The study addressed the effectiveness of AVE and CES apparatuses and the multisensory teaching approach within the defined scope of concrete nouns. As a starting point, the tangible nature of concrete nouns allowed us to control variables and establish a baseline of how these methods can aid in L2 vocabulary learning. Though the study is confined to a specific subset of vocabulary, which limits the generalizability of the findings, we believe that the underlying principles have the potential for broader applications. The application of this approach can even be extended to abstract concepts by using creative strategies that make the intangible more tangible. Different representations and cues evoke feelings or themes associated with the abstract concept. Moreover, since some of our words were low-frequency loanwords, it is suggested that future studies consider adopting alternative approaches, such as using pseudowords, to avoid this limitation. Besides, complementary experiments could be conducted to examine the effect of exposure to other photic stimulation colors on performances. In particular, the superior effect of the binaural beat over the monaural beat is speculative and requires further testing. Regarding the CES-related results, repetition of the session with other voltages is also recommended. In effect, replication of this study with repetitions in the AVE and CES sessions will expand the findings. In addition, conducting the study with a larger sample size will provide more reliable results. Finally, incorporating a task specifically designed to measure subjects' processing speed could enhance the validity of the data.

Conclusion

In conclusion, despite the myriad influences of AVE and CES on brain activities, they cannot rival the effect of senses on attention and retention mechanisms. Tracking the role of senses, we implicated that multisensory instruction can permit

a direct effect on the velocity of attention and retention mechanisms. In fact, the primary aim of the AVE and CES devices is to alter CBF and neurotransmitter responses to overcome the arousal and overarousal states emitted from cognitive challenges and to harness negative excitements. However, these modifications are in association with the challenge of slowing down learners' mental agility and vigilance. It is also concluded that the five-sense combination drives positive results in RT, as well as in RA. In other words, multisensory inputs are committed to the consolidation of new pieces of information, exerting effective influences on attention and L2 vocabulary retention. As such, for effective comprehension and successful retention, creating a learning situation based on the employment of enriched sensory inputs can be a solution to improve learners' attention and retention processes.

Acknowledgments. Nasim Boustani was partially supported by a grant from Ferdowsi University of Mashhad (No. FUM-61021).

References

- Addante, R., Yousif, M., Valencia, R., Greenwood, C., & Marino, R.** (2021). Boosting brain waves improves memory. *Frontiers for Young Minds*, *9*, 605677.
- Allan, D.** (1992). *The Oxford quick placement test*. Oxford: Oxford University Press.
- Alloway, T. P., & Alloway, R. G.** (2012). The impact of engagement with social networking sites (SNSs) on cognitive skills. *Computers in Human Behavior*, *28*(5), 1748–1754.
- Alpert, R., & Haber, R. N.** (1960). Anxiety in academic achievement situations. *The Journal of Abnormal and Social Psychology*, *61*(2), 207–215.
- Aparecido-Kanzler, S., Cidral-Filho, F. J., & Prediger, R. D.** (2021). Effects of binaural beats and isochronic tones on brain wave modulation: Literature review. *Revista Mexicana de Neurociencia*, *22*(6), 238–247.
- Aston-Jones, G., Chiang, C., & Alexinsky, T.** (1991). Discharge of noradrenergic locus coeruleus neurons in behaving rats and monkeys suggests a role in vigilance. In C. D. Barnes & O. Pomeiano (Eds.), *Progress in brain research* (pp. 501–519). Amsterdam: Elsevier Science.
- Bahri Roudposhti, R., & Al Abdwani, T.** (2024). Executive function and learning outcomes: A systematic review. *Journal of Cognition, Emotion and Education*, *2*(2), 39–50.
- Basu, S., & Banerjee, B.** (2020). Prospect of brainwave entrainment to promote well-being in individuals: A brief review. *Psychological Studies*, *65*(3), 296–306.
- Battich, L., Fairhurst, M., & Deroy, O.** (2020). Coordinating attention requires coordinated senses. *Psychonomic Bulletin and Review*, *27*(6), 1126–1138.
- Ben-Haim, M. S., Williams, P., Howard, Z., Mama, Y., Eidels, A., & Algom, D.** (2016). The emotional Stroop task: Assessing cognitive performance under exposure to emotional content. *Journal of Visualized Experiments*, *112*, e53720.
- Binder, K. S., Tremblay, K. A., & Joseph, A.** (2020). Vocabulary accessibility and acquisition: Do you get more from a finacestor or a sociophite? *Journal of Research in Reading*, *43*(4), 395–416.
- Bolognini, N., Frassinetti, F., Serino, A., & Ládavas, E.** (2005). “Acoustical vision” of below threshold stimuli: Interaction among spatially converging audiovisual inputs. *Experimental Brain Research*, *160*(3), 273–282.
- Borsipour, B.** (2016). *Emotioncy and willingness to read: A case of Iranian EFL learners*. Mashhad: Ferdowsi University of Mashhad.
- Boustani, N., & Al Abdwani, T.** (2023). The effect of music on the communication ability: A case of vocabularies. *Journal of Business, Communication and Technology*, *2*(1), 1–12.
- Boustani, N., Pishghadam, R., & Shayesteh, S.** (2021). Multisensory input modulates P200 and L2 sentence comprehension: a one-week consolidation phase. *Frontiers in Psychology*, *12*, 746813.
- Bremner, D.** (2002). *Does stress damage the brain?* New York: W. W. Norton & Company.

- Chen, Y., Jin, P., & Ding, N.** (2020). The influence of linguistic information on cortical tracking of words. *Neuropsychologia*, *148*, 107640.
- Collura, T. F., & Siever, D.** (2009). Auditory-visual entrainment in relation to mental health and EEG. In T. H. Budzinsky, H. K. Budzinski, J. R. Evans & A. A. Abarbanel (Eds.), *Introduction to QEEG and neurofeedback: Advanced theory and applications* (2nd ed., pp. 195–224). Cambridge, MA: Academic Press.
- Crespo, A., Recuero, M., Galvez, G., & Begoña, A.** (2013). Effect of binaural stimulation on attention and EEG. *Archives of Acoustics*, *38*(4), 517–528.
- Dunn, R. S., & Dunn, K. J.** (1993). *Teaching secondary students through their individual learning styles: Practical approaches for grades 7–12*. Boston: Allyn & Bacon.
- Engel, A. K., Fries, P., & Singer, W.** (2001). Dynamic predictions: Oscillations and synchrony in top-down processing. *Nature Reviews Neuroscience*, *2*(10), 704–716.
- Ernst, M. O.** (2008). Multisensory integration: A late bloomer. *Current Biology*, *18*(12), R519–R521.
- Fox, P. T., & Raichle, M. E.** (1985). Stimulus rate determines regional brain blood flow in striate cortex. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*, *17*(3), 303–305.
- Frassinetti, F., Bolognini, N., & Làdavas, E.** (2002). Enhancement of visual perception by crossmodal visuo-auditory interaction. *Experimental Brain Research*, *147*(3), 332–343.
- Frederick, J. A., Lubar, J. F., Rasey, H. W., Brim, S. A., & Blackburn, J.** (1999). Effects of 18.5 Hz auditory and visual stimulation on EEG amplitude at the vertex. *Journal of Neurotherapy*, *3*(3–4), 23–28.
- Jensen, E.** (2009). *Enriching the brain: How to maximize every learner's potential*. Hoboken, NJ: John Wiley & Sons.
- Jensen, O., Kaiser, J., & Lachaux, J. P.** (2007). Human gamma-frequency oscillations associated with attention and memory. *Trends in Neurosciences*, *30*(7), 317–324.
- Kennerly, R. C.** (2006). *Changes in quantitative EEG and low resolution tomography following cranial electrotherapy stimulation*. Denton, TX: University of North Texas.
- Lakens, D.** (2022). Sample size justification. *Collabra: Psychology*, *8*(1), 33267.
- Lane, J. D., Kasian, S. J., Owens, J. E., & Marsh, G. R.** (1998). Binaural auditory beats affect vigilance performance and mood. *Physiology and Behavior*, *63*(2), 249–252.
- Lee, J., Lee, H., & Park, W.** (2019). Effects of cranial electrotherapy stimulation on electroencephalogram. *Journal of International Academy of Physical Therapy Research*, *10*(1), 1687–1694.
- Li, J., & Deng, S. W.** (2023). Facilitation and interference effects of the multisensory context on learning: A systematic review and meta-analysis. *Psychological Research*, *87*(5), 1334–1352.
- Li, Z., Wu, S., & Liu, D.** (2023). The effects of context, story-type, and language proficiency on EFL word learning and retention from reading. *International Review of Applied Linguistics in Language Teaching*. <https://doi.org/10.1515/iral-2023-0155>.
- Liss, S., & Liss, B.** (1996). Physiological and therapeutic effects of high frequency electrical pulses. *Integrative Physiological and Behavioral Science*, *31*(2), 88–94.
- Lovelace, C. T., Stein, B. E., & Wallace, M. T.** (2003). An irrelevant light enhances auditory detection in humans: A psychophysical analysis of multisensory integration in stimulus detection. *Cognitive Brain Research*, *17*(2), 447–453.
- MacLeod, C. M.** (1991). Half a century of research on the Stroop effect: An integrative review. *Psychological Bulletin*, *109*(2), 163–203.
- Matthews, G., Warm, J. S., Reinerman-Jones, L. E., Langheim, L. K., Washburn, D. A., & Tripp, L.** (2010). Task engagement, cerebral blood flow velocity, and diagnostic monitoring for sustained attention. *Journal of Experimental Psychology: Applied*, *16*(2), 187–203.
- Oldfield, R. C.** (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113.
- On, F. R., Jailani, R., Norhazman, H., & Zaini, N. M.** (2013, March). *Binaural beat effect on brainwaves based on EEG*. IEEE 9th International Colloquium on Signal Processing and its Applications (pp. 339–343). IEEE. <http://doi.org/10.1109/CSPA.2013.6530068>
- Operators Manual** (2021). [Operator's manual] Usage and care of your AVE and CES device. <https://mindalive.com/manuals/>.

- Pahor, A., Collins, C., Smith-Peirce, R. N., Moon, A., Stavropoulos, T., Silva, I., Peng, E., Jaeggi, S. M., & Seitz, A. R. (2021). Multisensory facilitation of working memory training. *Journal of Cognitive Enhancement*, 5(3), 386–395.
- Park, J., Kwon, H., Kang, S., & Lee, Y. (2018, October). *The effect of binaural beat-based audiovisual stimulation on brain waves and concentration* [Conference presentation]. International Conference on Information and Communication Technology Convergence (ICTC) (pp. 420–423). IEEE. <http://doi.org/10.1109/ICTC.2018.8539512>
- Parris, B. A., Hasshim, N., Wadsley, M., Augustinova, M., & Ferrand, L. (2021). The loci of Stroop effects: A critical review of methods and evidence for levels of processing contributing to color-word Stroop effects and the implications for the loci of attentional selection. *Psychological Research*, 86, 1029–1053.
- Pishghadam, R. (2015, October). *Emotioncy in language education: From exvovement to involvement* [Conference presentation]. The 2nd Conference of Interdisciplinary Approaches to Language Teaching, Literature, and Translation Studies, Mashhad, Iran.
- Pishghadam, R., Al Abdwani, T., Al-Badi, A. H., Boustani, N., & Shayesteh, S. (2024). An investigation into online content's readability in light of a brain-based model: A case from the Sultanate of Oman. *Applied Research on English Language*, 13(3), 19–40.
- Pishghadam, R., Daneshvarfard, F., & Shayesteh, S. (2021). Oscillatory neuronal dynamics during L2 sentence comprehension: The effects of sensory enrichment and semantic incongruency. *Language, Cognition and Neuroscience*, 36(8), 903–920.
- Pishghadam, R., Jajarmi, H., & Shayesteh, S. (2021). Sense combinations influence the neural mechanism of L2 comprehension in semantically violated sentences: Insights from emotioncy. *Journal of Neurolinguistics*, 58, 100962.
- Pishghadam, R., Makiabadi, H., Zabetipour, M., Abbasnejad, H., Firoozian Pooresfahani, A., & Shayesteh, S. (2020). Development, validation and application of an inventory on emo-sensory intelligence. *Teaching English Language*, 14(2), 173–216.
- Pishghadam, R., Shakebaee, G., & Shayesteh, S. (2018). Introducing cultural weight as a tool of comparative analysis: An emotioncy-based study of social class. *Humanities Diliman*, 15(1), 139–158.
- Pishghadam, R., & Shayesteh, S. (2016). Emotioncy: A post-linguistic approach toward vocabulary learning and retention. *Sri Lanka Journal of Social Sciences*, 39(1), 27–36.
- Ralph, B. C., Thomson, D. R., Seli, P., Carriere, J. S., & Smilek, D. (2015). Media multitasking and behavioral measures of sustained attention. *Attention, Perception, and Psychophysics*, 77(2), 390–401.
- Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T., & Yiend, J. (1997). “Oops!”: Performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, 35(6), 747–758.
- Robison, M. K., Obulasetty, M., Blais, C., Wingert, K. M., & Brewer, G. A. (2022). The effect of binaural beat stimulation on sustained attention. *Psychological Research*, 86(3), 808–822.
- Rosenberg, L. (2015). *The effects of multisensory, explicit, and systematic instructional practices on elementary school students with learning impairments in encoding and oral reading*. Boston, MA: Northeastern University Library.
- Sansevere, K. S., Wooten, T., McWilliams, T., Peach, S., Hussey, E. K., Bruny , T. T., & Ward, N. (2022). Self-reported outcome expectations of non-invasive brain stimulation are malleable: A registered report that replicates and extends Rabipour et al. (2017). *Journal of Cognitive Enhancement*, 6, 496–513.
- Sappey-Marinier, D., Calabrese, G., Fein, G., Hugg, J. W., Biggins, C., & Weiner, M. W. (1992). Effect of photic stimulation on human visual cortex lactate and phosphates using 1H and 31P magnetic resonance spectroscopy. *Journal of Cerebral Blood Flow and Metabolism*, 12(4), 584–592.
- Segalowitz, N., & Frenkiel-Fishman, S. (2005). Attention control and ability level in a complex cognitive skill: Attention shifting and second-language proficiency. *Memory and Cognition*, 33(4), 644–653.
- Schroeder, M. J., & Barr, R. E. (2001). Quantitative analysis of the electroencephalogram during cranial electrotherapy stimulation. *Clinical Neurophysiology*, 112(11), 2075–2083.
- Shayesteh, S. (2019). *The neurocognitive effects of foreign language comprehension in response to the emotioncy-based language instruction (EBLI): Evidence from event-related brain potentials (ERPs) during semantic processing of a sentence*. Mashhad: Ferdowsi University of Mashhad.
- Shayesteh, S., Pishghadam, R., & Khodaverdi, A. (2020). FN400 and LPC responses to different degrees of sensory involvement: A study of sentence comprehension. *Advances in Cognitive Psychology*, 16(1), 45–58.

- Siever, D.** (2007). Audio-visual entrainment: History, physiology, and clinical studies. In J. R. Evans (Ed.), *Handbook of neurofeedback: Dynamics and clinical applications* (pp. 155–183). New York: Hayworth Medical.
- Siever, D.** (2008). Applying audio-visual entrainment technology for attention and learning (Part 3). *Biofeedback Magazine*, *31*(4), 1–15.
- Siever, D.** (2012). Audio-visual entrainment: A novel way of boosting grades and socialization while reducing stress in the typical college student. *Biofeedback*, *40*(3), 115–124.
- Southworth, S.** (1999). A study of the effects of cranial electrical stimulation on attention and concentration. *Integrative Physiological and Behavioral Science*, *34*(1), 43–53.
- Stein, B. E., Stanford, T. R., & Rowland, B. A.** (2009). The neural basis of multisensory integration in the midbrain: Its organization and maturation. *Hearing Research*, *258*(1–2), 4–15.
- Stroop, J. R.** (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, *18*, 643–662.
- Talsma, D., Senkowski, D., Soto-Faraco, S., & Woldorff, M. G.** (2010). The multifaceted interplay between attention and multisensory integration. *Trends in Cognitive Sciences*, *14*(9), 400–410.
- Wechsler, D.** (1981). *Wechsler adult intelligence scale: WAIS-R manual*. New York, NY: Psychological Corporation.
- Wuchrer, V.** (2009). *Study on memory and concentration*. Edmonton, Alberta, Canada: Mind Alive Inc.

Cite this article: Pishghadam, R., Boustani, N., Gerwien, J., Shayesteh, S., & Al Abdwani, T. (2024). Audio-visual entrainment, cranio-electro stimulation, and sensory involvement: rival effects on attention and L2 vocabulary retention. *Applied Psycholinguistics* *45*, 1067–1085. <https://doi.org/10.1017/S0142716424000419>