



# Some examples of noncommutative projective Calabi–Yau schemes

Yuki Mizuno

**Abstract.** In this article, we construct some examples of noncommutative projective Calabi–Yau schemes by using noncommutative Segre products and quantum weighted hypersurfaces. We also compare our constructions with commutative Calabi–Yau varieties and examples constructed in Kanazawa (2015, *Journal of Pure and Applied Algebra* 219, 2771–2780). In particular, we show that some of our constructions are essentially new examples of noncommutative projective Calabi–Yau schemes.

## 1 Introduction

Calabi–Yau varieties are rich objects and play an important role in mathematics and physics. In noncommutative algebraic geometry, (skew) Calabi–Yau algebras are often treated as noncommutative analogs of Calabi–Yau varieties. Calabi–Yau algebras have a deep relationship with quiver algebras (see [8, 31]). For example, many known Calabi–Yau algebras are constructed by using quiver algebras. They are also used to characterize Artin–Schelter regular algebras (see [24, 25]). In particular, a connected graded algebra  $A$  over a field  $k$  is Artin–Schelter regular if and only if  $A$  is skew Calabi–Yau.

On the other hand, a triangulated subcategory of the derived category of a cubic fourfold in  $\mathbb{P}^5$ , which is obtained by some semiorthogonal decompositions, has the 2-shift functor [2] as the Serre functor. Moreover, the structure of Hochschild (co)homology is the same as that of a projective K3 surface (see [13]). However, some such categories are not obtained as the derived categories of coherent sheaves of projective K3 surfaces and called noncommutative K3 surfaces.

Artin and Zhang constructed a framework of noncommutative projective schemes in [1], which are defined from noncommutative graded algebras. In this framework, we can think of Artin–Schelter regular algebras as noncommutative analogs of projective spaces, which are called quantum projective spaces. Our objective is to produce examples of noncommutative projective Calabi–Yau schemes that are not obtained from commutative Calabi–Yau varieties. In the future, it would be an interesting question to compare the derived category of a noncommutative projective Calabi–Yau scheme created in the framework of Artin–Zhang’s noncommutative projective

---

Received by the editors October 2, 2023; revised December 10, 2023; accepted February 3, 2024.

Published online on Cambridge Core February 8, 2024.

This work is supported by Grant-in-Aid for JSPS Fellows (Grant No. 22KJ2923)

AMS subject classification: 14A22, 14J32.

Keywords: Noncommutative algebraic geometry, Noncommutative projective schemes, Calabi–Yau varieties.



schemes with a noncommutative K3 surface obtained as a triangulated subcategory of the derived category of a cubic fourfold.

As the definition of noncommutative projective Calabi–Yau schemes, we adopt the definition introduced by Kanazawa [12]. His definition is a direct generalization of the definition of commutative Calabi–Yau varieties to noncommutative projective schemes. He also constructed the first examples of noncommutative projective Calabi–Yau schemes that are not isomorphic to commutative Calabi–Yau varieties as hypersurfaces of quantum projective spaces. Recently, some examples constructed by Kanazawa play an important role in noncommutative Donaldson–Thomas theory (see [14, 15]).

In this paper, we construct new examples of noncommutative projective Calabi–Yau schemes by using noncommutative Segre products and weighted hypersurfaces. There are many known examples of Calabi–Yau varieties in algebraic geometry. Some of them are complete intersections in products of projective spaces. Moreover, Reid gave a list of Calabi–Yau surfaces, which are hypersurfaces in weighted projective spaces (see [11, Table 1 in Section 13.3], [23, Theorem 4.5]). Motivated by these two facts, we construct noncommutative analogs of the two types of examples of Calabi–Yau varieties (Theorems 3.3 and 3.15) in Section 3.

In order to prove that a noncommutative projective scheme is Calabi–Yau, we use the methods of Kanazawa. However, they are not sufficient because the algebras we treat are more complicated than the ones he considered. In order to construct noncommutative projective Calabi–Yau schemes as noncommutative analogs of complete intersections in Segre products, we perform a more detailed analysis of noncommutative projective schemes defined by  $\mathbb{Z}^2$ -graded algebras, which were studied by Van Rompay [32]. A different approach to noncommutative Segre products is also studied in [9]. In order to construct noncommutative projective Calabi–Yau schemes as noncommutative analogs of weighted hypersurfaces, we consider quotients of weighted quantum polynomial rings. In commutative algebraic geometry, the category  $\text{Coh}(\text{Proj}(k[x_0, \dots, x_n]))$  of coherent sheaves on the projective spectrum  $\text{Proj}(k[x_0, \dots, x_n])$  of a weighted polynomial ring is not necessarily equivalent to  $\text{qgr}(k[x_0, \dots, x_n])$ , where  $\text{qgr}(k[x_0, \dots, x_n])$  is the quotient category associated with  $k[x_0, \dots, x_n]$  constructed in [1]. In fact,  $\text{qgr}(k[x_0, \dots, x_n])$  is equivalent to the category of coherent sheaves on a weighted projective space constructed as a Deligne–Mumford stack. Moreover,  $\text{qgr}(k[x_0, \dots, x_n])$  is thought of as a nonsingular model of  $\text{Proj}(k[x_0, \dots, x_n])$  (see [28, Example 4.9]). We use this idea to construct new noncommutative projective Calabi–Yau schemes. In addition, it should be noted that local structures of noncommutative projective schemes of quotients of weighted quantum polynomial rings are somewhat complicated. An analysis of the local structures was performed by Smith [28]. We show that the local structure obtained in [28] is described by the notion of quasi-Veronese algebras introduced by Mori [18].

In Section 4, we compare our constructions from weighted hypersurfaces in Section 3 with commutative Calabi–Yau varieties and the first examples constructed in [12], focusing on noncommutative projective Calabi–Yau schemes of dimensions 2. We show that some of our constructions in Section 3 are not isomorphic to any of the commutative Calabi–Yau varieties and the first examples constructed in [12] (Proposition 4.9). When we consider moduli spaces of point modules of noncommutative

projective schemes obtained from weighted hypersurfaces in Section 3, there is a problem, which is that in general weighted quantum polynomial rings are not generated in degree 1. So, the notion of point modules is not necessarily useful in this case. In this paper, we use theories of closed points studied in [19, 26, 27], etc. A different approach to closed points of weighted quantum polynomial rings is studied in [29]. The notion of point modules defined in [29] corresponds to those of ordinary and thin points in [19]. To show that some of our constructions are not isomorphic to the examples obtained in [12], we use Morita theory of noncommutative schemes, which is established in [5] (see also [1, Section 6]). In the theory, we need to calculate the centers of noncommutative rings. By using these calculations, we can do a detailed analysis and some classifications of noncommutative projective Calabi–Yau surfaces.

## 2 Preliminaries

**Notation and Terminology 2.1** In this article,  $k$  denotes an algebraically closed field of characteristic 0. We suppose  $\mathbb{N}$  contains 0. Let  $A$  be a  $k$ -algebra, let  $M$  be an  $A$ -bimodule, and let  $\psi, \phi$  be algebra automorphisms of  $A$ . Then, we denote the associated  $A$ -bimodule by  ${}^\psi M^\phi$ , i.e.,  ${}^\psi M^\phi = M$  as  $k$ -modules and the new bimodule structure is given by  $a * m * b := \psi(a)m\phi(b)$  for all  $a, b \in A$  and all  $m \in M$ . Let  $\mathcal{C}$  be a  $k$ -linear abelian category. We denote the global dimension of  $\mathcal{C}$  by  $\text{gl.dim}(\mathcal{C})$ . An  $\mathbb{N}$ -graded  $k$ -algebra  $A$  is connected if  $A_0 = k$ .

For any  $\mathbb{N}$ -graded  $k$ -algebra  $A = \bigoplus_{i=0}^{\infty} A_i$ , we denote the category of graded right  $A$ -modules (resp. finitely generated graded right  $A$ -modules) by  $\text{Gr}(A)$  (resp.  $\text{gr}(A)$ ). Let  $M \in \text{Gr}(A)$  and  $A^\circ$  be the opposite algebra of  $A$ . We define the Matlis dual  $M^* \in \text{Gr}(A^\circ)$  by  $M_i^* := \text{Hom}_k(M_{-i}, k)$  and the shift  $M(n) \in \text{Gr}(A)$  by  $M(n)_i := M_{i+n}$  ( $i, n \in \mathbb{Z}$ ). For  $M, N \in \text{Gr}(A)$ , we write  $\text{Hom}_A(M, N) := \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{\text{Gr}(A)}(M, N(n)) \in \text{Gr}(A)$ . For  $M \in \text{Gr}(A)$  and a homogeneous element  $m \in M$ , we denote the degree of  $m$  by  $\deg(m)$ . We define the truncation  $M_{\geq n} := \bigoplus_{i \geq n} M_i \in \text{Gr}(A)$  ( $n \in \mathbb{Z}$ ). An element  $m \in M$  is called torsion if  $mA_{\geq n} = 0$  for  $n \gg 0$ . We say  $M$  is a torsion module if any element of  $M$  is torsion. We denote the subcategory of torsion modules in  $\text{Gr}(A)$  (resp.  $\text{gr}(A)$ ) by  $\text{Tor}(A)$  (resp.  $\text{tor}(A)$ ).

**Definition 2.2** [1, Section 2] Let  $A$  be a right Noetherian  $\mathbb{N}$ -graded  $k$ -algebra. We define the quotient categories  $\text{QGr}(A) := \text{Gr}(A)/\text{Tor}(A)$  and  $\text{qgr}(A) := \text{gr}(A)/\text{tor}(A)$ . We denote the projection functor by  $\pi$  and its right adjoint functor by  $\omega$ . The general (resp. Noetherian) projective scheme of  $A$  is defined as  $\text{Proj}(A) := (\text{QGr}(A), \pi(A))$  (resp.  $\text{proj}(A) := (\text{qgr}(A), \pi(A))$ ).

**Definition 2.3** [1, Section 2], [26, Chapter 3] A quasi-scheme over  $k$  is a pair  $(\mathcal{C}, \mathcal{O})$ , where  $\mathcal{C}$  is a  $k$ -linear abelian category and  $\mathcal{O}$  is an object in  $\mathcal{C}$ . A morphism from a quasi-scheme  $(\mathcal{C}, \mathcal{O})$  to another quasi-scheme  $(\mathcal{C}', \mathcal{O}')$  is a pair  $(F, \varphi)$  consisting of a  $k$ -linear right exact functor  $F : \mathcal{C} \rightarrow \mathcal{C}'$  and an isomorphism  $\varphi : F(\mathcal{O}) \xrightarrow{\sim} \mathcal{O}'$ . We call  $(F, \varphi)$  an isomorphism if  $F$  is an equivalence.

When  $A$  is as in Definition 2.2, we think of  $\text{proj}(A) = (\text{qgr}(A), \pi(A))$  as a quasi-scheme. For any (commutative) Noetherian scheme  $X$ ,  $(\text{Coh}(X), \mathcal{O}_X)$  is also a quasi-scheme. From this observation, we regard  $X$  as a quasi-scheme.

**Definition 2.4** [30, Section 4], [35, Section 4] Let  $A, B$  be  $\mathbb{N}$ -graded  $k$ -algebras and let  $m_A$  be  $A_{\geq 1}$ . We define the torsion functor  $\Gamma_{m_A} : \text{Gr}(A \otimes_k B^\circ) \rightarrow \text{Gr}(A \otimes_k B^\circ)$  by  $\Gamma_{m_A}(M) := \{m \in M \mid m A_{\geq n} = 0 \text{ for some } n \in \mathbb{N}\}$ . We write  $H_{m_A}^i := R^i \Gamma_{m_A}$ .

**Definition 2.5** [30, Definitions 6.1 and 6.2], [35, Definitions 3.3 and 4.1] Let  $A$  be a right and left Noetherian connected  $\mathbb{N}$ -graded  $k$ -algebra and let  $A^e$  be the enveloping algebra of  $A$ . Let  $R$  be an object of  $D^b(\text{Gr}(A^e))$ . Then,  $R$  is called a dualizing complex of  $A$  if (1)  $R$  has finite injective dimension over  $A$  and  $A^\circ$ , (2) the cohomologies of  $R$  are finitely generated as both  $A$  and  $A^\circ$ -modules, (3) the natural morphisms  $A \rightarrow \text{RHom}_A(R, R)$  and  $A \rightarrow \text{RHom}_{A^\circ}(R, R)$  are isomorphisms in  $D^b(\text{Gr}(A^e))$ . Moreover,  $R$  is called balanced if  $\text{R}\Gamma_{m_A}(R) \simeq A^*$  and  $\text{R}\Gamma_{m_{A^\circ}}(R) \simeq A^*$  in  $D^b(\text{Gr}(A^e))$ .

### 3 Calabi–Yau conditions

**Definition 3.1** [12, Section 2.2] Let  $A$  be a connected right Noetherian  $\mathbb{N}$ -graded  $k$ -algebra. Then,  $\text{proj}(A)$  is a noncommutative projective Calabi–Yau scheme of dimension  $n$  if the global dimension of  $\text{qgr}(A)$  is  $n$  and the Serre functor of the derived category  $D^b(\text{qgr}(A))$  is the  $n$ -shift functor  $[n]$ .

**Remark 3.2** Actually, we do not need the condition that the global dimension of  $\text{qgr}(A)$  is  $n$ . If the Serre functor of the derived category  $D^b(\text{qgr}(A))$  is the  $n$ -shift functor  $[n]$ , then we can easily show that this condition holds. However, when we prove the existence of the Serre functor of  $D^b(\text{qgr}(A))$ , we essentially need the condition that the global dimension of  $\text{qgr}(A)$  is  $n$  (cf. [6, Theorem A.4 and Corollary A.5], Lemma 3.10).

#### 3.1 $\mathbb{Z}^2$ -graded algebras and Segre products

In commutative algebraic geometry, a smooth complete intersection  $X \subset \mathbb{P}^n \times \mathbb{P}^m$  of bidegrees  $(n+1, 0)$  and  $(0, m+1)$  provides a Calabi–Yau variety. We also have a little more complicated example that gives a Calabi–Yau variety. That is a smooth complete intersection of bidegrees  $(n, 0)$  (resp.  $(n+1, 0)$ ) and  $(1, n+1)$  in  $\mathbb{P}^n \times \mathbb{P}^n$  (resp.  $\mathbb{P}^{n+1} \times \mathbb{P}^n$ ). We construct noncommutative analogs of these examples.

Let  $C$  be an  $\mathbb{N}^2$ -graded  $k$ -algebra. We denote the category of  $\mathbb{Z}^2$ -graded right  $C$ -modules (resp. finitely generated  $\mathbb{Z}^2$ -graded right  $C$ -modules) by  $\text{BiGr}(C)$  (resp.  $\text{bigr}(C)$ ). Let  $M \in \text{BiGr}(C)$ . We denote by  $C^\circ$  (resp.  $C^e$ ) the opposite (resp. enveloping) algebra of  $C$ . We define the Matlis dual  $M^* \in \text{BiGr}(C^\circ)$  by  $M_{i,j}^* := \text{Hom}_k(M_{-i,-j}, k)$  and the shift  $M(n, m) \in \text{BiGr}(C)$  by  $M(m, n)_{i,j} := M_{i+m, j+n}(m, n, i, j \in \mathbb{Z})$ . For  $M, N \in \text{BiGr}(C)$ , we write  $\text{Hom}_C(M, N) := \bigoplus_{m, n \in \mathbb{Z}} \text{Hom}_{\text{BiGr}(C)}(M, N(m, n))$ . For a bihomogeneous element  $m \in M$ , we denote the bidegree of  $m$  by  $\text{bideg}(m)$ .

Let  $M \in \text{BiGr}(C)$ . We define the truncation  $M_{\geq n, \geq n} := \bigoplus_{i \geq n, j \geq n} M_{i,j} \in \text{BiGr}(C)$  ( $n \in \mathbb{Z}$ ). We say  $m \in M$  is torsion if  $m C_{\geq n, \geq n} = 0$  for  $n \gg 0$ . If all  $m \in M$  are torsion, then  $M$  is called a torsion  $C$ -module. We denote the category of  $\mathbb{Z}^2$ -graded torsion  $C$ -modules by  $\text{Tor}(C)$ . We also define  $\text{tor}(C)$  to be the intersection of  $\text{bigr}(C)$  and

$\text{Tor}(C)$ . When we assume that  $C$  is right Noetherian, we have the quotient categories  $\text{QBiGr}(C) := \text{BiGr}(C)/\text{Tor}(C)$  and  $\text{qbigr}(C) := \text{bigr}(C)/\text{tor}(C)$  (cf. [32, Section 2]). We denote the projection functor by  $\pi$  and its right adjoint functor by  $\omega$ . We can define the general (resp. Noetherian) projective scheme  $\text{Proj}(C)$  (resp.  $\text{proj}(C)$ ) associated with  $C$  and the notion of noncommutative projective Calabi–Yau schemes as in the case of  $\mathbb{N}$ -graded algebras.

Let  $D$  be an  $\mathbb{N}^2$ -graded algebra. We take the tensor product  $C \otimes_k D^\circ$  of  $C$  and  $D^\circ$  over  $k$ . We think of  $C \otimes_k D^\circ$  as an  $\mathbb{N}^2$ -graded algebra by  $(C \otimes_k D^\circ)_{i,j} := \bigoplus_{i_1+i_2=i, j_1+j_2=j} C_{i_1, j_1} \otimes_k D_{i_2, j_2}^\circ$ . We define  $m_{C_{++}} := C_{\geq 1, \geq 1}$  and the torsion functor  $\Gamma_{m_{C_{++}}} : \text{BiGr}(C \otimes_k D^\circ) \rightarrow \text{BiGr}(C \otimes_k D^\circ)$  by  $\Gamma_{m_{C_{++}}}(M) := \{m \in M \mid m_{C_{\geq n, \geq n}} = 0 \text{ for some } n \in \mathbb{N}\}$ . We write  $m_C := \bigoplus_{i+j \geq 1} C_{i,j}$  and define another torsion functor  $\Gamma_{m_C} : \text{BiGr}(C \otimes_k D^\circ) \rightarrow \text{BiGr}(C \otimes_k D^\circ)$  by  $\Gamma_{m_C}(M) := \{m \in M \mid m_{C_{\geq n}} = 0 \text{ for some } n \in \mathbb{N}\}$ , where  $C_{\geq n} := \bigoplus_{i+j \geq n} C_{i,j} \in \text{BiGr}(C)$ . See [25, Section 3] for details of  $\Gamma_{m_C}$ . We write  $H_{m_{C_{++}}}^i := R^i \Gamma_{m_{C_{++}}}$  and  $H_{m_C}^i := R^i \Gamma_{m_C}$ . The reason we define the functor  $\Gamma_{m_{C_{++}}}$  is that we can describe the Serre duality in  $\text{D}^b(\text{qbigr}(C))$  by using  $R\Gamma_{m_{C_{++}}}$  (cf. Lemma 3.10). However, it is not easy to calculate the functor  $R\Gamma_{m_{C_{++}}}$  directly. The reason we define the functor  $\Gamma_{m_C}$  is that we can use the theory of  $\mathbb{Z}$ -graded modules to calculate  $R\Gamma_{m_C}$  and we can reduce the calculation of  $R\Gamma_{m_{C_{++}}}$  to that of  $R\Gamma_{m_C}$  (cf. Lemma 3.6, the proof of Theorem 3.3).

**Theorem 3.3** *Let  $A := k\langle x_0, \dots, x_n \rangle / (x_j x_i - q_{ji} x_i x_j)_{i,j}$ ,  $B := k\langle y_0, \dots, y_m \rangle / (y_j y_i - q'_{ji} y_i y_j)_{i,j}$  and  $C := A \otimes_k B$ , where  $q_{ji}, q'_{ji} \in k^\times$  for all  $i, j$ . We regard  $C$  as an  $\mathbb{N}^2$ -graded algebra with  $\text{bideg}(x_i) = (1, 0)$  and  $\text{bideg}(y_i) = (0, 1)$  for all  $i$ .*

- (1) *Let  $f := \sum_{i=0}^n x_i^{n+1}$  and  $g := \sum_{i=0}^m y_i^{m+1}$ . We assume that (i)  $q_{ii} = q_{ij} q_{ji} = q_{ij}^{n+1} = 1$  for all  $i, j$ , (ii)  $q'_{ii} = q'_{ij} q'_{ji} = q'_{ij}^{m+1} = 1$  for all  $i, j$ .*

*Then,  $\text{proj}(C/(f, g))$  is a noncommutative projective Calabi–Yau scheme of dimension  $(n + m - 2)$  if and only if  $\prod_{i=0}^n q_{ij}$  and  $\prod_{i=0}^m q'_{ij}$  are independent of  $j$ , respectively.*

- (2) *Suppose that  $m = n + 1$  (resp.  $m = n$ ) and  $q'_{ij} = 1$  for all  $i, j$ . Let  $f := \sum_{i=0}^n x_i^{n+1} y_i$  and  $g := \sum_{i=0}^{n+1} y_i^{n+1}$  (resp.  $\sum_{i=0}^n y_i^n$ ). We assume that  $q_{ii} = q_{ij} q_{ji} = q_{ij}^{n+1} = 1$  for all  $i, j$ .*

*Then,  $\text{proj}(C/(f, g))$  is a noncommutative projective Calabi–Yau scheme of dimension  $(2n - 1)$  (resp.  $(2n - 2)$ ) if and only if  $\prod_{i=0}^n q_{ij}$  is independent of  $j$ .*

**Notation 3.4** For simplicity, we denote the bidegrees of  $f, g$  in the theorem by  $(d_0, d_1), (e_0, e_1)$ , respectively.

**Remark 3.5** •  $f, g$  are central elements in  $C$  because of the choices of  $\{q_{ij}\}, \{q'_{ij}\}$ .

- We have  $n + m - 2 = d_0 + d_1 + e_0 + e_1 - 4$  in (1). We have  $2n - 1$  (resp.  $2n - 2$ ) =  $d_0 + d_1 + e_0 + e_1 - 4$  also in (2).
- In (2) of the theorem, even if we do not assume  $q'_{ij} = 1$ , the condition for  $f, g$  to be central in  $C$  implies  $q'_{ij} = 1$  for all  $i, j$  after all.
- In the theorem, all equations appearing except for  $g$  of (2) are Fermat-type equations.

To prove the theorem, we need to show some lemmas. Perhaps some experts may know the following lemmas. However, to the best of the author's knowledge, there are no references written on those lemmas, so the proofs are given below. In addition, the following proofs do not depend on whether (1) or (2) in the theorem is considered (except for Lemma 3.8).

**Lemma 3.6** Let  $\mathcal{R} := \pi(\mathrm{R}\Gamma_{m_{C/(f,g)_{++}}}(C/(f,g))^*)$  and  $\mathcal{R}' := \pi(\mathrm{R}\Gamma_{m_{C/(f,g)}}(C/(f,g))^*)$ . Then, the functors  $-\otimes^{\mathbb{L}} \mathcal{R}$  and  $-\otimes^{\mathbb{L}} \mathcal{R}'[-1]$  between  $\mathrm{D}(\mathrm{QBiGr}(C/(f,g)))$  and itself are naturally isomorphic.

**Proof** Let  $I_1, I_2$  be the ideals generated by  $\{x_0, \dots, x_n\}, \{y_0, \dots, y_m\}$ , respectively. Then, we have  $m_{C/(f,g)_{++}} = I_1 \cap I_2$ ,  $m_{C/(f,g)} = I_1 + I_2$  and have the following long exact sequence in  $\mathrm{BiGr}(C/(f,g)^e)$

$$\cdots \rightarrow H_{m_{C/(f,g)}}^i(C/(f,g)) \rightarrow H_{I_1}^i(C/(f,g)) \oplus H_{I_2}^i(C/(f,g)) \rightarrow H_{m_{C/(f,g)_{++}}}^i(C/(f,g)) \rightarrow \cdots$$

by using the Mayer–Vietoris sequence in the sense of [4, Chapter 3], where  $\Gamma_{I_j} (j = 1, 2)$  is defined not by using the degrees of  $I_j$  but by using powers of  $I_j$  (i.e.,  $\Gamma_{I_j}(M) := \{m \in M \mid mI_j^n = 0 \text{ for some } n\}$ ). Note that we can use the Mayer–Vietoris sequence in our case because  $I_1, I_2$  are generated by normal elements and this implies that  $I_1, I_2$  satisfy Artin–Rees property. We also have the exact triangle in  $\mathrm{D}(\mathrm{BiGr}(C/(f,g)^e))$

$$\mathrm{R}\Gamma_{m_{C/(f,g)}}(C/(f,g)) \rightarrow \mathrm{R}\Gamma_{I_1}(C/(f,g)) \oplus \mathrm{R}\Gamma_{I_2}(C/(f,g)) \rightarrow \mathrm{R}\Gamma_{m_{C/(f,g)_{++}}}(C/(f,g)).$$

Moreover,  $H_{I_1}^i(C/(f,g))^*$  and  $H_{I_2}^i(C/(f,g))^*$  are torsion modules for  $m_{C/(f,g)_{++}}$  from Sub-Lemma 3.7. So, the cohomologies of  $\mathrm{R}\Gamma_{I_1}(C/(f,g))^* \oplus \mathrm{R}\Gamma_{I_2}(C/(f,g))^*$  are torsion. From this result, the above triangle gives an isomorphism between  $\mathcal{R}$  and  $\mathcal{R}'[-1]$  after taking the dual and applying  $\pi$ . Hence, we get the claim. ■

**Sub-Lemma 3.7** Let  $I_1, I_2$  be as in the proof of Lemma 3.6.  $H_{I_1}^i(C/(f,g))^*$  and  $H_{I_2}^i(C/(f,g))^*$  are torsion modules for  $m_{C/(f,g)_{++}}$  for any  $i$ .

**Proof** We only show that  $H_{I_1}^i(C/(f,g))^*$  are torsion modules for  $m_{C/(f,g)_{++}}$ . We can show that  $H_{I_2}^i(C/(f,g))^*$  are torsion in the same way.

First, we prove that  $H_{I_1}^i(C)^*$  is torsion. We have  $\Gamma_{I_1} = \Gamma_{I_1^{n+1}}$ . Moreover, if  $J_1$  is the ideal generated by  $x_0^{n+1}, \dots, x_n^{n+1}$ , then we have  $\Gamma_{I_1^{n+1}} = \Gamma_{J_1}$ . Note that  $x_0^{n+1}, \dots, x_n^{n+1}$  are central elements in  $C$  from the choice of  $\{q_{ij}\}$ .

Let  $M \in \mathrm{Gr}(C)$  be injective. Then, we have a surjective localization map  $M \rightarrow M[x_i^{-(n+1)}]$  for any  $i$  and  $\Gamma_{J_1}(M)$  is injective in  $\mathrm{Gr}(C)$  because  $J_1$  satisfies Artin–Rees property (cf. [34, Example 3.13], [7, Lemma A1.4]). When  $M'$  is injective in  $\mathrm{Gr}(C^e)$ , then  $M'$  is injective in  $\mathrm{Gr}(C)$ , where  $\mathrm{Res}_C : \mathrm{Gr}(C^e) \rightarrow \mathrm{Gr}(C)$  is the restriction functor (see [35, Lemma 2.1]). Thus, we can calculate  $\mathrm{Res}_C(H_{J_1}^i(C))$  by using a Čech complex  $\mathcal{C}(x_0^{n+1}, \dots, x_n^{n+1}; C)$  (cf. [17, Chapter 2, 3], [7, Theorem A1.3]). Then, we have  $\mathcal{C}(x_0^{n+1}, \dots, x_n^{n+1}; C) = \mathcal{C}(x_0^{n+1}, \dots, x_n^{n+1}; A) \otimes_k B$ . This induces that  $\mathrm{Res}_C(H_{J_1}^i(C)) \simeq H_{m_A}^i(A) \otimes_k B$ . Because  $H_{m_A}^i(A)_{>0} = 0$  (see [12, Proposition 2.4]),  $H_{J_1}^i(C)^* \simeq H_{I_1}^i(C)^*$  is torsion.

Finally, we consider the exact sequences of  $C$ -bimodules

$$(3.1.1) \quad 0 \rightarrow C(-d_0, -d_1) \xrightarrow{\times f} C \rightarrow C/(f) \rightarrow 0,$$

$$(3.1.2) \quad 0 \rightarrow C/(f)(-e_0, -e_1) \xrightarrow{\times g} C/(f) \rightarrow C/(f, g) \rightarrow 0.$$

Then, we take the long exact sequence for  $\Gamma_{I_1}$  and we get the claim since  $H_{I_1}^i(C)^*$  is torsion.  $\blacksquare$

**Lemma 3.8**  $\text{gl.dim}(\text{qbigr}(C/(f, g))) = d_0 + d_1 + e_0 + e_1 - 4.$

**Proof** We show the proposition only in (1) of the theorem. In (2) of the theorem, the proposition can be shown in the same way (cf. Remark 3.9). We consider a bigraded (commutative) algebra  $D := k[s_0, \dots, s_n, t_0, \dots, t_m]/(\sum_{i=0}^n s_i, \sum_{i=0}^m t_i)$  with  $s_i = x_i^{n+1}$ ,  $t_i = y_i^{m+1}$  and the projective spectrum  $\text{biProj}(D)$  in the sense of [10, Section 1]. Then,  $C/(f, g)$  is a finite  $D$ -module. So,  $\text{qbigr}(C/(f, g))$  can be thought of as the category of modules over a sheaf  $\mathcal{A}$  of  $\mathcal{O}_{\text{biProj}(D)}$ -algebras, where  $\mathcal{A}$  is the sheaf on  $\text{biProj}(D)$  which is locally defined by the algebra  $(k[x_0, \dots, x_n, y_0, \dots, y_m]/(f, g)_{x_i y_j})_{(0,0)}$  on each open affine scheme  $D_+(s_i t_j) \simeq \text{Spec}((D_{s_i t_j})_{(0,0)})$ . Hence, it is enough to prove that

$$\begin{aligned} \text{gl.dim}((k[x_0, \dots, x_n, y_0, \dots, y_m]/(f, g)_{x_i y_j})_{(0,0)}) &= d_0 + d_1 + e_0 + e_1 - 4 \\ &= n + m - 2. \end{aligned}$$

We can complete the rest of the proof in the same way as in [12, Section 2.3]. We give its sketch. For simplicity, we prove the claim when  $i = j = 0$ . We define a  $k$ -algebra  $E$  by

$$E := k[S_1, \dots, S_n, T_1, \dots, T_m] \left/ \left( 1 + \sum_{i=1}^n S_i, 1 + \sum_{i=1}^m T_i \right) \right.$$

with  $S_i = s_i/s_0$ ,  $T_i = t_i/t_0$ . We also define an  $E$ -algebra  $F$  by

$$F := k\langle X_1, \dots, X_n, Y_1, \dots, Y_m \rangle \left/ \left( \begin{aligned} &X_i X_j - (q_{0i} q_{ij} q_{j0}) X_j X_i, 1 + \sum_{l=1}^n X_l^{n+1}, 1 + \sum_{l=1}^m Y_l^{m+1} \\ &Y_i Y_j - (q'_{0i} q'_{ij} q'_{j0}) Y_j Y_i, \end{aligned} \right) \right|_{i,j}$$

with  $X_i = x_i/x_0$ ,  $Y_i = y_i/y_0$ . The module structure of  $F$  is given by the identifications  $S_i = X_i^{n+1}$ ,  $T_i = Y_i^{m+1}$ . Let  $F_{\tilde{m}}$  be the localization of  $F$  at a maximal ideal

$$\tilde{m} := (S_1 - a_1, \dots, S_n - a_n, T_1 - b_1, \dots, T_m - b_m)$$

of  $E$  with  $1 + \sum_{i=1}^n a_i = 1 + \sum_{i=1}^m b_i = 0$  ( $a_i, b_i \in k$ ). Then, it is enough to prove that the global dimension of  $F_{\tilde{m}}$  is  $n + m - 2$  (see [12, Lemma 2.6, 2.7]).

If all  $a_i, b_i$  are not 0, then  $F/\tilde{m}F$  is a twisted group ring and hence semisimple. Moreover,  $S_1 - a_1, \dots, S_n - a_n, T_1 - b_1, \dots, T_m - b_m$  is a regular sequence in  $F_{\tilde{m}}$ . This induces the claim (see [16, Theorem 7.3.7]).

On the other hand, assume that one of  $\{a_1, \dots, a_n, b_1, \dots, b_m\}$  is 0. For example, assume  $a_1 = 0$ . We consider  $F/(X_1)$ . Then, we can show that the global dimension of  $(F/(X_1))_{\tilde{m}} = n + m - 3$  because  $\text{pd}_F(S) = \text{pd}_{F/(X_1)}(S) + 1$  for any simple  $F$ -module



$S$  with  $\text{Ann}(S) = \widetilde{m}$  (see [16, Theorem 7.3.5]). If some other  $a_i, b_j$  are 0, we repeat taking quotients and can reduce to considering the global dimension of the algebra  $k[X, Y]/(X^{n+1} + 1, Y^{m+1} + 1)$ , which is 0. ■

**Remark 3.9** To prove Lemma 3.8 in (2) of the theorem, consider the projective spectrum  $X := \text{biProj}(k[s_0, \dots, s_n, t_0, \dots, t_{n+1}]/(\sum_{i=0}^n s_i t_i, \sum_{i=0}^{n+1} t_i^{n+1}))$  (resp.  $\text{biProj}(k[s_0, \dots, s_n, t_0, \dots, t_n]/(\sum_{i=0}^n s_i t_i, \sum_{i=0}^n t_i^n))$ ) and the sheaf  $\mathcal{A}$  of algebras on  $X$  associated with  $C/(f, g)$ .

**Proof of Theorem 3.3** First, we calculate  $\text{R}\Gamma_{m_{C/(f,g)}}(C/(f, g))^*$ . From [12, Proposition 2.4] (or [25, Example 5.5]) and the proof of [25, Lemma 6.1], we have

$$(3.1.3) \quad \begin{aligned} \text{R}\Gamma_{m_C}(C)^* &\simeq \text{R}\Gamma_{m_A}(A)^* \otimes \text{R}\Gamma_{m_B}(B)^* \\ &\simeq \phi A^1(-d_0 - e_0) \otimes_k \psi B^1(-d_1 - e_1)[d_0 + d_1 + e_0 + e_1], \end{aligned}$$

where  $\phi$  (resp.  $\psi$ ) is the graded automorphism of  $A$  (resp.  $B$ ) which maps  $x_j \mapsto \prod_{i=0}^n q_{ji} x_j$  (resp.  $y_j \mapsto \prod_{i=0}^m q'_{ji} y_j$ ). Then, we consider the distinguished triangles

$$(3.1.4) \quad \text{R}\Gamma_{m_C}(C(-d_0, -d_1)) \xrightarrow{\times f} \text{R}\Gamma_{m_C}(C) \longrightarrow \text{R}\Gamma_{m_{C/(f)}}(C/(f)),$$

$$(3.1.5)$$

$$\text{R}\Gamma_{m_{C/(f)}}((C/(f))(-e_0, -e_1)) \xrightarrow{\times g} \text{R}\Gamma_{m_{C/(f)}}(C/(f)) \longrightarrow \text{R}\Gamma_{m_{C/(f,g)}}(C/(f, g))$$

obtained from the exact sequences (3.1.1) and (3.1.2) of  $C$ -bimodules. Combining the formula (3.1.3) and the triangle (3.1.4), we have

$$(3.1.6) \quad \text{R}\Gamma_{m_{C/(f)}}(C/(f))^* \simeq \phi \otimes \psi (A \otimes_k B/(f))^1(-e_0, -e_1)[d_0 + d_1 + e_0 + e_1 - 1].$$

In addition, combining the triangle (3.1.5) and the formula (3.1.6), we have

$$(3.1.7) \quad \text{R}\Gamma_{m_{C/(f,g)}}(C/(f, g))^* \simeq \phi \otimes \psi (A \otimes_k B/(f, g))^1[d_0 + d_1 + e_0 + e_1 - 2].$$

On the other hand, we have the Serre duality in  $D^b(\text{qbigr}(C/(f, g)))$  from Lemma 3.10. Thus,  $-\otimes^{\mathbb{L}} \pi(\text{R}\Gamma_{m_{C/(f,g)++}}(C/(f, g))^*[-1])$  is the Serre functor of  $D^b(\text{qbigr}(C/(f, g)))$  because this functor induces an equivalence from Lemma 3.6 and the formula (3.1.7). Finally, the Serre functor  $-\otimes^{\mathbb{L}} \pi(\text{R}\Gamma_{m_{C/(f,g)++}}(C/(f, g))^*[-1])$  induces the  $[d_0 + d_1 + e_0 + e_1 - 4]$ -shift functor if and only if  $\prod_{i=0}^n q_{ij}$  and  $\prod_{i=0}^m q'_{ij}$  are independent of  $j$  (cf. [12, Remark 2.5]). This completes the proof. ■

The following lemma is well-known in the case of  $\mathbb{N}$ -graded algebras (for example, see [6, 37]).

**Lemma 3.10** (Local Duality and Serre Duality for  $\mathbb{N}^2$ -graded algebras) *Let  $D$  be a connected right Noetherian  $\mathbb{N}^2$ -graded  $k$ -algebra (connected means  $D_{0,0} = k$ ). Let  $E$  be a connected  $\mathbb{N}^2$ -graded  $k$ -algebra. We assume that  $\Gamma_{m_{D++}}$  has finite cohomological dimension.*

(1) *Let  $Q := \omega \circ \pi : \text{BiGr}(D) \rightarrow \text{BiGr}(D)$ . Let  $M \in D(\text{BiGr}(D \otimes_k E^\circ))$ . Then,*

$$(a) \quad \text{R}\Gamma_{m_{D++}}(M)^* \simeq \text{RHom}_D(M, \text{R}\Gamma_{m_{D++}}(D)^*),$$



$$(b) \quad RQ(M)^* \simeq R\mathrm{Hom}_D(M, RQ(D)^*)$$

in  $D(\mathrm{BiGr}(D^\circ \otimes_k E))$ , where we denote the natural extension of  $Q$  to a functor between  $\mathrm{BiGr}(D \otimes_k E^\circ)$  and itself by the same notation.

- (2) We assume that  $\mathrm{qbigr}(D)$  has finite global dimension. Let  $\mathcal{M} := \pi(M)$ ,  $\mathcal{N} := \pi(N)$  ( $M, N \in D^b(\mathrm{bigr}(D))$ ). Let  $\mathcal{R}_D := \pi(R\Gamma_{m_{D^{++}}}(D)^*) \in D^b(\mathrm{qbigr}(D^e))$ . Then,  $\mathcal{N} \otimes^{\mathbb{L}} \mathcal{R}_D \in D^b(\mathrm{qbigr}(D))$  and

$$\mathrm{Hom}_{D^b(\mathrm{qbigr}(D))}(\mathcal{N}, \mathcal{M}) \simeq \mathrm{Hom}_{D^b(\mathrm{qbigr}(D))}(\mathcal{M}, (\mathcal{N} \otimes^{\mathbb{L}} \mathcal{R}_D)[-1])',$$

which is functorial in  $\mathcal{M}$  and  $\mathcal{N}$ . Here,  $(-)'$  denotes the  $k$ -dual.

**Proof** Since  $R^i\Gamma_{m_{D^{++}}}(-) \simeq \lim_{n \rightarrow \infty} \mathrm{Ext}^i(D/D_{\geq n, \geq n}, -)$  and  $D$  is right Noetherian, one can check that  $R^i\Gamma_{m_{D^{++}}}(-)$  commutes with direct limits as in [36, Proposition 16.3.19]. In addition, if  $K$  is a complex of graded-free right  $D$ -modules and  $L$  is a complex of graded right  $D^e$ -modules, then  $\Gamma_{m_{D^{++}}}(K \otimes_D L) \simeq K \otimes_D \Gamma_{m_{D^{++}}}(L)$  (cf. [20, Lemma 6.10]). So, we can apply the argument of [30, Theorem 5.1] (or [22, Theorem 2.1]) to prove (a) of (1).

In order to prove (b) of (1), note that we have the canonical exact sequence and the isomorphism (see also [3, Lemmas 4.1.4 and 4.1.5])

$$0 \rightarrow \Gamma_{m_{D^{++}}}(M) \rightarrow M \rightarrow Q(M) \rightarrow \lim_{n \rightarrow \infty} \mathrm{Ext}^1(D/D_{\geq n, \geq n}, M) \rightarrow 0,$$

$$R^iQ(M) \simeq R^{i+1}\Gamma_{m_{D^{++}}}(M), \quad (1 \leq i, M \in \mathrm{BiGr}(D)).$$

So, from the previous paragraph,  $Q$  has finite cohomological dimension,  $R^iQ$  commutes with direct limits. We also have  $Q(K \otimes_D L) \simeq K \otimes_D Q(L)$ , where  $K, L$  are as above (cf. [21, Lemma 3.28]). Hence, we can also apply the argument of [30, Theorem 5.1] (or [21, Theorem 3.29]) to prove (b) of (1).

We can prove (2) in the same way as in [6, Lemma A.1 and Theorem A.4] by using (b) of (1). Note that we have a natural equivalence  $D^b(\mathrm{qbigr}(D)) \simeq D^b_f(\mathrm{QBiGr}(D))$ , where  $D^b_f(\mathrm{QBiGr}(D))$  is the full subcategory of  $D^b(\mathrm{QBiGr}(D))$  consisting of complexes with cohomology in  $\mathrm{qbigr}(D)$  (see [6, Lemma 2.2]). ■

As a corollary of Theorem 3.3, we construct examples of noncommutative projective Calabi–Yau schemes by using Segre products. Let  $A, B, f$  and  $g$  be as in Theorem 3.3.

- Definition 3.11** (1) The Segre product  $A \circ B$  of  $A$  and  $B$  is the  $\mathbb{N}$ -graded  $k$ -algebra with  $(A \circ B)_i = A_i \otimes_k B_i$ .
- (2) Let  $M \in \mathrm{bigr}(C)$ . We define a right graded  $A \circ B$ -module  $M_\Delta$  as the graded  $A \circ B$ -module with  $(M_\Delta)_i = M_{i,i}$ .

**Lemma 3.12** [32, Theorem 2.4] We have the following natural isomorphism:

$$\mathrm{qbigr}(C) \longrightarrow \mathrm{qgr}(A \circ B), \quad \pi(M) \longmapsto \pi(M_\Delta).$$

In addition, the functor defined by  $- \otimes_{A \circ B} C$  is the inverse of this equivalence.

**Remark 3.13** Let  $J := (f, g) \in \text{bigr}(C)$ . We similarly obtain an equivalence

$$\text{qbigr}(C/J) \simeq \text{qgr}(A \circ B/J_\Delta).$$

Combining Theorem 3.3 with Remark 3.13, we get the following.

**Corollary 3.14** Let  $J := (f, g) \in \text{bigr}(C)$ . Then,  $\text{proj}(A \circ B/J_\Delta)$  is a noncommutative projective Calabi–Yau scheme.

## 3.2 Weighted hypersurfaces

Reid produced the list of all commutative weighted Calabi–Yau hypersurfaces of dimensions 2 (for example, see [11, 23]). In this section, we construct noncommutative projective Calabi–Yau schemes from noncommutative weighted projective hypersurfaces. Let  $A$  be a right Noetherian  $\mathbb{N}$ -graded  $k$ -algebra. Then, the  $r$ th Veronese algebra  $A^{(r)}$  is the  $\mathbb{N}$ -graded  $k$ -algebra with  $A_i^{(r)} = A_{ri}$ . We consider the (commutative) weighted polynomial ring  $A = k[x_0, \dots, x_n]$  with  $\deg(x_i) = d_i$ . Then,  $\text{Coh}(\text{Proj}(A))$  is in general not equivalent to  $\text{qgr}(A)$ , but to  $\text{qgr}(A^{(n+1)\text{lcm}(d_0, \dots, d_n)})$ . However, we can think of  $\text{qgr}(A)$  as a resolution of singularities of  $\text{Coh}(\text{Proj}(A))$  (cf. [28, Example 4.9]). Moreover, we have  $\text{qgr}(A) \simeq \text{Coh}([\text{Spec}(A) \setminus \{0\}]/\mathbb{G}_m)$  and  $[(\text{Spec}(A) \setminus \{0\})/\mathbb{G}_m]$  is a smooth Deligne–Mumford stack whose coarse moduli space is  $\text{Proj}(A)$ .

**Theorem 3.15** Let  $(d_0, \dots, d_n) \in \mathbb{Z}_{>0}^{n+1}$  and  $d := \sum_{i=0}^n d_i$  such that  $d$  is divisible by  $d_i$  for all  $i$ . Let  $C := k\langle x_0, \dots, x_n \rangle / (x_j x_i - q_{ji} x_i x_j)_{i,j}$ , where  $q_{ji} \in k^\times$ ,  $\deg(x_i) = d_i$  for all  $i, j$ . Let  $f := \sum_{i=0}^n x_i^{h_i}$ , where  $h_i := d/d_i$ .

We assume that  $q_{ii} = q_{ij} q_{ji} = q_{ij}^{h_j} = q_{ji}^{h_i} = 1$  for all  $i, j$ . Then,  $\text{proj}(C/(f))$  is a noncommutative projective Calabi–Yau scheme of dimension  $(n-1)$  if and only if there exists  $c \in k$  such that  $c^{d_j} = \prod_{i=0}^n q_{ij}$  for all  $j$ .

**Remark 3.16** •  $f$  is a central element in  $C$  from the choice of  $\{q_{ij}\}$ .

• Theorem 3.15 is a generalization of [12, Theorem 1.1].

**Lemma 3.17** The balanced dualizing complex of  $C/(f)$  is isomorphic to  ${}^\phi(C/(f))^{[1]}$   $[n]$ , where  ${}^\phi$  is a graded automorphism of  $C$  which maps  $x_j \mapsto \prod_{i=0}^n q_{ji} x_j$ .

**Proof** Since  $C$  is Artin–Schelter regular,  $C$  is skew Calabi–Yau (see [25, Lemma 1.2]). This implies that the balanced dualizing complex of  $C$  is isomorphic to  ${}^\phi C^1(-d)[n+1]$ , where  ${}^\phi$  is the Nakayama automorphism of  $C$ . From [25, Example 5.5], the automorphism  ${}^\phi$  is the map which maps  $x_j \mapsto \prod_{i=0}^n q_{ji} x_j$ .

By using this result, we can obtain the claim in the same way as in the proof of Theorem 3.3 after Remark 3.9. ■

In general,  $C/(f)$  is not generated in degree 0 and 1. This fact prevents us from using the idea of the proof of Lemma 3.8 to calculate the global dimension of  $\text{qgr}(C/(f))$ . So, we need to find a right Noetherian  $\mathbb{N}$ -graded  $k$ -algebra  $R$  which

is generated in degree 0 and 1 and satisfies  $\text{qgr}(R) \simeq \text{qgr}(C/(f))$ . Quasi-Veronese algebras are effective in achieving this objective. We recall the notion of quasi-Veronese algebras below. In detail, see [18, Section 3].

**Definition 3.18** [18, Section 3] Let  $A$  be an  $\mathbb{N}$ -graded  $k$ -algebra. The  $l$ th quasi-Veronese algebra  $A^{[l]}$  of  $A$  is a graded  $k$ -algebra defined by

$$A^{[l]} := \bigoplus_{i \in \mathbb{N}} A_i^{[l]} := \bigoplus_{i \in \mathbb{N}} \begin{pmatrix} A_{li} & A_{li+1} & \cdots & A_{li+l-1} \\ A_{li-1} & A_{li} & \cdots & A_{li+l-2} \\ \vdots & \vdots & \ddots & \vdots \\ A_{li-l+1} & A_{li-l+2} & \cdots & A_{li} \end{pmatrix}.$$

**Remark 3.19** (1) We have  $\text{Gr}(A) \simeq \text{Gr}(A^{[l]})$  (see [18, Lemma 3.9]). The equivalence is obtained by the functor  $\Psi : \text{Gr}(A) \rightarrow \text{Gr}(A^{[l]})$ , which is defined by  $\Psi(M) := \bigoplus_{i \in \mathbb{Z}} \left( \bigoplus_{j=0}^{l-1} M_{li-j} \right)$ .

(2) When  $A$  is right Noetherian,  $A^{[l]} \simeq \bigoplus_{0 \leq i, j \leq n-1} A(j-i)^{(l)} \in \text{gr}(A^{(l)})$ , where  $A^{(l)}$  is the  $l$ th Veronese algebra of  $A$  and the  $A^{(l)}$ -module structure of  $A^{[l]}$  is given by the natural inclusion  $A^{(l)} \subset A^{[l]}$  (cf. the proof of [19, Proposition 4.11]). Then,  $A^{[l]}$  is also right Noetherian since  $A^{(l)}$  is right Noetherian. In this case,  $\Psi$  induces an equivalence between  $\text{qgr}(A)$  and  $\text{qgr}(A^{[l]})$ .

**Lemma 3.20** Let  $A$  be an  $\mathbb{N}$ -graded  $k$ -algebra which is generated by homogeneous elements  $y_0, \dots, y_h$  with  $\deg(y_i) > 0$  as an  $A_0$ -algebra. Let  $l \geq \max\{\deg(y_0), \dots, \deg(y_h)\}$ . Then,  $A^{[l]}$  is generated in degree 0 and 1.

**Proof** For any  $i \in \mathbb{N}$  and any  $a, b \in \{0, 1, \dots, l-1\}$ , it is enough to show that every homogeneous element  $m$  of the form

$$m = \begin{pmatrix} m_{0,0} & \cdots & m_{0,\beta} & \cdots & m_{0,l-1} \\ \vdots & & \vdots & & \vdots \\ m_{\alpha,0} & \cdots & m_{\alpha,\beta} & \cdots & m_{\alpha,l-1} \\ \vdots & & \vdots & & \vdots \\ m_{l-1,0} & \cdots & m_{l-1,\beta} & \cdots & m_{l-1,l-1} \end{pmatrix} \in A_i^{[l]}, \quad \begin{pmatrix} m_{\alpha,\beta} \in (A_i^{[l]})_{\alpha,\beta} := A_{li+\beta-\alpha} \\ m_{\alpha,\beta} = 0 \text{ when } (\alpha, \beta) \neq (a, b) \\ 0 \leq \alpha, \beta \leq l-1 \end{pmatrix}$$

is generated in degree 0 and 1. Moreover, we can assume that  $m_{a,b} = \prod_{j=0}^{n_1} y_{i_j}$  ( $i_j \in \{0, \dots, h\}, n_1 \in \mathbb{N}$ ).

If  $m_{a,b}$  is decomposed into  $\prod_{j=0}^{n_1} y_{i_j} = \prod_{j=0}^{n_2} y_{i_j} \prod_{j=n_2+1}^{n_1} y_{i_j}$  ( $n_2 \in \mathbb{N}$ ) such that  $l-a \leq \deg(\prod_{j=1}^{n_2} y_{i_j}) \leq 2l-a-1$ , then we have  $\prod_{j=0}^{n_2} y_{i_j} \in (A_1^{[l]})_{a,c} = A_{l+c-a}$  and  $\prod_{j=n_2+1}^{n_1} y_{i_j} \in (A_{l-1}^{[l]})_{c,b} = A_{l(i-1)+b-c}$  ( $0 \leq c \leq l-1$ ). In this case, we can show the claim by using induction on the degree of  $m$ . So, it is sufficient to show that we have such a decomposition for all  $m$ . Indeed, we can find at least one such decomposition from  $(2l-a-1) - (l-a) + 1 = l$  and the choice of  $l$ . In detail, we have  $l-a \leq \deg(y_{i_0}) \leq 2l-a-1$  or there exists  $n_3 \in \mathbb{N}$  such that  $\deg(y_{i_0} y_{i_1} \cdots y_{i_{n_3}}) < l-a$  and  $l-a \leq \deg(y_{i_0} y_{i_1} \cdots y_{i_{n_3}} y_{i_{n_3+1}}) \leq 2l-a-1$  since  $0 < \deg(y_i) \leq l$ . ■

**Lemma 3.21**  $\text{gl.dim}(\text{qgr}(C/(f))) = n-1$ .

**Proof** We use the idea of the proof of Lemma 3.8. We consider an  $\mathbb{N}$ -graded  $k$ -algebra  $B := k[s_0, \dots, s_n]/(\sum_{i=0}^n s_i)$  with  $s_i = x_i^{h_i}$ . Then,  $A^{[d]}$  is right Noetherian and  $\text{qgr}(C/(f)) \simeq \text{qgr}((C/(f))^{[d]})$  from Remark 3.19. So, it is enough to prove that  $\text{gl.dim}(\text{qgr}((C/(f))^{[d]})) = n - 1$ . Because  $C/(f)$  is finite over  $B$ , a  $B$ -submodule  $Z(C/(f))^{(d)}$  of  $C/(f)$  is finite over  $B$ . From [19, Propositions 4.10 and 4.11],  $(C/(f))^{[d]}$  is finite over  $Z(C/(f))^{(d)}$ . So,  $(C/(f))^{[d]}$  is finite over  $B$ . In addition,  $(C/(f))^{[d]}$  is generated in degrees 0 and 1 from Lemma 3.20. So,  $\text{qgr}((C/(f))^{[d]})$  is equivalent to the category of coherent modules over a sheaf  $\mathcal{A}$  of  $\mathcal{O}_{\text{Proj}(B)}$ -algebra, where  $\mathcal{A}$  is the sheaf on the projective spectrum  $\text{Proj}(B)$  which is locally defined by a tiled matrix algebra

$$N_i = \begin{pmatrix} E_{i,0} & E_{i,1} & \cdots & E_{i,d-1} \\ E_{i,-1} & E_{i,0} & \cdots & E_{i,d-2} \\ \vdots & \vdots & \cdots & \vdots \\ E_{i,-d+1} & E_{i,-d+2} & \cdots & E_{i,0} \end{pmatrix}$$

on each  $D_+(s_i)$ . Here,  $E_i := (C/(f))[x_i^{-1}]$  and  $E_{i,j}$  is the degree  $j$  part of  $E_i$ . As in the proof of Lemma 3.8, it is enough to show that the global dimension of  $N_i$  is  $n - 1$  for all  $i$ .

On the other hand, two graded algebras

$$\begin{aligned} R_1 &:= E_i \oplus E_i(1) \oplus \cdots \oplus E_i(d-2) \oplus E_i(d-1), \\ R_2 &:= E_i \oplus E_i(1) \oplus \cdots \oplus E_i(d_i-2) \oplus E_i(d_i-1) \end{aligned}$$

are progenerators in  $\text{Gr}(E_i)$ . So, the category of right  $\text{End}_{\text{gr}}(R_1)$ -modules and the category of right  $\text{End}_{\text{gr}}(R_2)$ -modules are equivalent because they are equivalent to the category of graded right  $E_i$ -modules (cf. [26, Lemma 4.8], [28, Remarks after Proposition 4.5]). We also have  $\text{End}_{\text{gr}}(R_1) \simeq N_i$  and

$$\text{End}_{\text{gr}}(R_2) \simeq M_i := \begin{pmatrix} E_{i,0} & E_{i,1} & \cdots & E_{i,d_i-1} \\ E_{i,-1} & E_{i,0} & \cdots & E_{i,d_i-2} \\ \vdots & \vdots & \cdots & \vdots \\ E_{i,-d_i+1} & E_{i,-d_i+2} & \cdots & E_{i,0} \end{pmatrix}.$$

So, it is sufficient to prove the global dimension of  $M_i$  is  $n - 1$  for each  $i$ .

For simplicity, we assume  $i = 0$ . When  $i \neq 0$ , we can show the claim in the same way. Let  $D = k[S_1, \dots, S_n]/(1 + \sum_{j=0}^n S_j)$  with  $S_j = s_j/s_0$ . We show that the global dimension of the  $D$ -algebra  $M_0$  is  $n - 1$ . The module structure of  $M_0$  is given by the identification  $S_j = (x_j^{h_j}/x_0^{h_0})I_{d_0} \in M_0$ , where  $I_{d_0}$  is the  $(d_0 \times d_0)$ -identity matrix. Let

$$\tilde{m} = (S_1 - a_1, \dots, S_n - a_n) \quad (a_j \in k)$$

be a maximal ideal of  $D$  with  $1 + \sum_{j=1}^n a_j = 0$ . It is sufficient to show that  $\text{gl.dim}((M_0)_{\tilde{m}}) = n - 1$ , where  $(M_0)_{\tilde{m}}$  is the localization of  $M_0$  at  $\tilde{m}$  (cf. the second paragraph of the proof of Lemma 3.8). We divide the proof of this claim into two cases.

*Case (a): all  $a_j$  are not 0.* Because  $S_1 - a_1, \dots, S_n - a_n$  is a regular sequence in  $(M_0)_{\tilde{m}}$ , we show that the global dimension of  $(M_0)_{\tilde{m}}/\tilde{m}(M_0)_{\tilde{m}} \simeq M_0/\tilde{m}M_0$  is 0 (cf. the third paragraph of the proof of Lemma 3.8).

First, the category of  $M_0/\widetilde{m}M_0$ -modules is equivalent to the category of graded  $E'_0$ -modules, where

$$E'_0 := E_0/(x_1^{h_1}/x_0^{h_0} - a_1, \dots, x_n^{h_n}/x_0^{h_0} - a_n)E_0.$$

This is a Morita equivalence obtained from the isomorphism  $\text{End}_{\text{gr}}(E'_0) \simeq M_0/\widetilde{m}M_0$  (cf. the three previous paragraph).

Next, we see that  $E'_0$  is strongly graded. Since

$$E_0 \simeq (C[x_0^{-1}])/(1 + (x_1^{h_1}/x_0^{h_0}) + \dots + (x_n^{h_n}/x_0^{h_0})),$$

we have

$$E'_0 \simeq (C[x_0^{-1}])/(x_1^{h_1}/x_0^{h_0} - a_1, \dots, x_n^{h_n}/x_0^{h_0} - a_n).$$

For any  $l \in \mathbb{Z}$ , if  $\widetilde{x} := x_0^{l_0} x_1^{l_1} \dots x_n^{l_n} \in (E'_0)_l$  ( $l_0 \in \mathbb{Z}, l_1, \dots, l_n \in \mathbb{N}$ ), then there exist  $k_1, \dots, k_n \in \mathbb{N}$  such that  $\widetilde{x}' := x_0^{(-\sum k_i)h_0 - l_0} x_1^{k_1 h_1 - l_1} \dots x_n^{k_n h_n - l_n} \in (E'_0)_{-l}$ . Because  $\widetilde{x} \widetilde{x}' \in k^*$ , we get  $1 \in (E'_0)_l (E'_0)_{-l}$  and  $E'_0$  is strongly graded.

Since  $E'_0$  is strongly graded, we have  $\text{Gr}(E'_0) \simeq \text{Mod}((E'_0)_0)$ . Then,  $(E'_0)_0$  is a twisted group algebra, where a  $k$ -basis of  $(E'_0)_0$  is

$$\left\{ x_0^{e_0} x_1^{e_1} x_2^{e_2} \dots x_n^{e_n} \in (E'_0)_0 \mid \sum_{j=0}^n e_j d_j = 0 \text{ and } 0 \leq e_j < h_j \ (\forall j = 1, 2, \dots, n) \right\}.$$

In particular,  $(E'_0)_0$  is semisimple. Hence, the graded global dimension of  $E'_0$  is 0 and  $\text{gl.dim}(M_0/\widetilde{m}M_0) = 0$ .

*Case (b): some of  $a_j$  are 0.* For example, we assume  $a_1 = 0$ . Then,  $(x_1^{h_1}/x_0^{h_0})I_{d_0}$  is an annihilator of any simple  $M_0$ -module  $N$ . On the other hand, we have a unique integer  $r_1$  such that  $0 \leq \deg(x_1/x_0^{r_1}) \leq d_0 - 1$ . If  $\deg(x_1/x_0^{r_1}) = 0$ , then  $J = x_1/x_0^{r_1}I_{d_0}$  annihilates  $N$ . Otherwise, the matrix

$$J = \begin{pmatrix} & & & x_1/x_0^{r_1} & & \\ & O & & \ddots & & \\ & & & & & x_1/x_0^{r_1} \\ x_1/x_0^{r_1+1} & & & & & \\ & \ddots & & & O & \\ & & x_1/x_0^{r_1+1} & & & \end{pmatrix} \in M_0$$

annihilates  $N$  because  $\exists n_J \in \mathbb{N}$  such that  $J^{n_J} = (x_1^{h_1}/x_0^{h_0})I_{d_0}$  (the reduction of  $N_i$  to  $M_i$  is used here). Thus, it is enough to prove that the global dimension of  $(M_0/JM_0)_{\widetilde{m}} = n - 2$  (cf. the fourth paragraph of the proof of Lemma 3.8). Note that we have

$$(3.2.1) \quad M_0/JM_0 \simeq \begin{pmatrix} F_{0,0} & F_{0,1} & \cdots & F_{0,d_0-1} \\ F_{0,-1} & F_{0,0} & \cdots & F_{0,d_0-2} \\ \vdots & \vdots & \cdots & \vdots \\ F_{0,-d_0+1} & F_{0,-d_0+2} & \cdots & F_{0,0} \end{pmatrix},$$

where

$$F_0 := E_0/x_1E_0 \simeq k\langle x_0, x_2, \dots, x_n \rangle / (x_j x_i - q_{ji} x_i x_j, x_0^{h_0} + x_2^{h_2} + \dots + x_n^{h_n})_{i,j} [x_0^{-1}]$$

and  $F_{0,j}$  is the degree  $j$  part of  $F_0$ .

If any of  $a_2, \dots, a_n$  is not 0, we can reduce to the case (a) from (3.2.1). If some of  $a_2, \dots, a_n$  are 0, repeat the above process until we can reduce to the case (a). ■

**Proof of Theorem 3.15**  $\text{gl.dim}(\text{qgr}(C/(f)))$  is finite. So, the balanced dualizing complex  ${}^\phi(C/(f))^{[n]}$  of  $C/(f)$  induces the Serre functor of  $\text{qgr}(C/(f))$  from [6, Theorem A.4]. We complete the proof as in the proof of Theorem 3.3. ■

## 4 Comparison and closed points

In this section, we calculate closed points of noncommutative projective Calabi–Yau schemes of dimensions 2 obtained in Section 3.2 and compare our examples with commutative Calabi–Yau varieties and the first examples constructed in [12]. In particular, we show that a noncommutative projective Calabi–Yau scheme in Section 3.2 gives essentially a new example of noncommutative projective Calabi–Yau schemes.

**Example 4.1** Any weight  $(d_0, d_1, d_2, d_3)$  of noncommutative projective Calabi–Yau schemes of dimensions 2 in Theorem 3.15 such that  $\gcd(d_0, d_1, d_2, d_3) = 1$  is one of the following (obtained by using a computer):

$$\begin{aligned}(d_0, d_1, d_2, d_3) = & (1, 1, 1, 1), (1, 1, 1, 3), (1, 1, 2, 2), (1, 1, 2, 4), (1, 1, 4, 6), \\ & (1, 2, 2, 5), (1, 2, 3, 6), (1, 2, 6, 9), (1, 3, 4, 4), (1, 3, 8, 12), \\ & (1, 4, 5, 10), (1, 6, 14, 21), (2, 3, 3, 4), (2, 3, 10, 15).\end{aligned}$$

From now, we focus on the closed points of noncommutative projective Calabi–Yau schemes of dimensions 2 in Theorem 3.15 whose weights are of type  $(1, 1, a, b)$ . We recall the notion of closed points of noncommutative projective schemes.

For simplicity, we often call an  $\mathbb{N}$ -graded  $k$ -algebra of the form  $k\langle z_0, \dots, z_m \rangle / (z_j z_i - p_{ji} z_i z_j)_{i,j}$  ( $p_{ji} \in k^\times$ ,  $m \in \mathbb{N}$ ) with  $\deg(z_i) > 0$  and  $p_{ji} p_{ij} = 1$  a weighted quantum polynomial ring. ( $p_{ji}$ ) is called the quantum parameter.

**Definition 4.2** [19, Section 3.1] Let  $A$  be a finitely generated right Noetherian connected  $\mathbb{N}$ -graded  $k$ -algebra. A closed point of  $\text{proj}(A)$  is an object of  $\text{qgr}(A)$  represented by a 1-critical module of  $A$ . We denote by  $|\text{proj}(A)|$  the set of closed points of  $\text{proj}(A)$ . For the definition of 1-critical modules, see [19, Definition 3.1].

**Remark 4.3** [19, Section 3.1] If  $A$  is a quotient of a weighted quantum polynomial ring, then every closed point of  $\text{proj}(A)$  is one of the following:

- (1) An ordinary point, which is represented by a finitely generated 1-critical module of multiplicity 1.
- (2) A fat point, which is represented by a finitely generated 1-critical module of multiplicity  $> 1$ .
- (3) A thin point, which is represented by a finitely generated 1-critical module of multiplicity  $< 1$ .

For the definition of multiplicities, see [19, Definition 3.10]. In addition, if  $A$  is generated in degree 1, the notion of ordinary points and that of point modules are the same, and there are no thin points.

Let  $C := k\langle x_0, x_1, x_2, x_3 \rangle / (x_j x_i - q_{ji} x_i x_j)_{i,j}$  whose weight is of type  $(d_0, d_1, d_2, d_3) = (1, 1, a, b)$  ( $0 < a \leq b$ ). We assume that  $q_{ij} q_{ji} = q_{ii} = 1$  for all  $i, j$ . Since  $d_0 = 1$ ,  $C[x_0^{-1}]$  is strongly graded. So, from [19, Theorem 4.20], we have

$$|\mathrm{proj}(C)| = |\mathrm{spec}(C[x_0^{-1}]_0)| \sqcup |\mathrm{proj}(C/(x_0))|,$$

where we denote by  $|\mathrm{spec}(C[x_0^{-1}]_0)|$  the set of simple modules of  $C[x_0^{-1}]_0$ . In this equality, the 1 (resp.  $n > 1$ )-dimensional simple modules of  $\mathrm{spec}(C[x_0^{-1}]_0)$  correspond to ordinary (resp. fat) points in  $\mathrm{proj}(C)$ . Similarly, we have

$$|\mathrm{proj}(C)| = |\mathrm{spec}(C[x_0^{-1}]_0)| \sqcup |\mathrm{spec}(C/(x_0)[x_1^{-1}]_0)| \sqcup |\mathrm{proj}(C/(x_0, x_1))|.$$

We have an isomorphism  $C[x_0^{-1}]_0 \xrightarrow{\sim} k\langle X_1, X_2, X_3 \rangle / (X_j X_i - q'_{ji} X_i X_j)_{i,j}$  which sends  $x_1 x_0^{-1}$ ,  $x_2 x_0^{-a}$  and  $x_3 x_0^{-b}$  to  $X_1, X_2$ , and  $X_3$ , respectively. Here,  $q'_{ji} := q_{0j}^{d_i} q_{ji} q_{i0}^{d_j}$  ( $i, j \neq 0$ ). In the same way,  $C/(x_0)[x_1^{-1}]_0$  is also isomorphic to  $k\langle Y_2, Y_3 \rangle / (Y_3 Y_2 - p_{32} Y_2 Y_3)$ , where  $p_{32} := q_{13}^a q_{32} q_{21}^b$ .

Let  $C_1 := k\langle x'_0, x'_1, x'_2, x'_3 \rangle / (x'_j x'_i - q'_{ji} x'_i x'_j)_{i,j}$ , where  $\deg(x'_i) = 1$ ,  $q'_{0i} = q'_{j0} = 1$  for all  $i, j$ . Let  $C_2 := k\langle y_1, y_2, y_3 \rangle / (y_j y_i - p_{ji} y_i y_j)_{i,j}$ , where  $\deg(y'_i) = 1$ ,  $p_{1i} = p_{j1} = 1$  for all  $i, j$ . Then, we can consider the point scheme of  $\mathrm{proj}(C_1)$  (resp.  $\mathrm{proj}(C_2)$ ), which is isomorphic to the set of ordinary points  $|\mathrm{proj}(C_1)|_{\mathrm{ord}}$  (resp.  $|\mathrm{proj}(C_2)|_{\mathrm{ord}}$ ) as sets. Thus, we regard  $|\mathrm{proj}(C_1)|_{\mathrm{ord}}$  (resp.  $|\mathrm{proj}(C_2)|_{\mathrm{ord}}$ ) as the point scheme of  $\mathrm{proj}(C_1)$  (resp.  $\mathrm{proj}(C_2)$ ).

Let  $|\mathrm{spec}(C[x_0^{-1}]_0)|_1$  (resp.  $|\mathrm{spec}(C/(x_0)[x_1^{-1}]_0)|_1$ ) be the set of one-dimensional simple modules of  $C[x_0^{-1}]_0$  (resp.  $C/(x_0)[x_1^{-1}]_0$ ). Because  $C_1[x_0'^{-1}]_0 \simeq C[x_0^{-1}]_0$  and  $C_2[y_1^{-1}]_0 \simeq C/(x_0)[x_1^{-1}]_0$ , we can think of  $|\mathrm{spec}(C[x_0^{-1}]_0)|_1$  (resp.  $|\mathrm{spec}(C/(x_0)[x_1^{-1}]_0)|_1$ ) as a locally closed subscheme of  $|\mathrm{proj}(C_1)|_{\mathrm{ord}}$  (resp.  $|\mathrm{proj}(C_2)|_{\mathrm{ord}}$ ) from [19, Theorem 4.20].

**Lemma 4.4** (1) If  $q'_{ji} \neq 1$  for all  $i, j \neq 0$ ,  $|\mathrm{spec}(C[x_0^{-1}]_0)|_1$  is a union of three affine lines.

(2) If  $p_{32} \neq 1$ ,  $|\mathrm{spec}(C/(x_0)[x_1^{-1}]_0)|_1$  is a union of two affine lines. Otherwise,  $|\mathrm{spec}(C/(x_0)[x_1^{-1}]_0)|_1 \simeq \mathbb{A}^2$ .

**Proof** (2) is well-known (for example, see [26, Section 4.3]). Regarding (1), under the assumption of the lemma,  $\mathrm{proj}(C_1)$  belongs to case (3) or case (4) in [33, Corollary 5.1]. This shows that  $|\mathrm{spec}(C_1[x_0'^{-1}]_0)|_1$  is isomorphic to  $\bigcup_{i \neq j} Z(X'_i, X'_j) \subset \mathbb{A}^3 = \mathrm{Spec}(k[X'_1, X'_2, X'_3])$  (cf. [33, Proposition 4.2] or [2, Theorem 1]). ■

**Remark 4.5** We consider the weights  $(1, 1, a, b)$  and the quantum parameters which give noncommutative projective Calabi–Yau schemes of dimensions 2 in Theorem 3.15. Then, we can check that if  $p_{32} \neq 1$ , then  $q'_{ji} \neq 1$  for all  $i, j \neq 0$  by using a computer. Moreover, if  $p_{32} = 1$ , then  $q'_{ji} = 1$  for all  $i, j \neq 0$ . In this case,  $|\mathrm{spec}(C[x_0^{-1}]_0)|_1 \simeq \mathbb{A}^3$ .



We consider  $C/(x_0, x_1) = k\langle x_2, x_3 \rangle / (x_3x_2 - q_{32}x_2x_3)$ . Then, it is known that a weighted quantum polynomial ring of two variables is a twisted algebra of a commutative weighted polynomial ring  $k[x, y]$  with  $\deg(x) = a > 0$ ,  $\deg(y) = b > 0$  (for example, see [29, Example 4.1] or [38, Example 3.6]). So, it is enough to consider the closed points of  $\text{proj}(k[x, y])$ . We want to study the closed points of  $\text{proj}(k[x, y])$  in the case of  $(a, b) = (2, 2)$ ,  $(2, 4)$  or  $(4, 6)$ . Note that when  $(a, b) = (1, 1)$  or  $(1, 3)$ , they are classified in [19, Theorem 3.16]. We treat a more general setting below.

**Lemma 4.6** *Let  $R = k[x, y]$  be a commutative weighted polynomial ring with  $\deg(x) = a > 0$ ,  $\deg(y) = b > 0$ . Let  $g := \gcd(a, b)$ ,  $a' := a/g$  and  $b' := b/g$ . Then, every closed point of  $\text{proj}(R)$  is one of the following:*

- (1)  $\pi R/(x)(-i)$ ,  $i = 0, \dots, b-1$ .
- (2)  $\pi R/(y)(-j)$ ,  $j = 0, \dots, a-1$ .
- (3)  $\pi R/(\beta x^{b'} - \alpha y^{a'})(-k)$ , where  $(\alpha, \beta) \in \mathbb{P}^1 \setminus \{(0, 1), (1, 0)\}$  and  $k = 0, \dots, g-1$ .

Moreover, all of them are not isomorphic in  $\text{proj}(R)$ .

**Proof** The proof is almost the same as the proof of [19, Lemma 3.15 and Theorem 3.16]. We give the sketch of the proof.

Firstly, every closed point of  $\text{proj}(R)$  is represented by a cyclic critical Cohen–Macaulay module of depth 1. Then,  $M \in \text{gr}(R)$  satisfies these conditions and is generated in degree 0 if and only if  $M$  is isomorphic to one of  $R/(x)$ ,  $R/(y)$  or  $R/(\beta x^{b'} - \alpha y^{a'})$  ( $\alpha, \beta \in k^\times$ ). Since being cyclic critical Cohen–Macaulay of depth 1 is invariant under shifting, any closed point is represented by some shifts of one of the above modules (that is,  $R/(x)(-l)$ ,  $R/(y)(-l)$ ,  $R/(\beta x^{b'} - \alpha y^{a'})(-l)$ ,  $l \in \mathbb{Z}$ ).

Finally, we classify the isomorphic classes of these modules in  $\text{proj}(R)$ . We have no isomorphisms between the three types of closed points by considering their Hilbert polynomials and multiplicities. Then, we have  $\pi R/(\beta x^{b'} - \alpha y^{a'}) \simeq \pi R/(\beta x^{b'} - \alpha y^{a'})(-gl)$ , ( $\forall l \in \mathbb{Z}$ ,  $\forall (\alpha, \beta) \in \mathbb{P}^1 \setminus \{(1, 0), (0, 1)\}$ ). We also have  $\pi R/(\beta x^{b'} - \alpha y^{a'}) \simeq \pi R/(\beta' x^{b'} - \alpha' y^{a'})$  if and only if  $(\alpha, \beta) = (\alpha', \beta')$  in  $\mathbb{P}^1 \setminus \{(1, 0), (0, 1)\}$ . In addition, we can show that  $\pi R/(x) \simeq \pi R/(x)(-i)$  (resp.  $\pi R/(y) \simeq \pi R/(y)(-j)$ ) if and only if  $i \equiv 0 \pmod{b}$  (resp.  $j \equiv 0 \pmod{a}$ ). From these discussions, we get the claim. ■

We can study ordinary and thin points of noncommutative projective Calabi–Yau schemes of dimensions 2 in Theorem 3.15 by using the above investigations. We give examples of noncommutative projective Calabi–Yau schemes whose moduli of ordinary closed points are different from those in [12, Proposition 3.4] and commutative Calabi–Yau varieties.

**Example 4.7** We consider the weight  $(1, 1, 2, 2)$  and the quantum parameter

$$\mathbf{q} = (q_{ij}) = \begin{pmatrix} 1 & 1 & 1 & \omega^2 \\ 1 & 1 & \omega^2 & 1 \\ 1 & \omega & 1 & 1 \\ \omega & 1 & 1 & 1 \end{pmatrix}, \quad \omega := \frac{-1 + i\sqrt{3}}{2}.$$

Then, we have

$$\mathbf{q}' = (q'_{ij}) = \begin{pmatrix} 1 & \omega^2 & \omega \\ \omega & 1 & \omega^2 \\ \omega^2 & \omega & 1 \end{pmatrix}, \quad q_{13}^2 q_{32} q_{21}^2 = \omega^2.$$

From Lemmas 4.4 and 4.6, the set of ordinary and thin points

$$|\mathrm{proj}(C/(f))|_{\mathrm{ord} \ \& \ \mathrm{thin}} = |\mathrm{spec}(C/(f)[x_0^{-1}]_0)|_1 \sqcup |\mathrm{spec}(C/(f, x_0)[x_1^{-1}]_0)|_1 \\ \sqcup |\mathrm{proj}(C/(f, x_0, x_1))|$$

is 24 points. To be more precise, we have  $|\mathrm{spec}(C/(f)[x_0^{-1}]_0)|_1 = \sqcup_{i \neq j} Z(X_i, X_j, 1 + X_1^6 + X_2^3 + X_3^3) \subset \mathbb{A}^3$ ,  $|\mathrm{spec}(C/(f, x_0)[x_1^{-1}]_0)|_1 = \sqcup_{i=1,2} Z(Y_i, 1 + Y_2^3 + Y_3^3)$  and  $|\mathrm{proj}(C/(f, x_0, x_1))| = \{3\mathrm{pts}\} \sqcup \{3\mathrm{pts}\}$ .

This calculation shows that for a fixed weight, if the set of ordinary and thin points of  $\mathrm{proj}(C/(f))$  is finite, then the cardinality is independent of the quantum parameters.

From the method in Example 4.7, Remark 4.5, and a direct computation, we have the following.

**Proposition 4.8** *For a weight  $(1, 1, a, b)$  in Example 4.1 and a quantum parameter  $\mathbf{q}$  which gives a noncommutative projective Calabi–Yau scheme, if the set of ordinary and thin points of  $\mathrm{proj}(C/(f))$  is finite, then the cardinality is always 24.*

The following proposition shows that some of noncommutative projective Calabi–Yau schemes of dimensions 2 in Theorem 3.15 are essentially new examples.

**Proposition 4.9** *There exists a noncommutative projective Calabi–Yau scheme of dimension 2 which is obtained in Theorem 3.15 and not isomorphic to either commutative Calabi–Yau surfaces or noncommutative projective Calabi–Yau schemes of dimensions 2 obtained in [12].*

**Proof** We divide the proof into four steps.

*Step 1.* We choose the weight  $(1, 1, a, b)$  and the quantum parameter  $\mathbf{q}$  as in Example 4.7. Then, the number of ordinary and thin points of  $\mathrm{proj}(C/(f))$  is finite. So,  $\mathrm{proj}(C/f)$  is not isomorphic to any commutative Calabi–Yau surfaces.

*Step 2.* We prove that  $\mathrm{proj}(C/(f))$  is not isomorphic to any noncommutative projective Calabi–Yau schemes of dimensions 2 in [12]. To prove this, we use the theory established in [5]. First, note that we can think of  $\mathrm{qgr}(C/(f))$  as the category of coherent modules of a sheaf  $\mathcal{A}$  of algebras on the projective spectrum  $\mathrm{Proj}(k[s_0, s_1, s_2, s_3]/(s_0 + s_1 + s_2 + s_3))$  (cf. the proof of Lemma 3.21). We define a sheaf  $\mathcal{Z}_{\mathcal{A}}$  to be the sheaf whose sections are

$$\Gamma(U, \mathcal{Z}_{\mathcal{A}}) = \{s \in \Gamma(U, \mathcal{A}) \mid s|_V \in Z(\Gamma(V, \mathcal{A})), \forall V \subset U : \text{open}\}$$

for all open subsets  $U$  (cf. [5, Proposition 2.11]). In particular, if  $U$  is affine,  $\Gamma(U, \mathcal{Z}_{\mathcal{A}}) = Z(\Gamma(U, \mathcal{A}))$ . Then, we show that  $\mathrm{Spec}(Z(\Gamma(D_+(s_i), \mathcal{A})))$  has four singular points when  $i = 0, 1$  and a one-dimensional singular locus when  $i = 2, 3$ .

In the following, we verify this claim for  $i = 0, 2$ . Similarly, the claim is proved for  $i = 1, 3$ . In the following, we write  $Z_i$  as  $Z(\Gamma(D_+(s_i), \mathcal{A}))$  for any  $i$ . We also use the notations in the proof of Lemma 3.21.

When  $i = 0$ , any  $m \in Z_0$  is of the form  $m = \begin{pmatrix} \mu_1 e & 0 \\ 0 & \mu_2 e \end{pmatrix} \in N_0$ , ( $e \in E_{0,0}$ ,  $\mu_1, \mu_2 \in k^\times$ ) from the definition of  $\mathcal{A}$ . We have

$$E_{0,0} \simeq k\langle X_1, X_2, X_3 \rangle (X_j X_i - q'_{ji} X_i X_j, 1 + X_1^6 + X_2^3 + X_3^3)_{i,j},$$

which is obtained from the identifications  $X_1 = x_1 x_0^{-1}$ ,  $X_2 = x_2 x_0^{-2}$  and  $X_3 = x_3 x_0^{-2}$ . Here, the  $q'_{ji}$  are as in Example 4.7. So, we have

$$Z(E_{0,0}) \simeq k[Y, Z, W, U] / (1 + Y^2 + Z + W, YZW - \lambda_1 U^3) \quad (\lambda_1 \in k^\times),$$

which is obtained from the identifications  $Y = (x_1 x_0^{-1})^3$ ,  $Z = (x_2 x_0^{-2})^3$ ,  $W = (x_3 x_0^{-2})^3$  and  $U = (x_1 x_0^{-1})(x_2 x_0^{-2})(x_3 x_0^{-2})$ . On the other hand, we define the inclusion  $\phi: Z(E_{0,0}) \rightarrow N_0$  in which  $Y, Z, W$  are mapped naturally and  $U$  to  $\begin{pmatrix} U & 0 \\ 0 & \omega U \end{pmatrix}$ . It is easy to see that  $\phi(Z(E_{0,0})) \subset Z_0$ . Because the choice of  $\mu_1$  determines  $\mu_2$  in the above form of  $m$ , the map  $\phi$  induces  $Z_0 \simeq Z(E_{0,0})$ . Thus, one can show that  $\text{Spec}(Z_0)$  has four singular points by using the Jacobi criterion.

When  $i = 2$ , any  $m \in Z_2$  is of the form  $m = \begin{pmatrix} \mu_1 e & 0 \\ 0 & \mu_2 e \end{pmatrix} \in N_2$ , ( $e \in E_{2,0}$ ,  $\mu_1, \mu_2 \in k^\times$ ) from the definition of  $\mathcal{A}$ . We also have

$$E_{2,0} \simeq k\langle X_0, X_1, X_2, X_3 \rangle / (X_j X_i - q''_{ji} X_i X_j, 1 + X_0^6 + X_1^6 + X_3^3, X_0 X_1 - \lambda_2 X_2^2)_{i,j} \quad (\lambda_2 \in k^\times),$$

which is obtained from the identifications  $X_0 = x_0^2 x_2^{-1}$ ,  $X_1 = x_1^2 x_2^{-1}$ ,  $X_2 = x_0 x_1 x_2^{-1}$  and  $X_3 = x_3 x_2^{-1}$ . Here, the  $q''_{ij}$  are defined by the matrix

$$(q''_{ij}) = \begin{pmatrix} 1 & \omega & \omega^2 & \omega \\ \omega^2 & 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 & 1 \\ \omega^2 & \omega & 1 & 1 \end{pmatrix}.$$

So, we have

$$Z(E_{2,0}) \simeq k[X, Y, W, U, V] / (X + Y + 1 + W, XY - \lambda_3 U^2, XYW - \lambda_4 V^2) \quad (\lambda_3, \lambda_4 \in k^\times),$$

which is obtained from the identifications  $X = (x_0^2 x_2^{-1})^3$ ,  $Y = (x_1^2 x_2^{-1})^3$ ,  $W = (x_3 x_2^{-1})^3$ ,  $U = (x_0 x_1 x_2^{-1})^3$ , and  $V = (x_0 x_1 x_2^{-1})(x_3 x_2^{-1})$ . On the other hand, we define the inclusion  $\phi: Z(E_{2,0}) \rightarrow N_2$  in which  $X, Y, W, U$  are mapped naturally and  $V$  to  $\begin{pmatrix} V & 0 \\ 0 & \omega V \end{pmatrix}$ . It is easy to see that  $\phi(Z(E_{2,0})) \subset Z_2$ . Because the choice of  $\mu_1$  determines  $\mu_2$  in the above form of  $m$ , the map  $\phi$  induces  $Z_2 \simeq Z(E_{2,0})$ . Thus, one can show that  $\text{Spec}(Z_2)$  has a one-dimensional singular locus by using the Jacobi criterion.

*Step 3.* We consider the weight  $(1, 1, 1, 1)$  and take a quantum parameter which gives a noncommutative projective Calabi–Yau scheme  $\text{proj}(C'/(f'))$  whose point scheme is finite.  $\text{qgr}(C'/(f'))$  is thought of as the category of coherent modules of a sheaf  $\mathcal{B}$  of algebras on the projective spectrum  $\text{Proj}(k[t_0, t_1, t_2, t_3]/(t_0 + t_1 + t_2 + t_3))$ .

The number of the choices of quantum parameters  $(q_{ij})$  which satisfy the conditions of Theorem 3.15 and give a noncommutative projective Calabi–Yau scheme

whose moduli space of point modules is finite is 20 up to permutating variables (we get the list below by using a computer and hand calculations):

$$\begin{aligned}
 &1. \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & 1 \end{pmatrix}, 2. \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -i & -i \\ 1 & -i & i & i \end{pmatrix}, 3. \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 \\ -1 & 1 & 1 & 1 \end{pmatrix}, 4. \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & i & -i & -i \\ -1 & i & -i & i \end{pmatrix}, \\
 &5. \begin{pmatrix} 1 & 1 & 1 & i \\ 1 & 1 & 1 & -i \\ -i & i & i & 1 \end{pmatrix}, 6. \begin{pmatrix} 1 & 1 & 1 & i \\ 1 & i & 1 & 1 \\ -i & -1 & 1 & 1 \end{pmatrix}, 7. \begin{pmatrix} 1 & 1 & 1 & -i \\ 1 & 1 & 1 & i \\ i & -i & -i & 1 \end{pmatrix}, 8. \begin{pmatrix} 1 & 1 & 1 & -i \\ 1 & i & 1 & 1 \\ i & i & 1 & 1 \end{pmatrix}, \\
 &9. \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -i & i \\ -1 & -i & -i & 1 \end{pmatrix}, 10. \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & i & 1 & 1 \\ -i & -1 & 1 & 1 \end{pmatrix}, 11. \begin{pmatrix} 1 & 1 & 1 & -i \\ 1 & 1 & 1 & -1 \\ -1 & i & 1 & 1 \end{pmatrix}, 12. \begin{pmatrix} 1 & 1 & 1 & -i \\ 1 & i & 1 & 1 \\ -i & i & 1 & 1 \end{pmatrix}, \\
 &13. \begin{pmatrix} 1 & 1 & i & -i \\ -i & i & 1 & 1 \\ -1 & -1 & 1 & 1 \end{pmatrix}, 14. \begin{pmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \end{pmatrix}, 15. \begin{pmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & 1 & -i \\ -1 & -1 & i & 1 \end{pmatrix}, 16. \begin{pmatrix} 1 & -1 & -1 & i \\ -1 & 1 & 1 & 1 \\ -1 & -1 & -i & i \end{pmatrix}, \\
 &17. \begin{pmatrix} 1 & -1 & -1 & -i \\ -1 & 1 & 1 & -i \\ -1 & -1 & 1 & 1 \end{pmatrix}, 18. \begin{pmatrix} 1 & -1 & i & i \\ -1 & 1 & i & i \\ -i & -i & 1 & 1 \end{pmatrix}, 19. \begin{pmatrix} 1 & i & i & i \\ -i & 1 & 1 & -i \\ -i & i & 1 & 1 \end{pmatrix}, 20. \begin{pmatrix} 1 & i & i & -i \\ -i & 1 & 1 & -i \\ i & i & -i & 1 \end{pmatrix}.
 \end{aligned}$$

When we choose one  $(q_{ij})$  of the above 20 quantum parameters, then for any  $l$ , we have

$$\Gamma(D_+(t_l), \mathcal{B}) \simeq k\langle Y_1, Y_2, Y_3 \rangle / (Y_i Y_j - q'_{ij} Y_j Y_i, Y_1^4 + Y_2^4 + Y_3^4 + 1)_{1 \leq i, j \leq 3},$$

where  $(q'_{ij})$  is represented by one of the following matrices (we can verify this with direct calculations):

$$(a). \begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix}, \quad (b). \begin{pmatrix} 1 & -i & i \\ i & 1 & -i \\ -i & i & 1 \end{pmatrix}.$$

We write  $Z'_l := Z(\Gamma(D_+(t_l), \mathcal{B}))$ . When  $(q'_{ij})$  is type (a),  $\text{Spec}(Z'_l)$  has six singular points because  $Z'_l$  is generated by  $Y_1^2, Y_2^2, Y_3^2$  and  $Y_1 Y_2 Y_3$  as a  $k$ -algebra. When  $(q'_{ij})$  is type (b),  $\text{Spec}(Z'_l)$  has three singular points because  $Z'_l$  is generated by  $Y_1^4, Y_2^4, Y_3^4$  and  $Y_1 Y_2 Y_3$  as a  $k$ -algebra. Moreover, for any  $(q_{ij})$  in the above table, if  $\mathcal{B}$  is type (a) (resp. (b)) on  $D_+(t_l)$  for some  $l$ , it is also type (a) (resp. (b)) on  $D_+(t_l)$  for any other  $l$ .

*Step 4.* If  $\text{qgr}(C/(f))$  is equivalent to  $\text{qgr}(C'/(f'))$  then, we must have an isomorphism of schemes between  $\text{Spec}(\mathcal{Z}_{\mathcal{A}})$  and  $\text{Spec}(\mathcal{Z}_{\mathcal{B}})$  by [5, Theorem 4.4] (cf. [1, Section 6]). Since  $\text{Spec}(\mathcal{Z}_{\mathcal{A}})$  has infinitely many singular points, but,  $\text{Spec}(\mathcal{Z}_{\mathcal{B}})$  has finitely many singular points, such a situation does not happen. Hence, we complete the proof. ■

**Acknowledgments** The author is grateful to his supervisor Professor Hajime Kaji for his encouragement. He is also grateful to Professor Atsushi Kanazawa for telling him the articles [14, 15]. He would like to thank Professor Izuru Mori, Professor Ryo Ohkawa, Professor Shinnosuke Okawa, Professor Balázs Szendrői, and Professor Kenta Ueyama for helpful comments. In addition, he thanks Niklas Lemcke for proofreading his English. Finally, he thanks the referee for his careful reading of the manuscript.

## References

- [1] M. Artin and J. J. Zhang, *Noncommutative projective schemes*. Adv. Math. **109**(1994), no. 2, 228–287.
- [2] P. Belmans, K. De Laet, and L. Le Bruyn, *The point variety of quantum polynomial rings*. J. Algebra **463**(2016), 10–22.
- [3] A. Bondal and M. Van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*. Mosc. Math. J. **3**(2003), no. 1, 1–36.
- [4] M. P. Brodmann and R. Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*. Vol. 136, Cambridge University Press, Cambridge, 2012.
- [5] I. Burban and Y. Drozd, *Morita theory for non-commutative Noetherian schemes*. Adv. Math. **399**(2022), 1–42.
- [6] K. De Naeghel and M. Van den Bergh, *Ideal classes of three-dimensional Sklyanin algebras*. J. Algebra **276**(2004), no. 2, 515–551.
- [7] D. Eisenbud, *The geometry of syzygies: a second course in commutative algebra and algebraic geometry*. Vol. 229, Springer, New York, 2005.
- [8] V. Ginzburg, *Calabi–Yau algebras*. Preprint, 2006. [arXiv:0612139](https://arxiv.org/abs/0612139).
- [9] J.-W. He and K. Ueyama, *Twisted Segre products*. J. Algebra **611**(2022), 528–560.
- [10] E. Hyry, *The diagonal subring and the Cohen–Macaulay property of a multigraded ring*. Trans. Amer. Math. Soc. **351**(1999), no. 6, 2213–2232.
- [11] A. R. Iano-Fletcher, *Working with weighted complete intersections*. In: A. Corti and M. Reid (eds.), *Explicit birational geometry of 3-folds*. Vol. 281, London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2000, pp. 101–174.
- [12] A. Kanazawa, *Non-commutative projective Calabi–Yau schemes*. J. Pure Appl. Algebra **219**(2015), no. 7, 2771–2780.
- [13] A. Kuznetsov, *Calabi–Yau and fractional Calabi–Yau categories*. J. Reine Angew. Math. **2019**(2019), no. 753, 239–267.
- [14] Y.-H. Liu, *Donaldson–Thomas theory of quantum Fermat quintic threefolds I*. Preprint, 2019. [arXiv:1911.07949](https://arxiv.org/abs/1911.07949).
- [15] Y.-H. Liu, *Donaldson–Thomas theory of quantum Fermat quintic threefolds II*. Preprint, 2020. [arXiv:2004.10346](https://arxiv.org/abs/2004.10346).
- [16] J. C. McConnell, J. C. Robson, and L. W. Small, *Noncommutative Noetherian rings*. Vol. 30, American Mathematical Society, Providence, RI, 2001.
- [17] R. Mckemey, *Relative local cohomology*. Ph.D. thesis, University of Manchester, 2012.
- [18] I. Mori, *B-construction and C-construction*. Comm. Algebra **41**(2013), no. 6, 2071–2091.
- [19] I. Mori, *Regular modules over 2-dimensional quantum Beilinson algebras of type S*. Math. Z. **279**(2015), no. 3, 1143–1174.
- [20] I. Mori and A. Nyman, *Local duality for connected  $\mathbb{Z}$ -algebras*. J. Pure Appl. Algebra **225**(2021), no. 9, 106676.
- [21] I. Mori and A. Nyman, *A categorical characterization of quantum projective  $\mathbb{Z}$ -spaces*. Preprint, 2023. [arXiv:2307.15253](https://arxiv.org/abs/2307.15253).
- [22] I. Mori and A. Nyman, *Corrigendum to “Local duality for connected  $\mathbb{Z}$ -algebras” [J. Pure Appl. Algebra 225 (9)(2019) 106676]*. J. Pure Appl. Algebra **228**(2024), no. 3, 107493.
- [23] M. Reid, *Canonical 3-folds*. Journées de géométrie algébrique, Angers/France 1979(1980), 273–310.
- [24] M. Reyes and D. Rogalski, *Graded twisted Calabi–Yau algebras are generalized Artin–Schelter regular*. Nagoya Math. J. **245**(2022), 100–153.
- [25] M. Reyes, D. Rogalski, and J. J. Zhang, *Skew Calabi–Yau algebras and homological identities*. Adv. Math. **264**(2014), 308–354.
- [26] S. P. Smith, *Noncommutative algebraic geometry*, Lecture Notes, University of Washington, 2000. <https://sites.math.washington.edu/~smith/Research/nag.pdf>.
- [27] S. P. Smith, *Subspaces of non-commutative spaces*. Trans. Amer. Math. Soc. **354**(2002), no. 6, 2131–2171.
- [28] S. P. Smith, *Maps between non-commutative spaces*. Trans. Amer. Math. Soc. **356**(2004), no. 7, 2927–2944.
- [29] D. R. Stephenson, *Quantum planes of weight  $(1, 1, n)$* . J. Algebra **225**(2000), no. 1, 70–92.
- [30] M. Van den Bergh, *Existence theorems for dualizing complexes over non-commutative graded and filtered rings*. J. Algebra **195**(1997), no. 2, 662–679.
- [31] M. Van den Bergh, *Calabi–Yau algebras and superpotentials*. Selecta Math. (N.S.) **21**(2015), no. 2, 555–603.

- [32] K. Van Rompay, *Segre product of Artin–Schelter regular algebras of dimension 2 and embeddings in quantum  $\mathbb{P}^3$ s*. J. Algebra 180(1996), no. 2, 483–512.
- [33] J. Vitoria, *Equivalences for noncommutative projective spaces*. Preprint, 2010. [arXiv:1001.4400](https://arxiv.org/abs/1001.4400).
- [34] R. Vyas and A. Yekutieli, *Weak proregularity, weak stability, and the noncommutative MGM equivalence*. J. Algebra 513(2018), 265–325.
- [35] A. Yekutieli, *Dualizing complexes over noncommutative graded algebras*. J. Algebra 153(1992), no. 1, 41–84.
- [36] A. Yekutieli, *Derived categories*. Vol. 183, Cambridge University Press, Cambridge, 2019.
- [37] A. Yekutieli and J. J. Zhang, *Serre duality for noncommutative projective schemes*. Proc. Amer. Math. Soc. 125(1997), no. 3, 697–707.
- [38] J. J. Zhang, *Twisted graded algebras and equivalences of graded categories*. Proc. Lond. Math. Soc. 3(1996), no. 2, 281–311.

*Department of Mathematics, School of Science and Engineering, Waseda University, Ohkubo 3-4-1, Shinjuku, Tokyo 169-8555, Japan*

*e-mail:* [m7d5932a72xxgx0@fuji.waseda.jp](mailto:m7d5932a72xxgx0@fuji.waseda.jp)