

pH-mediated inhibition of a bumble bee parasite by an intestinal symbiont

Evan C. Palmer-Young¹, Thomas R. Raffel² and Quinn S. McFrederick¹¹Department of Entomology, University of California Riverside, Riverside, CA, USA and ²Department of Biology, Oakland University, Rochester, MI, USA

Research Article

Cite this article: Palmer-Young EC, Raffel TR, McFrederick QS (2019). pH-mediated inhibition of a bumble bee parasite by an intestinal symbiont. *Parasitology* **146**, 380–388. <https://doi.org/10.1017/S0031182018001555>

Received: 28 June 2018
Revised: 10 August 2018
Accepted: 16 August 2018
First published online: 24 September 2018

Key words:

Cell-free supernatant; competition; Firmicutes; flagellates; lactic acid bacteria; microbiome; parasite strain variation; pH-dependent interactions; pollinator decline; Trypanosomatidae

Author for correspondence:

Evan C. Palmer-Young,
E-mail: ecp52@cornell.edu

Abstract

Gut symbionts can augment resistance to pathogens by stimulating host-immune responses, competing for space and nutrients, or producing antimicrobial metabolites. Gut microbiota of social bees, which pollinate many crops and wildflowers, protect hosts against diverse infections and might counteract pathogen-related bee declines. Bumble bee gut microbiota, and specifically abundance of *Lactobacillus* ‘Firm-5’ bacteria, can enhance resistance to the trypanosomatid parasite *Crithidia bombi*. However, the mechanism underlying this effect remains unknown. We hypothesized that the Firm-5 bacterium *Lactobacillus bombicola*, which produces lactic acid, inhibits *C. bombi* via pH-mediated effects. Consistent with our hypothesis, *L. bombicola* spent medium inhibited *C. bombi* growth via reduction in pH that was both necessary and sufficient for inhibition. Inhibition of all parasite strains occurred within the pH range documented in honey bees, though sensitivity to acidity varied among strains. Spent medium was slightly more potent than HCl, D- and L-lactic acids for a given pH, suggesting that other metabolites also contribute to inhibition. Results implicate symbiont-mediated reduction in gut pH as a key determinant of trypanosomatid infection in bees. Future investigation into *in vivo* effects of gut microbiota on pH and infection intensity would test the relevance of these findings for bees threatened by trypanosomatids.

Introduction

Both animals and plants associate with symbiotic bacterial communities that provide functional benefits to their hosts (Mueller and Sachs, 2015). The gut microbiota comprises some of the largest and most well-studied communities of host-associated symbionts (Kamada *et al.*, 2013). The bacterial communities that colonize the gut and skin epithelia interact directly with potential pathogens in the environment, and can influence infection by stimulation of the host-immune system, competition for space and nutrients and production of inhibitory substances such as organic acids and bacteriocins (Dillon and Dillon, 2004; Round and Mazmanian, 2009; Daskin *et al.*, 2014; Raymann and Moran, 2018). Interactions between the gut microbiota and pathogens of bees are an emerging area of research with both fundamental and applied importance (Gaggia *et al.*, 2018; Raymann and Moran, 2018). Elucidation of the antipathogenic potential of the bee microbiota may ultimately help to preserve the pollination services provided by both wild and managed bees, which improve yields of over two-thirds of common agricultural crops (Klein *et al.*, 2007) and contribute to the >\$150 billion per year in economic value supplied by animal pollination (Gallai *et al.*, 2009).

The gut microbiota of corbiculate (‘pollen basket’) bees, including honey and bumble bees, comprises a common core of 5 bacterial clades: *Snodgrassella* (Betaproteobacteria), *Gilliamella* (Gammaproteobacteria), *Bifidobacterium* and *Lactobacillus* clades ‘Firmicutes-4’ and ‘Firmicutes-5’ (Kwong *et al.*, 2017b). In addition to these core symbionts, bees may be infected by a variety of bacterial, fungal, protozoal and viral pathogens (Evans and Spivak, 2010), many of which are shared between wild and managed bees (Graystock *et al.*, 2016), can elevate mortality (Fürst *et al.*, 2014) and have been implicated in declines of bee populations on multiple continents (Cameron *et al.*, 2011; Schmid-Hempel *et al.*, 2014; Goulson *et al.*, 2015). Both core and non-core microbiota have been found to stimulate immunity and enhance bee resistance to pathogens (Evans and Lopez, 2004; Engel *et al.*, 2016; Kwong *et al.*, 2017a; Raymann and Moran, 2018). For example, depletion or perturbation of the gut microbiota increased the severity of bacterial, fungal and protozoal infections in honey bees (Schwarz *et al.*, 2016; Kwong *et al.*, 2017a; Raymann and Moran, 2018), whereas supplementation with core and hive-associated bacteria improved survival of infected larvae and adults (Forsgren *et al.*, 2010; Vásquez *et al.*, 2012; Kwong *et al.*, 2017a). Several studies have shown direct inhibitory effects of gut and hive-associated symbionts against common bee pathogens (Evans and Armstrong, 2006; Sabaté *et al.*, 2009; Praet *et al.*, 2018), which suggests a parsimonious explanation for the effects of symbionts on infection.

The gut microbiota of bumble bees (*Bombus* spp.) has been repeatedly associated with resistance to infection with the trypanosomatid gut parasite *Crithidia bombi* (Koch and Schmid-Hempel, 2012; Koch *et al.*, 2012; Cariveau *et al.*, 2014; Mockler *et al.*, 2018). This parasite has a variety of negative effects on bees. Symptoms include reduced rates of foraging, pollen collection and learning for worker bees (Shykoff and Schmid-Hempel, 1991; Gegear

et al., 2005, 2006); and reduced winter survival and spring nest-founding success for queen bees (Brown *et al.*, 2003). Infection with *C. bombi* can also exacerbate susceptibility to co-occurring stressors such as starvation, pesticides (Fauser-Misslin *et al.*, 2014) and nectar alkaloids (Palmer-Young *et al.*, 2017b). Trypanosomatid infections appear to be common in corbiculate bees, afflicting over half of individuals in some honey and bumble bee populations (Schmid-Hempel and Ebert, 2003; Cornman *et al.*, 2012), and have been correlated with honey bee colony collapses (Cornman *et al.*, 2012; Ravoet *et al.*, 2013) and native bumble bee declines (Schmid-Hempel *et al.*, 2014). Both the presence and composition of the bumble bee microbiota may improve resistance to *C. bombi*. For example, germ-free rearing conditions and treatment with antibiotics both resulted in higher infection intensity in *Bombus terrestris* (Koch and Schmid-Hempel, 2011, 2012). In contrast, absolute and relative abundance of select core gut bacteria were correlated with resistance to infection in both field surveys and fecal transplant experiments (Koch and Schmid-Hempel, 2011, 2012; Cariveau *et al.*, 2014; Mockler *et al.*, 2018). Just 3 taxa – *Snodgrassella*, *Gilliamella* and *Lactobacillus* Firm-5 – generally account for over 80% of the total gut bacteria in bumble bees (Koch and Schmid-Hempel, 2012; Billiet *et al.*, 2017; Mockler *et al.*, 2018). In both *B. terrestris* and *B. impatiens*, inoculation with microbiota rich in *Lactobacillus* Firm-5 resulted in resistance to *C. bombi* infection (Koch and Schmid-Hempel, 2012; Mockler *et al.*, 2018). However, no study has examined the mechanisms by which these microbes influence parasite growth.

Lactobacillus Firm-5 is a group of lactic acid-producing bacteria (Praet *et al.*, 2015). One way that they might increase host resistance to parasites is *via* modification of gut chemistry. Lactic acid fermentation results in production of organic acids that lower pH and inhibit growth of organisms that cause spoilage and infection (Adams and Hall, 1988; Lindgren and Dobrogosz, 1990; Glass *et al.*, 1992; Russell and Diez-Gonzalez, 1998). Indeed, lactic acid bacteria have a long history of use in food preservation in both human and insect societies (Salminen and von Wright, 2004; Anderson *et al.*, 2014). In the host intestine, lactic acid bacteria can inhibit enteric pathogens, such as *Salmonella* and *Escherichia coli* (Gorbach, 1990). This inhibition may reflect stimulation of the host-immune system (Presser *et al.*, 1997; Cox *et al.*, 2010), including that of insects (Evans and Lopez, 2004; Evans and Armstrong, 2006). However, *Lactobacillus*-mediated inhibition of pathogens is most simply explained by the direct antimicrobial activity of *Lactobacillus* metabolites. These metabolites, which include lactic acid and bacteriocins (Lindgren and Dobrogosz, 1990), may reduce the suitability of the gut environment for pathogens.

In the bee gut, *Lactobacillus* Firm-5 has been shown to have a disproportionately large effect on gut metabolomics. In honey bees, mono-inoculation with Firm-5 accounted for over 80% of the changes seen in bees inoculated with a full complement of gut microbes (Kešnerová *et al.*, 2017). Firm-5 isolates also showed *in vitro* inhibitory activity against the pathogens *Paenibacillus larvae* and *Melissococcus plutonius* (Praet *et al.*, 2018). The high relative abundance of the Firm-5 clade in bumble bees [often >30% total bacteria (Koch and Schmid-Hempel, 2012; Billiet *et al.*, 2017)], combined with its consistent association with resistance to trypanosomatid infection, suggests that *Lactobacillus* Firm-5 plays a major role in bumble bee resistance to trypanosomatid parasites.

We hypothesized that *Lactobacillus* Firm-5 enhances resistance to trypanosomatid parasites primarily by modifying the pH of the enteric environment. To test this hypothesis, we measured the inhibitory effects of spent medium from *Lactobacillus bombicola*, a member of the Firmicutes-5 clade that is ubiquitous in the gut

microbiota of corbiculate bees, including *Bombus* spp. (Billiet *et al.*, 2017; Kwong *et al.*, 2017b; Praet *et al.*, 2018), on *in vitro* growth of several strains of *C. bombi*. We predicted that spent medium from *L. bombicola* would inhibit *C. bombi* growth, that the acidity of spent medium would be both necessary and sufficient to account for parasite inhibition, and that *C. bombi* strains would vary in sensitivity to spent medium.

Materials and methods

Overview of experiments

Three experiments were conducted to evaluate effects of spent medium from *L. bombicola* cultures on growth of *C. bombi*. Spent medium was generated by growth of *L. bombicola* in MRS broth for 3 days, followed by sterile filtration to remove live cells. For *C. bombi* growth assays, the MRS-based spent medium (or MRS broth control) was diluted 1:1 in fresh, *Crithidia*-specific 'FPFB' medium (Salathé *et al.*, 2012). (1) The neutralization experiment tested whether spent medium would inhibit growth, and whether acidity of the spent medium was necessary or sufficient for inhibition. (2) The acidification experiment tested for variation in pH-dependent growth inhibition due to various sources of acidity. (3) The strain variation experiment tested for variation in sensitivity to spent medium among different parasite strains.

Cell cultures

Lactobacillus bombicola strain 70-3, isolated from *B. lapidarius* collected near Ghent Belgium [isolate '28288T' (Praet *et al.*, 2015)], was obtained from the DSMZ and grown in 2 mL screw-cap tubes in MRS broth (Research Products International, Mt. Prospect, IL) with 0.05% cysteine at 27 °C. *Crithidia bombi* cell cultures were isolated from bumble bee intestines by flow cytometry based single cell sorting as described previously (Salathé *et al.*, 2012). Cultures originated from wild infected bumble bees. Strains VT1 (Vermont, USA, 2013, courtesy Rebecca Irwin) and IL13.2 (Illinois, USA, 2013, courtesy Ben Sadd) originated from *B. impatiens* workers. Strains S08.1 (Switzerland, 2008, courtesy Ben Sadd) originated from *B. terrestris*. These same cell lines have been used to assess effects of phytochemicals on parasite growth (Palmer-Young *et al.*, 2017a). Briefly, cells from fecal samples were sorted into 96-well plates containing 'FPFB' culture medium with 10% heat-inactivated fetal bovine serum and incubated at 27 °C. Cultures with successful growth and absence of visible contamination were transferred to vented, 25 cm² tissue culture flasks, grown to high density and cryopreserved at –80 °C until several weeks before the experiments began (Salathé *et al.*, 2012). Culture identity was confirmed as *C. bombi* based on glyceraldehyde 3-phosphate dehydrogenase and cytochrome *b* gene sequences. Cultures were inspected weekly to verify absence of contamination.

The neutralization and acidification experiments were performed with a line of strain 'VT1' that had been in continuous culture for 2 months at the start of the experiments presented here, with transfers to fresh medium every 3–4 days. This line is referred to as 'VT1*' in the multi-strain experiment, to differentiate it from the more recently thawed line of the same strain. All other strains in the multi-strain experiment were thawed 20 days prior to the assay and transferred to fresh medium every 3–7 days, depending on growth.

Generation of spent medium

To generate spent medium, *L. bombicola* aliquots were transferred to 8 mL fresh medium and grown in screw-cap 14 mL

conical tubes at 27 °C for 3 days. The resulting spent medium (net OD_{630 nm} = 0.500–0.700) was sterile-filtered through a 0.22 µm membrane and stored at –20 °C until use in experiments (not more than 2 weeks). Fresh MRS medium, incubated under identical conditions, was used as a control.

Neutralization experiment

To evaluate the inhibitory effects of spent medium, neutralized spent medium and acidified fresh medium, growth of *C. bombi* was compared across 4 MRS-based treatments: spent medium ('Spent', initial pH 4.8), spent medium neutralized to pH 6.2 with 1 M NaOH ('Neutralized spent'), fresh MRS medium ('Fresh', pH 6.2), and fresh MRS medium acidified to pH 4.8 with 1 M HCl ('Acidified fresh'). The *C. bombi* culture was diluted to an OD of 0.020 in fresh FPFb medium (pH 5.88). The resulting cell suspension (100 µL) was added to wells of a 96-well plate containing an equal volume of the MRS-based treatment medium, resulting in a final net OD_{630 nm} of 0.010. Growth was measured twice daily by optical density (630 nm) over the ensuing 48 h on an EL-800 plate reader spectrophotometer (BioTek, Winooski, VT). Net optical density at each time point was computed by subtracting OD of wells containing the corresponding MRS-based treatment medium and FPFb medium without *C. bombi*; this controlled for any differences in optical density that occurred independent of *C. bombi* growth. The experiment included 18 replicate wells per treatment.

Acidification experiment

To compare variation in growth inhibition across different sources of acidity, growth of *C. bombi* was compared in dilutions of spent medium (initial pH 4.65), and in fresh medium acidified with D-lactic acid (pH 4.82), L-lactic acid (pH 4.77) or HCl (pH 4.73). Each base medium was diluted with fresh MRS medium to 0, 20, 40, 60, 80 or 100% of initial concentration. The final pH of each treatment was measured with a pH meter ('Orion Star', Thermo, Waltham, MA) after combination with an equal volume of FPFb medium. Growth of *C. bombi* (initial OD 0.010) was evaluated with a 96-well plate assay as in the neutralization experiment above. The experiment included 12 replicate wells per concentration of each acidification treatment. To verify the relative potency of spent vs acidified medium against different parasite strains, the experiment was repeated with 2 *C. bombi* strains (IL13.2 and VT1*) tested in parallel against 2 of the 4 acidification treatments (spent medium and HCl-acidified medium); results are shown in Supplementary Fig. 2.

Strain variation experiment

To compare susceptibility to spent medium across different *C. bombi* strains and degrees of acclimation to the culture environment, inhibitory concentration of spent medium was compared across 4 cell lines: the 'VT1*' line of strain VT1 that had been used for the above experiments, and by this time had been in continuous culture for 3 months; and strains VT1, IL13.2 and S08.1 that had been thawed 20 days prior to the experiment. Six dilutions of spent medium (0–100%) were prepared in fresh MRS medium; final pH of each treatment was measured after combination with an equal volume of FPFb medium. Growth of each line of *C. bombi* (initial OD 0.010) was evaluated with a 96-well plate assay as in the neutralization experiment above. The experiment included 12 replicate wells per cell line and spent medium concentration.

Statistical analysis

Statistical analysis was performed in R v3.4.3 for Windows (R Core Team, 2014). For the acidification and strain variation experiments, dose–response curves to relate growth rate to pH were estimated with the *drm* function in the R package *drc* (Ritz *et al.*, 2015). Under the conditions of the experiment (initial OD = 0.010, incubation temperature 32 °C), *C. bombi* growth was found to be exponential over the first 24 h of incubation (Supplementary Fig. 1). Therefore, the rate of growth ($\log_2(\text{OD}/\text{OD}_0)/t$) over the first incubation interval (0–21 h) was used as the response variable for the dose–response models. A log-logistic function, with the lower asymptote fixed at 0, was fitted to the growth measurements for each acidification treatment (acidification experiment) or cell line (strain variation experiment), using final pH of the treatment medium at the start of the experiment (i.e. after combination of the MRS-based treatment with an equal volume FPFb medium) as the predictor variable. The 95% confidence intervals (CIs) for EC₅₀ pH (i.e. the pH that inhibited growth by 50% relative to the unacidified control treatment), and 95% CIs for ratios of EC₅₀ pH among acidification treatments and strains, were estimated using the delta method [*drc* function *EDcomp* (Ritz *et al.*, 2015)]. Acidification treatments and strains were considered significantly different when the 95% CI for the ratio of their EC₅₀ pH values did not include 1.

Results

Neutralization experiment: acidity dependent inhibition of *C. bombi* by spent medium

We found that *L. bombicola* spent medium completely inhibited growth of *C. bombi* cell cultures (Fig. 1). However, neutralized spent medium had no inhibitory effect, indicating that acidity of the spent medium was necessary for inhibition (Fig. 1). Moreover, acidification of fresh (*Lactobacillus*-specific) MRS medium to pH 4.8 with L-lactic acid led to a level of growth inhibition that was comparable to that caused by pH 4.8 spent medium (Fig. 1). This demonstrated that spent medium from *L. bombicola* inhibited *C. bombi* growth, and that changes in pH were necessary and qualitatively sufficient to account for this inhibition.

Acidification experiment: pH-dependent inhibition of *C. bombi* with different sources of acidity

Having established pH-dependent growth inhibition, we conducted a follow-up experiment to quantify *C. bombi* growth rates across a range of pH values, and to compare the relative inhibitory effects of *L. bombicola* spent medium with that of 3 other sources of acidity: HCl, L-lactic acid [the form produced by animal cells (Ewaschuk *et al.*, 2002)] and D-lactic acid [the form produced by *L. bombicola* (Praet *et al.*, 2015)]. We used log-logistic models to estimate and compare the EC₅₀ pH (i.e. the pH that inhibited growth by 50% relative to the unacidified control treatment) for each source of acidity. All 4 sources of acidity resulted in qualitatively similar inhibition of *C. bombi* (Fig. 2A), with considerable inhibition achieved within the pH range previously measured in honey bee guts [vertical lines in Fig. 2A, from (Zheng *et al.*, 2017); no data are available on pH of bumble bee guts]. However, the EC pH varied somewhat across sources of acidity, with a hierarchy of pH-dependent inhibitory potency in the order HCl < L-lactic acid < D-lactic acid < spent medium (Fig. 2B, see Supplementary Table S1 for EC₅₀ values and Supplementary Data S1 for model parameters, CIs on EC₅₀ ratios and raw data). An additional experiment with a second *C. bombi* strain ('IL13.2') confirmed the greater potency of




Fig. 1. Spent medium from *Lactobacillus bombicola* inhibited growth of *Crithidia bombi* (strain VT1); acidity of the spent medium was necessary and sufficient for inhibition. Spent medium ('Spent', red solid line) completely inhibited parasite growth. However, spent medium neutralized to pH 6.2 with NaOH ('Neutralized spent', green dotted line) resulted in no inhibition relative to the fresh medium control ('Fresh', light green dashed line), demonstrating that acidity was necessary for inhibition. Fresh medium acidified to pH 4.8 with lactic acid ('Acidified fresh', blue dashed line) showed that acidity was sufficient for complete growth inhibition. Final pH of both spent medium and acidified fresh medium was 5.0 after combination with equal volume of fresh *Crithidia* medium (pH 5.9). Y-axis represents approximate number of parasite cell divisions, as measured by optical density (630 nm). Points and error bars show means and 95% CIs ($n = 36$ wells per treatment).

spent medium relative to HCl-acidified medium across multiple parasite genotypes (Supplementary Fig. 2).

Strain variation experiment: sensitivity to spent medium differs across *C. bombi* strains and rate of growth in culture (Fig. 3)

We tested the dose-dependent effects of spent medium across 3 *C. bombi* strains, including 2 lines of strain VT1 – the line that had been used for the above experiments and kept in continuous culture over the preceding 3 months ('VT1*'), and a second line that had been thawed 20 days prior to the experiment ('VT1'). Values for EC₅₀ pH showed statistically and biologically meaningful variation across cell lines (Fig. 3). The line most thoroughly acclimated to the culturing conditions, VT1*, exhibited both the fastest growth (Fig. 3A) and the lowest sensitivity to spent medium, as indicated by its low EC₅₀ pH value (Fig. 3B); strain IL13.2 exhibited the second-fastest growth and the second-lowest sensitivity to spent medium. Strains VT1 and S08.1 had similar growth rates in the absence of spent medium (Fig. 3A), but had significantly different EC₅₀ pH values (Fig. 3B; see Supplementary Table S1 for EC₅₀ values and Supplementary Data S1 for model parameters, CIs on EC₅₀ ratios and raw data). All *C. bombi* strains suffered growth inhibition within the pH range (5.2–6.0) documented in the honey bee hindgut (Zheng *et al.*, 2017) (Fig. 3A).

Discussion

Our results show that production of acids by the core bumble bee hindgut symbiont, *L. bombicola*, is both necessary and sufficient to inhibit growth of the widespread trypanosomatid parasite *C. bombi*. The growth-inhibitory effects of *L. bombicola*-acidified spent medium were qualitatively similar across *C. bombi* isolates and occurred within a pH range that is physiologically realistic for the bee hindgut where *C. bombi* establishes. Because lactobacilli and other acid-producing bacteria are dominant members of the bumble bee gut microbiota, our results suggest that the inhibitory effects of bumble bee gut microbiota on *C. bombi* infection intensity can be largely attributed to microbial acidification

of the gut. Our findings provide a mechanistic basis to understand how microbiota may affect trypanosomatid infection in corbiculate bees that share a core microbiome (Kwong *et al.*, 2017b) and can be infected with identical and related parasites, including trypanosomatids (McMahon *et al.*, 2015; Schwarz *et al.*, 2015; Tripodi *et al.*, 2018).

Inhibitory activity of *L. bombicola* spent medium is driven by production of acids

The effects of *L. bombicola* spent medium on *C. bombi* growth could be explained by the acidity of the spent medium. Environmental pH is a recognized driver of interactions in microbial communities, where species vary in how they alter the pH of their surroundings and in the pH range at which they can grow (Morton *et al.*, 2017; Ratzke and Gore, 2018). In honey bees, microbial colonization with core symbionts resulted in acidification of the hindgut lumen (Zheng *et al.*, 2017). In our cell culture experiments, a reduction in pH that corresponds to the difference between guts of germ-free (pH ~ 6.0) and normal, hive-reared honey bees [pH ~ 5.2 (Zheng *et al.*, 2017)] profoundly inhibited growth of *C. bombi*. Our findings indicate that symbiont-mediated gut acidification could act as an important filter that prevents trypanosomatid establishment and defends bees against infection.

Spent medium was more strongly inhibitory than expected based on pH alone

Whereas all 4 tested acids inhibited *C. bombi* in a dose-dependent fashion, the EC₅₀ pH varied across sources of acidity (Fig. 2). Both enantiomers of lactic acid were more inhibitory than HCl for a given pH. This finding is consistent with prior work on bacteria, which showed growth inhibition at relatively high pH when lactic acid, rather than HCl, was used as the acidulant (Adams and Hall, 1988; Glass *et al.*, 1992). For example, the inhibitory pH of *E. coli* was 0.1 pH units higher with lactic acid, rather than HCl, as the acidulant (Glass *et al.*, 1992). This increased activity reflects the fact that lactic acid and other weak acids are undissociated at low pH, which allows them to penetrate membranes of target cells with relative ease. Once inside the relatively alkaline cytoplasm of the cell, the acid dissociates, and disrupts the proton motive force necessary for energy production and homeostasis (Russell and Diez-Gonzalez, 1998).

We also found very slightly but significantly higher potency of D-lactic acid relative to L-lactic acid (Fig. 2). L-Lactic acid is the enantiomer more often produced by trypanosomatids, including by *Leishmania* species that are closely related to *C. bombi* (Bringaud *et al.*, 2006), and L-lactic acid dehydrogenase has been found in the genome of honey bee-infective trypanosomatids (Runckel *et al.*, 2014). It is possible that *C. bombi* can use oxidative phosphorylation (Bringaud *et al.*, 2006) to metabolize this enantiomer, and thereby reduce its toxicity, or that *C. bombi* have greater resistance to the enantiomer that they produce through their own metabolism. For example, *L. bulgaricus* produced D-lactic acid, and were less inhibited by this self-generated enantiomer than by L-lactic acid (Benthin and Villadsen, 1995).

Prior measurements of the pH range and sources of acidity in honey bee gut both indicate that pH-mediated inhibition of trypanosomatids is achievable for corbiculate bees. The pH of a symbiont-colonized honey bee hindgut [~5.2 (Zheng *et al.*, 2017)] was similar to the EC₅₀ pH for HCl (5.18) and lower than the EC₅₀ pH due to spent medium (5.29). *In vitro*, *L. bombicola* produced exclusively D-lactic acid (Praet *et al.*, 2015), which was the most inhibitory (i.e. highest EC₅₀ pH) of the pure acids tested here (Fig. 3). Moreover, the specific acids that are found




Fig. 2. Different sources of acidity varied in pH-dependent inhibitory potency against *C. bombi* (strain VT1). (A) Dose–response curves relating pH to growth rate. X-axis shows final pH of treatment medium after combination of *L. bombicola* spent medium or acidified MRS medium with an equal volume of *Crithidia*-specific FPF medium. Y-axis represents growth rate over first 21 h of incubation, measured as number of doublings per day by optical density (OD: 630 nm). Lines and shaded bands represent model predictions and standard errors for each source of acidity. Points show raw data for each replicate well ($n = 12$). Vertical lines correspond to pH values measured in ileum and rectum of microbe-colonized and germ-free honey bees (Zheng *et al.*, 2017), and are shown as an estimate of gut pH range in bumble bees. (B) 50% inhibitory concentrations (EC₅₀, i.e. the concentration that inhibits growth by 50%) for different sources of acidity. Point estimates and 95% CIs are derived from model fits shown in panel (A). Higher EC₅₀ pH estimates correspond to higher inhibitory potency for a given level of acidity. Different lower-case letters indicate statistically significant ($P < 0.05$) differences in pairwise comparisons of EC₅₀ pH by source of acidity.




Fig. 3. *Crithidia bombi* pH sensitivity varied according to strain identity and rate of growth in culture. (A) Dose–response curves relating pH to growth rate for different *C. bombi* strains. X-axis shows final pH of treatment medium after combination of *L. bombicola* spent MRS medium with an equal volume of *Crithidia*-specific FPF medium. Y-axis represents growth rate over first 20 h of incubation, measured as number of doublings per day by optical density (OD: 630 nm). Lines and shaded bands represent model predictions and standard errors for each strain. Points show raw data for each replicate well ($n = 12$). Vertical lines correspond to pH values measured in ileum and rectum of microbe-colonized and germ-free honey bees (Zheng *et al.*, 2017), and are shown as an estimate of gut pH range in bumble bees. Strains ‘VT1*’ and ‘VT1’ are the same strain, but ‘VT1*’ had been grown in continuous culture for 3 months prior to the experiment, whereas ‘VT1’ and all other strains had been thawed from cryopreserved stock 3 weeks prior. (B) 50% inhibitory concentrations (EC₅₀) for each strain. Point estimates and 95% CIs are derived from model fits shown in panel (A). Higher EC₅₀ pH estimates correspond to higher sensitivity to acidity. Different lower-case letters indicate statistically significant ($P < 0.05$) differences in pairwise comparisons of EC₅₀ pH by source of acidity. For all strains, inhibition occurred within the pH range measured in honey bee guts (data from Zheng *et al.*, 2017).

in the gut environment are generally more inhibitory for a given pH than those tested here. In the honey bee hindgut, the most abundant acids were acetic acid and succinic acid (Zheng *et al.*, 2017). Both acetic acid ($pK_a = 4.95$) and succinic acid ($pK_a = 4.2$ for first deprotonation and 5.6 for second deprotonation) have relatively high pK_a s relative to lactic acid ($pK_a = 3.86$) and HCl ($pK_a = -7$). These high pK_a values mean that at low pH, the acids are found in their more potent, undissociated form. As a result, these high- pK_a acids generally have antimicrobial effects that are even stronger than those of lactic acid for a given pH (Adams and Hall, 1988). Indeed, lactic and acetic acids can have synergistic effects against growth of *E. coli* (Adams and Hall, 1988), with lactic acid producing a low pH that increases the fraction of undissociated acetic acid (Adams and Hall, 1988).

Spent medium was slightly more inhibitory than all pure acids for a given pH (Fig. 3). Because neutralization completely removed the inhibitory activity of spent medium, it appears that some aspect of the spent medium – whether the existence of some metabolite or the relative lack of nutrients – is only inhibitory at low pH. In other words, something about the spent medium is potentiated by an acidic environment. For example, low pH could facilitate solubility or penetration of non-lactic acid components, such as bacteriocins; this type of synergy was seen in other studies of *Lactobacillus* spp. (Fayol-Messaoudi *et al.*, 2005; De Keersmaecker *et al.*, 2006). Within the bee gut, synergistic effects could also occur between organic acids and toxins produced by other members of the microbiota (Praet *et al.*, 2015; Steele *et al.*, 2017).

Strains varied in sensitivity to spent medium

We found that sensitivity to spent medium varied by *C. bombi* strain and rate of growth in culture. Strain VT1 that had been in continuous culture for 3 months and had the fastest growth rate was the least sensitive to spent medium, followed by the next-fastest strain IL13.2, the recently thawed line of strain VT1, and strain S08.1. The comparison between the recently thawed VT1 and S08.1 strains – which had similar maximal growth rates, but different levels of sensitivity to spent medium – indicates that pH sensitivity can have a genotypic basis and is not purely driven by the overall growth rate in culture. Strains of *C. bombi* have been shown to be both genetically and phenotypically diverse, and to vary in growth rate (Imhoof and Schmid-Hempel, 1998), infectivity (Barribeau *et al.*, 2014), and responses to host diet composition (Sadd, 2011), phytochemicals (Palmer-Young *et al.*, 2016) and microbiota (Koch and Schmid-Hempel, 2012). Our study documents variation in pH sensitivity within an ecologically relevant range of pH that is likely representative of the environment in the bumble bee gut.

Our results showed some growth inhibition of all strains – and complete inhibition of some strains – within the pH range measured in the gut of honey bees (Zheng *et al.*, 2017). The pH found in a germ-free gut (5.8–6.0) was favourable for growth, consistent with the high *C. bombi* infection intensities found in germ-free and antibiotic-treated bees (Koch and Schmid-Hempel, 2011), whereas pH of a symbiont-colonized gut (<5.2) would be expected to inhibit growth of all strains. Thus, pH sensitivity may constrain ability of strains to colonize certain host genotypes (Barribeau *et al.*, 2014) or enterotypes (Koch and Schmid-Hempel, 2012; Li *et al.*, 2015) characterized by low gut pH. For example, inoculation with a microbiota high in *Lactobacillus* Firm-5 resulted in lower overall infection intensity and favoured infection with a single parasite strain that was less successful in bees inoculated with microbiota low in Firm-5 (Koch and Schmid-Hempel, 2012). We hypothesize that low-pH conditions favour strains that are more tolerant, whereas high-pH conditions

favour strains that are strong competitors. Further experiments are needed to investigate the extent to which parasite populations are selected for pH tolerance, and possible trade-offs between growth rate, infectivity or tolerance to environmental stressors and insect immune factors. Environment-dependent selection for these other traits could maintain variation in pH tolerance within parasite populations.

Gut microbiota-driven changes in pH may explain patterns of trypanosomatid infection in bees

The pH-dependent inhibition demonstrated here is consistent with past surveys and experiments that showed negative correlations between the abundance of acid-producing gut symbionts and *C. bombi* infection intensity, and with associations between microbiota composition and relative infectivity of different parasite strains. The *Bombus* gut microbiota is dominated by 3 taxa – *Lactobacillus* Firm-5, *Gilliamella* and *Snodgrassella* – that made up over 80% of total gut bacteria (Koch and Schmid-Hempel, 2012; Billiet *et al.*, 2017). In fecal transplant experiments of *B. terrestris* and *B. impatiens*, relative abundances of *Lactobacillus* Firm-5 and *Gilliamella* were negatively correlated with *C. bombi* infection intensity (Koch and Schmid-Hempel, 2011; 2012; Koch *et al.*, 2012; Mockler *et al.*, 2018). Both *Lactobacillus* and *Gilliamella* ferment sugars to produce acids (Engel *et al.*, 2012; Kešnerová *et al.*, 2017).

In contrast, *Snodgrassella* abundance was not correlated with resistance to *C. bombi* in either *B. terrestris* or *B. impatiens* (Koch and Schmid-Hempel, 2012; Mockler *et al.*, 2018), and pre-inoculation with *Snodgrassella alvi* reduced resistance to trypanosomatid infection in *Apis mellifera* (Schwarz *et al.*, 2016). Whereas *Lactobacillus* and *Gilliamella* spp. produce acids from sugars, *Snodgrassella* consumes organic acids (Kwong and Moran, 2013; Kešnerová *et al.*, 2017). This metabolic activity could elevate gut pH to levels that are more hospitable to trypanosomatids. On the other hand, *Snodgrassella* abundance was negatively correlated with infection prevalence in a field survey (Cariveau *et al.*, 2014). This correlation might reflect the association of *Snodgrassella* with acid-producing *Gilliamella* (Kešnerová *et al.*, 2017), rather than inhibitory effects of *Snodgrassella per se*. *Snodgrassella* and *Gilliamella* form a biofilm that lines the ileum (Engel *et al.*, 2012) and could competitively inhibit trypanosomatid attachment to the gut wall (Gorbunov, 1996; Schwarz *et al.*, 2015). Further study would be needed to determine the relative contributions of gut acidification vs biofilm formation to microbiota-induced inhibition of parasites in bee guts. It would be intriguing to investigate how *Snodgrassella* abundance alters gut pH, and whether the effects of this symbiont on gut pH are outweighed by its direct competition with trypanosomatids for space along the gut wall, or by its contribution to anoxic gut environments (Zheng *et al.*, 2017) that can be toxic to insect gut trypanosomatids (Bringaud *et al.*, 2006). Because *Apis* and *Bombus* share a core gut microbiota (Martinson *et al.*, 2011; Kwong and Moran, 2016), we hypothesize that trypanosomatid parasites of honey bees (Schwarz *et al.*, 2015) and bumble bees interact with similar communities of gut symbionts, and that the same pH-altering symbiont taxa could govern parasite establishment in both host genera.

Do gut microbiota shape resistance to opportunistic infection via alteration of gut pH?

In the context of prior experiments, our results suggest a pH-mediated role of the bee microbiota in defence against opportunistic infection. Multiple studies have correlated lack of core gut bacteria, or abundance of non-core bacteria, with bee infection. In

North America and Europe, high diversity of non-core bacterial species correlated with higher prevalence of the pathogens *Crithidia* and *Nosema* (Koch *et al.*, 2012; Cariveau *et al.*, 2014). Similarly, in a survey of *Bombus* spp. in China, bees could be grouped into 2 microbial enterotypes (Li *et al.*, 2015). One enterotype was dominated by core symbionts *Snodgrassella* and *Gilliamella*, while the other enterotype comprised non-core *Hafnia*, Enterobacteriaceae and *Serratia*. The latter 2 are thought to be opportunistic pathogens (Kwong *et al.*, 2017a; Raymann and Moran, 2018). Finally, higher susceptibility to pathogens was found in germ-free honey and bumble bees (Koch and Schmid-Hempel, 2011; Kwong *et al.*, 2017a; Mockler *et al.*, 2018; Raymann and Moran, 2018).

Acidification of the bee gut lumen by symbionts (Zheng *et al.*, 2017) could explain how a strong core microbiota resists invasion by pathogens. This hypothesis is consistent with the dominant role of pH in determination of bacterial communities in soil (Morton *et al.*, 2017; Ratzke and Gore, 2018), and correlations between low gastric acidity and opportunistic infection in human subjects (Stark and Nylund, 2016). However, other explanations for the effects of core gut microbiota on parasites – such as physical competition for space and resources, enhancement of the immune response and improvement of host nutritional status (Raymann and Moran, 2018) – also deserve consideration. It is possible that different microbial assemblages are optimal for different functions. For example, a microbiome that produces abundant acids might be optimal for direct defence against parasites, whereas other community compositions might provide more benefits to immune system regulation, or to nutritional sufficiency that improves resistance or tolerance to infection (Scrimshaw *et al.*, 1959; Brown *et al.*, 2003). Further research is needed to investigate variation in gut pH within and across species, and possible causative relationships between microbiome composition and bee health under different environmental circumstances.

Conclusion

Our results build on prior associations between microbiome and infection intensity to provide mechanistic insights into how the bee gut microbiota are likely to influence trypanosomatid infection in bumble bees and possibly other corbiculate bee species. Our findings of pH-dependent parasite inhibition *in vitro* suggest that gut pH could be a critical determinant of trypanosomatid growth *in vivo*, a hypothesis that could be tested in both manipulative and observational studies that relate gut pH to trypanosomatid establishment. The role of gut pH in resistance to other enteric pathogens, and the selective forces acting on symbionts and parasites to create and tolerate different levels of acidity, warrants further study as part of continued investigations into the functional significance of the bee microbiome.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182018001555>

Acknowledgements. The authors thank Ben Sadd for providing strains IL13.2 and S08.1, Rebecca Irwin for providing the bees from which strain VT1 was established, Guang Xu and Ben Sadd for sharing DNA sequences and two anonymous reviewers for comments that improved the paper.

Financial support. This project was funded by a National Science Foundation Postdoctoral Research Fellowship to EPY (NSF-DBI-1708945); USDA NIFA Hatch funds (CA-R-ENT-5109-H), NIH (5R01GM122060-02) and NSF MSB-ECA (1638728) to QSM and an NSF-CAREER grant (IOS 1651888) to TRR. The funders had no role in study design, data collection and interpretation or the decision to submit the work for publication.

Conflicts of interest. The authors declare that they have no conflicts of interest.

Ethical standards. Not applicable.

Data availability. All data are supplied in Supplementary Information, Data S1.

References

- Adams MR and Hall CJ (1988) Growth inhibition of food-borne pathogens by lactic and acetic acids and their mixtures. *International Journal of Food Science & Technology* **23**, 287–292.
- Anderson KE, Carroll MJ, Sheehan T, Mott BM, Maes P and Corby-Harris V (2014) Hive-stored pollen of honey bees: many lines of evidence are consistent with pollen preservation, not nutrient conversion. *Molecular Ecology* **23**, 5904–5917.
- Barribeau SM, Sadd BM, du Plessis L and Schmid-Hempel P (2014) Gene expression differences underlying genotype-by-genotype specificity in a host–parasite system. *Proceedings of the National Academy of Sciences* **111**, 3496–3501.
- Benthin S and Villadsen J (1995) Different inhibition of *Lactobacillus delbrueckii* subsp. *bulgaricus* by D- and L-lactic acid: effects on lag phase, growth rate and cell yield. *Journal of Applied Bacteriology* **78**, 647–654.
- Billiet A, Meeus I, Van Nieuwerburgh F, Deforce D, Wäckers F and Smaghe G (2017) Colony contact contributes to the diversity of gut bacteria in bumblebees (*Bombus terrestris*). *Insect Science* **24**, 270–277.
- Bringaud F, Rivière L and Coustou V (2006) Energy metabolism of trypanosomatids: adaptation to available carbon sources. *Molecular and Biochemical Parasitology* **149**, 1–9.
- Brown MJF, Schmid-Hempel R and Schmid-Hempel P (2003) Strong context-dependent virulence in a host–parasite system: reconciling genetic evidence with theory. *Journal of Animal Ecology* **72**, 994–1002.
- Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF and Griswold TL (2011) Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences* **108**, 662–667.
- Cariveau DP, Elijah Powell J, Koch H, Winfree R and Moran NA (2014) Variation in gut microbial communities and its association with pathogen infection in wild bumble bees (*Bombus*). *The ISME Journal* **8**, 2369–2379.
- Cornman RS, Tarpy DR, Chen Y, Jeffreys L, Lopez D, Pettis JS, vanEngelsdorp D and Evans JD (2012) Pathogen webs in collapsing honey bee colonies. *PLoS ONE* **7**, e43562.
- Cox AJ, Pyne DB, Saunders PU and Fricker PA (2010) Oral administration of the probiotic *Lactobacillus fermentum* VRI-003 and mucosal immunity in endurance athletes. *British Journal of Sports Medicine* **44**, 222–226.
- Daskin JH, Bell SC, Schwarzkopf L and Alford RA (2014) Cool temperatures reduce antifungal activity of symbiotic bacteria of threatened amphibians – implications for disease management and patterns of decline. *PLoS ONE* **9**, e100378.
- De Keersmaecker SCJ, Verhoeven TLA, Desair J, Marchal K, Vanderleyden J and Nagy I (2006) Strong antimicrobial activity of *Lactobacillus rhamnosus* GG against *Salmonella typhimurium* is due to accumulation of lactic acid. *FEMS Microbiology Letters* **259**, 89–96.
- Dillon RJ and Dillon VM (2004) The gut bacteria of insects: nonpathogenic interactions. *Annual Review of Entomology* **49**, 71–92.
- Engel P, Martinson VG and Moran NA (2012) Functional diversity within the simple gut microbiota of the honey bee. *Proceedings of the National Academy of Sciences* **109**, 11002–11007.
- Engel P, Kwong WK, McFrederick Q, Anderson KE, Barribeau SM, Chandler JA, Cornman RS, Dainat J, Miranda JR, de, Doublet V, Emery O, Evans JD, Farinelli L, Flenniken ML, Granberg F, Grasis JA, Gauthier L, Hayer J, Koch H, Kocher S, Martinson VG, Moran N, Munoz-Torres M, Newton I, Paxton RJ, Powell E, Sadd BM, Schmid-Hempel P, Schmid-Hempel R, Song SJ, Schwarz RS, vanEngelsdorp D and Dainat B (2016) The bee microbiome: impact on bee health and model for evolution and ecology of host–microbe interactions. *mBio* **7**, e02164–15.
- Evans JD and Armstrong T-N (2006) Antagonistic interactions between honey bee bacterial symbionts and implications for disease. *BMC Ecology* **6**, 4.
- Evans JD and Lopez DL (2004) Bacterial probiotics induce an immune response in the honey bee (Hymenoptera: Apidae). *Journal of Economic Entomology* **97**, 752–756.
- Evans JD and Spivak M (2010) Socialized medicine: individual and communal disease barriers in honey bees. *Journal of Invertebrate Pathology* **103** (Suppl.), S62–S72.

- Ewaschuk JB, Zello GA, Naylor JM and Brocks DR (2002) Metabolic acidosis: separation methods and biological relevance of organic acids and lactic acid enantiomers. *Journal of Chromatography B* **781**, 39–56.
- Fausser-Misslin A, Sadd BM, Neumann P and Sandrock C (2014) Influence of combined pesticide and parasite exposure on bumblebee colony traits in the laboratory. *Journal of Applied Ecology* **51**, 450–459.
- Fayol-Messaoudi D, Berger CN, Coconnier-Polter M-H, Moal VL-L and Servin AL (2005) pH-, lactic acid-, and non-lactic acid-dependent activities of probiotic Lactobacilli against *Salmonella enterica* serovar Typhimurium. *Applied and Environmental Microbiology* **71**, 6008–6013.
- Forsgren E, Olofsson TC, Vázquez A and Fries I (2010) Novel lactic acid bacteria inhibiting *Paenibacillus* larvae. *Apidologie* **41**, 99–108.
- Fürst MA, McMahon DP, Osborne JL, Paxton RJ and Brown MJF (2014) Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* **506**, 364–366.
- Gaggia F, Baffoni L and Alberoni D (2018) Probiotics for honeybees' health. In Di Gioia D and Biavati B (eds), *Probiotics and Prebiotics in Animal Health and Food Safety*. Cham: Springer, pp. 219–245. doi: 10.1007/978-3-319-71950-4_9.
- Gallai N, Salles J-M, Settele J and Vaissière BE (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* **68**, 810–821.
- Gegeer RJ, Otterstatter MC and Thomson JD (2005) Does parasitic infection impair the ability of bumblebees to learn flower-handling techniques? *Animal Behaviour* **70**, 209–215.
- Gegeer RJ, Otterstatter MC and Thomson JD (2006) Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proceedings of Biological Sciences/The Royal Society* **273**, 1073–1078.
- Glass KA, Loeffelholz JM, Ford JP and Doyle MP (1992) Fate of *Escherichia coli* O157:H7 as affected by pH or sodium chloride and in fermented, dry sausage. *Applied and Environmental Microbiology* **58**, 2513–2516.
- Gorbach SL (1990) Lactic acid bacteria and human health. *Annals of Medicine* **22**, 37–41.
- Gorbunov PS (1996) Peculiarities of life cycle in flagellate *Crithidia bombi* (protozoa, trypanosomatidae). *Zoologicheskii Zhurnal* **75**, 808–810.
- Goulson D, Nicholls E, Botías C and Rotheray EL (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* **347**, 1255957.
- Graystock P, Blane EJ, McFrederick QS, Goulson D and Hughes WOH (2016) Do managed bees drive parasite spread and emergence in wild bees? *International Journal for Parasitology: Parasites and Wildlife* **5**, 64–75.
- Imhoof B and Schmid-Hempel P (1998) Single-clone and mixed-clone infections versus host environment in *Crithidia bombi* infecting bumblebees. *Parasitology* **117**, 331–336.
- Kamada N, Chen GY, Inohara N and Núñez G (2013) Control of pathogens and pathobionts by the gut microbiota. *Nature Immunology* **14**, 685–690.
- Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U and Engel P (2017) Disentangling metabolic functions of bacteria in the honey bee gut. *PLoS Biology* **15**, e2003467.
- Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham S, Kremen C and Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303–313.
- Koch H and Schmid-Hempel P (2011) Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 19288–19292.
- Koch H and Schmid-Hempel P (2012) Gut microbiota instead of host genotype drive the specificity in the interaction of a natural host–parasite system. *Ecology Letters* **15**, 1095–1103.
- Koch H, Cisarovsky G and Schmid-Hempel P (2012) Ecological effects on gut bacterial communities in wild bumblebee colonies. *The Journal of Animal Ecology* **81**, 1202–1210.
- Kwong WK and Moran NA (2013) Cultivation and characterization of the gut symbionts of honey bees and bumble bees: description of *Snodgrassella alvi* gen. nov., sp. nov., a member of the family Neisseriaceae of the Betaproteobacteria, and *Gilliamella apicola* gen. nov., sp. nov., a member of Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order 'Enterobacteriales' of the Gammaproteobacteria. *International Journal of Systematic and Evolutionary Microbiology* **63**, 2008–2018.
- Kwong WK and Moran NA (2016) Gut microbial communities of social bees. *Nature Reviews Microbiology* **14**, 374–384.
- Kwong WK, Mancenido AL and Moran NA (2017a) Immune system stimulation by the native gut microbiota of honey bees. *Open Science* **4**, 170003.
- Kwong WK, Medina LA, Koch H, Sing K-W, Soh EJY, Ascher JS, Jaffé R and Moran NA (2017b) Dynamic microbiome evolution in social bees. *Science Advances* **3**, e1600513.
- Li J, Powell JE, Guo J, Evans JD, Wu J, Williams P, Lin Q, Moran NA and Zhang Z (2015) Two gut community enterotypes recur in diverse bumblebee species. *Current Biology* **25**, R652–R653.
- Lindgren SE and Dobrogosz WJ (1990) Antagonistic activities of lactic acid bacteria in food and feed fermentations. *FEMS Microbiology Letters* **87**, 149–164.
- Martinson VG, Danforth BN, Mincley RL, Rueppell O, Tingek S and Moran NA (2011) A simple and distinctive microbiota associated with honey bees and bumble bees. *Molecular Ecology* **20**, 619–628.
- McMahon DP, Fürst MA, Caspar J, Theodorou P, Brown MJF and Paxton RJ (2015) A sting in the spit: widespread cross-infection of multiple RNA viruses across wild and managed bees. *Journal of Animal Ecology* **84**, 615–624.
- Mockler BK, Kwong WK, Moran NA and Koch H (2018) Microbiome structure influences infection by the parasite *Crithidia bombi* in bumble bees. *Applied and Environmental Microbiology* **84**, e02335–17.
- Morton JT, Sanders J, Quinn RA, McDonald D, Gonzalez A, Vázquez-Baeza Y, Navas-Molina JA, Song SJ, Metcalf JL, Hyde ER, Lladser M, Dorrestein PC and Knight R (2017) Balance trees reveal microbial niche differentiation. *mSystems* **2**, e00162–16.
- Mueller UG and Sachs JL (2015) Engineering microbiomes to improve plant and animal health. *Trends in Microbiology* **23**, 606–617.
- Palmer-Young EC, Sadd BM, Stevenson PC, Irwin RE and Adler LS (2016) Bumble bee parasite strains vary in resistance to phytochemicals. *Scientific Reports* **6**, 37087.
- Palmer-Young EC, Sadd BM, Irwin RE and Adler LS (2017a) Synergistic effects of floral phytochemicals against a bumble bee parasite. *Ecology and Evolution* **7**, 1836–1849.
- Palmer-Young EC, Hogeboom A, Kaye AJ, Donnelly D, Andicoechea J, Cannon SJ, Weston I, Skyrn K, Irwin RE and Adler LS (2017b) Context-dependent medicinal effects of anabasine and infection-dependent toxicity in bumble bees. *PLoS ONE* **12**, e0183729.
- Praet J, Meeus I, Cnockaert M, Houf K, Smaghe G and Vandamme P (2015) Novel lactic acid bacteria isolated from the bumble bee gut: *Convivina intestini* gen. nov., sp. nov., *Lactobacillus bombicola* sp. nov., and *Weissella bombi* sp. nov. *Antonie van Leeuwenhoek* **107**, 1337–1349.
- Praet J, Parmentier A, Schmid-Hempel R, Meeus I, Smaghe G and Vandamme P (2018) Large-scale cultivation of the bumblebee gut microbiota reveals an underestimated bacterial species diversity capable of pathogen inhibition. *Environmental Microbiology* **20**, 214–227.
- Presser KA, Ratkowsky DA and Ross T (1997) Modelling the growth rate of *Escherichia coli* as a function of pH and lactic acid concentration. *Applied and Environmental Microbiology* **63**, 2355–2360.
- R Core Team (2014) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ratzke C and Gore J (2018) Modifying and reacting to the environmental pH can drive bacterial interactions. *PLoS Biology* **16**, e2004248.
- Ravoet J, Maharramov J, Meeus I, De Smet L, Wenseleers T, Smaghe G and de Graaf DC (2013) Comprehensive bee pathogen screening in Belgium reveals *Crithidia mellificae* as a new contributory factor to winter mortality. *PLoS ONE* **8**, e72443.
- Raymann K and Moran NA (2018) The role of the gut microbiome in health and disease of adult honey bee workers. *Current Opinion in Insect Science* **26**, 97–104.
- Ritz C, Baty F, Streibig JC and Gerhard D (2015) Dose-response analysis using R. *PLoS ONE* **10**, e0146021.
- Round JL and Mazmanian SK (2009) The gut microbiota shapes intestinal immune responses during health and disease. *Nature Reviews Immunology* **9**, 313–323.
- Runckel C, DeRisi J and Flenniken ML (2014) A draft genome of the honey bee trypanosomatid parasite *Crithidia mellificae*. *PLOS ONE* **9**, e95057.
- Russell JB and Diez-Gonzalez F (1998) The effects of fermentation acids on bacterial growth. *Advances in Microbial Physiology* **39**, 205–234.
- Sabaté DC, Carrillo L and Carina Audisio M (2009) Inhibition of *Paenibacillus* larvae and *Ascosphaera apis* by *Bacillus subtilis* isolated from honeybee gut and honey samples. *Research in Microbiology* **160**, 193–199.

- Sadd BM** (2011) Food-environment mediates the outcome of specific interactions between a bumblebee and its trypanosome parasite. *Evolution* **65**, 2995–3001.
- Salathé R, Tognazzo M, Schmid-Hempel R and Schmid-Hempel P** (2012) Probing mixed-genotype infections I: extraction and cloning of infections from hosts of the trypanosomatid *Crithidia bombi*. *PLoS ONE* **7**, e49046.
- Salminen S and von Wright A** (2004) *Lactic Acid Bacteria: Microbiological and Functional Aspects*, 3rd Edn. Boca Raton, FL: CRC Press.
- Schmid-Hempel P and Ebert D** (2003) On the evolutionary ecology of specific immune defence. *Trends in Ecology & Evolution* **18**, 27–32.
- Schmid-Hempel R, Eckhardt M, Goulson D, Heinzmann D, Lange C, Plischuk S, Escudero LR, Salathé R, Scriven JJ and Schmid-Hempel P** (2014) The invasion of Southern South America by imported bumblebees and associated parasites. *Journal of Animal Ecology* **83**, 823–837.
- Schwarz RS, Bauchan GR, Murphy CA, Ravoet J, de Graaf DC and Evans JD** (2015) Characterization of two species of trypanosomatidae from the honey bee *Apis mellifera*: *Crithidia mellificae* Langridge and McGhee, and *Lotmaria passim* n. gen., n. sp. *Journal of Eukaryotic Microbiology* **62**, 567–583.
- Schwarz RS, Moran NA and Evans JD** (2016) Early gut colonizers shape parasite susceptibility and microbiota composition in honey bee workers. *Proceedings of the National Academy of Sciences* **113**, 9345–9350.
- Scrimshaw NS, Taylor CE and Gordon JE** (1959) Interactions of nutrition and infection. *American Journal of Medical Sciences* **237**, 367–403.
- Shykoff JA and Schmid-Hempel P** (1991) Incidence and effects of four parasites in natural populations of bumble bees in Switzerland. *Apidologie* **22**, 117–125.
- Stark CM and Nylund CM** (2016) Side effects and complications of proton pump inhibitors: a pediatric perspective. *The Journal of Pediatrics* **168**, 16–22.
- Steele MI, Kwong WK, Whiteley M and Moran NA** (2017) Diversification of type VI secretion system toxins reveals ancient antagonism among bee gut microbes. *mBio* **8**, e01630–17.
- Tripodi AD, Szalanski AL and Strange JP** (2018) Novel multiplex PCR reveals multiple trypanosomatid species infecting North American bumble bees (Hymenoptera: Apidae: Bombus). *Journal of Invertebrate Pathology* **153**, 147–155.
- Vásquez A, Forsgren E, Fries I, Paxton RJ, Flaberg E, Szekely L and Olofsson TC** (2012) Symbionts as Major modulators of insect health: lactic acid Bacteria and honeybees. *PLoS ONE* **7**, e33188.
- Zheng H, Powell JE, Steele MI, Dietrich C and Moran NA** (2017) Honeybee gut microbiota promotes host weight gain via bacterial metabolism and hormonal signaling. *Proceedings of the National Academy of Sciences* **114**, 4775–4780.