

Adhesion and volume filling in one-dimensional population dynamics under Dirichlet boundary condition

Hyung Jun Choi 

School of Liberal Arts, Korea University of Technology and Education,
Cheonan 31253, Republic of Korea (email: hjchoi@koreatech.ac.kr)

Seonghak Kim 

Department of Mathematics, College of Natural Sciences, Kyungpook
National University, Daegu 41566, Republic of Korea
(email: shkim17@knu.ac.kr)

Youngwoo Koh 

Department of Mathematics Education, Kongju National University,
Kongju 32588, Republic of Korea (email: ywkoh@kongju.ac.kr)

(Received 21 May 2023; accepted 12 December 2023)

We generalize the one-dimensional population model of Anguige & Schmeiser [1] reflecting the cell-to-cell adhesion and volume filling and classify the resulting equation into the six types. Among these types, we fix one that yields a class of advection-diffusion equations of forward-backward-forward type and prove the existence of infinitely many global-in-time weak solutions to the initial-Dirichlet boundary value problem when the maximum value of an initial population density exceeds a certain threshold. Such solutions are extracted from the method of convex integration by Müller & Šverák [12]; they exhibit fine-scale density mixtures over a finite time interval, then become smooth and identical, and decay exponentially and uniformly to zero as time approaches infinity. TE check: Please check the reference citation in abstract.

Keywords: Population model; forward-backward-forward type; adhesion and volume filling; partial differential inclusion; convex integration

2020 *Mathematics Subject Classification:* Primary: 35M13
Secondary: 35K59, 35D30, 92D25

1. Introduction

The evolution process for spatial distribution of cells or animals in one-dimensional homogeneous habitat can be modelled by quasilinear advection-diffusion equations of the form,

$$u_t = (\rho(u))_{xx}, \quad (1.1)$$

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Royal Society of Edinburgh

where $u = u(x, t)$ denotes the population density of a single species at position x and time t , and the derivative $\sigma(s) = \rho'(s)$ of a given flux function $\rho(s)$ represents the diffusivity of equation (1.1).

In zoological studies, as an improvement to the Δ -model by Taylor & Taylor [15] for adhesive movement, Turchin [16] proposed equation (1.1), based on a random walk approach [13], as a model of individual movement, which is not only reflecting adhesion or repulsion between conspecific organisms but also avoiding some defects in their model. In his model, the flux function ρ is given by

$$\rho(s) = \frac{2k_0}{3\omega} s^3 - k_0 s^2 + \frac{\mu}{2} s, \tag{1.2}$$

where $k_0 > 0$ is the maximum degree of gregariousness, $\omega > 0$ is the critical density at which movement switches from adhesive to repulsive, and $\mu \in (0, 1]$ is the motility rate. With this flux function ρ , he considered the initial-Dirichlet boundary value problem

$$\begin{cases} u_t = (\rho(u))_{xx} & \text{in } \Omega \times (0, \infty), \\ u = u_0 & \text{on } \Omega \times \{t = 0\}, \\ u(0, t) = u(L, t) = \delta_0 & \text{for } t > 0, \end{cases} \tag{1.3}$$

where $\Omega = (0, L) \subset \mathbb{R}$ is a favourable habitat of size $L > 0$, $u_0 = u_0(x)$ is the initial density of a single species, and $\delta_0 \geq 0$ is a constant for the Dirichlet (or absorbing) boundary condition. Here, the constant δ_0 reflects the intensity of hostility of the surrounding habitat and/or the migration rate towards there. For example, when $\delta_0 = 0$, animals touching the border $\partial\Omega = \{0, L\}$ are permanently lost to the population, either because they move away from the habitat Ω or because they are killed by predators residing in the very hostile surrounding area. In this case, it is expected that the total population eventually vanishes as time approaches infinity.

In mathematical biology, Anguige & Schmeiser [1] obtained equation (1.1), based also on the random walk approach, as a model of cell motility which incorporates the effects of cell-to-cell adhesion and volume filling. In their model, the flux function ρ is given by

$$\rho(s) = \alpha s^3 - 2\alpha s^2 + s, \tag{1.4}$$

where $\alpha \in [0, 1]$ is the adhesion constant. With this flux function ρ , they studied the initial-Neumann boundary value problem

$$\begin{cases} u_t = (\rho(u))_{xx} & \text{in } \Omega \times (0, \infty), \\ u = u_0 & \text{on } \Omega \times \{t = 0\}, \\ u_x(0, t) = u_x(L, t) = 0 & \text{for } t > 0, \end{cases} \tag{1.5}$$

where $\Omega = (0, 1) \subset \mathbb{R}$ is a habitat of unit size. Here, the Neumann (or reflecting) boundary condition should imply that the total population remains the same for all time.

In both models of [1, 16], when the species in question are highly adhesive (that is, $k_0\omega > \mu$ in (1.2) and $\alpha > \frac{3}{4}$ in (1.4), resp.), the diffusivity $\sigma(s) = \rho'(s)$ admits a nonempty open interval in which it is negative so that the equation $u_t = (\rho(u))_{xx}$ becomes *backward parabolic* for the population density values $s = u$ lying in that

interval. So when the range of the initial population density u_0 overlaps with the interval of backward regime, problems (1.3) and (1.5) may be ill-posed. This awkward situation was regarded as a clue for the authors of [1, 16] to expect a certain pattern formation in the population density u as time goes by. However, there have not been any existence results on the highly adhesive random walk models since the usual methods of parabolic theory are no longer applicable to this case.

Regardless of the expectation stated above, there has been scepticism on the random walk models of adhesive population dynamics due to their limited practicality and inability of displaying complicated behaviour such as sorting [5, 8]. Meanwhile, many nonlocal adhesion models have been suggested and studied actively to reflect various phenomena in population dynamics. For an overview, one may refer to the review article [5].

Nonetheless, in this paper, we generalize the adhesion population model of Anguige & Schmeiser [1] and classify the resulting equation into the six types. Among such types, we fix one that yields a class of advection-diffusion equations of forward-backward-forward type and prove the existence of infinitely many global-in-time weak solutions to the initial-Dirichlet boundary value problem when the maximum value of the initial population density u_0 exceeds a certain threshold. Although our solutions may not exhibit expected behaviour of adhesion such as clustering and sorting, they capture some interesting phenomena: fine-scale density mixtures between the high and low density regimes, smoothing after a finite time, and smooth extinction of the species at an exponential rate due to the absorbing boundary condition. On the other hand, in the same initial value problem under the Neumann boundary condition, we expect to obtain global weak solutions showing quite different behaviours from the Dirichlet case. This will be explored in the subsequent paper as a sequel.

Plan of the paper: In the rest of this section, we derive a continuum model of adhesion, which is a generalization of [1], and then justify the definition of a global weak solution to the corresponding initial-Dirichlet boundary value problem. In § 2, depending on the choice of a pair of adhesion and volume filling constants, we classify the resulting equation from continuum modelling into the six types. Then the main result of the paper, theorem 3.2, is sophisticated in § 3 along with a strategy to obtain solutions. As an independent section, in § 4, we formulate a differential inclusion problem and present its special solvability result as theorem 4.2, which serves as an essential ingredient for a proof of theorem 3.2 in § 5. Sections 6 and 7 carry out a long proof of theorem 4.2, which can be regarded as the core analysis of the paper.

Notations: Hereafter, let $L > 0$ denote the size of a favourable habitat $\Omega = (0, L) \subset \mathbb{R}$. For $\tau \in (0, \infty]$, we write

$$\Omega_\tau := \Omega \times (0, \tau) \subset \mathbb{R}^2.$$

Let k , m , and n be positive integers, let $0 < a < 1$, and let $U \subset \mathbb{R}^n$ be an open set.

- (i) We denote by $C^k(U)$ the space of functions $u : U \rightarrow \mathbb{R}$ whose partial derivatives of order up to k exist and are continuous in U .

- (ii) Let $C^k(\bar{U})$ be the space of functions $u \in C^k(U)$ whose partial derivatives of order up to k are uniformly continuous in every bounded subset of U .
- (iii) Let $n = 2$; so $U \subset \mathbb{R}^2 = \mathbb{R}_x \times \mathbb{R}_t$. We define $C^{2,1}(U)$ to be the space of functions $u : U \rightarrow \mathbb{R}$ such that u_x, u_{xx} , and u_t exist and are continuous in U . Also, let $C^{2,1}(\bar{U})$ denote the space of functions $u \in C^{2,1}(U)$ such that u_x, u_{xx} , and u_t are uniformly continuous in every bounded subset of U .
- (iv) We denote by $C^{2+a}(\bar{\Omega})$ the space of functions $u \in C^2(\bar{\Omega})$ with

$$\sup_{x,y \in \Omega, x \neq y} \frac{|u_{xx}(x) - u_{xx}(y)|}{|x - y|^a} < \infty.$$

- (v) Let $0 < T < \infty$. We define $C^{2+a,1+\frac{a}{2}}(\bar{\Omega}_T)$ to be the space of functions $u \in C^{2,1}(\bar{\Omega}_T)$ whose quantities

$$\begin{aligned} & \sup_{x,y \in \Omega, x \neq y, 0 < t < T} \frac{|u_{xx}(x,t) - u_{xx}(y,t)|}{|x - y|^a}, \\ & \sup_{x \in \Omega, 0 < s, t < T, s \neq t} \frac{|u_{xx}(x,s) - u_{xx}(x,t)|}{|s - t|^{\frac{a}{2}}}, \\ & \sup_{x,y \in \Omega, x \neq y, 0 < t < T} \frac{|u_t(x,t) - u_t(y,t)|}{|x - y|^a}, \quad \sup_{x \in \Omega, 0 < s, t < T, s \neq t} \frac{|u_t(x,s) - u_t(x,t)|}{|s - t|^{\frac{a}{2}}} \end{aligned}$$

are all finite.

- (vi) We denote by $\mathbb{M}^{m \times n}$ the space of $m \times n$ real matrices.
- (vii) For a Lebesgue measurable set $E \subset \mathbb{R}^n$, its n -dimensional measure is denoted by $|E| = |E|_n$ with subscript n omitted if it is clear from the context.
- (viii) We use $W^{k,p}(U)$ to denote the space of functions $u \in L^p(U)$ whose weak partial derivatives of order up to k exist in U and belong to $L^p(U)$.
- (ix) A sequence $\{U_\ell\}_{\ell \in \mathbb{N}}$ of disjoint open subsets of U is called a *Vitali cover* of U if $|U \setminus \cup_{\ell \in \mathbb{N}} U_\ell| = 0$; in this case, $\{U_\ell\}_{\ell \in \mathbb{N}}$ is said to cover U in the sense of Vitali.

1.1. Continuum model

For any fixed $n \in \mathbb{N}$ with $n \geq 2$, let us consider a discrete distribution of homogeneous cells residing on the endpoints of the 2^n uniform subintervals

$$[0, h_n], [h_n, 2h_n], [2h_n, 3h_n], \dots, [(2^n - 1)h_n, L]$$

of the spatial domain $\bar{\Omega} = [0, L]$, where $h_n := L/2^n$. Assume that $N_{cap} \in \mathbb{N}$ is the maximum number of cells that can stay at each position $x_{ni} := ih_n$ ($i = 0, 1, \dots, 2^n$). For $i = 0, 1, \dots, 2^n$, let $\tilde{u}_{ni}(t) \in \{0, 1, \dots, N_{cap}\}$ denote the number

of cells at position x_{ni} and time $t \geq 0$, and write

$$u_{ni}(t) := \frac{\tilde{u}_{ni}(t)}{N_{cap}} \in [0, 1],$$

called the *discrete cell density* at position x_{ni} and time $t \geq 0$. We assume that the dynamics of such cells is governed by the system of ordinary differential equations,

$$\frac{du_{ni}}{dt} = \mathcal{T}_{n(i-1)}^+ u_{n(i-1)} + \mathcal{T}_{n(i+1)}^- u_{n(i+1)} - (\mathcal{T}_{ni}^+ + \mathcal{T}_{ni}^-) u_{ni}, \tag{1.6}$$

where $i = 2, 3, \dots, 2^n - 2$. Here, for $i = 1, 2, \dots, 2^n - 1$, we denote by \mathcal{T}_{ni}^\pm the *transitional probabilities* per unit time of a one-step jump from ih_n to $(i \pm 1)h_n$ that are given by

$$\begin{aligned} \mathcal{T}_{ni}^+ &= \frac{(1 - \beta u_{n(i+1)})(1 - \alpha u_{n(i-1)})}{h_n^2}, \\ \mathcal{T}_{ni}^- &= \frac{(1 - \alpha u_{n(i+1)})(1 - \beta u_{n(i-1)})}{h_n^2}, \end{aligned} \tag{1.7}$$

where $\alpha, \beta \in [0, 1]$ are the *adhesion* and *volume filling* constants, respectively. Note here that only the maximal volume filling constant $\beta = 1$ is considered by Anguige & Schmeiser [1]. We also write $I_n := \{ih_n \mid i = 0, 1, \dots, 2^n\}$, the n th discrete habitat; then $I_2 \subset I_3 \subset I_4 \subset \dots$.

Inserting (1.7) into (1.6), we obtain that for $i = 2, 3, \dots, 2^n - 2$,

$$\begin{aligned} \frac{du_{ni}}{dt} &= \frac{1}{h_n} \left(\frac{R_{n(i+1)} - R_{ni}}{h_n} - \frac{R_{ni} - R_{n(i-1)}}{h_n} + \frac{Q_{n(i+1)} - Q_{ni}}{h_n} \right. \\ &\quad \left. - \frac{Q_{ni} - Q_{n(i-1)}}{h_n} + \frac{Q_{ni} - Q_{n(i-1)}}{h_n} - \frac{Q_{n(i-1)} - Q_{n(i-2)}}{h_n} \right), \end{aligned} \tag{1.8}$$

where

$$R_{nj} := u_{nj}(1 - \alpha\beta + \alpha\beta(1 - u_{n(j-1)})(1 - u_{n(j+1)}))$$

for $j = 1, 2, \dots, 2^n - 1$, and

$$Q_{nk} := \alpha(\beta - 1)u_{nk}u_{n(k+1)}$$

for $k = 0, 1, \dots, 2^n - 1$.

To derive the continuum equation, assume that there exists a function $u = u(x, t) \in C^{2,1}(\Omega_\infty; [0, 1]) \cap C(\bar{\Omega}_\infty; [0, 1])$ such that

$$u(ih_n, t) = u_{ni}(t)$$

for all $n \in \mathbb{N}$, $i = 0, 1, \dots, 2^n$, and $t \geq 0$.

Now, fix any $n \in \mathbb{N}$ with $n \geq 2$. Then for $j = 1, 2, \dots, 2^n - 1$, $k = 0, 1, \dots, 2^n - 1$, and $t \geq 0$, we have

$$R_{nj}(t) = u(jh_n, t)(1 - \alpha\beta + \alpha\beta(1 - u((j - 1)h_n, t))(1 - u((j + 1)h_n, t))),$$

$$Q_{nk}(t) = \alpha(\beta - 1)u(kh_n, t)u((k + 1)h_n, t).$$

In this regard, for $t \geq 0$, define

$$R_n(x, t) = u(x, t)(1 - \alpha\beta + \alpha\beta(1 - u(x - h_n, t))(1 - u(x + h_n, t)))$$

for $h_n \leq x \leq L - h_n$ and

$$Q_n(x, t) = \alpha(\beta - 1)u(x, t)u(x + h_n, t)$$

for $0 \leq x \leq L - h_n$; then

$$R_n \in C^{2,1}((h_n, L - h_n) \times (0, \infty)) \cap C([h_n, L - h_n] \times [0, \infty)),$$

$$Q_n \in C^{2,1}((0, L - h_n) \times (0, \infty)) \cap C([0, L - h_n] \times [0, \infty)),$$

$$(R_n)_{xx}(x, t) = u_{xx}(x, t)(1 - \alpha\beta + \alpha\beta(1 - u(x - h_n, t))(1 - u(x + h_n, t)))$$

$$- 2\alpha\beta u_x(x, t)u_x(x - h_n, t)(1 - u(x + h_n, t))$$

$$- 2\alpha\beta u_x(x, t)u_x(x + h_n, t)(1 - u(x - h_n, t))$$

$$- \alpha\beta u(x, t)u_{xx}(x - h_n, t)(1 - u(x + h_n, t))$$

$$+ 2\alpha\beta u(x, t)u_x(x - h_n, t)u_x(x + h_n, t)$$

$$- \alpha\beta u(x, t)u_{xx}(x + h_n, t)(1 - u(x - h_n, t)),$$

$$(Q_n)_{xx}(x, t) = \alpha(\beta - 1)u_{xx}(x, t)u(x + h_n, t) + 2\alpha(\beta - 1)u_x(x, t)u_x(x + h_n, t)$$

$$+ \alpha(\beta - 1)u(x, t)u_{xx}(x + h_n, t).$$

Next, fix any $x_0 \in (\cup_{n \geq 2} I_n) \setminus \{0, L\} \subset \Omega$ and $t_0 > 0$. Let n_0 denote the smallest positive integer with $x_0 \in I_{n_0}$; then

$$x_0 = i_0 h_{n_0}$$

for some odd integer $i_0 \in \{1, 3, \dots, 2^{n_0} - 1\}$. Choose any integer $n > n_0$. Then

$$x_0 = i_{0n} h_n,$$

where $i_{0n} := 2^{n-n_0} i_0 \in \{2, 4, \dots, 2^n - 2\}$. From (1.8) and the mean value theorem, we have

$$u_t(x_0, t_0) = u_t(i_{0n} h_n, t_0) = \frac{du_{ni_{0n}}(t_0)}{dt}$$

$$= (R_n)_{xx}(x_{1n}, t_0) + (Q_n)_{xx}(x_{2n}, t_0) + (Q_n)_{xx}(x_{3n}, t_0)$$

for some $x_{1n}, x_{2n} \in ((i_{0n} - 1)h_n, (i_{0n} + 1)h_n) = (x_0 - h_n, x_0 + h_n)$ and $x_{3n} \in ((i_{0n} - 2)h_n, i_{0n}h_n) = (x_0 - 2h_n, x_0)$. Thus,

$$\begin{aligned}
 u_t(x_0, t_0) &= u_{xx}(x_{1n}, t_0)(1 - \alpha\beta + \alpha\beta(1 - u(x_{1n} - h_n, t_0))(1 - u(x_{1n} + h_n, t_0))) \\
 &\quad - 2\alpha\beta u_x(x_{1n}, t_0)u_x(x_{1n} - h_n, t_0)(1 - u(x_{1n} + h_n, t_0)) \\
 &\quad - 2\alpha\beta u_x(x_{1n}, t_0)u_x(x_{1n} + h_n, t_0)(1 - u(x_{1n} - h_n, t_0)) \\
 &\quad - \alpha\beta u(x_{1n}, t_0)u_{xx}(x_{1n} - h_n, t_0)(1 - u(x_{1n} + h_n, t_0)) \\
 &\quad + 2\alpha\beta u(x_{1n}, t_0)u_x(x_{1n} - h_n, t_0)u_x(x_{1n} + h_n, t_0) \\
 &\quad - \alpha\beta u(x_{1n}, t_0)u_{xx}(x_{1n} + h_n, t_0)(1 - u(x_{1n} - h_n, t_0)) \\
 &\quad + \alpha(\beta - 1)u_{xx}(x_{2n}, t_0)u(x_{2n} + h_n, t_0) \\
 &\quad + 2\alpha(\beta - 1)u_x(x_{2n}, t_0)u_x(x_{2n} + h_n, t_0) \\
 &\quad + \alpha(\beta - 1)u(x_{2n}, t_0)u_{xx}(x_{2n} + h_n, t_0) \\
 &\quad + \alpha(\beta - 1)u_{xx}(x_{3n}, t_0)u(x_{3n} + h_n, t_0) \\
 &\quad + 2\alpha(\beta - 1)u_x(x_{3n}, t_0)u_x(x_{3n} + h_n, t_0) \\
 &\quad + \alpha(\beta - 1)u(x_{3n}, t_0)u_{xx}(x_{3n} + h_n, t_0) \\
 &\longrightarrow u_{xx}(x_0, t_0)(3\alpha\beta u^2(x_0, t_0) - 4\alpha u(x_0, t_0) + 1) \\
 &\quad + u_x^2(x_0, t_0)(6\alpha\beta u(x_0, t_0) - 4\alpha)
 \end{aligned}$$

as $n \rightarrow \infty$; that is,

$$u_t = (\sigma(u)u_x)_x \text{ at } (x, t) = (x_0, t_0),$$

where $\sigma(s) = \sigma_{\alpha\beta}(s) := 3\alpha\beta s^2 - 4\alpha s + 1$ ($s \in \mathbb{R}$). By continuity, we conclude that

$$u_t = (\sigma(u)u_x)_x = (\rho(u))_{xx} \text{ in } \Omega_\infty, \tag{1.9}$$

where $\rho(s) = \rho_{\alpha\beta}(s) := \alpha\beta s^3 - 2\alpha s^2 + s$ ($s \in \mathbb{R}$). For later use, let us denote

$$\begin{aligned}
 Z_\sigma &= Z_{\sigma, \alpha\beta} := \{s \in [0, 1] \mid \sigma(s) = 0\}, \\
 Z_\rho &= Z_{\rho, \alpha\beta} := \{s \in (0, 1] \mid \rho(s) = 0\}.
 \end{aligned}$$

In this paper, we study equation (1.9), coupled with the initial condition,

$$u = u_0 \text{ on } \Omega \times \{t = 0\} \tag{1.10}$$

and the Dirichlet boundary condition,

$$u = 0 \text{ on } \partial\Omega \times (0, \infty), \tag{1.11}$$

where $u_0 \in L^\infty(\Omega; [0, 1])$ is a given initial population density.

1.2. Global weak solutions

To derive a natural definition of a weak solution to problem (1.9)–(1.11), let $u_0 \in C^2(\bar{\Omega}; [0, 1])$ be such that

$$u_0(0) = u_0(L) = 0, \quad 4\alpha(u'_0(0))^2 = u''_0(0), \quad \text{and} \quad 4\alpha(u'_0(L))^2 = u''_0(L).$$

Assume that $u \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ is a global classical solution to problem (1.9)–(1.11). Fix any $T > 0$, and choose a test function $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ such that

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}).$$

Then from the integration by parts,

$$\begin{aligned} 0 &= \int_0^T \int_0^L (u_t - (\rho(u))_{xx})\varphi \, dx \, dt \\ &= - \int_0^L u_0(x)\varphi(x, 0) \, dx - \int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt. \end{aligned}$$

Conversely, assume that $u \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ is a function satisfying that

$$\int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx = 0$$

for each $T > 0$ and each $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}).$$

We will check below that u is a global classical solution to problem (1.9)–(1.11) provided that

$$u \notin Z_\rho \text{ on } \partial\Omega \times (0, \infty). \tag{1.12}$$

To show that (1.9) holds, fix any $\varphi \in C_c^\infty(\Omega_\infty)$. Choose a $T = T_\varphi > 0$ so large that $\text{spt}(\varphi) \subset\subset \Omega \times (0, T)$. Then from the integration by parts,

$$\begin{aligned} 0 &= \int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx \\ &= \int_0^T \int_0^L (-u_t + (\rho(u))_{xx})\varphi \, dx \, dt = \int_0^\infty \int_0^L (-u_t + (\rho(u))_{xx})\varphi \, dx \, dt. \end{aligned}$$

Thus, (1.9) is satisfied.

Next, to check that (1.10) holds, fix any $\psi \in C_c^\infty(\Omega)$. Choose a function $\omega \in C^\infty(\mathbb{R})$ such that

$$\omega = 1 \text{ on } (-\infty, 0] \quad \text{and} \quad \omega = 0 \text{ on } [2, \infty),$$

and define $\varphi(x, t) = \psi(x)\omega(t)$ for $(x, t) \in \bar{\Omega}_\infty$. Then with $T = 2$, it follows from (1.9) that

$$\begin{aligned} 0 &= \int_0^2 \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx \\ &= \int_0^L (u(x, 2)\varphi(x, 2) - u(x, 0)\varphi(x, 0)) \, dx - \int_0^2 \int_0^L u_t\varphi \, dx \, dt \\ &\quad + \int_0^2 (\rho(u(L, t))\varphi_x(L, t) - \rho(u(0, t))\varphi_x(0, t)) \, dt \\ &\quad - \int_0^2 ((\rho(u))_x(L, t)\varphi(L, t) - (\rho(u))_x(0, t)\varphi(0, t)) \, dt \\ &\quad + \int_0^2 \int_0^L (\rho(u))_{xx}\varphi \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx \\ &= \int_0^L (u_0(x) - u(x, 0))\psi(x) \, dx. \end{aligned}$$

Thus, (1.10) is true.

Finally, to see that (1.11) holds, fix any $\omega \in C_c^\infty((0, \infty))$. Let $T = T_\omega > 0$ be chosen so large that $\text{spt}(\omega) \subset\subset (0, T)$. Choose two functions $\psi_0, \psi_1 \in C^\infty(\mathbb{R})$ such that

$$\psi_0(x) = \begin{cases} x & \text{for } x \leq \frac{1}{4}L \\ 0 & \text{for } x \geq \frac{3}{4}L \end{cases} \quad \text{and} \quad \psi_1(x) = \begin{cases} 0 & \text{for } x \leq \frac{1}{4}L \\ -x + L & \text{for } x \geq \frac{3}{4}L. \end{cases}$$

For $i = 0, 1$, define $\varphi_i(x, t) = \psi_i(x)\omega(t)$ for $(x, t) \in \bar{\Omega}_\infty$. Then we have from (1.9) that for $i = 0, 1$,

$$\begin{aligned} 0 &= \int_0^L (u(x, T)\varphi_i(x, T) - u(x, 0)\varphi_i(x, 0)) \, dx - \int_0^T \int_0^L u_t\varphi_i \, dx \, dt \\ &\quad + \int_0^T (\rho(u(L, t))(\varphi_i)_x(L, t) - \rho(u(0, t))(\varphi_i)_x(0, t)) \, dt \\ &\quad - \int_0^T ((\rho(u))_x(L, t)\varphi_i(L, t) - (\rho(u))_x(0, t)\varphi_i(0, t)) \, dt \\ &\quad + \int_0^T \int_0^L (\rho(u))_{xx}\varphi_i \, dx \, dt + \int_0^L u_0(x)\varphi_i(x, 0) \, dx \\ &= - \int_0^T \rho(u(iL, t))\omega(t) \, dt = - \int_0^\infty \rho(u(iL, t))\omega(t) \, dt; \end{aligned}$$

that is, $\rho(u(x, t)) = 0$ for all $(x, t) \in \partial\Omega \times (0, \infty)$. Thus, (1.11) follows from (1.12) and the definition of $\rho(s)$.

Summarizing the previous discussion, we have the following.

PROPOSITION 1.1. *Let $u_0 \in C^2(\bar{\Omega}; [0, 1])$ satisfy the compatibility conditions,*

$$u_0(0) = u_0(L) = 0, \quad 4\alpha(u'_0(0))^2 = u''_0(0), \quad \text{and} \quad 4\alpha(u'_0(L))^2 = u''_0(L).$$

Assume that $u \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ is such that

$$u \notin Z_\rho \text{ on } \partial\Omega \times (0, \infty).$$

Then u is a global classical solution to problem (1.9)–(1.11) if and only if

$$\int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx = 0$$

for each $T > 0$ and each $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}).$$

Motivated by this observation, we fix the definition of a global weak solution to problem (1.9)–(1.11) as follows.

DEFINITION 1.2. *Let $u_0 \in L^\infty(\Omega; [0, 1])$ and $u \in L^\infty(\Omega_\infty; [0, 1])$.*

- (i) *Assume $Z_\rho = \emptyset$. Then we say that u is a global weak solution to problem (1.9)–(1.11) provided that for each $T > 0$ and each $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with*

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}),$$

one has

$$\int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt + \int_0^L u_0(x)\varphi(x, 0) \, dx = 0. \tag{1.13}$$

- (ii) *Assume $Z_\rho \neq \emptyset$ so that $|Z_\rho| \in \{1, 2\}$. Then we say that u is a global weak solution to (1.9)–(1.11) provided that for each $T > 0$ and each $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with*

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}),$$

(1.13) holds and that there exists a number $0 < \delta < \frac{L}{2}$ with $\min Z_\rho - \delta > 0$ such that for each $s_z \in Z_\rho$,

$$u \notin [s_z - \delta, s_z + \delta] \text{ a.e. in } ((0, \delta) \cup (L - \delta, L)) \times (0, \infty).$$

Although

$$\{(\alpha, \beta) \in [0, 1]^2 \mid Z_\rho = Z_{\rho_{\alpha\beta}} \neq \emptyset\} \neq \emptyset,$$

the adhesion-volume filling pairs $(\alpha, \beta) \in [0, 1]^2$ that we mainly consider in this paper satisfy that $Z_\rho = Z_{\rho_{\alpha\beta}} = \emptyset$ (see § 3). Thus, we only have to keep in mind definition 1.2(i) even if definition 1.2(ii) is included for the sake of completeness.

2. Classification of equation

In this section, we classify the types of equation (1.9) as follows. To do so, for any fixed adhesion-volume filling pair $(\alpha, \beta) \in [0, 1]^2$, let us write

$$I_{\alpha\beta}^+ := \{s \in [0, 1] \mid \sigma(s) > 0\} \quad \text{and} \quad I_{\alpha\beta}^- := \{s \in [0, 1] \mid \sigma(s) < 0\}.$$

Case $\beta = 1$: In this case,

$$\sigma(s) = 3\alpha s^2 - 4\alpha s + 1 = 3\alpha \left(s - \frac{2}{3}\right)^2 + 1 - \frac{4}{3}\alpha.$$

- (a) If $0 \leq \alpha < \frac{3}{4}$, then $\sigma(s) \geq 1 - \frac{4}{3}\alpha > 0$ for all $s \in [0, 1]$; hence, equation (1.9) is *forward parabolic* on $[0, 1]$. We may call this case as (F).
- (b) If $\alpha = \frac{3}{4}$, then $\sigma(s) > 1 - \frac{4}{3}\alpha = 0$ for all $s \in [0, 1] \setminus \{\frac{2}{3}\}$ and $\sigma(\frac{2}{3}) = 1 - \frac{4}{3}\alpha = 0$; that is, equation (1.9) is forward parabolic on $[0, 1] \setminus \{\frac{2}{3}\}$ and *degenerate* at $s = \frac{2}{3}$. We refer to this case as (FDF).
- (c) If $\frac{3}{4} < \alpha < 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in \left[0, \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}\right) \cup \left(\frac{2\alpha + \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}, 1\right] = I_{\alpha 1}^+, \\ < 0 & \text{for } s \in \left(\frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}, \frac{2\alpha + \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}\right) = I_{\alpha 1}^-, \\ = 0 & \text{for } s = \frac{2\alpha \pm \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha 1}^+$, *backward parabolic* on $I_{\alpha 1}^-$, and degenerate at $s = \frac{2\alpha \pm \sqrt{4\alpha^2 - 3\alpha}}{3\alpha}$. We refer to this case as (FDBDF).

- (d) If $\alpha = 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in \left[0, \frac{1}{3}\right) = I_{11}^+, \\ < 0 & \text{for } s \in \left(\frac{1}{3}, 1\right) = I_{11}^-, \\ = 0 & \text{for } s \in \left\{\frac{1}{3}, 1\right\}; \end{cases}$$

that is, equation (1.9) is forward parabolic on I_{11}^+ , *backward parabolic* on I_{11}^- , and degenerate at $s = \frac{1}{3}, 1$. We refer to this case as (FDBD).

Case $\frac{2}{3} < \beta < 1$: In this case,

$$\sigma(s) = 3\alpha\beta s^2 - 4\alpha s + 1 = 3\alpha\beta \left(s - \frac{2}{3\beta}\right)^2 + 1 - \frac{4\alpha}{3\beta}.$$

- (a) If $0 \leq \alpha < \frac{3}{4}\beta$, then $\sigma(s) \geq 1 - \frac{4\alpha}{3\beta} > 0$ for all $s \in [0, 1]$; hence, equation (1.9) is forward parabolic on $[0, 1]$. We refer to this case as (F).
- (b) If $\alpha = \frac{3}{4}\beta$, then $\sigma(s) > 1 - \frac{4\alpha}{3\beta} = 0$ for all $s \in [0, 1] \setminus \{\frac{2}{3\beta}\}$ and $\sigma(\frac{2}{3\beta}) = 1 - \frac{4\alpha}{3\beta} = 0$; that is, equation (1.9) is forward parabolic on $[0, 1] \setminus \{\frac{2}{3\beta}\}$ and degenerate at $s = \frac{2}{3\beta}$. We refer to this case as (FDF).
- (c) If $\frac{3}{4}\beta < \alpha < \frac{1}{4-3\beta}$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in [0, \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}) \cup (\frac{2\alpha + \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}, 1] = I_{\alpha\beta}^+, \\ < 0 & \text{for } s \in (\frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}, \frac{2\alpha + \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}) = I_{\alpha\beta}^-, \\ = 0 & \text{for } s = \frac{2\alpha \pm \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha\beta}^+$, backward parabolic on $I_{\alpha\beta}^-$, and degenerate at $s = \frac{2\alpha \pm \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}$. We refer to this case as (FDBDF).

- (d) If $\alpha = \frac{1}{4-3\beta}$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in [0, \frac{1}{4\alpha - 1}) = I_{\alpha\beta}^+, \\ < 0 & \text{for } s \in (\frac{1}{4\alpha - 1}, 1) = I_{\alpha\beta}^-, \\ = 0 & \text{for } s \in \{\frac{1}{4\alpha - 1}, 1\}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha\beta}^+$, backward parabolic on $I_{\alpha\beta}^-$, and degenerate at $s = \frac{1}{4\alpha - 1}, 1$. We refer to this case as (FDBD).

- (e) If $\frac{1}{4-3\beta} < \alpha \leq 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in [0, \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}) = I_{\alpha\beta}^+, \\ < 0 & \text{for } s \in (\frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}, 1] = I_{\alpha\beta}^-, \\ = 0 & \text{for } s = \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha\beta}^+$, backward parabolic on $I_{\alpha\beta}^-$, and degenerate at $s = \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}$. We refer to this case as (FDB).

Case $\beta = \frac{2}{3}$: In this case,

$$\sigma(s) = 2\alpha s^2 - 4\alpha s + 1 = 2\alpha(s - 1)^2 + 1 - 2\alpha.$$

- (a) If $0 \leq \alpha < \frac{1}{2}$, then $\sigma(s) \geq 1 - 2\alpha > 0$ for all $s \in [0, 1]$; hence, equation (1.9) is forward parabolic on $[0, 1]$. We refer to this case as (F).
- (b) If $\alpha = \frac{1}{2}$, then $\sigma(s) > 1 - 2\alpha = 0$ for all $s \in [0, 1)$ and $\sigma(1) = 0$; that is, equation (1.9) is forward parabolic on $[0, 1)$ and *degenerate* at $s = 1$. We refer to this case as (FD).
- (c) If $\frac{1}{2} < \alpha \leq 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in \left[0, \frac{2\alpha - \sqrt{4\alpha^2 - 2\alpha}}{2\alpha}\right) = I_{\frac{\alpha}{3}}^+, \\ < 0 & \text{for } s \in \left(\frac{2\alpha - \sqrt{4\alpha^2 - 2\alpha}}{2\alpha}, 1\right] = I_{\frac{\alpha}{3}}^-, \\ = 0 & \text{for } s = \frac{2\alpha - \sqrt{4\alpha^2 - 2\alpha}}{2\alpha}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\frac{\alpha}{3}}^+$, backward parabolic on $I_{\frac{\alpha}{3}}^-$, and degenerate at $s = \frac{2\alpha - \sqrt{4\alpha^2 - 2\alpha}}{2\alpha}$. We refer to this case as (FDB).

Case $0 < \beta < \frac{2}{3}$: In this case,

$$\sigma(s) = 3\alpha\beta s^2 - 4\alpha s + 1 = 3\alpha\beta\left(s - \frac{2}{3\beta}\right)^2 + 1 - \frac{4\alpha}{3\beta}.$$

- (a) If $0 \leq \alpha < \frac{1}{4-3\beta}$, then $\sigma(s) \geq 3\alpha\beta - 4\alpha + 1 > 0$ for all $s \in [0, 1]$; hence, equation (1.9) is forward parabolic on $[0, 1]$. We refer to this case as (F).
- (b) If $\alpha = \frac{1}{4-3\beta}$, then $\sigma(s) > 3\alpha\beta - 4\alpha + 1 = 0$ for all $s \in [0, 1)$ and $\sigma(1) = 0$; that is, equation (1.9) is forward parabolic on $[0, 1)$ and degenerate at $s = 1$. We refer to this case as (FD).
- (c) If $\frac{1}{4-3\beta} < \alpha \leq 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in \left[0, \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}\right) = I_{\alpha\beta}^+, \\ < 0 & \text{for } s \in \left(\frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}, 1\right] = I_{\alpha\beta}^-, \\ = 0 & \text{for } s = \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha\beta}^+$, backward parabolic on $I_{\alpha\beta}^-$, and degenerate at $s = \frac{2\alpha - \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta}$. We refer to this case as (FDB).

Case $\beta = 0$: In this case,

$$\sigma(s) = -4\alpha s + 1.$$

- (a) If $0 \leq \alpha < \frac{1}{4}$, then $\sigma(s) \geq -4\alpha + 1 > 0$ for all $s \in [0, 1]$; hence, equation (1.9) is forward parabolic on $[0, 1]$. We refer to this case as (F).
- (b) If $\alpha = \frac{1}{4}$, then $\sigma(s) > -4\alpha + 1 = 0$ for all $s \in [0, 1)$ and $\sigma(1) = 0$; that is, equation (1.9) is forward parabolic on $[0, 1)$ and degenerate at $s = 1$. We refer to this case as (FD).
- (c) If $\frac{1}{4} < \alpha \leq 1$, then

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in \left[0, \frac{1}{4\alpha}\right) = I_{\alpha 0}^+, \\ < 0 & \text{for } s \in \left(\frac{1}{4\alpha}, 1\right] = I_{\alpha 0}^-, \\ = 0 & \text{for } s = \frac{1}{4\alpha}; \end{cases}$$

that is, equation (1.9) is forward parabolic on $I_{\alpha\beta}^+$, backward parabolic on $I_{\alpha\beta}^-$, and degenerate at $s = \frac{1}{4\alpha}$. We refer to this case as (FDB).

In short, we can summarize the classification of equation (1.9) as in Fig. 1.




Figure 1. Classification of equation (1.9).

3. Main result

From this section, we mainly study problem (1.9)–(1.11) of type (FDBDF); that is, the initial-Dirichlet boundary value problem in one space dimension,

$$\begin{cases} u_t = (\sigma(u)u_x)_x = (\rho(u))_{xx} & \text{in } \Omega_\infty, \\ u = u_0 & \text{on } \Omega \times \{t = 0\}, \\ u = 0 & \text{on } \partial\Omega \times (0, \infty), \end{cases} \tag{3.1}$$

where $u_0 : \Omega \rightarrow [0, 1]$ is a given initial population density, $u(x, t) \in [0, 1]$ represents the population density at a space-time point $(x, t) \in \Omega_\infty$, the diffusivity $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$\sigma(s) = 3\alpha\beta s^2 - 4\alpha s + 1 \quad (s \in \mathbb{R})$$

for some constants $\frac{2}{3} < \beta \leq 1$ and $\frac{3}{4}\beta < \alpha < \frac{1}{4-3\beta}$, and

$$\rho(s) = \alpha\beta s^3 - 2\alpha s^2 + s \quad (s \in \mathbb{R}).$$

Letting

$$s_0^\pm = \frac{2\alpha \pm \sqrt{4\alpha^2 - 3\alpha\beta}}{3\alpha\beta},$$

we observe that $0 < s_0^- < s_0^+ < 1$ and that

$$\sigma(s) \begin{cases} > 0 & \text{for } s \in I_{\alpha\beta}^+ = [0, s_0^-] \cup (s_0^+, 1], \\ < 0 & \text{for } s \in I_{\alpha\beta}^- = (s_0^-, s_0^+), \\ = 0 & \text{for } s \in Z_\sigma = \{s_0^+, s_0^-\}. \end{cases} \tag{3.2}$$

We check below that

$$\rho(s) > 0 \quad \forall s \in (0, 1]; \tag{3.3}$$

hence, our global weak solutions to problem (3.1) should be as in definition 1.2(i). Note from $\frac{2}{3} < \beta \leq 1$ that

$$3\beta^2 - 4\beta + 1 = (3\beta - 1)(\beta - 1) \leq 0$$

so that

$$\alpha < \frac{1}{4 - 3\beta} \leq \beta;$$

thus, $\alpha^2 - \alpha\beta = \alpha(\alpha - \beta) < 0$. This implies that the equation $\rho(s) = s(\alpha\beta s^2 - 2\alpha s + 1) = 0$ has precisely one zero $s = 0$ in \mathbb{R} ; hence, inequality (3.3) holds.




Figure 2. Two possible graphs of $r = \rho(s)$ ($0 \leq s \leq 1$).

Let

$$r^* = \min\{\rho(s_0^-), \rho(1)\};$$

then from (3.2) and (3.3), we have

$$r^* > \rho(s_0^+) > 0.$$

For each $r \in [\rho(s_0^+), r^*]$, let $s^+(r) \in [s_0^+, 1]$ and $s^-(r) \in (0, s_0^-]$ denote the unique numbers with

$$\rho(s^\pm(r)) = r.$$

Let us write

$$s_1^\pm := s^\pm(\rho(s_0^+)) \quad \text{and} \quad s_2^\pm := s^\pm(r^*);$$

then

$$0 < s_1^- < s_2^- \leq s_0^- < s_0^+ = s_1^+ < s_2^+ \leq 1,$$

and

$$s_2^- = s_0^- \quad \text{or} \quad s_2^+ = 1$$

(see Fig. 2).

Initial population density: We assume that the initial population density u_0 to problem (3.1) fulfils the regularity condition,

$$u_0 \in C^{2+a}(\bar{\Omega}; [0, 1]) \text{ for some } 0 < a < 1$$

and the compatibility conditions,

$$u_0(0) = u_0(L) = 0, \quad 4\alpha(u_0'(0))^2 = u_0''(0), \quad \text{and} \quad 4\alpha(u_0'(L))^2 = u_0''(L).$$

We write

$$M_0 := \max_{\bar{\Omega}} u_0 \in [0, 1].$$

From the classical parabolic theory, we can handle the following case that the maximum value M_0 of the initial population density u_0 is strictly less than the infimum s_0^- of the backward regime (s_0^-, s_0^+) .

THEOREM 3.1 Case $M_0 < s_0^-$: smooth extinction. Assume $M_0 < s_0^-$. Then there exists a unique global classical solution $u \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ to problem (3.1) such that

$$u \in C^{2+a, 1+\frac{a}{2}}(\bar{\Omega}_T; [0, 1])$$

for all $T > 0$ and that $0 \leq u \leq M_0$ in Ω_∞ . Moreover, it follows that

$$0 \leq u(x, t) \leq \max_{\Omega} u(\cdot, s)$$

for all $x \in \Omega$ and $0 \leq s \leq t$ and that there exist two constants $C \geq 0$ and $\gamma > 0$, depending only on M_0, L, α , and β , such that

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} \leq Ce^{-\gamma t}$$

for all $t \geq 0$.

Proof. Let $M_1 = \frac{M_0 + s_0^-}{2}$. By elementary calculus, we can choose a function $\rho^* \in C^3(\mathbb{R})$ such that

$$\rho^* = \rho \text{ on } (-\infty, M_1] \quad \text{and} \quad (\rho^*)' \geq c_0 \text{ in } \mathbb{R},$$

for some constant $c_0 > 0$. Then from [11, Theorem 12.14], the modified problem,

$$\begin{cases} u_t^* = (\rho^*(u^*))_{xx} & \text{in } \Omega_\infty, \\ u^* = u_0 & \text{on } \Omega \times \{t = 0\}, \\ u^* = 0 & \text{on } \partial\Omega \times (0, \infty), \end{cases}$$

admits a unique global classical solution $u^* \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ with

$$u^* \in C^{2+a, 1+\frac{a}{2}}(\bar{\Omega}_T; [0, 1]) \quad \forall T > 0.$$

Also, it follows from [10, Theorem 1.1] that

$$0 \leq u^*(x, t) \leq \|u^*(\cdot, s)\|_{L^\infty(\Omega)} \text{ for all } x \in \bar{\Omega} \quad \text{and} \quad t \geq s \geq 0$$

and that

$$\|u^*(\cdot, t)\|_{L^\infty(\Omega)} \leq Ce^{-\gamma t} \text{ for all } t \geq 0,$$

for some constants $C \geq 0$ and $\gamma > 0$, depending only on M_0, L, α , and β . In particular, $0 \leq u^* \leq M_0 < M_1$ in Ω_∞ ; thus, from the choice of ρ^* , we see that $u := u^*$ is a global classical solution to problem (3.1).

Suppose $\tilde{u} \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ is a unique global classical solution to problem (3.1) such that

$$\tilde{u} \in C^{2+a, 1+\frac{a}{2}}(\bar{\Omega}_T; [0, 1])$$

for all $T > 0$ and that $0 \leq \tilde{u} \leq M_0$ in Ω_∞ . Then from the choice of ρ^* , \tilde{u} is also a global classical solution to the modified problem above. By uniqueness of the modified problem, we have $\tilde{u} = u^* = u$ in Ω_∞ . □

As the main result of the paper, we present the following theorem on the case that the maximum value M_0 of the initial population density u_0 exceeds the threshold s_1^- . A proof of this theorem is given in § 5.

THEOREM 3.2 Case $M_0 > s_1^-$: density mixtures and smooth extinction. Assume $M_0 > s_1^-$. Let $r_1 \in [\rho(s_0^+), r^*)$ be any number such that

$$s^-(r_1) < M_0,$$

and let $r_2 \in (r_1, r^*)$. Then there exist a function $u^* \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ with

$$u^* \in C^{2+a, 1+\frac{a}{2}}(\bar{\Omega}_T; [0, 1]) \quad \forall T > 0,$$

a nonempty bounded open set $Q \subset \Omega_\infty$ with

$$\bar{Q} \subset \Omega \times [0, \infty) \quad \text{and} \quad \bar{Q} \cap (\Omega \times \{0\}) \neq \emptyset,$$

and infinitely many global weak solutions $u \in L^\infty(\Omega_\infty; [0, 1])$ to problem (3.1) satisfying the following:

(a) Smoothing in finite time:

$$u = u^* \text{ in } \Omega_\infty \setminus \bar{Q};$$

(b) Density mixtures:

$$u \in [s^-(r_1), s^-(r_2)] \cup [s^+(r_1), s^+(r_2)] \text{ a.e. in } Q;$$

(c) Fine-scale oscillations: for any nonempty open set $O \subset Q$,

$$\text{ess\,osc}_O u := \text{ess\,sup}_O u - \text{ess\,inf}_O u \geq s^+(r_1) - s^-(r_2) > 0;$$

(d) Nonincreasing total population:

$$\int_\Omega u(x, t) \, dx = \int_\Omega u^*(x, t) \, dx \quad \forall t \geq 0,$$

and the function $t \mapsto \int_\Omega u^*(x, t) \, dx$ is nonincreasing on $[0, \infty)$;

(e) Maximum principle:

$$0 \leq u^*(x, t) \leq \|u^*(\cdot, s)\|_{L^\infty(\Omega)} \leq 1 \quad \forall x \in \bar{\Omega}, \quad \forall t \geq s \geq 0;$$

(f) Exponential and smooth extinction: there exists two constants $C > 0$ and $\gamma > 0$ such that

$$\|u^*(\cdot, t)\|_{L^\infty(\Omega)} \leq Ce^{-\gamma t} \quad \forall t \geq 0.$$

Since $Q \neq \emptyset$ is a bounded subset of Ω_∞ , it follows from (a) that the solutions u are smooth and identical on $\bar{\Omega} \times [t^*, \infty)$, where

$$0 < t^* := \sup_{(x,t) \in Q} t < \infty.$$

So properties (e) and (f) are valid for the solutions u when $t \geq t^*$. Since $\bar{Q} \subset \Omega \times [0, \infty)$, observe from (a) that the solutions u are smooth and identical near the

cylindrical boundary $\partial\Omega \times [0, \infty)$; thus, $u = 0$ pointwise on $\partial\Omega \times [0, \infty)$. From (b), (c), and $\bar{Q} \cap (\Omega \times \{0\}) \neq \emptyset$, the solutions u experience immediate fine-scale density mixtures in Q between low density regime $[s^-(r_1), s^-(r_2)]$ and high density regime $[s^+(r_1), s^+(r_2)]$. Observe also from (d), (e), and (f) that the total population in Ω decreases exponentially in time:

$$0 \leq \int_{\Omega} u(x, t) \, dx = \int_{\Omega} u^*(x, t) \, dx \leq CLe^{-\gamma t} \quad \forall t \geq 0.$$

Instead of the initial-Dirichlet boundary value problem (3.1), one may consider the Cauchy problem with the diffusivity $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ as in (3.1) and initial population density $u_0 \in C_c^{2+a}(\mathbb{R}; [0, 1])$. In this case, it is essential to study first the existence and properties of a global classical solution to the modified Cauchy problem as in (5.3) with Ω replaced by \mathbb{R} . On the other hand, in a biological viewpoint, it is also interesting to consider in problem (3.1) a Dirac delta distribution as the initial population density u_0 . This may be regarded as the case of describing population dynamics of a species, concentrated initially at a single point. Likewise, one then needs to establish first the existence and properties of a global classical solution to the modified (initial-boundary value or Cauchy) problem under the initial Dirac delta distribution u_0 .

Approach by differential inclusion: Let us take a moment here to explain our approach to prove theorem 3.2. To solve the equation in (3.1), we formally put $v_x = u$ in Ω_{∞} for some function $v : \Omega_{\infty} \rightarrow \mathbb{R}$; so we consider the equation,

$$v_t = (\rho(v_x))_x \text{ in } \Omega_{\infty}.$$

To solve the previous equation in the sense of distributions in Ω_{∞} , we may try to find a vector function $z = (v, w) \in W^{1,\infty}(\Omega_{\infty}; \mathbb{R}^{1+1})$ with $v_x \in L^{\infty}(\Omega_{\infty}; [0, 1])$ such that

$$w_x = v \quad \text{and} \quad w_t = \rho(v_x) \text{ a.e. in } \Omega_{\infty}. \tag{3.4}$$

If there is such a function $z = (v, w)$, we take $u = v_x \in L^{\infty}(\Omega_{\infty}; [0, 1])$; then from the integration by parts, for each $\varphi \in C_c^{\infty}(\Omega_{\infty})$,

$$\begin{aligned} \int_0^{\infty} \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt &= \int_0^{T_{\varphi}} \int_0^L (v_x\varphi_t + \rho(v_x)\varphi_{xx}) \, dx \, dt \\ &= \int_0^{T_{\varphi}} \int_0^L (-v\varphi_{tx} + w_t\varphi_{xx}) \, dx \, dt \\ &= \int_0^{T_{\varphi}} \int_0^L (-v\varphi_{tx} + w_x\varphi_{xt}) \, dx \, dt = 0, \end{aligned}$$

where $T_{\varphi} := \sup_{(x,t) \in \text{spt}(\varphi)} t + 1$. Hence, u is a global weak solution of the equation in (3.1) in the sense of distributions in Ω_{∞} .

On the other hand, for each $b \in \mathbb{R}$, define

$$\Sigma(b) = \Sigma_\rho(b) = \left\{ \begin{pmatrix} s & c \\ b & \rho(s) \end{pmatrix} \in \mathbb{M}^{2 \times 2} \mid 0 \leq s \leq 1, c \in \mathbb{R} \right\};$$

then system (3.4) is equivalent to the inhomogeneous partial differential inclusion,

$$\nabla z = \begin{pmatrix} v_x & v_t \\ w_x & w_t \end{pmatrix} \in \Sigma(v) \text{ a.e. in } \Omega_\infty,$$

where $\nabla = (\partial_x, \partial_t)$ is the space-time gradient operator. In this regard, utilizing the method of convex integration by Müller & Šverák [12], we aim at solving this inclusion for certain sets $K(b) \subset \Sigma(b)$ ($b \in \mathbb{R}$) in a generic setup (section 4) while reflecting the initial and Dirichlet boundary conditions in (3.1).

After the successful understanding of homogeneous partial differential inclusions in the study of crystal microstructures by Ball & James [2] and Chipot & Kinderlehrer [6], the methods of convex integration in differential inclusions have been extensively applied to many important problems; see, e.g., elliptic systems [12], the Euler equations and Onsager’s conjecture [7, 9], the porous media equation [3], active scalar equations [14], and the Muskat problem [4].

4. Generic problem

In this section that is independent of the previous sections, we develop a generic inclusion problem that can be applied to the main problem (3.1) as a special case. Since the core analysis part is essentially the same for both the Dirichlet problem and Neumann problem, we take the generic approach instead of studying the convex integration in a special setup to avoid repetition when we deal with the Neumann problem in a subsequent paper.

4.1. Two-wall inclusions

As a setup, we fix some generic notations and introduce a two-wall partial differential inclusion of inhomogeneous type.

4.1.1. *Related sets.* Let $r_1 < r_2$, and let $\omega_1, \omega_2 \in C([r_1, r_2])$ be any two functions such that

$$\max_{[r_1, r_2]} \omega_1 < \min_{[r_1, r_2]} \omega_2.$$

For each $b \in \mathbb{R}$, define the matrix sets

$$\begin{aligned} K^+(b) &= K_{\omega_2}^+(b) = \left\{ \begin{pmatrix} \omega_2(r) & c \\ b & r \end{pmatrix} \in \mathbb{M}^{2 \times 2} \mid r \in [r_1, r_2], c \in \mathbb{R} \right\}, \\ K^-(b) &= K_{\omega_1}^-(b) = \left\{ \begin{pmatrix} \omega_1(r) & c \\ b & r \end{pmatrix} \in \mathbb{M}^{2 \times 2} \mid r \in [r_1, r_2], c \in \mathbb{R} \right\}, \\ U(b) &= U_{\omega_1, \omega_2}(b) = \left\{ \begin{pmatrix} s & c \\ b & r \end{pmatrix} \in \mathbb{M}^{2 \times 2} \mid r \in (r_1, r_2), \omega_1(r) < s < \omega_2(r), c \in \mathbb{R} \right\}, \end{aligned}$$

and $K(b) = K_{\omega_1, \omega_2}(b) = K^+(b) \cup K^-(b)$. Let

$$K^+ = K_{\omega_2}^+ = \{(\omega_2(r), r) \mid r \in [r_1, r_2]\},$$

$$K^- = K_{\omega_1}^- = \{(\omega_1(r), r) \mid r \in [r_1, r_2]\},$$

and $K = K_{\omega_1, \omega_2} = K^+ \cup K^-$. Also, let

$$U = \{(s, r) \mid r \in (r_1, r_2), \omega_1(r) < s < \omega_2(r)\}.$$

4.1.2. *Two-wall inclusions.* Let

$$\Omega_{t_1}^{t_2} = \Omega \times (t_1, t_2) = (0, L) \times (t_1, t_2) \subset \mathbb{R}^2,$$

where $t_1 < t_2$ are any two fixed real numbers, and let $Q \subset \Omega_{t_1}^{t_2}$ be a nonempty open set. Consider the inhomogeneous partial differential inclusion,

$$\nabla z \in K(v) \text{ in } Q, \tag{4.1}$$

where $z = (v, w) : Q \rightarrow \mathbb{R}^2$. Regarding this, we fix some terminologies.

DEFINITION 4.1. *Let $z = (v, w) \in W^{1, \infty}(Q; \mathbb{R}^2)$. Then the function z is called a solution of inclusion (4.1) if*

$$\nabla z \in K(v) \text{ a.e. in } Q,$$

a subsolution of (4.1) if

$$\nabla z \in K(v) \cup U(v) \text{ a.e. in } Q,$$

and a strict subsolution of (4.1) if

$$\nabla z \in U(v) \text{ a.e. in } Q,$$

respectively.

Observe that if $z = (v, w) \in W^{1, \infty}(Q; \mathbb{R}^2)$ is a solution of (4.1), then

$$(v_x, w_t) \in K = K^+ \cup K^- \text{ a.e. in } Q;$$

that is, (v_x, w_t) lies either in the ‘right wall’ K^+ or in the ‘left wall’ K^- almost everywhere in Q (see Fig. 3).

4.2. Special solutions to generic problem

Continuing the previous setup, we present an important existence result on inclusion (4.1) that will serve as the main ingredient for proving theorem 3.2.




Figure 3. The right wall K^+ and left wall K^- .

Assume that $z^* = (v^*, w^*) \in C^1(\bar{\Omega}_{t_1}^{t_2}; \mathbb{R}^2)$ is a function such that in Q ,

$$w_x^* = v^*, \quad r_1 < w_t^* < r_2, \quad \text{and} \quad \omega_1(w_t^*) < v_x^* < \omega_2(w_t^*). \tag{4.2}$$

From the definition of $U(b)$ ($b \in \mathbb{R}$),

$$\nabla z^* = \begin{pmatrix} v_x^* & v_t^* \\ w_x^* & w_t^* \end{pmatrix} \in U(v^*) \text{ in } Q;$$

that is, z^* is a strict subsolution of inclusion (4.1). In particular, we have

$$(v_x^*, w_t^*) \in U \text{ in } Q. \tag{4.3}$$

Assume further that

$$(v_x^*, w_t^*)(Q) \cap \{(s, r) \in U \mid \text{dist}((s, t), \partial U) < \delta\} \neq \emptyset \tag{4.4}$$

for all sufficiently small $\delta > 0$.

We are now ready to state the main result of this section whose proof is given in § 6.

THEOREM 4.2. *Let $\epsilon > 0$. Then there exists a function $z = (v, w) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$ satisfying the following:*

- (i) z is a solution of inclusion (4.1),
- (ii) $z = z^*$ on $\bar{\Omega}_{t_1}^{t_2} \setminus Q$,
- (iii) $\nabla z = \nabla z^*$ a.e. on $\Omega_{t_1}^{t_2} \cap \partial Q$,
- (iv) $\|z - z^*\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} < \epsilon$,
- (v) $\|v_t - v_t^*\|_{L^\infty(\Omega_{t_1}^{t_2})} < \epsilon$,

(vi) for any nonempty open set $O \subset Q$,

$$\operatorname{ess\,osc}_O v_x \geq d_0,$$

where $d_0 := \min_{[r_1, r_2]} \omega_2 - \max_{[r_1, r_2]} \omega_1 > 0$.

5. Application of theorem 4.2: proof of main result

In this section, we turn back to § 3 and prove the main result of the paper, theorem 3.2, by applying theorem 4.2. For the reader's convenience, we underline the arguments in theorem 3.2 that are proved along the way.

To start the proof, assume

$$M_0 > s_1^-,$$

and fix any two numbers $r_1 < r_2$ in $[\rho(s_0^+), r^*]$ such that

$$s^-(r_1) < M_0. \tag{5.1}$$

In order to fit into the setup in § 4, for $r_1 \leq r \leq r_2$, define

$$\omega_1(r) = s^-(r) \quad \text{and} \quad \omega_2(r) = s^+(r);$$

then $\omega_1, \omega_2 \in C([r_1, r_2])$, and

$$\max_{[r_1, r_2]} \omega_1 = s^-(r_2) < s^+(r_1) = \min_{[r_1, r_2]} \omega_2.$$

Next, using elementary calculus, we can choose a function $\rho^* \in C^3(\mathbb{R})$ such that

$$\begin{cases} \rho^* = \rho \text{ on } (-\infty, s^-(r_1)] \cup [s^+(r_2), \infty), \\ (\rho^*)' > 0 \text{ on } [s^-(r_1), s^+(r_2)], \\ \rho^* < \rho \text{ on } (s^-(r_1), s^-(r_2)], \quad \text{and} \quad \rho^* > \rho \text{ on } [s^+(r_1), s^+(r_2)). \end{cases} \tag{5.2}$$

Define $\sigma^* = (\rho^*)' \in C^2(\mathbb{R})$; then $\exists c_0 > 0$ such that $\sigma^* \geq c_0$ in \mathbb{R} . Now, from [11, Theorem 12.14] and [10, Theorem 1.1], the modified problem,

$$\begin{cases} u_t^* = (\sigma^*(u^*)u_x^*)_x = (\rho^*(u^*))_{xx} & \text{in } \Omega_\infty, \\ u^* = u_0 & \text{on } \Omega \times \{t = 0\}, \\ u^* = 0 & \text{on } \partial\Omega \times (0, \infty), \end{cases} \tag{5.3}$$

possesses a unique global solution $u^* \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$ with $u^* \in C^{2+a, 1+\frac{a}{2}}(\bar{\Omega}_T; [0, 1])$ for each $T > 0$ such that

$$0 \leq u^*(x, t) \leq \|u^*(\cdot, s)\|_{L^\infty(\Omega)} \quad \text{for all } x \in \bar{\Omega} \quad \text{and} \quad t \geq s \geq 0 \tag{5.4}$$

and that

$$\|u^*(\cdot, t)\|_{L^\infty(\Omega)} \leq Ce^{-\gamma t} \quad \text{for all } t \geq 0, \tag{5.5}$$

for some constants $C > 0$ and $\gamma > 0$, depending only on M_0, L, σ^* , and $(\sigma^*)'$. Thus, (e) and (f) in theorem 3.2 are satisfied. Note also from (5.2) and (5.3) that for each

$t \geq 0$,

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u^*(x, t) \, dx &= \int_{\Omega} u_t^*(x, t) \, dx = \sigma^*(u^*(L, t))u_x^*(L, t) - \sigma^*(u^*(0, t))u_x^*(0, t) \\ &= u_x^*(L, t) - u_x^*(0, t) \leq 0 \end{aligned}$$

as $u^*(x, t) \geq 0$ for $0 < x < L$. So the second of (d) in theorem 3.2 holds.

We define

$$v^*(x, t) = \int_0^x u^*(y, t) \, dy + \int_0^t u_x^*(0, s) \, ds \quad \forall (x, t) \in \bar{\Omega}_\infty; \tag{5.6}$$

then from (5.3) and the choice of $\rho^*, v^* \in C^{3,1}(\bar{\Omega}_\infty)$ satisfies that for all $(x, t) \in \Omega_\infty$,

$$\begin{aligned} v_t^*(x, t) &= \int_0^x u_t^*(y, t) \, dy + u_x^*(0, t) = \int_0^x (\sigma^*(u^*)u_x^*)_x(y, t) \, dy + u_x^*(0, t) \\ &= \sigma^*(u^*(x, t))u_x^*(x, t) - \sigma^*(u^*(0, t))u_x^*(0, t) + u_x^*(0, t) \\ &= \sigma^*(v_x^*(x, t))v_{xx}^*(x, t) = (\rho^*(v_x^*))_x(x, t). \end{aligned}$$

Hence, v^* is a global solution to the problem,

$$\begin{cases} v_t^* = (\rho^*(v_x^*))_x & \text{in } \Omega_\infty, \\ v^* = v_0 & \text{on } \Omega \times \{t = 0\}, \\ v_x^* = 0 & \text{on } \partial\Omega \times (0, \infty), \end{cases} \tag{5.7}$$

where

$$v_0(x) := \int_0^x u_0(y) \, dy \quad \forall x \in \bar{\Omega}.$$

In turn, we define

$$w^*(x, t) = \int_0^t \rho^*(v_x^*(x, s)) \, ds + \int_0^x v_0(y) \, dy \quad \forall (x, t) \in \bar{\Omega}_\infty;$$

then from (5.6) and (5.7),

$$\begin{cases} w_x^* = v^* \\ w_t^* = \rho^*(v_x^*) \end{cases} \quad \text{in } \Omega_\infty, \tag{5.8}$$

$v_x^* = u^* \in C^{2,1}(\bar{\Omega}_\infty; [0, 1])$, and $z^* := (v^*, w^*) \in (C^{3,1} \times C^{4,2})(\bar{\Omega}_\infty)$.

Define

$$Q = \{(x, t) \in \Omega_\infty \mid s^-(r_1) < v_x^*(x, t) < s^+(r_2)\}.$$

Fix a number $T_0 > 0$ so large that $Ce^{-\gamma T_0} < s^-(r_1)$. Then from (5.4) and (5.5),

$$v_x^*(x, t) = u^*(x, t) < s^-(r_1) \quad \forall (x, t) \in \Omega \times [T_0, \infty);$$

thus,

$$Q \subset \Omega_0^{T_0} = \Omega \times (0, T_0).$$

Since $u_0(0) = u_0(L) = 0$ and $M_0 > s^-(r_1)$, we can take a point $x_0 \in \Omega$ such that

$$s^-(r_1) < v_x^*(x_0, 0) = u_0(x_0) < s^+(r_2).$$

Then by continuity, we can take an $r_0 \in (0, T_0)$ with $r_0 < \min\{x_0, L - x_0\}$ so small that

$$s^-(r_1) < v_x^*(x, t) < s^+(r_2) \quad \forall (x, t) \in (\Omega \times [0, \infty)) \cap B_{r_0}(x_0, 0);$$

thus, $\Omega_\infty \cap B_{r_0}(x_0, 0) \subset Q \neq \emptyset$ so that

$$(x_0 - r_0, x_0 + r_0) \times \{0\} \subset \underline{\bar{Q}} \cap (\Omega \times \{0\}) \neq \emptyset.$$

Note here that Q is a nonempty bounded open subset of Ω_∞ . Observe also that

$$\bar{Q} \subset \Omega \times [0, \infty)$$

as \bar{Q} is compact and $v_x^*(0, t) = v_x^*(L, t) = 0$ for all $t \geq 0$.

Following the notations in § 4, note from the choice of ρ^* , the definition of Q , (5.1), (5.7), and (5.8) that (4.2) holds in Q and (4.4) holds for all sufficiently small $\delta > 0$. Thus, for any fixed $\epsilon > 0$, we can apply theorem 4.2 (with $t_1 = 0$ and $t_2 = T_0$ in the current case) to obtain a function $z = z_\epsilon \in W^{1,\infty}(\Omega_0^{T_0}; \mathbb{R}^2)$ satisfying the following, where $z = (v, w)$:

- (i) z is a solution of inclusion (4.1);
- (ii) $z = z^*$ on $\bar{\Omega}_0^{T_0} \setminus Q$;
- (iii) $\nabla z = \nabla z^*$ a.e. on $\Omega_0^{T_0} \cap \partial Q$;
- (iv) $\|z - z^*\|_{L^\infty(\Omega_0^{T_0}; \mathbb{R}^2)} < \epsilon$;
- (v) $\|v_t - v_t^*\|_{L^\infty(\Omega_0^{T_0})} < \epsilon$;
- (vi) for any nonempty open set $O \subset Q$,

$$\text{ess osc}_O v_x \geq d_0,$$

where $d_0 = \min_{[r_1, r_2]} \omega_2 - \max_{[r_1, r_2]} \omega_1 = s^+(r_1) - s^-(r_2) > 0$.

For each $(x, t) \in \bar{\Omega} \times [T_0, \infty)$, define

$$z(x, t) = z^*(x, t); \tag{5.9}$$

then from (ii), $z = (v, w) \in W^{1,\infty}(\Omega_T; \mathbb{R}^2)$ for all $T > 0$.

For each $t \geq 0$, define

$$u(\cdot, t) = v_x(\cdot, t) \text{ a.e. in } \Omega; \tag{5.10}$$

then $u \in L^\infty(\Omega_T)$ for all $T > 0$. First, from (vi), (c) in theorem 3.2 is fulfilled. Also, note from (ii) and (5.6) that for every $t \geq 0$,

$$\begin{aligned} \int_{\Omega} u(x, t) \, dx &= \int_{\Omega} v_x(x, t) \, dx = v(L, t) - v(0, t) \\ &= v^*(L, t) - v^*(0, t) = \int_{\Omega} v_x^*(x, t) \, dx = \int_{\Omega} u^*(x, t) \, dx; \end{aligned}$$

hence, the first of (d) in theorem 3.2 holds. From (ii), (5.6), and (5.9), we have

$$u = v_x = v_x^* = u^* \in [0, 1] \text{ in } \Omega_\infty \setminus \bar{Q},$$

and from (iii) and (5.6), we get

$$u = v_x = v_x^* = u^* \in [0, 1] \text{ a.e. in } \Omega_\infty \cap \partial Q.$$

In particular, (a) in theorem 3.2 is satisfied. From (i), we have

$$\nabla z \in K(v) \text{ a.e. in } Q;$$

that is, a.e. in Q ,

$$\begin{cases} u = v_x \in [s^-(r_1), s^-(r_2)] \cup [s^+(r_1), s^+(r_2)], \\ w_t = \rho(v_x), \\ w_x = v. \end{cases} \tag{5.11}$$

In particular, we see that $u \in [0, 1]$ a.e. in Ω_∞ , that is, $u \in L^\infty(\Omega_\infty; [0, 1])$ and that (b) in theorem 3.2 holds.

Now, we check that u is a global weak solution to problem (3.1). To do so, fix any $T > 0$ and any test function $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$ with

$$\varphi = 0 \text{ on } (\partial\Omega \times [0, T]) \cup (\Omega \times \{t = T\}). \tag{5.12}$$

Observe from (ii), (5.7), (5.9)–(5.12), the choice of ρ^* , and the integration by parts that

$$\begin{aligned} & \int_0^T \int_0^L (u\varphi_t + \rho(u)\varphi_{xx}) \, dx \, dt \\ &= \int_0^T \int_0^L (u\varphi_t + w_t\varphi_{xx}) \, dx \, dt \\ &= \int_0^T w_t^*(L, t)\varphi_x(L, t) \, dt - \int_0^T w_t^*(0, t)\varphi_x(0, t) \, dt + \int_0^L w_x^*(x, 0)\varphi_x(x, 0) \, dx \\ &= \int_0^T \rho^*(v_x^*(L, t))\varphi_x(L, t) \, dt - \int_0^T \rho^*(v_x^*(0, t))\varphi_x(0, t) \, dt + \int_0^L v_0(x)\varphi_x(x, 0) \, dx \\ &= - \int_0^L v_0'(x)\varphi(x, 0) \, dx + v_0(L)\varphi(L, 0) - v_0(0)\varphi(0, 0) = - \int_0^L u_0(x)\varphi(x, 0) \, dx. \end{aligned}$$

Thus, according to definition 1.2(i), u is a global weak solution to (3.1).

Since u^* itself is not a global weak solution to problem (3.1), it follows from (iv) that there are infinitely many global weak solutions to (3.1) that satisfy properties (a)–(f) in theorem 3.2.

The proof of theorem 3.2 is now complete.

6. Proof of theorem 4.2

Following § 4, this section presents a proof of theorem 4.2 with the help of the key lemma, lemma 6.1, to be proved in § 7.

6.1. Selection of an in-approximation to K

For each $b \in \mathbb{R}$ and $0 \leq \lambda < \frac{1}{2}$, define the matrix set $U^\lambda(b) = U_{\omega_1, \omega_2}^\lambda(b) \subset \mathbb{M}^{2 \times 2}$ by

$$U^\lambda(b) = \left\{ \begin{pmatrix} s & c \\ b & r \end{pmatrix} \mid c \in \mathbb{R}, (1 - \lambda)r_1 + \lambda r_2 < r < \lambda r_1 + (1 - \lambda)r_2, \right. \\ \left. (1 - \lambda)\omega_1(r) + \lambda\omega_2(r) < s < \lambda\omega_1(r) + (1 - \lambda)\omega_2(r) \right\},$$

and let $U^\lambda = U_{\omega_1, \omega_2}^\lambda \subset \mathbb{R}^2$ be given by

$$U^\lambda = \left\{ (s, r) \mid (1 - \lambda)r_1 + \lambda r_2 < r < \lambda r_1 + (1 - \lambda)r_2, \right. \\ \left. (1 - \lambda)\omega_1(r) + \lambda\omega_2(r) < s < \lambda\omega_1(r) + (1 - \lambda)\omega_2(r) \right\}.$$

Observe that $U^0(b) = U(b)$ for every $b \in \mathbb{R}$ and that $U^0 = U$.

Since $|Q| < \infty$, we can select a sequence $\{\lambda_i\}_{i \in \mathbb{N}}$ in \mathbb{R} with

$$\frac{1}{(2i + 1)^2} < \lambda_i < \frac{1}{(2i)^2} < \frac{1}{2} \quad \forall i \in \mathbb{N} \tag{6.1}$$

such that for every $i \in \mathbb{N}$,

$$\left| \{(x, t) \in Q \mid (v_x^*(x, t), w_t^*(x, t)) \in \partial U^{\lambda_i}\} \right| = 0.$$

Thanks to (4.4), we may assume

$$Q_i := \{(x, t) \in Q \mid (v_x^*(x, t), w_t^*(x, t)) \in U^{\lambda_i} \setminus \bar{U}^{\lambda_{i-1}}\} \neq \emptyset \quad \forall i \in \mathbb{N},$$

where $U^{\lambda_0} := \emptyset$; then from (4.3), $\{Q_i\}_{i \in \mathbb{N}}$ is a sequence of disjoint open subsets of Q whose union has measure $|Q|$.

For each $i \in \mathbb{N}$, let $\lambda'_i = \frac{\lambda_i + \lambda_{i+1}}{2}$. Observe from (6.1) that for all $i \in \mathbb{N}$,

$$\begin{aligned} \frac{\lambda_i - \lambda'_{i+1}}{\lambda_{i+1} - \lambda'_{i+1}} &= \frac{\lambda_i - \frac{\lambda_{i+1} + \lambda_{i+2}}{2}}{\lambda_{i+1} - \frac{\lambda_{i+1} + \lambda_{i+2}}{2}} = \frac{2\lambda_i - \lambda_{i+1} - \lambda_{i+2}}{\lambda_{i+1} - \lambda_{i+2}} \\ &\leq \frac{\frac{2}{(2i)^2} - \frac{1}{(2(i+1)+1)^2} - \frac{1}{(2(i+2)+1)^2}}{\frac{1}{(2(i+1)+1)^2} - \frac{1}{(2(i+2))^2}} \\ &= \frac{2(64i^3 + 308i^2 + 480i + 225)(i + 2)^2}{i^2(2i + 5)^2(4i + 7)} \leq 36 \end{aligned}$$

and that

$$\begin{aligned} \frac{\lambda_i - \lambda'_{i+1}}{1 - 2\lambda'_{i+1}} &= \frac{\lambda_i - \frac{\lambda_{i+1} + \lambda_{i+2}}{2}}{1 - \lambda_{i+1} - \lambda_{i+2}} = \frac{1}{2} \cdot \frac{2\lambda_i - \lambda_{i+1} - \lambda_{i+2}}{1 - \lambda_{i+1} - \lambda_{i+2}} \\ &\leq \frac{1}{2} \cdot \frac{\frac{2}{(2i)^2} - \frac{1}{(2(i+1)+1)^2} - \frac{1}{(2(i+2)+1)^2}}{1 - \frac{1}{(2(i+1))^2} - \frac{1}{(2(i+2))^2}} \\ &= \frac{(i + 1)^2(i + 2)^2(64i^3 + 308i^2 + 480i + 225)}{i^2(2i + 3)^2(2i + 5)^2(4i^4 + 24i^3 + 50i^2 + 42i + 11)}. \end{aligned}$$

From these observations, we deduce that for each $i \in \mathbb{N}$,

$$0 < \alpha_i := \frac{\lambda_i - \lambda'_{i+1}}{1 - 2\lambda'_{i+1}} \leq 36 \frac{\lambda_{i+1} - \lambda'_{i+1}}{1 - 2\lambda'_{i+1}} \tag{6.2}$$

and that

$$0 < \ell_0 := \sum_{i=1}^{\infty} \alpha_i < 1.$$

Fix a number $\kappa_0 \in (1, 1/\ell_0)$, and define

$$\beta_1 = 1 - \frac{1}{\kappa_0};$$

then

$$\ell_0 < \kappa_0 \ell_0 < 1 \quad \text{and} \quad 0 < \beta_1 = 1 - 1/\kappa_0 < 1 - \ell_0 < 1.$$

In turn, for each $i \in \mathbb{N}$, define

$$\beta_{i+1} = \beta_i + \frac{\alpha_i}{\kappa_0 \ell_0}; \tag{6.3}$$

then the sequence $\{\beta_i\}_{i \in \mathbb{N}}$ fulfils that

$$0 < \beta_1 < \beta_2 < \dots < 1, \quad \lim_{i \rightarrow \infty} \beta_i = 1,$$

and

$$\frac{\lambda_i - \lambda'_{i+1}}{1 - 2\lambda'_{i+1}} = \kappa_0 \ell_0 (\beta_{i+1} - \beta_i) \leq 36 \frac{\lambda_{i+1} - \lambda'_{i+1}}{1 - 2\lambda'_{i+1}} \quad \forall i \in \mathbb{N}.$$

For each $i \in \mathbb{N}$, let

$$U_i^+ = \left\{ (s, r) \left| \begin{array}{l} (1 - \lambda_{i+1})r_1 + \lambda_{i+1}r_2 < r < \lambda_{i+1}r_1 + (1 - \lambda_{i+1})r_2, \\ \lambda_i \omega_1(r) + (1 - \lambda_i)\omega_2(r) < s < \lambda_{i+1}\omega_1(r) + (1 - \lambda_{i+1})\omega_2(r) \end{array} \right. \right\},$$

$$U_i^- = \left\{ (s, r) \left| \begin{array}{l} (1 - \lambda_{i+1})r_1 + \lambda_{i+1}r_2 < r < \lambda_{i+1}r_1 + (1 - \lambda_{i+1})r_2, \\ (1 - \lambda_{i+1})\omega_1(r) + \lambda_{i+1}\omega_2(r) < s < (1 - \lambda_i)\omega_1(r) + \lambda_i\omega_2(r) \end{array} \right. \right\}.$$

Next, for each $i \in \mathbb{N}$, let

$$I_{i+1} = [(1 - \lambda_{i+1})r_1 + \lambda_{i+1}r_2, \lambda_{i+1}r_1 + (1 - \lambda_{i+1})r_2] \subset \mathbb{R},$$

$$\eta_i^{1,+} = \min_{r, \bar{r} \in I_{i+1}} |(\lambda'_i \omega_1(r) + (1 - \lambda'_i)\omega_2(r), r) - (\lambda_i \omega_1(\bar{r}) + (1 - \lambda_i)\omega_2(\bar{r}), \bar{r})|,$$

$$\eta_i^{2,+} = \min_{r, \bar{r} \in I_{i+1}} |(\lambda'_i \omega_1(r) + (1 - \lambda'_i)\omega_2(r), r) - (\lambda_{i+1}\omega_1(\bar{r}) + (1 - \lambda_{i+1})\omega_2(\bar{r}), \bar{r})|,$$

$$\eta_i^{1,-} = \min_{r, \bar{r} \in I_{i+1}} |((1 - \lambda'_i)\omega_1(r) + \lambda'_i\omega_2(r), r) - ((1 - \lambda_i)\omega_1(\bar{r}) + \lambda_i\omega_2(\bar{r}), \bar{r})|,$$

$$\eta_i^{2,-} = \min_{r, \bar{r} \in I_{i+1}} |((1 - \lambda'_i)\omega_1(r) + \lambda'_i\omega_2(r), r) - ((1 - \lambda_{i+1})\omega_1(\bar{r}) + \lambda_{i+1}\omega_2(\bar{r}), \bar{r})|,$$

and

$$\eta_i = \min\{\eta_i^{1,+}, \eta_i^{2,+}, \eta_i^{1,-}, \eta_i^{2,-}, (\lambda_i - \lambda_{i+1})(r_2 - r_1)\} > 0.$$

6.2. Main lemma

Define

$$\zeta_0^\pm(s, r) = \theta_0(\text{dist}((s, r), K^\pm)) \quad ((s, r) \in \mathbb{R}^2),$$

where $\theta_0 \in C^\infty([0, \infty))$ is a cutoff function such that $0 \leq \theta_0 \leq 1$ on $[0, \infty)$, $\theta_0 = 1$ on $[0, d_0/4]$, and $\theta_0 = 0$ on $[d_0/2, \infty)$. Then we can choose an integer $i_0 \geq 2$ so large that for all $i \geq i_0$,

$$\zeta_0^\pm(s, r) = 1 \quad \forall (s, r) \in U_{i-1}^\pm,$$

respectively.

We present here the key lemma to finish the proof of theorem 4.2. A proof of this lemma is provided in § 7.

LEMMA 6.1. *Let $\epsilon > 0$. Then there exist a sequence $\{z_i\}_{i \in \mathbb{N}} = \{(v_i, w_i)\}_{i \in \mathbb{N}}$ in $W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$, three sequences $\{\mathcal{F}_i\}_{i \in \mathbb{N}}$ and $\{\mathcal{F}_i^\pm\}_{i \in \mathbb{N}}$ as follows, and three positive constants c_1, c_2 , and c_3 such that for each $i \in \mathbb{N}$, the following are satisfied:*

- (a) $\mathcal{F}_i = \{D_{ij} \mid j \in \mathbb{N}\}$ is a countable collection of disjoint rhombic domains in $Q_1 \cup \dots \cup Q_i$;
- (b) $\mathcal{F}_i^+ = \{T_{ij} \mid j \in \mathbb{N}\}$ is a countable collection of disjoint triangular domains in $Q_1 \cup \dots \cup Q_i$;
- (c) $\mathcal{F}_i^- = \{R_{ij} \mid j \in \mathbb{N}\}$ is a countable collection of disjoint rhombic domains in $Q_1 \cup \dots \cup Q_i$;
- (d) $(\cup_{j \in \mathbb{N}} T_{ij}) \cap (\cup_{j \in \mathbb{N}} R_{ij}) = \emptyset$ and

$$|Q_1 \cup \dots \cup Q_i| = |\cup_{j \in \mathbb{N}} D_{ij}| = |\cup_{j \in \mathbb{N}} (T_{ij} \cup R_{ij})|;$$

- (e) for each $D \in \mathcal{F}_i$, there are four disjoint triangular domains $T_D^1, T_D^2, T_D^3, T_D^4 \in \mathcal{F}_i^+$ and one rhombic domain $R_D \in \mathcal{F}_i^-$ such that

$$T_D^1 \cup T_D^2 \cup T_D^3 \cup T_D^4 \cup R_D \subset D \text{ and } |D| = |T_D^1 \cup T_D^2 \cup T_D^3 \cup T_D^4 \cup R_D|;$$

- (f) one has

$$\sup_{j \in \mathbb{N}} \text{diam } D_{ij} \leq \frac{1}{2^i};$$

- (g) one has

$$\begin{cases} \text{for each } T \in \mathcal{F}_i^+, z_i \in C^1(\bar{T}; \mathbb{R}^2), \text{ and } ((v_i)_x, (w_i)_t) \in U_i^+ \text{ in } T, \\ \text{for each } R \in \mathcal{F}_i^-, z_i \in C^1(\bar{R}; \mathbb{R}^2), \text{ and } ((v_i)_x, (w_i)_t) \in U_i^- \text{ in } R, \\ z_i = z_0 \text{ on } \Omega_{t_1}^{t_2} \setminus (Q_1 \cup \dots \cup Q_i), \\ \nabla z_i = \nabla z_0 \text{ a.e. on } \Omega_{t_1}^{t_2} \cap \partial Q, \end{cases}$$

where $z_0 = (v_0, w_0) := z^*$;

- (h) one has

$$\begin{cases} (w_i)_x = v_i \text{ a.e. in } Q; \\ \|(v_i)_t - (v_{i-1})_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \leq \frac{\epsilon}{2^{i+1}}; \\ \|z_i - z_{i-1}\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \leq \frac{\epsilon}{2^{i+1}}; \\ |z_i(x, t) - z_i(y, s)| \leq c_1 |(x, t) - (y, s)| \quad \forall (x, t), (y, s) \in \Omega_{t_1}^{t_2}; \end{cases}$$

- (i) if $i \geq 2$, then

$$\int_{\Omega_{t_1}^{t_2}} |\nabla z_i - \nabla z_{i-1}| \, dx \, dt \leq c_2 ((\beta_i - \beta_{i-1})|Q| + |Q_i|);$$

- (j) if $i \geq i_0$ and $D \in \mathcal{F}_{i-1}$, then

$$\begin{aligned} \int_D \zeta_0^\pm((v_i)_x, (w_i)_t) \, dx \, dt &\geq c_3(\beta_i - \beta_{i-1})|D| \quad \text{and} \\ \int_D \zeta_0^\pm((v_i)_x, (w_i)_t) \, dx \, dt &\geq (1 - (\beta_i - \beta_{i-1})) \int_D \zeta_0^\pm((v_{i-1})_x, (w_{i-1})_t) \, dx \, dt. \end{aligned}$$

6.3. Completion of proof

Utilizing lemma 6.1, we now complete the proof of theorem 4.2.

From the third of (h) in lemma 6.1, it follows that for all $i > j \geq 1$,

$$\|z_i - z_j\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \leq \epsilon \left(\frac{1}{2^{i+1}} + \dots + \frac{1}{2^{j+2}} \right) \leq \frac{\epsilon}{2^{j+1}};$$

thus, $\{z_i\}_{i \in \mathbb{N}}$ is a Cauchy sequence in $L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$ so that from the fourth of (h) in lemma 6.1,

$$z_i \rightarrow z \text{ in } L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2) \text{ as } i \rightarrow \infty, \tag{6.4}$$

for some $z = (v, w) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$ satisfying

$$|z(x, t) - z(y, s)| \leq c_1 |(x, t) - (y, s)| \quad \forall (x, t), (y, s) \in \Omega_{t_1}^{t_2}.$$

Convergence (6.4) together with the third of (g) and third of (h) in lemma 6.1 implies that

$$z = z_0 = z^* \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q$$

and

$$\|z - z^*\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} = \|z - z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \leq \frac{\epsilon}{2} < \epsilon;$$

that is, (ii) and (iv) in theorem 4.2 hold.

From (i) in lemma 6.1, we have

$$\|\nabla z_i - \nabla z_j\|_{L^1(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} \leq c_2 \left(|\beta_i - \beta_j| |Q| + \sum_{k=\min\{i,j\}+1}^{\infty} |Q_k| \right) \rightarrow 0$$

as $i, j \rightarrow \infty$; thus, after passing to a subsequence if necessary, for a.e. $(x, t) \in \Omega_{t_1}^{t_2}$,

$$\nabla z_i(x, t) \rightarrow \nabla z(x, t) \text{ in } \mathbb{M}^{2 \times 2} \text{ as } i \rightarrow \infty. \tag{6.5}$$

In particular, this pointwise convergence holds for a.e. $(x, t) \in \Omega_{t_1}^{t_2} \cap \partial Q$. Thus, (iii) in theorem 4.2 follows from the fourth of (g) in lemma 6.1.

From (6.4), (6.5), and the first and second of (h) in lemma 6.1, we have

$$w_x = v \text{ a.e. in } Q \tag{6.6}$$

and

$$\|v_t - v_t^*\|_{L^\infty(\Omega_{t_1}^{t_2})} = \|v_t - (v_0)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \leq \frac{\epsilon}{2} < \epsilon;$$

that is, (v) in theorem 4.2 is fulfilled.

For each $i \in \mathbb{N}$, observe from (b), (c), (d), and the first and second of (g) in lemma 6.1 and from the definition of U_i^\pm and $K = K^+ \cup K^-$ that

$$\begin{aligned} \int_Q \text{dist}(((v_i)_x, (w_i)_t), K) \, dx \, dt &= \left(\sum_{\ell=1}^i + \sum_{\ell=i+1}^\infty \right) \int_{Q_\ell} \text{dist}(((v_i)_x, (w_i)_t), K) \, dx \, dt \\ &\leq S_M |Q| \lambda_i + d_1 \sum_{\ell=i+1}^\infty |Q_\ell|, \end{aligned}$$

where $d_1 := \text{diam } U$ and

$$S_M := \max_{[r_1, r_2]} (\omega_2 - \omega_1) > 0.$$

Here, letting $i \rightarrow \infty$, we obtain from (6.5) that

$$\int_Q \text{dist}((v_x, w_t), K) \, dx \, dt = 0;$$

that is,

$$(v_x, w_t) \in K \text{ a.e. in } Q \tag{6.7}$$

as $K \subset \mathbb{R}^2$ is compact. This inclusion together with (6.6) and the definition of $K(b)$ ($b \in \mathbb{R}$) implies that

$$\nabla z \in K(v) \text{ a.e. in } Q;$$

hence, (i) in theorem 4.2 holds.

Finally, to verify (vi) in theorem 4.2, choose an integer $i_1 \geq i_0$ so large that for all $i \geq i_1$,

$$\lim_{n \rightarrow \infty} (1 - (\beta_n - \beta_{n-1}))(1 - (\beta_{n-1} - \beta_{n-2})) \cdots (1 - (\beta_{i_1+1} - \beta_{i_1})) \geq \frac{1}{2}.$$

Fix any integer $i \geq i_1$, and let $D \in \mathcal{F}_{i-1}$. Then note from (6.5) and (j) in lemma 6.1 that

$$\begin{aligned} \int_D \zeta_0^\pm(v_x, w_t) \, dx \, dt &= \lim_{n \rightarrow \infty} \int_D \zeta_0^\pm((v_n)_x, (w_n)_t) \, dx \, dt \\ &\geq \limsup_{n \rightarrow \infty} (1 - (\beta_n - \beta_{n-1})) \int_D \zeta_0^\pm((v_{n-1})_x, (w_{n-1})_t) \, dx \, dt \\ &\geq \limsup_{n \rightarrow \infty} (1 - (\beta_n - \beta_{n-1}))(1 - (\beta_{n-1} - \beta_{n-2})) \\ &\quad \times \int_D \zeta_0^\pm((v_{n-2})_x, (w_{n-2})_t) \, dx \, dt \\ &\quad \vdots \\ &\geq \limsup_{n \rightarrow \infty} (1 - (\beta_n - \beta_{n-1})) \cdots (1 - (\beta_{i+1} - \beta_i)) \\ &\quad \times \int_D \zeta_0^\pm((v_i)_x, (w_i)_t) \, dx \, dt \\ &\geq \frac{c_3}{2} (\beta_i - \beta_{i-1}) |D| > 0. \end{aligned}$$

Next, let $O \subset Q$ be any nonempty open set. Since $\{Q_i\}_{i \in \mathbb{N}}$ is a Vitali cover of Q , we have $O \cap Q_{i_2} \neq \emptyset$ for some $i_2 \in \mathbb{N}$. Thus, from (a), (d), and (f) in lemma 6.1, there exist an $i_3 > \max\{i_1, i_2\}$ and a $D_3 \in \mathcal{F}_{i_3-1}$ such that

$$D_3 \subset O \cap Q_{i_2}.$$

This inclusion and the above positivity estimate imply that

$$\int_O \zeta_0^\pm(v_x, w_t) \, dx \, dt \geq \int_{D_3} \zeta_0^\pm(v_x, w_t) \, dx \, dt > 0.$$

So from the definition of ζ_0^\pm , there are two disjoint sets $O^\pm \subset O$ of positive measure such that

$$\text{dist}((v_x, w_t), K^\pm) < \frac{d_0}{2} \text{ in } O^\pm;$$

thus, with (6.7), we conclude that

$$(v_x, w_t) \in K^\pm \text{ a.e. in } O^\pm.$$

Therefore, (vi) in theorem 4.2 follows from the definition of K^\pm .

The proof of theorem 4.2 is now complete.

7. Proof of lemma 6.1

This final section is entirely devoted to a proof of lemma 6.1. First, we begin with an elementary lemma in subsection 7.1 that provides us with a building block for our constructions. Then we perform inductive surgeries in subsection 7.2 that complete the proof of lemma 6.1.

7.1. Piecewise affine maps with boundary trace 0

Although the following theorem is essential in our constructions, its proof is so elementary that we omit it (see [17] and Fig. 4).

LEMMA 7.1. *Let τ^\pm and δ be any three positive numbers, and let $D = D_\delta$ be the interior of the convex hull of the four points $(\pm\delta, 0)$ and $(0, \pm 1)$ in \mathbb{R}^2 ; that is,*

$$D = \text{int co}\{(\delta, 0), (-\delta, 0), (0, 1), (0, -1)\}.$$

Let $\varphi = \varphi_{\tau^+, \tau^-, \delta} : \bar{D} \rightarrow \mathbb{R}$ be the piecewise affine map defined as follows: for each $0 \leq t \leq 1$, define

$$\varphi(x, t) = \begin{cases} \tau^+(x + \delta - \delta t), & -\delta + \delta t \leq x \leq \frac{\tau^+}{\tau^+ + \tau^-}(-\delta + \delta t), \\ -\tau^-x, & \frac{\tau^+}{\tau^+ + \tau^-}(-\delta + \delta t) \leq x \leq \frac{\tau^+}{\tau^+ + \tau^-}(\delta - \delta t), \\ \tau^+(x - \delta + \delta t), & \frac{\tau^+}{\tau^+ + \tau^-}(\delta - \delta t) \leq x \leq \delta - \delta t; \end{cases}$$

and for each $(x, t) \in \bar{D}$ with $t \leq 0$, define $\varphi(x, t) = \varphi(x, -t)$. Then φ satisfies the following:

- (i) $\varphi \in W_0^{1,\infty}(D)$;
- (ii) there are five disjoint domains $T^1, T^2, T^3, T^4, R \subset D$ which cover D in the sense of Vitali, and φ is affine in each of T^1, T^2, T^3, T^4, R ;
- (iii) after a proper ordering of T^1, T^2, T^3, T^4, R ,

$$\nabla \varphi = \begin{cases} (\tau^+, \delta\tau^+) & \text{in } T^1 \cup T^2, \\ (\tau^+, -\delta\tau^+) & \text{in } T^3 \cup T^4, \\ (-\tau^-, 0) & \text{in } R; \end{cases}$$

- (iv) for each $t \in [-1, 1]$,

$$\int_{-\delta + \delta|t|}^{\delta - \delta|t|} \varphi(x, t) \, dx = 0;$$

- (v) $\|\varphi\|_{L^\infty(D)} = \frac{\tau^+ \tau^-}{\tau^+ + \tau^-} \delta$;

$$(vi) \begin{cases} |\{(x, t) \in D \mid \varphi_x(x, t) = \tau^+\}| = \frac{\tau^-}{\tau^+ + \tau^-} |D|, \\ |\{(x, t) \in D \mid \varphi_x(x, t) = -\tau^-\}| = \frac{\tau^+}{\tau^+ + \tau^-} |D|. \end{cases}$$

7.2. Inductive surgeries

In this subsection, we prove lemma 6.1 by utilizing affine maps in lemma 7.1.

Fix any $\epsilon \in (0, 1]$. Then we perform the first two surgeries in order that fulfil (a)–(h) for $i = 1$ and (a)–(j) for $i = 2$. We do not proceed with the n th surgery




Figure 4. Five disjoint domains $T^1, T^2, T^3, T^4, R \subset D$ in lemma 7.1.

under the assumption that we have performed the surgeries up to the $(n - 1)$ th one as it is essentially the repetition of the second one under the fulfilment of the first surgery.

7.2.1. *The first surgery.* In this first step, we construct a function $z_1 = (v_1, w_1) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$, three countable collections \mathcal{F}_1 and \mathcal{F}_1^\pm , and a constant $c_1 > 0$ satisfying (a)–(h) for $i = 1$, where c_1 is independent of the index i .

Let $\delta_1 \in (0, 1]$ be such that

$$\delta_1 < \min \left\{ \frac{\epsilon}{2^{1+1}\sqrt{1 + L^2}S_M}, \frac{\eta_1}{4} \right\}, \tag{7.1}$$

where the constant $S_M > 0$ is as in subsection 6.3. Since ∇z_0 is uniformly continuous in Q_1 , there exists a number $\gamma_1 \in (0, \frac{1}{2^{1+1}}]$ such that

$$(x, t), (y, s) \in Q_1, |(x, t) - (y, s)| \leq \gamma_1 \implies |\nabla z_0(x, t) - \nabla z_0(y, s)| \leq \delta_1. \tag{7.2}$$

Consider the diamond D_{δ_1} . From the Vitali covering lemma, we can choose a sequence $\{(x_{1j}, t_{1j})\}_{j \in \mathbb{N}}$ in Q_1 and a sequence $\{\nu_{1j}\}_{j \in \mathbb{N}}$ in $(0, \gamma_1]$ such that the sequence $\{\tilde{D}_{1j}\}_{j \in \mathbb{N}}$ forms a Vitali cover of Q_1 , where $\tilde{D}_{1j} := (x_{1j}, t_{1j}) + \nu_{1j}D_{\delta_1} \subset \subset Q_1$ ($j \in \mathbb{N}$).

For each $j \in \mathbb{N}$, let

$$(s_{1j}, r_{1j}) = ((v_0)_x(x_{1j}, t_{1j}), (w_0)_t(x_{1j}, t_{1j})) \in U^{\lambda_1}, \tag{7.3}$$

$$\tau_{1j}^+ = \lambda'_1 \omega_1(r_{1j}) + (1 - \lambda'_1) \omega_2(r_{1j}) - s_{1j} > 0, \tag{7.4}$$

$$\tau_{1j}^- = s_{1j} - (1 - \lambda'_1) \omega_1(r_{1j}) - \lambda'_1 \omega_2(r_{1j}) > 0, \tag{7.5}$$

and

$$\varphi_{1j}(x, t) = \begin{cases} \nu_{1j} \varphi_{\tau_{1j}^+, \tau_{1j}^-, \delta_1} \left(\frac{1}{\nu_{1j}} (x - x_{1j}, t - t_{1j}) \right), & (x, t) \in \tilde{D}_{1j}, \\ 0, & (x, t) \in \bar{\Omega}_{t_1}^{t_2} \setminus \tilde{D}_{1j}. \end{cases} \tag{7.6}$$

Also, for each $j \in \mathbb{N}$, let $T_{1j}^1, T_{1j}^2, T_{1j}^3, T_{1j}^4 \subset \tilde{D}_{1j}$ denote the four disjoint triangular domains in each of which φ_{1j} is affine and has spatial derivative τ_{1j}^+ , and let $R_{1j}^5 \subset \tilde{D}_{1j}$ denote the rhombic domain in which φ_{1j} is affine and has spatial derivative $-\tau_{1j}^-$; then

$$|\tilde{D}_{1j}| = |T_{1j}^1 \cup T_{1j}^2 \cup T_{1j}^3 \cup T_{1j}^4 \cup R_{1j}^5|. \tag{7.7}$$

We write

$$\mathcal{F}_1 = \{\tilde{D}_{1j} \mid j \in \mathbb{N}\}, \quad \mathcal{F}_1^+ = \{T_{1j}^1, T_{1j}^2, T_{1j}^3, T_{1j}^4 \mid j \in \mathbb{N}\}, \quad \text{and} \quad \mathcal{F}_1^- = \{R_{1j}^5 \mid j \in \mathbb{N}\};$$

here, let $\{D_{1j}\}_{j \in \mathbb{N}}$, $\{T_{1j}\}_{j \in \mathbb{N}}$, and $\{R_{1j}\}_{j \in \mathbb{N}}$ be enumerations of \mathcal{F}_1 , \mathcal{F}_1^+ , and \mathcal{F}_1^- , respectively.

Since \mathcal{F}_1 is a Vitali cover of Q_1 , it follows from the definition of \mathcal{F}_1^\pm , (7.7), and $2\nu_{1j} \leq 2\gamma_1 \leq \frac{1}{2\tau}$ ($j \in \mathbb{N}$) that (a), (b), (c), (d), (e), and (f) for $i = 1$ hold.

To check the rest, define

$$\varphi_1 = \sum_{j=1}^{\infty} \varphi_{1j} \text{ on } \bar{\Omega}_{t_1}^{t_2}.$$

Also, let $N \in \mathbb{N}$ and

$$\varphi_1^N = \sum_{j=1}^N \varphi_{1j} \text{ on } \bar{\Omega}_{t_1}^{t_2}.$$

Then note from the definition of φ_{1j} ($j \in \mathbb{N}$) and (iii) in lemma 7.1 that

$$\begin{aligned} |\varphi_1^N(x, t) - \varphi_1^N(y, s)| &\leq |(x, t) - (y, s)| \max_{1 \leq j \leq N} \max\{|\tau_{1j}^+, \delta_1 \tau_{1j}^+|, \tau_{1j}^-\} \\ &\leq \sqrt{2}S_M |(x, t) - (y, s)| \end{aligned} \tag{7.8}$$

for all $(x, t), (y, s) \in \Omega_{t_1}^{t_2}$ and from $\tilde{D}_{1j} \subset \subset Q_1$ ($j \in \mathbb{N}$) that

$$\varphi_1^N = 0 \quad \text{and} \quad \nabla \varphi_1^N = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q_1. \tag{7.9}$$

Observe from (v) in lemma 7.1 that

$$\|\varphi_1^N - \varphi_1\|_{L^1(\Omega_{t_1}^{t_2})} \leq \sum_{j=N+1}^{\infty} |\tilde{D}_{1j}| \sup_{j \geq N+1} \frac{\nu_{1j} \tau_{1j}^+ \tau_{1j}^-}{\tau_{1j}^+ + \tau_{1j}^-} \delta_1 \leq S_M \sum_{j=N+1}^{\infty} |\tilde{D}_{1j}| \rightarrow 0 \tag{7.10}$$

as $N \rightarrow \infty$; thus, it follows from (7.8) that

$$|\varphi_1(x, t) - \varphi_1(y, s)| \leq \sqrt{2}S_M |(x, t) - (y, s)| \quad \forall (x, t), (y, s) \in \Omega_{t_1}^{t_2},$$

that is, $\varphi_1 \in W^{1,\infty}(\Omega_{t_1}^{t_2})$, and from (7.9) that

$$\varphi_1 = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q_1. \tag{7.11}$$

From (iii) in lemma 7.1,

$$\begin{aligned} \|\nabla\varphi_1^{N_2} - \nabla\varphi_1^{N_1}\|_{L^1(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} &\leq \sum_{j=N_1+1}^{N_2} |\tilde{D}_{1j}| \max_{N_1+1 \leq j \leq N_2} \max\{|\tau_{1j}^+, \delta_1 \tau_{1j}^+|, \tau_{1j}^-\} \\ &\leq \sqrt{2} S_M \sum_{j=N_1+1}^{N_2} |\tilde{D}_{1j}| \rightarrow 0 \end{aligned}$$

as $N_2 > N_1 \rightarrow \infty$; thus, from (7.10),

$$\nabla\varphi_1^N \rightarrow \nabla\varphi_1 \text{ in } L^1(\Omega_{t_1}^{t_2}; \mathbb{R}^2) \text{ as } N \rightarrow \infty$$

so that (7.9) implies that

$$\nabla\varphi_1 = 0 \text{ a.e. in } \Omega_{t_1}^{t_2} \setminus Q_1. \tag{7.12}$$

Next, for every $(x, t) \in \bar{\Omega}_{t_1}^{t_2}$, define

$$\psi_1(x, t) = \int_0^x \varphi_1(y, t) \, dy$$

and

$$\psi_1^N(x, t) = \int_0^x \varphi_1^N(y, t) \, dy \quad (N \in \mathbb{N}).$$

Then $\psi_1 \in W^{1,\infty}(\Omega_{t_1}^{t_2})$,

$$(\psi_1)_x = \varphi_1 \text{ in } \Omega_{t_1}^{t_2}, \tag{7.13}$$

from (iv) and (v) in lemma 7.1,

$$\begin{aligned} \|\psi_1^N - \psi_1\|_{L^1(\Omega_{t_1}^{t_2})} &= \int_{\Omega_{t_1}^{t_2}} \left| \int_0^x (\varphi_1^N(y, t) - \varphi_1(y, t)) \, dy \right| \, dx \, dt \\ &= \sum_{j=N+1}^{\infty} \int_{\tilde{D}_{1j}} \left| \int_0^x \varphi_{1j}(y, t) \, dy \right| \, dx \, dt \\ &\leq \sum_{j=N+1}^{\infty} |\tilde{D}_{1j}| \frac{\nu_{1j}^2 \delta_1^2 \tau_{1j}^+ \tau_{1j}^-}{2(\tau_{1j}^+ + \tau_{1j}^-)} \leq S_M \sum_{j=N+1}^{\infty} |\tilde{D}_{1j}| \rightarrow 0 \end{aligned} \tag{7.14}$$

as $N \rightarrow \infty$, and from (iii) in lemma 7.1,

$$\begin{aligned} \|\nabla\psi_1^{N_2} - \nabla\psi_1^{N_1}\|_{L^1(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} &= \int_{\Omega_{t_1}^{t_2}} \left| \int_0^x (\nabla\varphi_1^{N_2}(y, t) - \nabla\varphi_1^{N_1}(y, t)) dy \right| dx dt \\ &= \sum_{j=N_1+1}^{N_2} \int_{\tilde{D}_{1j}} \left| \int_0^x \nabla\varphi_{1j}(y, t) dy \right| dx dt \\ &\leq \max_{N_1+1 \leq j \leq N_2} 2\nu_{1j}\delta_1 \max\{|\tau_{1j}^+, \delta_1\tau_{1j}^+|, \tau_{1j}^-\} \sum_{j=N_1+1}^{N_2} |\tilde{D}_{1j}| \\ &\leq S_M \sum_{j=N_1+1}^{N_2} |\tilde{D}_{1j}| \rightarrow 0 \end{aligned}$$

as $N_2 > N_1 \rightarrow \infty$; thus,

$$\nabla\psi_1^N \rightarrow \nabla\psi_1 \text{ in } L^1(\Omega_{t_1}^{t_2}; \mathbb{R}^2) \text{ as } N \rightarrow \infty. \tag{7.15}$$

Note from the definition of ψ_1^N ($N \in \mathbb{N}$) and (iii) and (iv) in lemma 7.1 that for each $N \in \mathbb{N}$,

$$\psi_1^N = 0 \quad \text{and} \quad \nabla\psi_1^N = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q_1;$$

thus, letting $N \rightarrow \infty$, it follows from (7.14) and (7.15) that

$$\psi_1 = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q_1 \quad \text{and} \quad \nabla\psi_1 = 0 \text{ a.e. in } \Omega_{t_1}^{t_2} \setminus Q_1. \tag{7.16}$$

In turn, define

$$z_1 = (v_1, w_1) = z_0 + (\varphi_1, \psi_1) \text{ on } \bar{\Omega}_{t_1}^{t_2}; \tag{7.17}$$

then $z_1 \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$ as $(\varphi_1, \psi_1) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$. From (7.11) and (7.16), we have

$$z_1 = z_0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus Q_1;$$

hence, the third of (g) for $i = 1$ holds. Also, the fourth of (g) for $i = 1$ follows from (7.12), (7.16), and $Q_1 \subset Q$. The first of (h) for $i = 1$ is a consequence of (4.2) and (7.13).

From (7.2)–(7.6), (7.17), the definition of U_1^\pm , λ_1' , η_1 , and \mathcal{F}_1^\pm , and lemma 7.1, the first and second of (g) for $i = 1$ are fulfilled.

From (7.1), (7.17), and (iii) in lemma 7.1, we have

$$\|(v_1)_t - (v_0)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} = \|(\varphi_1)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \leq \delta_1 \sup_{j \in \mathbb{N}} \tau_{1j}^+ \leq \delta_1 S_M \leq \frac{\epsilon}{2^{1+1}}; \tag{7.18}$$

hence, the second of (h) for $i = 1$ holds.

From (7.1), (7.6), (7.17), $\nu_{1j} \leq \gamma_1 \leq \frac{1}{2^{1+j}}$ ($j \in \mathbb{N}$), and (v) in lemma 7.1, we have

$$\begin{aligned} \|z_1 - z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} &= \|(\varphi_1, \psi_1)\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \\ &\leq (1 + L^2)^{\frac{1}{2}} \delta_1 \sup_{j \in \mathbb{N}} \frac{\nu_{1j} \tau_{1j}^+ \tau_{1j}^-}{\tau_{1j}^+ + \tau_{1j}^-} \leq (1 + L^2)^{\frac{1}{2}} \delta_1 S_M \leq \frac{\epsilon}{2^{1+j}}; \end{aligned}$$

thus, the third of (h) for $i = 1$ is fulfilled.

Finally, to check the fourth of (h) for $i = 1$, note from the first and third of (h) for $i = 1$, (7.12), (7.16), (7.17), and (7.18) that

$$\begin{aligned} \|\nabla z_1\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} &\leq \|((v_1)_x, (w_1)_t)\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} + \|(v_1)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} + \|(w_1)_x\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\leq \|((v_0)_x, (w_0)_t)\|_{L^\infty(\Omega_{t_1}^{t_2} \setminus Q_1; \mathbb{R}^2)} + \|((v_1)_x, (w_1)_t)\|_{L^\infty(Q_1; \mathbb{R}^2)} \\ &\quad + \|(v_0)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\quad + \|(\varphi_1)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} + \|v_1\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\leq 2\|\nabla z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} + C_U + \|z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} + \frac{2}{2^{1+j}} \leq c_1 \end{aligned}$$

where $C_U := \sup_{(s,r) \in U} |(s, r)|$ and

$$c_1 := 2\|\nabla z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} + C_U + \|z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} + 1 > 0.$$

Therefore, the fourth of (h) for $i = 1$ follows.

The first step is now finished.

The second surgery: In the second step, we construct a function $z_2 = (v_2, w_2) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$, three countable collections \mathcal{F}_2 and \mathcal{F}_2^\pm , and two positive constants c_2 and c_3 satisfying (a)–(j) for $i = 2$, where the constant $c_1 > 0$ is as in the first step, and the two constants c_2 and c_3 are independent of the index i .

Let $\delta_2 \in (0, 1]$ be such that

$$\delta_2 < \min \left\{ \frac{\epsilon}{2^{2+j} \sqrt{1 + L^2} S_M}, \frac{\eta_2}{4} \right\}. \tag{7.19}$$

Since ∇z_1 is uniformly continuous in Q_2 , there exists a number $\gamma_2 \in (0, \frac{1}{2^{2+j}}]$ such that

$$(x, t), (y, s) \in Q_2, |(x, t) - (y, s)| \leq \gamma_2 \implies |\nabla z_1(x, t) - \nabla z_1(y, s)| \leq \delta_2. \tag{7.20}$$

For each $j \in \mathbb{N}$, since ∇z_1 is uniformly continuous in T_{1j} and in R_{1j} , there exists a number $\gamma_{1j} \in (0, \frac{1}{2^{2+j}}]$ such that

$$(x, t), (y, s) \in T_{1j}, |(x, t) - (y, s)| \leq \gamma_{1j} \implies |\nabla z_1(x, t) - \nabla z_1(y, s)| \leq \delta_2 \tag{7.21}$$

and that

$$(x, t), (y, s) \in R_{1j}, |(x, t) - (y, s)| \leq \gamma_{1j} \implies |\nabla z_1(x, t) - \nabla z_1(y, s)| \leq \delta_2. \tag{7.22}$$

Consider the diamond D_{δ_2} . From the Vitali covering lemma, we can choose a sequence $\{(x_{2j}, t_{2j})\}_{j \in \mathbb{N}}$ in Q_2 and a sequence $\{\nu_{2j}\}_{j \in \mathbb{N}}$ in $(0, \gamma_2]$ such that the

sequence $\{\tilde{D}_{2j}\}_{j \in \mathbb{N}}$ forms a Vitali cover of Q_2 , where $\tilde{D}_{2j} := (x_{2j}, t_{2j}) + \nu_{2j}D_{\delta_2} \subset \subset Q_2$ ($j \in \mathbb{N}$).

Let $j \in \mathbb{N}$. Also, from the Vitali covering lemma, we can choose a sequence $\{(x_{1jk}^+, t_{1jk}^+)\}_{k \in \mathbb{N}}$ in T_{1j} and a sequence $\{\nu_{1jk}^+\}_{k \in \mathbb{N}}$ in $(0, \gamma_{1j}]$ such that the sequence $\{\tilde{D}_{1jk}^+\}_{k \in \mathbb{N}}$ forms a Vitali cover of T_{1j} , where $\tilde{D}_{1jk}^+ := (x_{1jk}^+, t_{1jk}^+) + \nu_{1jk}^+D_{\delta_2} \subset \subset T_{1j}$ ($k \in \mathbb{N}$). Likewise, we can choose a sequence $\{(x_{1jk}^-, t_{1jk}^-)\}_{k \in \mathbb{N}}$ in R_{1j} and a sequence $\{\nu_{1jk}^-\}_{k \in \mathbb{N}}$ in $(0, \min\{\gamma_{1j}, \lambda_1 S_M\}]$ such that the sequence $\{\tilde{D}_{1jk}^-\}_{k \in \mathbb{N}}$ forms a Vitali cover of R_{1j} , where $\tilde{D}_{1jk}^- := (x_{1jk}^-, t_{1jk}^-) + \nu_{1jk}^-D_{\delta_2} \subset \subset R_{1j}$ ($k \in \mathbb{N}$).

For each $j \in \mathbb{N}$, let

$$(s_{2j}, r_{2j}) = ((v_1)_x(x_{2j}, t_{2j}), (w_1)_t(x_{2j}, t_{2j})) \in U^{\lambda_2} \setminus \bar{U}^{\lambda_1}, \tag{7.23}$$

$$\tau_{2j}^+ = \lambda'_2 \omega_1(r_{2j}) + (1 - \lambda'_2) \omega_2(r_{2j}) - s_{2j} > 0, \tag{7.24}$$

$$\tau_{2j}^- = s_{2j} - (1 - \lambda'_2) \omega_1(r_{2j}) - \lambda'_2 \omega_2(r_{2j}) > 0, \tag{7.25}$$

and

$$\varphi_{2j}(x, t) = \begin{cases} \nu_{2j} \varphi_{\tau_{2j}^+, \tau_{2j}^-, \delta_2} \left(\frac{1}{\nu_{2j}}(x - x_{2j}, t - t_{2j}) \right), & (x, t) \in \tilde{D}_{2j}, \\ 0, & (x, t) \in \bar{\Omega}_{t_1} \setminus \tilde{D}_{2j}. \end{cases} \tag{7.26}$$

Also, for each $j \in \mathbb{N}$, let $T_{2j}^1, T_{2j}^2, T_{2j}^3, T_{2j}^4 \subset \tilde{D}_{2j}$ denote the four disjoint triangular domains in each of which φ_{2j} is affine and has spatial derivative τ_{2j}^+ , and let $R_{2j}^5 \subset \tilde{D}_{2j}$ denote the rhombic domain in which φ_{2j} is affine and has spatial derivative $-\tau_{2j}^-$; then

$$|\tilde{D}_{2j}| = |T_{2j}^1 \cup T_{2j}^2 \cup T_{2j}^3 \cup T_{2j}^4 \cup R_{2j}^5|. \tag{7.27}$$

Let $j, k \in \mathbb{N}$, and let

$$(s_{1jk}^+, r_{1jk}^+) = ((v_1)_x(x_{1jk}^+, t_{1jk}^+), (w_1)_t(x_{1jk}^+, t_{1jk}^+)) \in U_1^+, \tag{7.28}$$

$$\tau_{1jk}^{++} = \lambda'_2 \omega_1(r_{1jk}^+) + (1 - \lambda'_2) \omega_2(r_{1jk}^+) - s_{1jk}^+ > 0, \tag{7.29}$$

$$\tau_{1jk}^{+-} = s_{1jk}^+ - (1 - \lambda'_2) \omega_1(r_{1jk}^+) - \lambda'_2 \omega_2(r_{1jk}^+) > 0, \tag{7.30}$$

and

$$\varphi_{1jk}^+(x, t) = \begin{cases} \nu_{1jk}^+ \varphi_{\tau_{1jk}^{++}, \tau_{1jk}^{+-}, \delta_2} \left(\frac{1}{\nu_{1jk}^+}(x - x_{1jk}^+, t - t_{1jk}^+) \right), & (x, t) \in \tilde{D}_{1jk}^+, \\ 0, & (x, t) \in \bar{\Omega}_{t_1} \setminus \tilde{D}_{1jk}^+. \end{cases} \tag{7.31}$$

Observe from (7.28) that

$$\begin{aligned} \frac{\tau_{1jk}^{++}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} &= \frac{\lambda'_2 \omega_1(r_{1jk}^+) + (1 - \lambda'_2) \omega_2(r_{1jk}^+) - s_{1jk}^+}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))} \\ &\leq \frac{(\lambda_1 - \lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))} = \frac{\lambda_1 - \lambda'_2}{1 - 2\lambda'_2} \end{aligned} \tag{7.32}$$

and that

$$\frac{\tau_{1jk}^{++}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \geq \frac{\lambda_2 - \lambda'_2}{1 - 2\lambda'_2}. \tag{7.33}$$

Note also that

$$\begin{aligned} \frac{\tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} &= \frac{s_{1jk}^+ - (1 - \lambda'_2)\omega_1(r_{1jk}^+) - \lambda'_2\omega_2(r_{1jk}^+)}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))} \\ &\geq \frac{(1 - \lambda_1 - \lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^+) - \omega_1(r_{1jk}^+))} = \frac{1 - \lambda_1 - \lambda'_2}{1 - 2\lambda'_2}. \end{aligned} \tag{7.34}$$

Here, we claim that

$$\frac{1 - \lambda_1 - \lambda'_2}{1 - 2\lambda'_2} > 1 - (\beta_2 - \beta_1). \tag{7.35}$$

To check this, we start from the inequality $\lambda_1 > \lambda'_2$. Since $\ell_0 < \kappa_0\ell_0 < 1$, we have $\frac{1}{\kappa_0\ell_0} - 1 > 0$ so that

$$\left(\frac{1}{\kappa_0\ell_0} - 1\right)\lambda_1 > \left(\frac{1}{\kappa_0\ell_0} - 1\right)\lambda'_2;$$

that is,

$$1 - \lambda_1 - \lambda'_2 > 1 - 2\lambda'_2 - \frac{1}{\kappa_0\ell_0}(\lambda_1 - \lambda'_2).$$

As $1 - 2\lambda'_2 > 0$, we now have

$$\frac{1 - \lambda_1 - \lambda'_2}{1 - 2\lambda'_2} > 1 - \frac{1}{\kappa_0\ell_0} \cdot \frac{\lambda_1 - \lambda'_2}{1 - 2\lambda'_2} = 1 - \frac{\alpha_1}{\kappa_0\ell_0} = 1 - (\beta_2 - \beta_1);$$

hence the claim holds.

Let $T_{1jk}^{+,1}, T_{1jk}^{+,2}, T_{1jk}^{+,3}, T_{1jk}^{+,4} \subset \tilde{D}_{1jk}^+$ denote the four disjoint triangular domains in each of which φ_{1jk}^+ is affine and has spatial derivative τ_{1jk}^{++} , and let $R_{1jk}^{+,5} \subset \tilde{D}_{1jk}^+$ denote the rhombic domain in which φ_{1jk}^+ is affine and has spatial derivative $-\tau_{1jk}^{+-}$; then

$$|\tilde{D}_{1jk}^+| = |T_{1jk}^{+,1} \cup T_{1jk}^{+,2} \cup T_{1jk}^{+,3} \cup T_{1jk}^{+,4} \cup R_{1jk}^{+,5}|. \tag{7.36}$$

Let $j, k \in \mathbb{N}$, and let

$$(s_{1jk}^-, r_{1jk}^-) = ((v_1)_x(x_{1jk}^-, t_{1jk}^-), (w_1)_t(x_{1jk}^-, t_{1jk}^-)) \in U_1^-, \tag{7.37}$$

$$\tau_{1jk}^{+-} = \lambda'_2\omega_1(r_{1jk}^-) + (1 - \lambda'_2)\omega_2(r_{1jk}^-) - s_{1jk}^- > 0, \tag{7.38}$$

$$\tau_{1jk}^{--} = s_{1jk}^- - (1 - \lambda'_2)\omega_1(r_{1jk}^-) - \lambda'_2\omega_2(r_{1jk}^-) > 0, \tag{7.39}$$

and

$$\varphi_{1jk}^-(x, t) = \begin{cases} \nu_{1jk}^- \varphi_{\tau_{1jk}^{+-}, \tau_{1jk}^{--}, \delta_2} \left(\frac{1}{\nu_{1jk}^-} (x - x_{1jk}^-, t - t_{1jk}^-) \right), & (x, t) \in \tilde{D}_{1jk}^-, \\ 0, & (x, t) \in \tilde{\Omega}_{t_1}^{t_2} \setminus \tilde{D}_{1jk}^-. \end{cases} \tag{7.40}$$

Observe from (7.37) that

$$\begin{aligned} \frac{\tau_{1jk}^-}{\tau_{1jk}^+ + \tau_{1jk}^-} &= \frac{s_{1jk}^- - (1 - \lambda'_2)\omega_1(r_{1jk}^-) - \lambda'_2\omega_2(r_{1jk}^-)}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^-) - \omega_1(r_{1jk}^-))} \\ &\leq \frac{(\lambda_1 - \lambda'_2)(\omega_2(r_{1jk}^-) - \omega_1(r_{1jk}^-))}{(1 - 2\lambda'_2)(\omega_2(r_{1jk}^-) - \omega_1(r_{1jk}^-))} = \frac{\lambda_1 - \lambda'_2}{1 - 2\lambda'_2} \end{aligned} \tag{7.41}$$

and that

$$\frac{\tau_{1jk}^-}{\tau_{1jk}^+ + \tau_{1jk}^-} \geq \frac{\lambda_2 - \lambda'_2}{1 - 2\lambda'_2}. \tag{7.42}$$

As above, we also have

$$\frac{\tau_{1jk}^-}{\tau_{1jk}^+ + \tau_{1jk}^-} \geq \frac{1 - \lambda_1 - \lambda'_2}{1 - 2\lambda'_2} > 1 - (\beta_2 - \beta_1).$$

Let $T_{1jk}^{-,1}, T_{1jk}^{-,2}, T_{1jk}^{-,3}, T_{1jk}^{-,4} \subset \tilde{D}_{1jk}^-$ denote the four disjoint triangular domains in each of which φ_{1jk}^- is affine and has spatial derivative τ_{1jk}^- , and let $R_{1jk}^{-,5} \subset \tilde{D}_{1jk}^-$ denote the rhombic domain in which φ_{1jk}^- is affine and has spatial derivative $-\tau_{1jk}^-$; then

$$|\tilde{D}_{1jk}^-| = |T_{1jk}^{-,1} \cup T_{1jk}^{-,2} \cup T_{1jk}^{-,3} \cup T_{1jk}^{-,4} \cup R_{1jk}^{-,5}|. \tag{7.43}$$

We write

$$\begin{aligned} \mathcal{F}_2 &= \{\tilde{D}_{2j} \mid j \in \mathbb{N}\} \cup \{\tilde{D}_{1jk}^+ \mid j, k \in \mathbb{N}\} \cup \{\tilde{D}_{1jk}^- \mid j, k \in \mathbb{N}\}, \\ \mathcal{F}_2^+ &= \{T_{2j}^1, T_{2j}^2, T_{2j}^3, T_{2j}^4 \mid j \in \mathbb{N}\} \cup \{T_{1jk}^{+,1}, T_{1jk}^{+,2}, T_{1jk}^{+,3}, T_{1jk}^{+,4} \mid j, k \in \mathbb{N}\} \\ &\quad \cup \{T_{1jk}^{-,1}, T_{1jk}^{-,2}, T_{1jk}^{-,3}, T_{1jk}^{-,4} \mid j, k \in \mathbb{N}\}, \end{aligned}$$

and

$$\mathcal{F}_2^- = \{R_{2j}^5 \mid j \in \mathbb{N}\} \cup \{R_{1jk}^{+,5} \mid j, k \in \mathbb{N}\} \cup \{R_{1jk}^{-,5} \mid j, k \in \mathbb{N}\};$$

here, let $\{D_{2j}\}_{j \in \mathbb{N}}, \{T_{2j}\}_{j \in \mathbb{N}}$, and $\{R_{2j}\}_{j \in \mathbb{N}}$ be enumerations of $\mathcal{F}_2, \mathcal{F}_2^+$, and \mathcal{F}_2^- , respectively.

Since \mathcal{F}_2 is a Vitali cover of $Q_1 \cup Q_2$, it follows from the definition of \mathcal{F}_2^\pm , (7.27), (7.36), (7.43), $2\nu_{2j} \leq 2\gamma_2 \leq \frac{1}{2^2}$ ($j \in \mathbb{N}$), and $2\nu_{1jk}^\pm \leq 2\gamma_{1j} \leq \frac{1}{2^2}$ ($j, k \in \mathbb{N}$) that (a), (b), (c), (d), (e), and (f) for $i = 2$ hold.

To check the rest, define

$$\varphi_2 = \sum_{j=1}^\infty \varphi_{2j} + \sum_{j=1}^\infty \sum_{k=1}^\infty \varphi_{1jk}^+ + \sum_{j=1}^\infty \sum_{k=1}^\infty \varphi_{1jk}^- \text{ on } \tilde{\Omega}_{t_1}^{t_2}; \tag{7.44}$$

then as in the first step, we can see that

$$|\varphi_2(x, t) - \varphi_2(y, s)| \leq \sqrt{2}S_M|(x, t) - (y, s)| \quad \forall (x, t), (y, s) \in \Omega_{t_1}^{t_2},$$

that is, $\varphi_2 \in W^{1,\infty}(\Omega_{t_1}^{t_2})$,

$$\varphi_2 = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus (Q_1 \cup Q_2), \tag{7.45}$$

and

$$\nabla \varphi_2 = 0 \text{ a.e. in } \Omega_{t_1}^{t_2} \setminus (Q_1 \cup Q_2). \tag{7.46}$$

Next, for every $(x, t) \in \bar{\Omega}_{t_1}^{t_2}$, define

$$\psi_2(x, t) = \int_0^x \varphi_2(y, t) dy;$$

then as in the first step, we can check that $\psi_2 \in W^{1,\infty}(\Omega_{t_1}^{t_2})$,

$$(\psi_2)_x = \varphi_2 \text{ in } \Omega_{t_2}^{t_1}, \tag{7.47}$$

$$\psi_2 = 0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus (Q_1 \cup Q_2), \quad \text{and} \quad \nabla \psi_2 = 0 \text{ a.e. in } \Omega_{t_1}^{t_2} \setminus (Q_1 \cup Q_2). \tag{7.48}$$

In turn, define

$$z_2 = (v_2, w_2) = z_1 + (\varphi_2, \psi_2) \text{ on } \bar{\Omega}_{t_1}^{t_2}; \tag{7.49}$$

then $z_2 \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$ as $(\varphi_2, \psi_2) \in W^{1,\infty}(\Omega_{t_1}^{t_2}; \mathbb{R}^2)$. From (7.45), (7.48), and the third of (g) for $i = 1$, we have

$$z_2 = z_0 \text{ on } \bar{\Omega}_{t_1}^{t_2} \setminus (Q_1 \cup Q_2);$$

hence, the third of (g) for $i = 2$ holds. Also, the fourth of (g) for $i = 2$ follows from (7.46), (7.48), and $Q_1 \cup Q_2 \subset Q$. The first of (h) for $i = 2$ is implied by (7.47) and the first of (h) for $i = 1$.

From (7.20)–(7.26), (7.28)–(7.31), (7.37)–(7.40), (7.49), the definition of U_2^\pm , λ'_2 , η_2 , and \mathcal{F}_2^\pm , and lemma 7.1, the first and second of (g) for $i = 2$ are satisfied.

From (7.19), (7.49), and (iii) in lemma 7.1, we have

$$\begin{aligned} \|(v_2)_t - (v_1)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} &= \|(\varphi_2)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\leq \delta_2 \sup_{j,k \in \mathbb{N}} \max\{\tau_{2j}^+, \tau_{1jk}^{++}, \tau_{1jk}^{--}\} \leq \delta_2 S_M \leq \frac{\epsilon}{2^{2+1}}; \end{aligned} \tag{7.50}$$

hence, the second of (h) for $i = 2$ holds.

From (7.19), (7.26), (7.31), (7.40), (7.49), $\nu_{2j} \leq \gamma_2 \leq \frac{1}{2^{2+1}}$ ($j \in \mathbb{N}$), and $\nu_{1jk}^\pm \leq \gamma_{1j} \leq \frac{1}{2^{2+1}}$ ($j, k \in \mathbb{N}$), we have

$$\begin{aligned} \|z_2 - z_1\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} &= \|(\varphi_2, \psi_2)\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \\ &\leq (1 + L^2)^{\frac{1}{2}} \delta_2 \sup_{j,k \in \mathbb{N}} \max \left\{ \frac{\nu_{2j} \tau_{2j}^+ \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-}, \frac{\nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}}, \frac{\nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{--}}{\tau_{1jk}^{-+} + \tau_{1jk}^{--}} \right\} \\ &\leq (1 + L^2)^{\frac{1}{2}} \delta_2 S_M \leq \frac{\epsilon}{2^{2+1}}; \end{aligned}$$

thus, the third of (h) for $i = 2$ is satisfied.

To check the fourth of (h) for $i = 2$, note from the first and third of (h) for $i = 2$, (7.18), (7.46), (7.48), (7.49), and (7.50) that

$$\begin{aligned} \|\nabla z_2\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} &\leq \|((v_2)_x, (w_2)_t)\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} + \|(v_2)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\quad + \|(w_2)_x\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\leq \|((v_0)_x, (w_0)_t)\|_{L^\infty(\Omega_{t_1}^{t_2} \setminus (Q_1 \cup Q_2); \mathbb{R}^2)} \\ &\quad + \|((v_2)_x, (w_2)_t)\|_{L^\infty(Q_1 \cup Q_2; \mathbb{R}^2)} \\ &\quad + \|(v_0)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} + \|(\varphi_1)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\quad + \|(\varphi_2)_t\|_{L^\infty(\Omega_{t_1}^{t_2})} + \|v_2\|_{L^\infty(\Omega_{t_1}^{t_2})} \\ &\leq 2\|\nabla z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{M}^{2 \times 2})} + C_U + \|z_0\|_{L^\infty(\Omega_{t_1}^{t_2}; \mathbb{R}^2)} \\ &\quad + 2\left(\frac{1}{2^{1+1}} + \frac{1}{2^{2+1}}\right) \leq c_1. \end{aligned}$$

Hence, the fourth of (h) for $i = 2$ is true.

We now verify (i) for $i = 2$. Note from (7.44), (7.46), (7.48), and (7.49) that

$$\begin{aligned} \int_{\Omega_{t_1}^{t_2}} |\nabla z_2 - \nabla z_1| \, dx \, dt &= \int_{\Omega_{t_1}^{t_2}} |\nabla(\varphi_2, \psi_2)| \, dx \, dt = \left(\int_{Q_1} + \int_{Q_2} \right) |\nabla(\varphi_2, \psi_2)| \, dx \, dt \\ &= \bigcup_{j,k \in \mathbb{N}} \int_{\bar{D}_{1jk}^+} |\nabla(\varphi_{1jk}^+, \psi_2)| \, dx \, dt \\ &\quad + \bigcup_{j,k \in \mathbb{N}} \int_{\bar{D}_{1jk}^-} |\nabla(\varphi_{1jk}^-, \psi_2)| \, dx \, dt \\ &\quad + \bigcup_{j \in \mathbb{N}} \int_{\bar{D}_{2j}} |\nabla(\varphi_{2j}, \psi_2)| \, dx \, dt. \end{aligned}$$

Let $j, k \in \mathbb{N}$. Then from (6.1), (6.3), (7.31), (7.32), (7.47), and (iii), (v), and (vi) in lemma 7.1,

$$\begin{aligned} \int_{\bar{D}_{1jk}^+} |\nabla(\varphi_{1jk}^+, \psi_2)| \, dx \, dt &= \int_{T_{1jk}^{+,1} \cup T_{1jk}^{+,2} \cup T_{1jk}^{+,3} \cup T_{1jk}^{+,4} \cup R_{1jk}^{+,5}} |\nabla(\varphi_{1jk}^+, \psi_2)| \, dx \, dt \\ &= \int_{T_{1jk}^{+,1} \cup T_{1jk}^{+,2} \cup T_{1jk}^{+,3} \cup T_{1jk}^{+,4}} ((1 + \delta_2^2)(\tau_{1jk}^{++})^2 \\ &\quad + (\varphi_{1jk}^+)^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} \, dx \, dt \end{aligned}$$

$$\begin{aligned}
 & + \int_{R_{1jk}^{+,5}} ((\tau_{1jk}^{+-})^2 + (\varphi_{1jk}^+)^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} dx dt \\
 & \leq \left((1 + \delta_2^2)(\tau_{1jk}^{++})^2 + \left(\frac{\delta_2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right. \\
 & \quad \left. + \left(\frac{\delta_2^2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right)^{\frac{1}{2}} \frac{\tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \\
 & \quad + \left((\tau_{1jk}^{+-})^2 + \left(\frac{\delta_2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right. \\
 & \quad \left. + \left(\frac{\delta_2^2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right)^{\frac{1}{2}} \frac{\tau_{1jk}^{++}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \\
 & = \left((1 + \delta_2^2)(\tau_{1jk}^{+-})^2 + \left(\frac{\delta_2 \nu_{1jk}^+ (\tau_{1jk}^{+-})^2}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right. \\
 & \quad \left. + \left(\frac{\delta_2^2 \nu_{1jk}^+ (\tau_{1jk}^{+-})^2}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right)^{\frac{1}{2}} \frac{\tau_{1jk}^{++}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \\
 & \quad + \left((\tau_{1jk}^{+-})^2 + \left(\frac{\delta_2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right. \\
 & \quad \left. + \left(\frac{\delta_2^2 \nu_{1jk}^+ \tau_{1jk}^{++} \tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} \right)^2 \right)^{\frac{1}{2}} \frac{\tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \\
 & \leq \left(\left(2S_M^2 + \frac{2S_M^4}{(1 - 2\lambda_2')^2 S_m^2} \right)^{\frac{1}{2}} + \sqrt{3} S_M \right) (\beta_2 - \beta_1) |\tilde{D}_{1jk}^+| \\
 & \leq \left(\sqrt{2} \left(1 + \frac{64}{49} \left(\frac{S_M}{S_m} \right)^2 \right)^{\frac{1}{2}} + \sqrt{3} \right) S_M (\beta_2 - \beta_1) |\tilde{D}_{1jk}^+|,
 \end{aligned}$$

where $S_m := \min_{[r_1, r_2]}(\omega_2 - \omega_1) > 0$. Similarly, from (6.1), (6.3), (7.31), (7.41), (7.47), and (iii), (v), and (vi) in lemma 7.1,

$$\begin{aligned}
 \int_{\tilde{D}_{1jk}^-} |\nabla(\varphi_{1jk}^-, \psi_2)| dx dt & = \int_{T_{1jk}^{-,1} \cup T_{1jk}^{-,2} \cup T_{1jk}^{-,3} \cup T_{1jk}^{-,4} \cup R_{1jk}^{-,5}} |\nabla(\varphi_{1jk}^-, \psi_2)| dx dt \\
 & = \int_{T_{1jk}^{-,1} \cup T_{1jk}^{-,2} \cup T_{1jk}^{-,3} \cup T_{1jk}^{-,4}} ((1 + \delta_2^2)(\tau_{1jk}^{--})^2 \\
 & \quad + (\varphi_{1jk}^-)^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} dx dt \\
 & \quad + \int_{R_{1jk}^{-,5}} ((\tau_{1jk}^{--})^2 + (\varphi_{1jk}^-)^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} dx dt
 \end{aligned}$$

$$\begin{aligned}
 &\leq \left((1 + \delta_2^2)(\tau_{1jk}^{-+})^2 + \left(\frac{\delta_2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \right) \\
 &\quad + \left(\frac{\delta_2^2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \frac{\tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} |\tilde{D}_{1jk}^-| \\
 &\quad + \left((\tau_{1jk}^{-})^2 + \left(\frac{\delta_2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \right) \\
 &\quad + \left(\frac{\delta_2^2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \frac{\tau_{1jk}^{-+}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} |\tilde{D}_{1jk}^-| \\
 &= \left((1 + \delta_2^2)(\tau_{1jk}^{-+})^2 + \left(\frac{\delta_2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \right) \\
 &\quad + \left(\frac{\delta_2^2 \nu_{1jk}^- \tau_{1jk}^{-+} \tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \frac{\tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} |\tilde{D}_{1jk}^-| \\
 &\quad + \left((\tau_{1jk}^{-+})^2 + \left(\frac{\delta_2 \nu_{1jk}^- (\tau_{1jk}^{-+})^2}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \right) \\
 &\quad + \left(\frac{\delta_2^2 \nu_{1jk}^- (\tau_{1jk}^{-+})^2}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} \right)^2 \frac{\tau_{1jk}^{-}}{\tau_{1jk}^{-+} + \tau_{1jk}^{-}} |\tilde{D}_{1jk}^-| \\
 &\leq \left(2S_M + \left(S_M^2 + \frac{2S_M^4}{(1 - 2\lambda_2')^2 S_m^2} \right)^{\frac{1}{2}} \right) (\beta_2 - \beta_1) |\tilde{D}_{1jk}^-| \\
 &\leq \left(2 + \left(1 + \frac{128}{49} \left(\frac{S_M}{S_m} \right)^2 \right)^{\frac{1}{2}} \right) S_M (\beta_2 - \beta_1) |\tilde{D}_{1jk}^-|.
 \end{aligned}$$

Also,

$$\begin{aligned}
 \int_{\tilde{D}_{2j}} |\nabla(\varphi_{2j}, \psi_2)| \, dx \, dt &= \int_{T_{2j}^1 \cup T_{2j}^2 \cup T_{2j}^3 \cup T_{2j}^4 \cup R_{2j}^5} |\nabla(\varphi_{2j}, \psi_2)| \, dx \, dt \\
 &= \int_{T_{2j}^1 \cup T_{2j}^2 \cup T_{2j}^3 \cup T_{2j}^4} ((1 + \delta_2^2)(\tau_{2j}^+)^2 \\
 &\quad + (\varphi_{2j})^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} \, dx \, dt \\
 &\quad + \int_{R_{2j}^5} ((\tau_{2j}^-)^2 + (\varphi_{2j})^2 + ((\psi_2)_t)^2)^{\frac{1}{2}} \, dx \, dt \\
 &\leq \left((1 + \delta_2^2)(\tau_{2j}^+)^2 + \left(\frac{\delta_2 \nu_{2j} \tau_{2j}^+ \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} \right)^2 \right) \\
 &\quad + \left(\frac{\delta_2^2 \nu_{2j} \tau_{2j}^+ \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} \right)^2 \frac{\tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} |\tilde{D}_{2j}|
 \end{aligned}$$

$$\begin{aligned}
 &+ \left((\tau_{2j}^-)^2 + \left(\frac{\delta_2 \nu_{2j} \tau_{2j}^+ \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} \right)^2 \right. \\
 &+ \left. \left(\frac{\delta_2^2 \nu_{2j} \tau_{2j}^+ \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} \right)^2 \right)^{\frac{1}{2}} \frac{\tau_{2j}^+}{\tau_{2j}^+ + \tau_{2j}^-} |\tilde{D}_{2j}| \\
 &\leq \frac{2S_M \tau_{2j}^-}{\tau_{2j}^+ + \tau_{2j}^-} |\tilde{D}_{2j}| + \frac{\sqrt{3} S_M \tau_{2j}^+}{\tau_{2j}^+ + \tau_{2j}^-} |\tilde{D}_{2j}| \leq 2S_M |\tilde{D}_{2j}|.
 \end{aligned}$$

Combining these estimates, we obtain that

$$\int_{\Omega_{t_1}^{t_2}} |\nabla z_2 - \nabla z_1| \, dx \, dt \leq c_2((\beta_2 - \beta_1)|Q_1| + |Q_2|) \leq c_2((\beta_2 - \beta_1)|Q| + |Q_2|),$$

where

$$c_2 := \frac{6S_M^2}{S_m} > 0.$$

Thus, (i) for $i = 2$ holds.

In this final stage, we show that if $i_0 = 2$, then (j) for $i = 2$ holds. So we assume $i_0 = 2$. Hence, for all $i \geq i_0 = 2$,

$$\zeta_0^\pm(s, r) = 1 \quad \forall (s, r) \in U_{i-1}^\pm. \tag{7.51}$$

Let $D \in \mathcal{F}_1$. Then from (b), (c), and (e) for $i = 1$, there are five numbers $j_1, \dots, j_5 \in \mathbb{N}$ with

$$T_{1j_1} \cup T_{1j_2} \cup T_{1j_3} \cup T_{1j_4} \cup R_{1j_5} \subset D$$

such that

$$|D| = |T_{1j_1} \cup T_{1j_2} \cup T_{1j_3} \cup T_{1j_4} \cup R_{1j_5}|.$$

We now observe that

$$\begin{aligned}
 \int_D \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt &= \int_{T_{1j_1} \cup T_{1j_2} \cup T_{1j_3} \cup T_{1j_4} \cup R_{1j_5}} \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt \\
 &= \sum_{k \in \mathbb{N}} \left(\int_{\tilde{D}_{1j_1k}^+} + \int_{\tilde{D}_{1j_2k}^+} + \int_{\tilde{D}_{1j_3k}^+} + \int_{\tilde{D}_{1j_4k}^+} \right. \\
 &\quad \left. + \int_{\tilde{D}_{1j_5k}^-} \right) \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt.
 \end{aligned}$$

Let $k \in \mathbb{N}$. Then from (7.28), (7.51), the first of (g) for $i = 2$, and (vi) in lemma 7.1,

$$\int_{\tilde{D}_{1jk}^+} \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt = \frac{\tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \geq \frac{\tau_{1jk}^{++}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+|$$

for $j = j_1, \dots, j_4$, and

$$\int_{\tilde{D}_{1j_5k}^-} \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt = \frac{\tau_{1j_5k}^{--}}{\tau_{1j_5k}^{--} + \tau_{1j_5k}^{--}} |\tilde{D}_{1j_5k}^-|.$$

Thus, from (6.2), (6.3), (7.33), and (7.42),

$$\int_D \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt \geq \frac{\lambda_2 - \lambda_2'}{1 - 2\lambda_2'} |D| \geq c_3(\beta_2 - \beta_1) |D|,$$

where $c_3 := \frac{\kappa_0 \ell_0}{36} > 0$. Likewise, one can check that

$$\int_D \zeta_0^-((v_2)_x, (w_2)_t) \, dx \, dt \geq c_3(\beta_2 - \beta_1) |D|.$$

Hence, the first inequalities in (j) for $i = 2$ hold.

Next, for $k \in \mathbb{N}$ and $j = j_1, \dots, j_4$, it follows from (7.34) and (7.35) that

$$\int_{\tilde{D}_{1jk}^+} \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt = \frac{\tau_{1jk}^{+-}}{\tau_{1jk}^{++} + \tau_{1jk}^{+-}} |\tilde{D}_{1jk}^+| \geq (1 - (\beta_2 - \beta_1)) |\tilde{D}_{1jk}^+|;$$

thus,

$$\int_D \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt \geq (1 - (\beta_2 - \beta_1)) |T_{1j_1} \cup T_{1j_2} \cup T_{1j_3} \cup T_{1j_4}|.$$

On the other hand, from (7.51) and the first of (g) for $i = 1$,

$$\int_D \zeta_0^+((v_1)_x, (w_1)_t) \, dx \, dt = |T_{1j_1} \cup T_{1j_2} \cup T_{1j_3} \cup T_{1j_4}|$$

so that

$$\int_D \zeta_0^+((v_2)_x, (w_2)_t) \, dx \, dt \geq (1 - (\beta_2 - \beta_1)) \int_D \zeta_0^+((v_1)_x, (w_1)_t) \, dx \, dt.$$

Similarly, one can check that

$$\int_D \zeta_0^-((v_2)_x, (w_2)_t) \, dx \, dt \geq (1 - (\beta_2 - \beta_1)) \int_D \zeta_0^-((v_1)_x, (w_1)_t) \, dx \, dt.$$

Hence, the second inequalities in (j) for $i = 2$ hold.

The second step is now complete.

Acknowledgements

The authors would like to thank the anonymous referees for valuable suggestions that greatly improved the presentation of the paper. H. J. Choi was supported by the National Research Foundation of Korea (grant RS-2023-00280065). S. Kim was supported by the National Research Foundation of Korea (grant NRF-2022R1F1A1063379, RS-2023-00217116) and Korea Institute for Advanced Study(KIAS) grant funded by the Korea government (MSIP).

Y. Koh was supported by the National Research Foundation of Korea (grant NRF-2022R1F1A1061968) and Korea Institute for Advanced Study(KIAS) grant funded by the Korea government (MSIP).

Ethical statement

The manuscript has not been submitted to more than one journal for simultaneous consideration. The manuscript has not been published previously.

Competing interest

None.

References

- 1 K. Anguige and C. Schmeiser. A one-dimensional model of cell diffusion and aggregation, incorporating volume filling and cell-to-cell adhesion. *J. Math. Biol.* **58** (2009), 395–427.
- 2 J. M. Ball and R. D. James. Fine phase mixtures as minimizers of energy. *Arch. Ration. Mech. Anal.* **100** (1987), 13–52.
- 3 D. Córdoba, D. Faraco and F. Gancedo. Lack of uniqueness for weak solutions of the incompressible porous media equation. *Arch. Ration. Mech. Anal.* **200** (2011), 725–746.
- 4 A. Castro, D. Córdoba and D. Faraco. Mixing solutions for the Muskat problem. *Invent. Math.* **226** (2021), 251–348.
- 5 L. Chen, K. Painter, C. Surulescu and A. Zhigun. Mathematical models for cell migration: a non-local perspective. *Phil. Trans. R. Soc. B.* **375** (2020), 20190379.
- 6 M. Chipot and D. Kinderlehrer. Equilibrium configurations of crystals. *Arch. Ration. Mech. Anal.* **103** (1988), 237–277.
- 7 C. De Lellis and L. Székelyhidi Jr.. The Euler equations as a differential inclusion. *Ann. Math. (2)*. **170** (2009), 1417–1436.
- 8 T. Hillen, K. Painter and M. Winkler. Global solvability and explicit bounds for non-local adhesion models. *Eur. J. Appl. Math.* **29** (2018), 645–684.
- 9 P. Isett. A proof of Onsager’s conjecture. *Ann. Math. (2)*. **188** (2018), 871–963.
- 10 S. Kim. Rate of convergence for one-dimensional quasilinear parabolic problem and its applications. *J. Differ. Equ.* **264** (2018), 82–97.
- 11 G. M. Lieberman. *Second order parabolic differential equations* (World Scientific Publishing Co., Inc., River Edge, NJ, 1996).
- 12 S. Müller and V. Šverák. Convex integration for Lipschitz mappings and counterexamples to regularity. *Ann. Math. (2)*. **157** (2003), 715–742.
- 13 A. Okubo. *Diffusion and ecological problems: mathematical models* (Springer-Verlag, Berlin-New York, 1980).
- 14 R. Shvydkoy. Convex integration for a class of active scalar equations. *J. Amer. Math. Soc.* **24** (2011), 1159–1174.
- 15 L. R. Taylor and R. A. J. Taylor. Aggregation, migration and population dynamics. *Nature* **265** (1977), 415–421.
- 16 P. Turchin. Population consequences of aggregative movement. *J. Anim. Ecol.* **58** (1989), 75–100.
- 17 K. Zhang. On existence of weak solutions for one-dimensional forward-backward diffusion equations. *J. Differ. Equ.* **220** (2006), 322–353.