

Short Note

Life history of the Antarctic sea star *Labidiaster annulatus* (Asteroidea: Labidiasteridae) revealed by DNA barcodingALEXIS M. JANOSIK¹, ANDREW R. MAHON¹, RUDOLF S. SCHELTEMA² and KENNETH M. HALANYCH^{1,*}¹Department of Biological Sciences, Auburn University, 101 Rouse Life Sciences Building, Auburn, AL 36849, USA²Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

*corresponding author: ken@auburn.edu

Received 25 June 2008, accepted 18 July 2008

Labidiaster annulatus, Sladen (1889) is a multi-rayed (9–50) voracious Antarctic sea star with numerous large, conspicuous crossed pedicellariae. An active and opportunistic predator, it commonly preys upon euphausiids, amphipods, and small fish in the water column (Dearborn *et al.* 1991). *Labidiaster annulatus* is distributed around the Antarctic, Kerguelen, South Orkney, South Sandwich Islands, South Georgia, and Shag Rocks, at recorded depths of 30–440 m (Fisher 1940, unpublished data).

Nothing is reported on the mode of reproduction in *Labidiaster*. Furthermore, the recognized family Labidiasteridae, composed of *Labidiaster*, *Coronaster Rathbunaster*, and *Plazaster*, is unlikely to be monophyletic, and the closest extant relative to *Labidiaster* remains unknown (Mah 2000, Foltz *et al.* 2007). In such a case larval identification by barcoding can be an important tool for examining life history (Webb *et al.* 2006). Here we use DNA barcoding techniques on partial mitochondrial 16S sequences, which serendipitously matched adults of *L. annulatus* to an unknown asteroid larvae collected along the western Antarctic Peninsula and Bransfield Strait region.

Larvae and adult specimens were collected during two five week Antarctic voyages aboard the RV *Laurence M. Gould* from 23 November–22 December 2004 and 12 May–13 June 2006 (Table I). Larval specimens were collected using a conical 75 cm plankton net and with a 250 micron mesh towed for 20 min in a slow oblique decent to a depth of *c.* 180 m and then similarly returned to the surface. Benthic samples were collected using a Blake trawl, wire dredge, or epibenthic sled. Adult voucher specimens have been deposited at The Smithsonian Institution National Museum of Natural History (USNM 1115369 and 1115370).

Individual asteroid larvae (19 bipinnaria and eight brachiolaria) were subjected to whole genome amplification using the GenomiPhi Kit following the manufacturer's recommendations (GE Healthcare) without prior DNA extraction because the protocol's first heating step lyses cells. DNA of adult specimens was extracted using the DNeasy® Tissue Kit (Qiagen). An approximately 500 bp region of the 16S gene was amplified using the 16SarL and 16SbrH primers following Palumbi (1991). Purified products were sequenced in both directions on a CEQ 8000 Genetic

Analysis System (Beckman Coulter). Novel sequences are deposited under Genbank accession numbers EU248958–EU248964. Edited sequences were compared to Genbank sequences using “blastn” (Altschul *et al.* 1990). Genetic distances (uncorrected *p*-distance values) were calculated using PAUP*4.0 (Swofford 2003). To objectively confirm that all the sequences probably represented a single species, sequences were analysed using TCS 1.21 (gaps treated as missing) to create a parsimony network with a 95% connection limit between haplotypes (Clement *et al.* 2000).

Of the 27 larvae examined, four (three bipinnaria and one brachiolaria) from the 2004 voyage showed > 99% sequence similarity to the *L. annulatus* sequence reported by Foltz *et al.* (2007; Genbank accession AY706154). All other larvae sampled were *Odontaster validus*. To confirm the result, we sequenced three adult *L. annulatus* and found uncorrected *p*-distances of > 0.378% when compared to larval samples and the Foltz *et al.* sample (representing four unique haplotypes). The parsimony network found all samples to be within a single network with a maximum distance of three nucleotide changes in the network (data not shown).

Thus, *Labidiaster annulatus* has an indirect mode of development with planktonic bipinnaria and brachiolaria larvae. The less than 0.4% uncorrected distance values recovered among *L. annulatus* individuals are considerably lower than the 5–7% interspecific mtDNA sequence divergences generally found in echinoderms (Foltz 1997, Hart *et al.* 1997, Waters & Roy 2003, Waters *et al.* 2004). Furthermore, 16S sequence data are known to be informative and variable in intraspecific studies for Antarctic marine invertebrates (Raupach & Wagele 2006, Wilson *et al.* 2007, Hunter & Halanych 2008, Mahon *et al.* 2008) as well as within asteroids (Waters *et al.* 2004).

Unfortunately, we cannot determine with certainty the morphology of the *L. annulatus* larvae. Larval samples were destroyed in data collection and no photographs could be taken of live larvae. Larvae were examined under a dissecting microscope and *L. annulatus* larvae seem to be superficially similar to those of *Odontaster*. Substantial morphological differences were not immediately obvious, and as such it is most likely that these are feeding (planktotrophic) larvae, but future research should further explore this issue.

Table I. Collection information for *Labidiaster annulatus* and *Odontaster validus*.

Species/ Station number	Stage	Depth (m)	Latitude/Longitude (S/W)	No. of samples
<i>L. annulatus</i>				
LMG 06-13	A	132	63°24.96'; 61°50.48'	1
LMG 06-14	A	132	62°56.004'; 61°28.751'	1
LMG 06-45	A	195	67°43.60'; 69°18.10'	1
LMG 04-68	L	0–180	63°28.02'; 62°23.97'	1
LMG 04-47	L	0–180	62°51.00'; 59°27.07'	3
-	A	-	60°58.08'; 55°6.85'	1 ^a
<i>O. validus</i>				
LMG 04-45	L	0–180	62°15.80'; 58°16.70'	7
LMG 04-47	L	0–180	62°51.00'; 59°27.07'	4
LMG 04-60	L	0–180	62°58.07'; 61°35.46'	5
LMG 04-68	L	0–180	63°28.02'; 62°23.97'	5
LMG 06-10	L	0–180	64°23.54'; 62°59.82'	1
LMG 06-30	L	0–180	65°50.51'; 66°59.83'	1

Stage: A = adult, L = larva.

^afrom Foltz *et al.* 2007.

Linking larval and adult forms together via DNA barcoding raises some interesting issues about *L. annulatus*. We now know that this organism has planktonic development, but the duration of larvae in the water column is unknown because length of larval development can vary greatly and there can be a long delay in settlement after reaching competence if cues for metamorphosis are lacking (e.g. Strathmann & Strathmann 2007). Moreover, studies of modes of reproduction in Antarctic marine invertebrates indicate that larvae often spend very long periods of time in the plankton (Pearse *et al.* 1991). Larvae of *L. annulatus* were present in the summer 2004 but not in May/June 2006. Although our numbers are low, the fact that more bipinnaria were found suggest that larvae were in the water column well past the May/June time frame during which we sampled. Our results suggest that the South American sister species, *Labidiaster radiosus* may also have planktonic larval development. Future efforts should expand the temporal and spatial sampling of larvae so that a better understanding of the life history of this conspicuous predatory sea star can be obtained.

Acknowledgements

We gratefully acknowledge all the help we received aboard the RV *Laurence M. Gould*. Helpful comments were provided to this manuscript by Scott Santos and Rebecca Hunter. This work was supported by a National Science Foundation grants (OPP9910164, OPP0338087, and OPP0338218) to R.S. Scheltema and K.M. Halanych. This work is AU Marine Biology Program contribution #42.

References

- ALTSCHUL, S.F., GISH, W., MILLER, W., MYERS, E.W. & LIPMAN, D.J. 1990. Basic local alignment search tool. *Journal of Molecular Biology*, **215**, 403–410.
- CLEMENT, M., POSADA, D. & CRANDALL, K.A. 2000. TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, **9**, 1657–1659.
- DEARBORN, J.H., EDWARDS, K.C. & FRATT, D.B. 1991. Diet, feeding behavior, and surface morphology of the multi-armed Antarctic sea star *Labidiaster annulatus* (Echinodermata: Asteroidea). *Marine Ecology Progress Series*, **77**, 65–84.
- FISHER, W.K. 1940. Asteroidea. *Discovery Reports*, **20**, 69–306.
- FOLTZ, D.W. 1997. Hybridization frequency is negatively correlated with divergence time of mitochondrial DNA haplotypes in a sea star (*Leptasterias spp.*) species complex. *Evolution*, **51**, 283–288.
- FOLTZ, D.W., BOLTON, M.T., KELLEY, S.P., KELLEY, B.D. & NGUYEN, A.T. 2007. Combined mitochondrial and nuclear sequences support the monophyly of forcipulatacean sea stars. *Molecular Phylogenetics and Evolution*, **43**, 627–634.
- HART, M.W., BYRNE, M. & SMITH, M.J. 1997. Molecular phylogenetic analysis of life-history evolution in Asterinid starfish. *Evolution*, **51**, 1848–1861.
- HUNTER, R.L. & HALANYCH, K.M. 2008. Evaluating connectivity in the brooding brittle star *Astrotoma agassizii* across the Drake Passage. *Journal of Heredity*, **99**, 137–148.
- MAH, C.L. 2000. Preliminary phylogeny of the forcipulatacean Asteroidea. *American Zoologist*, **40**, 375–381.
- MAHON, A.R., ARANGO, C.P. & HALANYCH, K.M. 2008. Genetic diversity of *Nymphon* (Arthropoda: Pycnogonida: Nymphonidae) along the Antarctic Peninsula with a focus on *Nymphon australe*. *Marine Biology*, **155**, 315–323.
- PALUMBI, S.R., MARTIN, A.P., ROMANO, S., McMILLAN, W.O., STICE, L. & GRABOWSKI, G. 1991. The simple fool's guide to PCR. *Special Publication of the Department of Zoology*. Honolulu, HI: University of Hawaii, 47 pp.
- PEARSE, J.S., McCLINTOCK, J.B. & BOSCH, I. 1991. Reproduction of Antarctic benthic marine invertebrates: tempos, modes and timing. *American Zoologist*, **31**, 65–80.
- RAUPACH, M.J. & WAGELE, J.W. 2006. Distinguishing cryptic species in Antarctic Asellota (Crustacea: Isopoda) - a preliminary study of mitochondrial DNA in *Acanthaspidia drygalskii*. *Antarctic Science*, **18**, 191–198.
- STRATHMANN, M.F. & STRATHMANN, R.R. 2007. An extraordinarily long larval duration of 4.5 years from hatching to metamorphosis for teleplanic veligers of *Fusitriton oregonensis*. *Biological Bulletin*, **213**, 152–159.
- SWOFFORD, D.L. 2003. *PAUP*: phylogenetic analysis using parsimony (*and other methods)*. Version 4.0. Sunderland, MA: Sinauer Associates.
- WATERS, J.M., O'LOUGHLIN, P.M. & ROY, M.S. 2004. Cladogenesis in a starfish species complex from southern Australia: evidence for vicariant speciation? *Molecular Phylogenetics and Evolution*, **32**, 236–245.
- WATERS, J.M. & ROY, M.S. 2003. Marine biogeography of southern Australia: phylogeographical structure in a temperate sea-star. *Journal of Biogeography*, **30**, 1787–1796.
- WILSON, N.G., HUNTER, R.L., LOCKHART, S.J. & HALANYCH, K.M. 2007. Multiple lineages and absence of panmixia in the Antarctic 'circumpolar' crinoid *Promachocrinus kerguelensis* from the Atlantic sector of Antarctica. *Marine Biology*, **152**, 895–904.
- WEBB, K.E., BARNES, K.A., CLARK, M.S. & BOWDEN, D.A. 2006. DNA barcoding: a molecular tool to identify Antarctic marine larvae. *Deep-Sea Research II*, **53**, 1053–1060.