

# Prevalence of obesity and associated sociodemographic and lifestyle factors in Morocco

K El Rhazi<sup>1,\*</sup>, C Nejjari<sup>1</sup>, A Zidouh<sup>2</sup>, R Bakkali<sup>2</sup>, M Berraho<sup>1</sup> and P Barberger Gateau<sup>3</sup>

<sup>1</sup>Laboratoire d'Epidémiologie, Recherche Clinique et Santé Communautaire, Faculté de Médecine et de Pharmacie, Km2.2 Route de Sidi Harazem, BP 1893, 30000 Fès, Maroc: <sup>2</sup>Association Lalla Salma de Lutte contre le Cancer, Rabat, Maroc: <sup>3</sup>Inserm, Univ Victor Segalen Bordeaux 2, Bordeaux, France

Submitted 16 December 2009; Accepted 13 May 2010; First published online 6 July 2010

## Abstract

**Objective:** In Morocco, the association between obesity/overweight and socio-demographic and lifestyle factors is poorly understood. The present study aimed to investigate this association in a representative sample of the Moroccan population aged 18 years and above.

**Design:** This is a cross-sectional study using a questionnaire including demographic, socio-economic and physical activity items. Height and weight were measured and BMI was computed. The association between obesity ( $\text{BMI} \geq 30.0 \text{ kg/m}^2$ ) or overweight ( $25.0 \leq \text{BMI} < 29.9 \text{ kg/m}^2$ ) and the other variables was analysed using multiple binomial logistic regression, separately in men and women.

**Setting:** The whole Moroccan territory.

**Subjects:** A total of 2891 subjects took part in the survey (1430 men and 1461 women).

**Results:** The prevalence of obesity was 20.9% in women and 6.0% in men ( $P < 0.0001$ ). The prevalence of overweight was 32.9% in women *v.* 26.8% in men ( $P < 0.0001$ ). In women, the risk of obesity and overweight increased with age, with the highest risk being in individuals aged 45–54 years (OR = 3.02, 95% CI 2.06, 4.44) compared to individuals <35 years old. Married women were more prone to obesity and overweight (OR = 2.42, 95% CI 1.50, 3.91) than single women. In men, the risk of obesity and overweight increased with average family income (OR = 2.62, 95% CI 1.40, 4.87 for family income  $\geq 5000$  MAD/month compared to <2000 MAD/month) and in married persons (OR = 3.75, 95% CI 1.78, 7.81) compared to single individuals.

**Conclusions:** These results contribute to target groups in whom prevention programmes could be implemented.

## Keywords

Obesity  
Sociodemographic factors  
Morocco

Obesity and overweight constitute a major public health problem and their prevalence is increasing at an alarming rate worldwide<sup>(1)</sup>. Obesity is a major risk factor for the development of certain chronic diseases and has been related to some hormone-dependent cancers<sup>(2,3)</sup>. Many environmental factors such as lifestyle and poor diet<sup>(4,5)</sup>, age patterns<sup>(6)</sup>, socio-economic factors<sup>(7)</sup> and a lack of physical activity<sup>(8)</sup> play an important role in the onset of obesity<sup>(9)</sup>. Changes in diet and inactivity patterns in transitional societies are fuelling the obesity epidemic<sup>(10,11)</sup>.

Socio-economic status (SES) as a marker for lifestyle behaviours is inversely associated with overweight and obesity in developed societies<sup>(12)</sup> and positively associated in developing countries<sup>(13)</sup>. However, the relationship between SES and obesity is complex and poorly understood in many countries.

In North African countries, obesity is a growing problem, especially as female fatness is a cultural symbol of beauty, fertility and prosperity<sup>(13,14)</sup>. These countries are undergoing a demographic transition<sup>(15)</sup> and economic

development, which have some implications for lifestyle, diet and related health conditions, especially obesity. However, there are some exceptions, such as Tunisia, where the prevalence of obesity becomes quite similar to that found in European countries, especially in women with higher SES, who prefer a slimmer body such as that in Western societies<sup>(16,17)</sup>. The aetiology of obesity in North Africa is still not well understood and few studies shed any light on its development among women<sup>(14)</sup>.

Morocco is a North African country with Mediterranean climate and a population of 30 million. During the past 50 years, the Moroccan population has been undergoing an epidemiological transition<sup>(18)</sup>. A marked shift was also documented in dietary patterns and lifestyle as part of the nutritional transition. This was associated with an increase in overweight and obesity<sup>(19)</sup>. Many studies have been interested in the prevalence of obesity in the general population and in particular groups<sup>(20–23)</sup>. In the most recent national survey in the year 2000, overweight and obesity were found to occur among 39.3% and 13.2%,

\*Corresponding author: Email elrhazikarima@gmail.com

respectively, of individuals aged 20 years and above<sup>(23)</sup> but their association with SES remains poorly understood. The objective of the present study was to estimate the prevalence of overweight and obesity in a national representative Sample of Moroccan adults and describe its relationship to SES and the lifestyle of subjects.

## Population and methods

### Study design

The present cross-sectional study was carried out in May 2008 on a national random sample of the Moroccan population aged 18 years and above. Sampling procedures were as follows: sample size was calculated on the basis of 15% risk factor prevalence, 2% precision, 95% CI and a cluster effect of 2; and sample size was estimated at 2448 and rounded to 3000 persons to compensate for people refusing to take part or being absent during the survey. The people to be surveyed were selected at random from 150 communes, in clusters of twenty persons per commune. A cluster was defined as a neighbourhood in an urban area and a locality in a rural area. One cluster was selected at random from each commune included in the survey and one person aged 20 years or above from each household of the cluster was selected at random. The total cluster selection was done proportionally to the distribution of the Moroccan population in urban and rural areas (53 and 47%, respectively)<sup>(24)</sup>. Pregnant women and bed-ridden terminally ill people, as well as the mentally ill, were considered ineligible. A person selected at random was not replaced in the case of absence or refusal to take part. Exclusion criteria included individuals unable to provide the requested information or written consent to participate in the study.

### Survey procedure

The interviewers were first trained to the survey in order to standardize the data collected and the anthropometric measurements. They were teamed up in pairs (one man and one woman) and were supervised by members of the survey planning team. The data were collected in the subjects' respective homes during a personal interview. The questionnaire included questions on demographics (age, sex, place of residence, educational level), SES (declared total monthly income of all the members of the household, profession, etc.) and physical activity, which were assessed by means of a detailed frequency questionnaire, including weekdays and weekends. Usual physical activity in work, method of going to and from work, leisure time and sports practice were considered in the questionnaire.

### Variables studied

#### Anthropometric measurements

The following measurements were performed: body weight, height, BMI ( $\text{kg}/\text{m}^2$ ). Body weight was measured

using bathroom scales, with the person wearing light clothes and no shoes. Body weight was expressed in kilograms. The bathroom scales were calibrated before and during the study. Size was measured using a height gauge with the subject standing barefoot. Size was expressed in centimetres.

BMI, corresponding to the person's weight divided by the square of the person's height ( $\text{kg}/\text{m}^2$ ), was used to define underweight ( $\text{BMI} < 18.5 \text{ kg}/\text{m}^2$ ), normal weight ( $\text{BMI} \geq 18.5$  and  $< 25.0 \text{ kg}/\text{m}^2$ ), overweight ( $\text{BMI} \geq 25.0$  and  $< 30.0 \text{ kg}/\text{m}^2$ ) and obesity ( $\text{BMI} \geq 30.0 \text{ kg}/\text{m}^2$ ) according to WHO recommendations<sup>(1)</sup>.

#### Sociodemographic and lifestyle variables

Among determinants of SES, monthly income was used as a class variable into the following categories:  $< 2000$  MAD corresponding about to the guaranteed minimum wage,  $2000$ – $4999$  MAD,  $5000$ – $9999$  MAD and  $\geq 10\,000$  MAD (equivalent currency exchange is:  $1 \text{ MAD} = 0.09 \text{ €}$ ). Given the small number of the two latter classes ( $5000$ – $9999$  and  $\geq 10\,000 \text{ MAD}/\text{month}$ ), they were grouped in the univariate analysis. Profession was categorized into five classes: active, retired, unemployed, housewife and student. We also grouped the highest level of education achieved into the following categories: university, secondary level, primary level, Koranic school or informal education and illiterates. In the multivariate analysis, educational level was grouped into three categories (illiterates, 0 school years; primary level and Koranic school or informal education,  $\leq 6$  school years; university and secondary level,  $> 6$  school years). Ages were grouped into 10-year segments ranging as follows:  $< 25$ ,  $25$ – $34$ ,  $35$ – $44$ ,  $45$ – $54$ ,  $55$ – $64$  and  $\geq 65$  years or older. Marital status was also grouped into three classes: married, single, divorced or widowed. Physical activity was defined according to WHO guidelines, which recommended that at least 30 min of regular, moderate or intense physical activity on most days reduces the risk of CVD and diabetes, colon cancer and breast cancer<sup>(1)</sup>. A person was therefore considered as active if he/she performed at least 30 min of physical activity per day. Housing group was classified according to criteria used previously<sup>(25)</sup> into five categories: luxurious, modern, new medina, old medina, poor housing or slums and rural housing.

#### Statistical analysis

Summary statistics were used to describe the study population. Results were reported as percentage and mean and standard deviation. The association between obesity and the determinant factors considered was researched through univariate analysis and then through multiple binomial logistic regression with being obese as the dependent variable in the first model (model 1) and being obese or overweight as the dependent variable in the second model (model 2). Factors associated with inclusion at  $\alpha < 10\%$  in univariate analysis were included

**Table 1** Main demographic and socio-economic characteristics according to gender

	All		Men		Women	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
<b>Age groups (years)</b>						
<25	420	14.5	176	12.3	244	16.7
25–34	663	22.9	293	20.5	370	25.3
35–44	632	21.9	270	18.9	362	24.8
45–54	573	19.8	295	20.6	278	19.0
55–64	347	12.0	216	15.1	131	9.0
≥65	256	8.9	180	12.6	76	5.2
Total	2891	100.0	1430	100.0	1461	100.0
<b>Marital status</b>						
Single	667	23.2	368	25.9	299	20.6
Married	1986	69.2	1028	72.2	958	66.1
Divorced or widowed	219	7.6	27	1.9	192	13.3
Total	2872	100.0	1423	100.0	1449	100.0
<b>Educational level</b>						
Illiterate	1250	43.4	427	30.1	823	56.4
Primary school	588	20.4	339	23.9	249	17.1
Secondary	569	19.8	342	24.1	227	15.5
University	219	7.6	143	10.1	76	5.2
Informal education or Koranic school	252	8.8	167	11.8	85	5.8
Total	2878	100.0	1418	–	1460	100.0
<b>Occupational activity</b>						
Active	1250	43.6	1053	73.8	197	13.7
Retired	173	6.0	160	11.2	13	0.9
Unemployed	212	7.4	156	10.9	56	3.9
Housewife	1114	38.9	–	–	1113	77.3
Student	117	4.1	57	4.0	60	4.2
Total	2866	100.0	1426	100.0	1439	100.0
<b>Average family income (MAD/month)</b>						
<2000	1456	52.5	819	58.6	637	46.3
2000–4999	714	25.7	389	27.8	325	23.6
5000–9999	159	5.7	88	6.3	71	5.2
≥10000	89	3.2	55	3.9	34	2.5
Don't know	355	12.8	47	3.4	308	22.4
Total	2773	100.0	1398	100.0	1375	100.0
<b>Housing</b>						
Luxurious or modern	450	15.5	234	16.3	216	14.8
New medina	416	14.4	201	14.0	215	14.7
Old medina	546	18.9	263	18.4	283	19.3
Poor housing or slums	303	10.5	145	10.1	158	10.8
Rural housing	1181	40.8	590	41.2	591	40.4
Total	2896	100.0	1433	100.0	1463	100.0

Sample of the adult Moroccan population, 2008.

in the initial multivariate logistic model. Manual descending multivariate logistic regressions were carried out. Factors associated with inclusion at  $\alpha < 5\%$  remained in the final multivariate logistic model. Dummy categorical variables were used in logistic regressions. Univariate and multivariate regression analyses were carried out separately for men and women. Adjusted OR and 95% CI are presented for both sex groups. All statistical analyses were performed using the Statistical Package for Social Sciences statistical software package version 17.0 (SPSS Inc., Chicago, IL, USA).

## Results

In total, out of the 3000 people selected, 2891 (96.5%) took part in the survey, including 1430 men (49.5%) and 1461 women (50.5%). People who are not included in the survey

had refused to participate or were absent (3.5%). The sample description according to sex is shown in Table 1.

The overall prevalence of obesity was 13.5%, but was much higher in women than in men (20.9% *v.* 6.0%,  $P < 0.0001$ ; Table 2). This prevalence was significantly higher in urban areas (14.9%) than rural areas (11.7%;  $P < 0.001$ ), but the difference was significant only in women. The overall prevalence of overweight was 29.9% (32.9% in women *v.* 26.8% in men;  $P < 0.0001$ ).

In univariate analyses (Table 2), the prevalence of obesity increased with age until 64 years in women, but in men the highest prevalence was observed between 45 and 54 years of age. The prevalence of overweight (not including obesity) increased with age until the age of 64 years in men and until the age of 54 years in women. According to educational level, the prevalence of obesity and overweight decreased from illiterate to university level in women, whereas in men the prevalence of obesity varied little with

**Table 2** BMI distribution by sociodemographic and lifestyle characteristics and gender

BMI (kg/m <sup>2</sup> )	All				Men				Women			
	n	<25.0	25.0–29.9	≥30.0	n	<25.0	25.0–29.9	≥30.0	n	<25.0	25.0–29.9	≥30.0
<b>Origin area</b>												
All	2881	56.8	29.6	13.5	1425	67.3	26.7	6.0	1456	46.6	32.5	20.9
Rural	1201	61.1	27.2	11.7	595	69.1	25.2	5.7	606	53.3	29.2	17.5
Urban	1680	53.8	31.4	14.9	830	66.0	27.8	6.1	850	41.8	35.6	23.4
P		0.0001				NS				0.0001		
<b>Age groups (years)</b>												
All	2876	56.8	29.7	13.5	1422	67.3	26.8	5.9	1454	46.6	32.5	21.0
<25	417	82.5	14.6	2.9	175	92.0	8.0	–	242	75.6	19.4	5.0
25–34	661	60.4	28.3	11.3	292	70.5	24.0	5.5	369	52.3	32.2	16.0
35–44	628	51.3	32.8	15.9	267	67.0	27.7	5.2	361	39.6	37.4	23.8
45–54	568	43.0	36.4	20.6	293	56.0	34.5	9.6	275	29.1	38.5	32.4
55–64	346	48.8	35.8	15.3	215	58.6	36.7	4.7	131	32.8	35.1	32.8
≥65	256	60.9	26.6	12.5	180	67.2	23.9	8.9	76	46.6	32.9	21.1
P		0.001				0.001				0.0001		
<b>Educational level</b>												
All	2863	56.7	29.8	13.5	1410	67.2	26.9	6.0	1453	46.6	32.6	20.9
Informal education or Koranic school	250	52.0	31.2	16.8	166	58.4	30.7	10.8	84	39.3	32.1	28.6
Illiterate	1242	51.0	32.0	16.9	422	66.4	28.0	5.7	820	43.2	34.1	22.7
Primary	587	59.5	30.0	10.6	338	68.6	26.6	4.7	249	47.0	34.5	18.5
Secondary	567	66.7	23.1	10.2	341	73.9	20.2	5.9	226	55.8	27.4	16.8
University	217	61.3	31.8	6.9	143	60.1	35.7	4.2	74	63.5	24.3	12.2
P		0.0001				0.006				0.002		
<b>Marital status</b>												
All	2857	56.7	29.7	13.6	1415	67.2	26.8	6.0	1442	46.5	32.5	21.0
Single	664	76.8	18.7	4.5	367	81.5	16.9	1.6	297	71.0	20.9	8.1
Married	1974	51.5	32.7	15.8	1021	61.9	30.6	7.5	953	40.3	35.0	24.7
Divorced or widowed	219	43.4	35.6	21.0	27	74.1	18.5	7.4	192	39.1	38.0	22.9
P		0.0001				0.0001				0.0001		
<b>Occupational activity</b>												
All	2850	56.7	29.7	13.5	1418	67.3	26.7	6.0	1432	46.3	32.7	21.0
Active	1244	62.8	28.7	8.5	1046	66.5	27.5	5.9	198	42.9	34.8	22.2
Retired	172	49.4	39.0	11.6	160	50.0	40.6	9.4	12	41.7	16.7	41.7
Unemployed	212	75.9	17.9	6.1	156	80.1	14.7	5.1	56	64.3	26.8	8.9
Housewife	1108	43.9	34.1	22.0	–	–	–	–	1108	43.9	34.1	22.0
Student	114	91.2	6.1	2.6	56	94.6	5.4	–	58	87.9	06.9	5.2
P		0.0001				0.0001				0.0001		
<b>Average family income (MAD/month)</b>												
All	2760	57.3	29.5	13.2	1391	67.4	26.5	6.0	1369	47.0	32.6	20.4
<2000	1449	60.8	27.8	11.4	814	71.5	24.1	4.4	635	47.1	32.6	20.3
2000–5000	710	53.4	32.0	14.6	387	62.3	29.7	8.0	323	42.7	34.7	22.6
≥5000	247	48.6	34.4	17.0	143	57.3	31.5	11.2	104	36.5	38.5	25.0
Don't know	354	57.1	28.2	14.7	47	70.2	27.7	2.1	307	55.0	28.3	16.6
P		0.002				0.001				0.02		
<b>Housing</b>												
All	2881	56.8	29.6	13.5	1425	67.3	26.7	6.0	1456	46.6	32.5	20.9
Luxurious or modern	449	50.1	37.0	12.9	234	56.8	37.2	6.0	215	42.8	36.7	20.5
New medina	415	52.0	28.9	19.0	201	66.2	26.4	7.5	214	38.8	31.3	29.9
Old medina	543	54.3	30.4	15.3	262	70.2	23.7	6.1	281	39.5	36.7	23.8
Poor housing or slums	299	60.2	28.8	11.0	143	71.3	23.8	4.9	156	50.0	33.3	16.7
Rural housing	1175	61.4	27.0	11.7	585	69.6	24.8	5.6	590	53.2	29.2	17.6
P		0.0001				0.02				0.0001		
<b>Physical activity (≥30 min/d)</b>												
All	2620	58.4	29.2	12.4	1359	67.8	26.3	5.9	1261	48.2	32.4	19.4
Yes	1606	61.2	28.4	10.4	918	70.0	24.8	5.1	688	49.4	33.1	17.4
No	1014	53.8	30.6	15.6	441	63.0	29.5	7.5	573	46.8	31.4	21.8
P		0.0001				0.02				0.15		

Sample of adult Moroccan population 2008.  
P for global  $\chi^2$  tests.

educational level. The prevalence of overweight increased up to primary level of schooling in women. In men, overweight was much more frequent in those with a university level of education and in persons attending only informal education or Koranic school. Married women and men had the highest prevalence of obesity. In both

genders, single persons had the lowest prevalence of both overweight and obesity.

Within the five occupational groups and according to sex, retired subjects were more often obese or overweight than those with other occupational activities (total prevalence 50.0% and 58.4% in men and women, respectively), which

**Table 3** Adjusted OR for obesity or overweight in women

	Model 1			Model 2		
	OR	95% CI	<i>P</i>	OR	95% CI	<i>P</i>
Age groups (years)						
<35	1.00	–	<0.0001	1.00	–	<0.0001
35–44	1.93	1.33, 2.78		1.80	1.35, 2.40	
45–54	3.02	2.06, 4.44		2.67	1.90, 3.75	
≥55	2.85	1.83, 4.44		1.94	1.32, 2.86	
Marital status						
Single	1.00	–	0.0001	1.00	–	0.0001
Married	2.42	1.50, 3.91		2.45	1.77, 3.38	
Divorced or widowed	1.58	0.86, 2.88		2.00	1.27, 3.16	
Housing						
Rural housing	1.00	–	0.0001	1.00	–	0.0001
Luxurious or modern	1.16	0.77, 1.75		1.99	1.35, 2.91	
New medina	2.31	1.58, 3.37		2.59	1.79, 3.74	
Old medina	1.57	1.09, 2.24		2.09	1.51, 2.90	
Poor housing or slums	0.99	0.62, 1.62		1.38	0.94, 2.03	
Educational level						
Illiterate	–	–		1.00	–	0.0120
Primary or informal education or Koranic school	–	–		1.04	0.77, 1.41	
Secondary or university	–	–		0.63	0.44, 0.90	

Sample of adult Moroccan population, 2008 (*n* 1440).

Model 1: obesity as a dependent variable.

Model 2: obesity or overweight as a dependent variable.

may be explained by their older age: 83.7% of the retired subjects were 55 years old and above *v.* 16.2% of the active persons ( $P < 0.0001$ ; Table 2).

A positive relationship was found between the prevalence of obesity and total family income in both genders; as income of the family increased, the prevalence of obesity increased.

According to housing, the prevalence of obesity and overweight increased with the quality of housing. A higher prevalence was observed among individuals who lived in new or old medina or in luxurious or modern houses than in those who lived in poor housing or in slums. The same trend was observed in the whole population and separately in men and women.

The number of persons living in the same house, as an indirect indicator of SES, was not different between obese (mean 5.9; SD 3.0) and non-obese persons (mean 6.1; SD 3.0). The same results were found separately in men (mean 6.2; SD 3.0) and in women (mean 5.7; SD 2.9) and also when we adjusted for the type of housing or on average family income.

The prevalence of obesity appeared to be lower among subjects who practised at least 30 min of physical activity per day than in the others (10.4% *v.* 15.6%,  $P < 0.0001$ ). The same trend was observed separately in men and in women even if it was not significant in women. But this association disappeared after adjusting for average family income and for the other socio-economic factors.

After correction for possible confounding factors including all demographic and socio-economic factors in multivariate regression models separately in men and women, the retained variables in the final manual descending

multivariate logistic regression models differed between men and women. However, in women as in men, physical activity did not remain a significant correlate of obesity or overweight in the multivariate models.

In women, physical activity and occupation were no longer associated with obesity in the final model (Table 3). Age remained strongly associated with obesity with the highest risk in individuals aged 45–54 years compared to those aged <35 years. According to marital status, married women remained more prone to obesity than single and divorced or widowed women. However, both married and divorced or widowed women were at higher risk of overweight including obesity. Housing was another independent correlate of obesity with a higher risk in women living in new medina. Similar results were observed in the estimation of adjusted OR for obesity or overweight with, in addition to the previous factors, a significantly decreased risk in women with a higher educational level.

Given the small number of divorced or widowed men (*n* 27) this category was grouped with the singles in multivariate analyses (Table 4). Age, education and occupation were no longer associated with obesity in men in the final multivariate model. Married men were more prone to obesity than single, divorced or widowed individuals. Average family income was a strong correlate with increasing risk with higher family income, the highest risk being associated with an income of ≥5000 MAD/month. When the dependent variable was 'obesity or overweight', age group and occupation were both independently associated with obesity or overweight with the highest risk in retired men. Conversely, risk was significantly decreased in unemployed men.

**Table 4** Adjusted OR for obesity or overweight in men

	Model 1			Model 2		
	OR	95 % CI	<i>P</i>	OR	95 % CI	<i>P</i>
Age groups (years)						
<35	–	–		1.00	–	0.010
35–44	–	–		1.24	0.83, 1.85	
45–54	–	–		1.91	1.28, 2.85	
≥55	–	–		1.34	0.87, 2.05	
Marital status						
Single or divorced or widowed	1.00	–	0.0001	1.00	–	0.009
Married	3.75	1.78, 7.81		1.66	1.13, 2.43	
Average family income (MAD/month)						
<2000	1.00	–	0.002	1.00	–	0.001
2000–4999	2.02	1.23, 3.33		1.57	1.20, 2.06	
≥5000	2.62	1.40, 4.87		1.67	1.14, 2.43	
Profession						
Active or student	–	–		1.00	–	0.001
Retired	–	–		1.67	1.12, 2.50	
Unemployment	–	–		0.53	0.32, 0.88	

Sample of adult Moroccan population, 2008 (*n* 1325).

Model 1: obesity as a dependent variable.

Model 2: obesity or overweight as a dependent variable.

## Discussion

In the present study, we investigated the relationship between obesity and socio-economic factors within a sample from the Moroccan population aged 18 years and above. The overall prevalence of obesity and overweight was 13.5% and 29.9%, respectively, and higher in women and urban areas compared to men and rural areas. The prevalence of obesity in Morocco has increased from 1.6% in 1984 to 4.3% in 1998 in men and from 6.4% in 1984 to 16.0% in 1998 in women aged 20 years and above. In addition, it increased from 5.5% in 1984 to 12.2% in 1998 in the urban population and from 2.6% in 1984 to 7.4% in 1998 in the rural population<sup>(19)</sup>.

Gender difference in the prevalence of obesity was found to be highly significant in our population and there are similar findings in previous studies in Morocco<sup>(23)</sup> and in other countries<sup>(26,27)</sup>.

Age was strongly associated with obesity in women, as previously reported<sup>(28)</sup>. This association between obesity and age can be explained, in part, by parity and post-menopausal status<sup>(29)</sup>. The reduced risk of obesity in younger women may reflect a possible shift in the burden of obesity in women, from the positive association observed in most studies from sub-Saharan Africa<sup>(20,21)</sup> to the inverse association reported from developed countries. Therefore, younger women, who are more likely to be educated, adopt lifestyles that are less prone to obesity in response to their exposure to the cultures of more developed countries.

Obesity was inversely associated with educational level in our study, which remained after adjustment for confounding variables in women. This may be due to a healthier lifestyle in individuals who have been educated to a higher level and is in accordance with other recent studies<sup>(30)</sup>.

Several studies have underlined the importance of socio-economic factors and physical activity<sup>(31,32)</sup>. Our finding showed that family income was strongly and positively associated with obesity, especially in men. In women, a strong association between housing and obesity or overweight was observed. Obesity and overweight were more common among the women living in luxurious or modern housing and in new or old medina. This may be explained by lifestyle factors including level of physical activity. Women living in more comfortable houses and urban areas probably spend less time walking and doing heavy housework than those living in poor housing or in rural areas. Similarly, higher income may provide access to more abundant dietary resources and lifestyle choices. Therefore, wealthier men may be consuming larger portions of food because they can afford to do so. Concerning physical activity, our study showed that it decreased with the average family income (73.1% in <2000, 67.5% in 2000–4999 and 44.9% in ≥5000 MAD/month; *P* < 0.0001). Therefore, the wealthier men walked less often because they owned and used cars frequently. Men with higher income did not do chores that were probably physically demanding because they could afford to employ others. Therefore, they were probably expending less energy. They were also spending greater time at work, which was often sedentary, and hence had less time to participate in leisure-time physical activity. People in the highest socio-economic classes were more prone to adopting a Western way of life, including more meals taken in restaurants or fast foods, and more time spent watching television or working on computers. However, physical activity was not significantly correlated with obesity or overweight in the multivariate models, perhaps because of a lack of accuracy of its measurement in two classes only. As in other studies<sup>(33,34)</sup>, we evidenced a significant difference in the

prevalence of obesity with marital status in univariate analysis, with more obesity in married, widowed and divorced individuals compared to single persons. This association remained significant in the multivariate analysis in both genders. This may be explained by access to more abundant food and rich dishes. Furthermore, married, widowed or divorced persons were older than the single persons (24.7% and 47.0% among married and widowed or divorced, respectively, were 55 years old and above *v.* 1.1% among single persons;  $P < 0.0001$ ). Moreover, it is very likely that the cultural preference for female plumpness coupled with physical and cultural barriers to physical activity makes the difference<sup>(20)</sup>.

Housewives and retired persons were the most frequently obese group in our study compared to those who were employed, unemployed and students. These results are concordant with the results of previous studies<sup>(26)</sup>. But this association disappeared after correction for confounding variables, in particular older age.

The main limitation of the present study is its cross-sectional design. We cannot therefore demonstrate causality of the associations with the factors evidenced in the present study. The second limitation was a recall bias, especially for income and education level that might be over- or underestimated because data were collected by direct interview. Furthermore, given the lack of data on number of pregnancies and engagement in sport in the present study, we could not ascertain the impact of these variables on overweight and obesity. In particular, a more precise evaluation of the time spent on various professional, household and recreational activities would have allowed a better estimation of physical activity. However, the strength of this nationwide study was that it was representative of the whole Moroccan population since the prevalence of sociodemographic factors was similar to that reported in the last general census of habitat and population in 2004<sup>(24)</sup>. Our findings are in accordance with two previous studies in Morocco, one with a sociological approach of Moroccan women's household roles in relation to obesity<sup>(35)</sup> and the other one conducted in a specific group of Sahraoui women<sup>(36)</sup>. Another strength was the direct measurement of anthropometric data.

In conclusion, the present study contributes to understanding the relationship between the socio-economic factors and obesity and overweight in the Moroccan population. Higher income and better living conditions associated with higher SES are the main risk factors of overweight and obesity. Recognizing these factors may facilitate planning of preventive programmes for reducing obesity in the targeted groups in this society.

### Acknowledgements

The present study was funded by 'Association Lalla Salma de Lutte contre le Cancer – Maroc'. The authors have no

conflict of interest. K.E.R. contributed to the conception and design, acquisition of data, analysis and interpretation of data and was involved in drafting the manuscript; C.N. contributed to the conception and design, interpretation of data, was involved in revising the manuscript critically and gave final approval of the version to be published. A.Z. contributed to the conception and design of data; R.B. contributed to the conception and design of data. M.B. made substantial contributions to conception and design of data; P.B.G. was involved in drafting the manuscript and gave final approval of the version to be published. Special thanks to Dr T. Norat (Department of Epidemiology and Public Health, Imperial College London, UK) for reviewing the manuscript.

### References

1. World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation on Obesity. WHO Technical Report Series no. 894*. Geneva: WHO.
2. Pi-Sunyer FX (1993) Medical hazards of obesity. *Ann Intern Med* **119**, 655–660.
3. Khan LK & Bowman BA (1999) Obesity: a major global public health problem. *Am Rev Nutr* **19**, 13–17.
4. Chen M, You Y, Zhao W *et al.* (2002) Influences of diet and nutrition on obesity of pre-school children. *Wei Sheng Yan Jiu* **31**, 370–372.
5. Shepard TY, Weil KM, Sharp TA *et al.* (2001) Occasional physical inactivity combined with a high-fat diet may be important in the development and maintenance of obesity in human subjects. *Am J Clin Nutr* **73**, 703–708.
6. Bartali B, Benvenuti E, Corsi AM *et al.* (2002) Changes in anthropometric measures in men and women across the life-span: findings from the InCHIANTI study. *Soz Präventivmed* **47**, 336–348.
7. Heitmann BL (1999) Occurrence and development of overweight and obesity among adult Danes aged 30–60 years. *Ugeskr Laeger* **161**, 4380–4384.
8. Lindstrom M, Isacson SO & Merlo J (2003) Increasing prevalence of overweight, obesity and physical inactivity: two population based studies 1986 and 1994. *Eur J Public Health* **13**, 306–312.
9. Bouchard C (1996) Genetics of obesity in humans: current issues. In *The Origin and Consequences of Obesity, CIBA Foundation Symposium*, pp. 108–117 [DJ Chadwick and GC Cardew, editors]. Chichester: Wiley.
10. Popkin BM (2001) The nutrition transition and obesity in the developing world. *J Nutr* **131**, 871S–873S.
11. Popkin BM & Gordon-Larsen P (2004) The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord* **28**, Suppl. 3, S2–S9.
12. McLaren L (2007) Socioeconomic status and obesity. *Epidemiol Rev* **29**, 29–48.
13. Monteiro CA, Moura EC, Conde WL *et al.* (2004) Socio-economic status and obesity in adult populations of developing countries: a review. *Bull World Health Organ* **82**, 940–946.
14. Mokhtar N, Elati J, Chabir R *et al.* (2001) Diet culture and obesity in northern Africa. *J Nutr* **131**, 887S–892S.
15. Popkin BM (1993) Nutritional patterns and transitions. *Popul Dev Rev* **19**, 138–157.
16. Kamoun M, Hajem S, Imen S *et al.* (2008) Prevalence of obesity and overweight in Tunisia on 2001. *Tunis Med* **86**, 649–652.

17. Tlili F, Mahjoub A, Lefèvre P *et al.* (2008) Tunisian women's perceptions of desirable body size and chronic disease risk. *Ecol Food Nutr* **47**, 399–414.
18. Jroundi I (2007) Epidemiological transition in Morocco in the past 50 years. *Epidemiology* **18**, S12.
19. Benjelloun S (2002) Nutrition transition in Morocco. *Public Health Nutr* **5**, 135–140.
20. Rguibi M & Belahsen R (2007) Prevalence of obesity in Morocco. *Obes Rev* **8**, 11–13.
21. Lahmam A, Baali A, Hilali MK *et al.* (2008) Obesity, overweight and body-weight perception in a High Atlas Moroccan population. *Obes Rev* **9**, 93–99.
22. Belahsen R, Mziwira M & Fertat F (2004) Anthropometry of women of childbearing age in Morocco: body composition and prevalence of overweight and obesity. *Public Health Nutr* **7**, 523–530.
23. Tazi MA, Abir-Khalil S, Chaouki N *et al.* (2003) Prevalence of the main cardiovascular risk factors in Morocco: results of a national survey 2000. *J Hypertens* **21**, 897–903.
24. Haut Commissariat au Plan (2004) Recensement Général de la Population et de l'Habitat 2004. [http://www.hca.ma/pubData/Demographie/RGPH/RGPH\\_Rapport\\_National.pdf](http://www.hca.ma/pubData/Demographie/RGPH/RGPH_Rapport_National.pdf) (accessed June 2010).
25. Direction de la planification et des ressources financières, Division de la planification et des études, Service des études et de l'information sanitaire (2003) Enquête sur la population et la Santé Familiale –2003. Protocole <http://doc.abhatoo.net.ma/doc/IMG/pdf/Protocole-PAPFAM-web.pdf> (accessed June 2010).
26. Hajian-Tilaki KO & Heidari B (2007) Prevalence of obesity, central obesity and the associated factors in urban population aged 20–70 years, in the north of Iran: a population-based study and regression approach. *Obes Rev* **8**, 3–10.
27. Xu F, Yin XM, Zhang M *et al.* (2004) Family average income and body mass index above the healthy weight range among urban and rural residents in regional Mainland China. *Public Health Nutr* **8**, 47–51.
28. Peixoto MD, Benício MH & Jardim PC (2007) The relationship between body mass index and lifestyle in a Brazilian adult population: a cross-sectional survey. *Cad Saude Publica* **23**, 2694–2740.
29. Tufano A, Marzo P, Enrini R *et al.* (2004) Anthropometric, hormonal and biochemical differences in lean and obese women before and after menopause. *J Endocrinol Invest* **27**, 648–653.
30. Hajian-Tilaki KO & Heidari B (2010) Association of educational level with risk of obesity and abdominal obesity in Iranian adults. *J Public Health (Oxf)* **32**, 202–209.
31. Martinez JA, Kearney JM, Kafatos A *et al.* (1999) Variables independently associated with self-reported obesity in the European Union. *Public Health Nutr* **2**, 125–133.
32. Aal-Isa AN (1999) Dietary and socio-economic factors associated with obesity among Kuwaiti college men. *Br J Nutr* **82**, 369–374.
33. Kahn HS, Williamson DF & Stevens JA (1991) Race and weight change in US women: the roles of socioeconomic and marital status. *Am J Public Health* **81**, 319–323.
34. Sobal J, Rauschenbach B & Frongillo EA (2003) Marital status changes and body weight changes: a US longitudinal analysis. *Soc Sci Med* **56**, 1543–1555.
35. Batnitzky A (2008) Obesity and household roles: gender and social class in Morocco. *Sociol Health Illn* **30**, 445–462.
36. Rguibi M & Belahsen R (2006) Body size preferences and sociocultural influences on attitudes towards obesity among Moroccan Sahraoui women. *Body Image* **3**, 395–400.