

## Review Paper

†This article is dedicated to the great academician Nikolay Ivanovich Vavilov (25 November 1887 to 26 January 1943), who had envisioned a hunger-free world but, ironically, died of hunger and malnutrition.

**Cite this article:** Rangan P (2020). Endosperm variability: from endoreduplication within a seed to higher ploidy across species, and its competence. *Seed Science Research* 30, 173–185. <https://doi.org/10.1017/S0960258520000148>

Received: 3 October 2019

Revised: 23 April 2020

Accepted: 22 May 2020

First published online: 17 June 2020

**Key words:**

apomixis; cereals; dosage level; endoreduplication; endosperm variability; plant genetic resources; single-cell sequencing

**Author for Correspondence:**

Parimalan Rangan, E-mail: [r.parimalan@icar.gov.in](mailto:r.parimalan@icar.gov.in); [p.rangan@uq.edu.au](mailto:p.rangan@uq.edu.au)

# Endosperm variability: from endoreduplication within a seed to higher ploidy across species, and its competence<sup>†</sup>

Parimalan Rangan<sup>1,2</sup> 

<sup>1</sup>Division of Genomic Resources, ICAR-National Bureau of Plant Genetic Resources, New Delhi 110012, India and

<sup>2</sup>Queensland Alliance for Agriculture and Food Innovation, University of Queensland, St Lucia, QLD 4072, Australia

**Abstract**

Endosperm tissue that nourishes the embryo during seed development, upon maturity, nourishes the global population with special reference to cereal crops like maize, wheat and rice. In about 70% of the angiosperms, endosperm genome content is '3n' with 2:1 (maternal:paternal) contribution, as a result of the second fertilization event. However, angiosperms evolution also documents diversity in endosperm genome content from '2n' to '15n', in scale with the corresponding maternal genome dosage variability ('1n' to '14n'), whereas paternal contribution is invariable. In apomicts, due to lack of fertilization, or pseudogamy (fertilization of the central cell for endosperm formation), endosperm genome dosage (m:p) has been reported to range between 1:1 and 8:3. Exceptionally, the central cell with one unreduced nucleus and fused with a reduced sperm cell, with 2:1 normal genome dosage, has been reported in *Panicum*. Altered genome dosage levels are reportedly correlative with eccentricities among maternal and paternal contribution to seed resource allocation. Besides endosperm ploidy variability between species of angiosperms, the present review gives an overview of the ploidy variability in endosperm cells within a seed, up to '690n'. In addition to genome-scale variability in the endosperm, some taxa of angiosperms exhibit chlorophyllous endosperms and some chlorophyllous embryos. Also, endosperm cell number during seed development is reported to have a strong association with grain weight at maturity. Genes underlying these traits of variability are unknown, and the present review underscores the variability and highlights the potential of the single-cell sequencing techniques towards understanding the genetic mechanisms associated with these variable traits.

**Introduction**

Endosperm is not just an embryo-nourishing tissue; being the major component of the cereal grain, it nourishes the global population. From an evolutionary perspective, the origin of endosperm (resultant of the second fertilization event) in angiosperms is predicted to have evolved from multiple fertilization events in gymnosperms, leading to polyembryony (Friedman, 1992). In 70% of the angiosperms genome dosage level of the embryo is '2n' with equal genetic contribution from the female (m, maternal) and male (p, paternal) parent, while in the endosperm, it is '3n' (2m:1p) (Raghavan, 2006). Diversity in endosperm ploidy level of angiosperms broadly ranges from 2n to 15n, especially due to the variability in the genetic contribution of the female parent from 1n to 14n (Friedman et al., 2008). Key genes regulating the cell cycle, mega-sporogenesis or -gametogenesis or endosperm developmental process, including ontogeny and cell numbers are well documented (Motamayor et al., 2000; Berger et al., 2006; Huh et al., 2007; Fiume and Fletcher, 2012; Batista et al., 2019; Huang et al., 2019; Kirkbride et al., 2019). However, the variability underlying these traits between species and its regulatory mechanism that flexibly accommodates and accepts the variable endosperm dosage level from 2n to 15n, and endosperm cell number in angiosperms is not yet fully understood. The present review highlights the endosperm variability patterns in angiosperms, from variable genome dosage to chlorophyllous endosperm involving carbon fixation; and the potentialities of the genomics tools and techniques to underscore the genetic mechanism for better scientific understanding.

**Endosperm variability and development**

Endosperm is a tissue in transition across dicot families, while it is the major source of reserves in monocots. However, early endosperm development is remarkably conserved across monocots and dicots (Becraft, 2001). Endosperm development is a dynamic and intricate process involving genome imprinting, dosage balance, positional cues, cell-cycle modifications and programmed cell death (PCD) for a normal ontogeny (Wangenheim, 1957; Becraft, 2001;

Gehring and Satyaki, 2017; Satyaki and Gehring, 2019). In monocots, where the endosperm is persistent, it is comprised of three cell types, namely starchy endosperm, aleurone layer and transfer cell layers; among these, the aleurone layer retains its viability at seed maturity (Lopes and Larkins, 1993; Becraft, 2001; Becraft and Gutierrez-Marcos, 2012). In addition to cell-type variability, the endosperm exhibits variability of *inter alia* ploidy (both across species and within the seed, as well), chlorophyll and polyendospermy.

### Endosperm ploidy variability in sexual and apomictic species

Around 70% of the angiosperms exhibit an endosperm ploidy level of '3n' with two-thirds of the genome dosage contribution from the maternal parent, and the remaining one-third through the paternal parent due to the second fertilization event (Raghavan, 2006). The maternal to paternal genome ratio of 2:1 is the most balanced dosage levels among parental genomes during sexual reproduction and is postulated as endosperm balance number (EBN) for normal endosperm and seed development (Johnston et al., 1980). However, in several angiosperms, endosperm ploidy deviates from '3n'. A central cell with 1 to 14 nuclei is the source of variability for the endosperm ploidy level, resulting in a genome dosage level of 2n to 15n upon the second fertilization event, reflecting altered maternal dosage levels, ranging from 1m:1p to 14m:1p (Friedman et al., 2008). Genera exhibiting variable endosperm ploidy level due to altered genome dosage levels of maternal parents include *Oenothera* (Von Wangenheim, 1962; Haig and Westoby, 1991), *Nuphar* (Williams and Friedman, 2002), *Manekia* (Arias and Williams, 2008), *Gagea* (Greilhuber et al., 2000), *Penea*, *Plumbago*, *Fritillaria*, *Plumbagella* and *Peperomia* (Friedman et al., 2008). Invariably, in all these species with seed development through a fertilization event, the paternal genome dosage contribution is uniformly '1n' only.

However, in aposporous apomicts, the paternal genome dosage level could be higher than '1n', at instances where the paternal ploidy level is higher than the maternal one (Haig and Westoby, 1991; Quarin, 1999; Alves et al., 2001; Šarhanová et al., 2012). In diplosporous apomicts, the maternal dosage alone is inherited with 4n (4m:0p) genome content without the fertilization event (Kollmann et al., 2000). In certain diplosporous apomictic grass species, such as *Elymus*, *Poa*, *Eragrostis* and *Tripsacum*, pollination is a necessary event for successful endosperm development – one of the three components of apomictic seed development (Bashaw and Hanna, 1990; Kaushal et al., 2019). For example, *Tripsacum* with diploids and tetraploids reported to exhibit genome dosage (maternal to paternal) of 8:1 and 8:2 ratios in apomictic plants, but 2:1, 4:1 and 4:2 ratios observed in sexual plants (Grimanelli et al., 1997). *Paspalum* with di-, tri-, tetra-, penta-, hexa- and octaploid species, the endosperm genome dosage level varies widely from 1:1, 2:1, 3:1, 4:1, 4:3, 8:1 to 8:3, with 4:1 genome dosage exhibiting the maximum reproductive efficiency (Burton, 1948; Quarin et al., 1984; Quarin, 1999; Ortiz et al., 2013; Felitti et al., 2015). An AFLP Marker associated with apomictic trait loci in *Paspalum* has been identified (Labombarda et al., 2002), its cross-transferability in other related species might be of advantage in differentiating the apomictic from sexual plants. Exceptionally, the occurrence of pseudogamy resulting in the normal EBN genome dosage level of 2:1 is possible only when the reduced sperm cell is fused with the central cell containing a single unreduced polar nucleus as reported in *Panicum maximum* (Warmke, 1954). However, in general, due

to pseudogamy, 4:1 is formed when the two unreduced polar nuclei are fused with one reduced sperm nucleus (Grimanelli et al., 1997; Quarin, 1999; Felitti et al., 2015). Polyploidy and apomixis are reported to be strongly associated, although gametophytic apomixis is reported in diploids (Sharbel et al., 2009; Ortiz et al., 2013).

The occurrence of apomixis (gametophytic: apospory and diplospory; sporophytic: adventitious embryony type) is documented in at least 300 species, predominantly from four families, namely Gramineae, Compositae, Rosaceae and Rutaceae, but for more than 35 families in total (Koltunow, 1993; Khush, 1994). As a rule of thumb, gametophytic apomicts (apospory – *Cenchrus*, *Dichanthium*, *Panicum* and *Heracium*; diplospory – *Taraxacum*, *Ixeris* and *Antennaria*) are polyploid in nature, while the sporophytic ones (adventitious embryony – *Citrus*) exhibit diploid behaviour (Knox, 1967; Young et al., 1979; Asker and Jerling, 1992).

From an evolutionary perspective, polyploidization and hybridization events are the drivers of genetic modifications at the genome level for apomixis development (Barke et al., 2018). Photoperiod (natural) induced apomixis in *Dichanthium* was the first report documenting the role of the environment, in addition to genetic factors, leading to apomixis (Knox, 1967; Rodrigo et al., 2017). Moreover, experimentally induced apomixis through gamma-ray irradiation has also been documented for maize (Yudin, 1966). Chemically induced apomixis has employed *inter alia* nitrous oxide, dimethyl sulfoxide, gibberellic acid, 6-benzyl aminopurine, 2,4-dichlorophenoxy acetic acid and zeatin in *Datura*, *Solanum*, *Ficus*, *Gossypium* and *Zea* (Montezuma-de-Carvalho, 1967; Arendt, 1970; Zhou, 1980; Hu et al., 1991).

Facultative apomixis – coexistence of apomixis and sexual reproduction – have been demonstrated in *Hieracium* and *Sorghum* (Tang, 1977; Bicknell and Koltunow, 2004; Carman et al., 2011). Probably due to the apomictic nature of *Hieracium*, Gregor Mendel in 1869 obtained contrasting results as compared to *Pisum*, which he noted as 'almost opposed behaviour' at a time when the phenomenon of apomixis was unknown (Savidan, 2000; Bicknell and Koltunow, 2004). Inheritance of aposporous apomixis has been studied in *Pennisetum*, *Panicum* and *Brachiaria* and reported to be determined by a single dominant locus (Sherwood et al., 1994; Valle et al., 1994; Savidan, 2000). Genes (protein coding and lncRNA) and epigenetic mechanisms regulating apomixis are well documented (Guerin et al., 2000; Albertini et al., 2005; Laspina et al., 2008; Garcia-Aguilar et al., 2010; Polegri et al., 2010; Ortiz et al., 2013; Hand and Koltunow, 2014; Podio et al., 2014a,b; Felitti et al., 2015; Ortiz et al., 2017; Selva et al., 2017; Tang et al., 2017; Bocchini et al., 2018; Ochogavía et al., 2018; Tang et al., 2019; de Oliveira et al., 2020). However, complete molecular regulatory mechanisms are yet to be revealed and would help fertilize agricultural crops innovatively through fixing the hybrid heterotic vigour in order to propagate indefinitely with enhanced productivity (Dujardin and Hanna, 1983; Khush, 1994; Ramulu et al., 1999; Spillane et al., 2001, 2004; Ortiz et al., 2013; Brukhin, 2017).

Besides, in an inter-specific cross combination, when the ploidy level of the maternal plant (e.g. diploid) is less than that of the paternal plant (e.g. tetraploid); the genome proportion of the paternal plant in the resultant endosperm tissues is greater than '1n', and as per the example, it is '2n' (Quarin, 1999; Felitti et al., 2015; Batista et al., 2019). Evolutionarily, the occurrence of the character – endosperm development in angiosperms – resembles poly-embryonic events (more than one fertilization

event leading to embryo formation) in gymnosperms, and later evolved to form endosperm in angiosperms. The genome dosage level variability in the endosperm, from 2n to 15n, might be termed ‘radicals’ or ‘variables’ for the said character, as defined by Vavilov (1922).

### *Polyendospermy and chlorophyllous endosperm*

Besides endosperm ploidy level variability (2n to 15n), *Arceuthobium americanum* Nutt. ex Engelm. has been reported to exhibit polyendospermy – formation of multiple ‘3n’ endosperms (Friedman and Sumner, 2009). Non-endospermic species have also been reported, for example in Podostemaceae due to the lack of a central cell (Battaglia, 1971; Baroux et al., 2002) and in Trapaceae and Orchidaceae, due to either suppression of primary endosperm nucleus’ cell division or disintegration of nuclei after few cell divisions (Johri et al., 1992). Very few taxa have been reported to exhibit chlorophyllous endosperm; Amaryllidaceae members are the notable ones (Meerow and Snijman, 2001). However, taxa from more than 70 families exhibit chlorophyllous embryos (Johri et al., 1992) during seed development. However, at maturity, the chlorophyllous nature disappears. The nature or type of photosynthesis, the quantity of carbon being fixed and its relative contribution with respect to the carbon translocated from leaves are open for further scientific research to assess the importance of chlorophyllous seed tissues. Irrespective of the statistical significance of the quantum of carbon fixed through endosperm or embryo, in comparison with translocated carbon, it will provide a greater understanding from a biological and evolutionary viewpoint, if the nature of photosynthesis is different from leaves. Answers from such studies will form a niche for further research questions, especially when the nature of photosynthesis in endosperm and/or embryo is different from leaves, as reported in wheat grain pericarp (Rangan et al., 2016). C<sub>4</sub> photosynthesis in wheat grain pericarp is yet an unsettled issue with arguments for and against (Henry et al., 2017). Species exhibiting variability in photosynthesis type, between leaves and endosperm/embryo parts of a plant might be of much help to throw light on this issue. The importance of non-foliar photosynthesis for yield improvement in crop plants is an untapped potential and is a current prioritized issue globally, on implementing C<sub>4</sub> photosynthesis in rice for improved productivity (Normile, 2008; Ermakova et al., 2020; Simkin et al., 2020). Carbon fixation in non-foliar (reproductive or sink) tissues of crop plants might, potentially, help accelerate grain filling to keep pace with the increase in cell size due to ploidy effects in endosperm tissues and its variability with respect to the cell position in the seed as detailed in the following section.

### *Endosperm ploidy variability within a seed*

The variability in ploidy level of endosperm across species or hybrids derived between parents at different ploidy levels is prominent in crossable species wherein fertile seeds are produced, involving parents differing in their ploidy (Tomaszewska and Kosina, 2018). Remarkably, the variability of endosperm ploidy within a seed was reported well before the DNA-double helix structure (Duncan and Ross, 1950; Swift, 1950). First, report to point the variability in the size of cells and nuclei of the young endosperm cells present in the central portion of the endosperm, was made as early as 1931 in maize (Lampe, 1931). However, it was only in 1950 that increased cell and nuclei size were

associated with the increased DNA content (Swift, 1950) or ploidy levels for those specific cells of enlarged size (Duncan and Ross, 1950) through quantified DNA content, and identified to be due to the endomitotic process. Genes (like *Rh11*) associated with cell size enlargement and higher ploidy were reported much later (Sugimoto-Shirasu et al., 2005). An upsurge of around 64 or 125 and even up to 1000 times the volume of nuclei from the central portion of endosperm, when compared to its peripheral nuclei, was documented in maize (Duncan and Ross, 1950); the ploidy correlation with DNA content has been reported recently (Santeramo et al., 2020). The ploidy level variation of endosperm nucleus within a maize seed, when compared in the aleurone and central regions of the endosperm, was reported to be up to ‘6n’ and ‘24n’, respectively (Swift, 1950). Also, the variability in endosperm ploidy among the centrally located endosperm cells was as high as ‘690n’ in maize (Kowles and Phillips, 1985). In addition to central region cells of the endosperm, suspensor cells too were reported to undergo endoreduplication events (Lee et al., 2009).

Apparently, the cells present in the central region of the endosperm undergo cell cycle only between S and G phases (endoreduplication) interrupted by gap periods and a doubling time of 24 h (Kowles et al., 1990, 1992b; Schweizer et al., 1995). The Cyclin A gene (*CYCA2;3*) is one of the key genes regulating the endoreduplication event (Imai et al., 2006), and the importance of this functionality has been viewed from a broader perspective on overall physiology and development (De Veylder et al., 2011). Besides increased mRNA and protein formation through endoreduplication, it has also been suggested that storing nucleotides by this mechanism might anticipate its use during embryogenesis and germination (Lee et al., 2009). It has been estimated that roughly 3% of the total endosperm cells (positioned in the central region) in maize seed exhibit the variability in ploidy from the normal triploid endosperm cells, and a few cells (approximately 1%) contained around 90 chromosomes (Lin, 1977). Ploidy variability, like in maize, has been reported in oats, as well (Tomaszewska and Kosina, 2018). Later, endoreduplication in plants was found in cotyledons, roots, cell suspensions, anthers, developing fruits (tomato) and leaves, as well (Joubès and Chevalier, 2000).

Defective kernel (*Dek*) mutants in maize exhibit reduced mitotic activity and, in turn, the endoreduplication, especially in the central regions of endosperm cell, was found to be controlled by a recessive gene (Kowles et al., 1992a) and also lacking an aleurone layer (Becraft and Yi, 2010). Water deficit (Artlip et al., 1995), abscisic acid (Mambelli and Setter, 1998), high temperature (Engelen-Eigles et al., 2000), parental dosage effect (Kowles et al., 1997; Leblanc et al., 2002; Tomaszewska and Kosina, 2018) and post-translational modification (Zhao and Grafi, 2000) may alter endoreduplication through regulating mitotic cycles in the endosperm, thereby suggesting multiple checkpoints, besides some recessive genes. Key molecular mechanisms involved in endoreduplication and increased ploidy level within the seed were identified as primarily due to the loss of activity of M-phase cyclin-dependent kinase and alterations in S-phase cyclin-dependent kinase (Larkins et al., 2001). Post-translational modifications such as hypophosphorylation on high mobility group I/Y protein by CDC2 kinase have been suggested to be associated with endoreduplication events in maize endosperm by alleviating the transcriptional repression by Histone H1 (Zhao and Grafi, 2000). Cyclin-dependent kinase inhibitors KRP1 and KRP2 inhibit the cell cycle, leading to the onset of endoreduplication events in the maize endosperm (Coelho et al., 2005).

Tolerance to altered genome dosage levels is evident in angiosperms, albeit in a few species only, and in different cells of a tissue, depending on the position (Kowles and Phillips, 1985; Joubès and Chevalier, 2000; Friedman et al., 2008). Understanding the linkage between the genetic nature of embryosac variability and endosperm ploidy might help shed light on the genetic regulatory mechanism underlying the ploidy syndrome. The following section associates endosperm ploidy with embryosac variability, using the common genetic factor. However, how far they could be influenced by environmental factors on ploidy variability with respect to the different cell positions within a tissue is yet to be revealed.

### Are endosperm ploidy and embryosac variability genetically linked?

Cells destined for endosperm formation are determined during embryosac formation. Later, with the onset of fertilization, the determined central cell, upon second fertilization, gets differentiated into endosperm tissue. To understand the variability in endosperm ploidy level, primarily, the megaspore mother cell (MMC) formation process – its root, needs to be understood. Identification of the diversity in embryosac development might shed some light on the endosperm variability. Normal meiosis of MMC leads to monosporic functional megaspores. The absence of cytokinesis during the second meiotic division or both meiotic divisions yields bi- (two haploid nuclei) or tetra-sporic (four haploid nuclei) functional megaspores. Evolutionarily, bi- and tetra-sporic trait types are derived from the monosporic trait (Arias and Williams, 2008). The single functional megaspore undergoes three mitotic division to yield an eight-nucleate (seven cellular) mature embryosac (ES), classified as *Polygonum* type – the most common one across angiosperms (Friedman, 1998). However, mega-gametogenesis devoid of the second or third mitosis, in combination with mono-, bi- or tetra-sporic megaspore, potentially generates 4 or 16 nucleate ES instead of the most frequently found eight-nucleate ES. Structured arrangement of 4 or 8 or 16 nuclei within ES in different species exhibits diverse forms of ES. Structurally (and the number of nuclei as well) differing ES were named upon the taxa from which they were described and reported at first. Documented ES types are *Polygonum*- (most common), *Oenothera*-, *Allium*-, *Peperomia*-, *Penaea*-, *Drusa*-, *Fritillaria*-, *Plumbagella*-, *Plumbago*-, *Adoxa*-, *Butomopsis*-, *Acalypha indica*-, *Peperomia hispidula*-, *Apinagia*- and *Dicraea* types (Maheshwari, 1950; Battaglia, 1971; Friedman et al., 2008). Due to the variability in the number of nuclei of the central cell of ES, accordingly the endosperm ploidy level gets modulated from the normal '3n' endosperm during the second fertilization event.

To obtain an overall understanding on such variable ES for number of nuclei, and, in turn, endosperm genome dosage levels, ES classified on genetic basis can be grouped broadly into seven types (Friedman et al., 2008), namely monosporic 2n (1m:1p), monosporic 3n (2m:1p), bi-sporic 3n (2m:1p), tetra-sporic 3n (2m:1p), tetra-sporic 5n (4m:1p), tetra-sporic 9n (8m:1p) and tetra-sporic 15n (14m:1p). Taxonomically, the order *Piperales* alone accommodates six of these seven genetic types (Arias and Williams, 2008) excluding the monosporic 2n type, suggesting the possibility for better understanding of the genetic variability of embryosac or endosperm and its development. The law of homologous variation, described by Vavilov (1922), might help foresee the endosperm variability in allied taxa as well for the

genera reported with variable endosperm ploidy levels. Most of the understanding of the process of organogenesis of MMC and flower or seed development has been reported only in few plant models like *Arabidopsis*, *Medicago*, *Zea* and *Nicotiana* (Table 1). With the present understanding from these model plants, the genetic mechanism underlying endosperm or ES variability may be revealed.

### Tolerance of altered genome dosage

Several reports have highlighted the importance of balanced genome dosage levels, positional cues, PCD and cell-cycle modifications primarily involved in normal endosperm development and, in turn, seed development and viability (Wangenheim, 1957; Becraft, 2001; Satyaki and Gehring, 2019). Whenever there is a deviation from the 2m:1p genome dosage ratio during endosperm formation, either premature cellularization (increased maternal dosage) or lack of cellularization (increased paternal dosage), seed abortion is the most probable result (Scott et al., 1998). However, the evolution (on parsimony basis) of 2m:1p from 1m:1p further from 1m:0p (Cailleau et al., 2010), and the existence of species accommodating the altered genome dosage, indicates nature's tolerance of genome dosage levels deviating from 2m:1p (Satyaki and Gehring, 2019). Likely, this tolerance mechanism is responsible for the 30% of angiosperms exhibiting altered genome dosage levels, from 1m:1p up to 14m:1p (Raghavan, 2006; Friedman et al., 2008). It is essential to underpin the genetic regulatory mechanisms of the tolerance of the deviation from the most common 2m:1p, especially so because understanding the genetic regulation and genes involved in endosperm development have been reported mostly in the model plant *Arabidopsis* (Table 1). However, in well-studied cereal crops like rice, wheat and maize, the initial phase of endosperm development (cell determination phase) is largely unknown (Li and Song, 2020; Olsen, 2020). Information gained from model plant species may help understand the endosperm development and the genome dosage balance in the species deviating from 2m:1p ratio. Also, methylation patterns involved in parental imprinting to functionalize parent-of-origin effects, when neutralized in both the parents, the impact of the nature of imprinting no longer hinders or alters normal endosperm development (Adams et al., 2000). Hence, this potentially gives clues on the possible tolerance for the nature's acceptance of deviation from the 2m:1p dosage level.

### Roadmap for understanding the endosperm genome variability

Gametophytic mutations (from completion of meiosis until fertilization and maternal control of seed development) affecting mega-gametogenesis follows non-Mendelian segregation (Yadegari and Drews, 2004). However, sporophytic mutations affecting mega-gametogenesis (from MMC till completion of meiosis) are yet to be studied. Hence, to sufficiently address this, two approaches may be followed. Firstly, working with the existing knowledge on the reproductive ontogeny in model flowering plants, potential candidate genes could be identified, and further, its validation in the taxa of interest, either through forward or reverse genetics approaches (Jankowicz-Cieslak and Till, 2015). Alternatively, differential gene expression (RNA-seq) between the two nearest taxa variable for endosperm and embryosac formation may be used to identify and validate the genes or genetic mechanism underlying variability. Irrespective of the

**Table 1.** Potential candidate genes involved in regulating plant reproductive development (based on reports of genes/metabolites associated with cell cycle)

Gene/metabolite/pathway	Name	Species	Function	Reference
Auxin	–	<i>Arabidopsis</i>	Prevents endosperm cellularization and leads to seed arrest	Batista et al. (2019)
<i>CYCD</i>	D-type cyclin	<i>Arabidopsis</i> <i>Medicago</i>	Cell cycle entry	den Boer and Murray (2000) and citations therein
<i>CDC2A</i>	Cyclin-dependent kinase ( <i>cdk</i> )	<i>Arabidopsis</i> <i>Zea</i>	Cell cycle regulation	
<i>CAK</i>	CDK-activating kinase	<i>Arabidopsis</i>	Phosphorylation of CDK to regulate cell cycle	
<i>RML</i>	Root meristemless	<i>Arabidopsis</i> <i>Nicotiana</i>	Encodes for Glutathione (GSH) biosynthesis linking cell cycle	
<i>CDC26</i>	Cell division cycle protein 26 (uORF)	<i>Arabidopsis</i>	Regulates cell cycle at anaphase	Lorenzo-Orts et al. (2019)
TOR pathway	Target of Rapamycin pathway	<i>Arabidopsis</i>	Integrates cytoplasm growth, cell expansion and cell cycle. Also, cell size checkpoint	Sablowski and Carnier Dornelas (2014) and citations therein
<i>ARP6</i> , <i>E2FA</i> , <i>E2FB</i> and <i>E2FC</i>	Actin-related protein 6, elongation factor complex	<i>Arabidopsis</i>	Distinguishing MMC identity to take meiosis; while other cells to take mitosis	Pinto et al. (2019) and citations therein
<i>KNU</i>	Knuckles	<i>Arabidopsis</i>	Drives expression specifically in MMC (even at tetrad) – the primary germline cell	
<i>THO/TREX</i> complex	Includes <i>Tex1</i> , <i>Hpr1</i> and <i>Tho6</i>	<i>Arabidopsis</i>	Leads to multiple MMC-like cells through involvement of tasiRNA that represses ARF3	Lora et al. (2019) and citations therein
Auxin cytokinin ratio	–	<i>Arabidopsis</i>	Positional information with auxin at distal and cytokinin at proximal polarity in ovule primordia development. Possibly, the absence of auxin leads to multiple MMC-like cells?	
<i>ASF1</i>	Anti-silencing function 1	<i>Arabidopsis</i>	Involved in gametophyte (both male and female) development and acquiring fertilization competency	Min et al. (2019)
F-box, E3 ligases	–	<i>Boecheira</i>	Distinguishes sexual and apomictic germline	Zühl et al. (2019)
<i>FRK3</i>	Fertilization-related kinase3	<i>Solanum chacoense</i>	Gametophyte (male and female) development	Daigle et al. (2019)

approaches taken to identify the candidate genes, validating the identified genes is an important part of the studies for understanding the genome dosage variability. A schematic overview of the roadmap towards understanding the genetic mechanism underlying endosperm or embryosac variability is provided in Fig. 1 and detailed in the following section.


### Regulators of the cell cycle as potential candidates

With the brief highlights of the genetic or genomic variation in endosperm and embryosac formation, as explained in previous sections, first a framework with identified potential target taxonomical groups for better understanding of the underlying genetic mechanism needs to be made. Secondly, potential candidate genes regulating the cell cycle (mitotic and meiotic) and cytokinetic checkpoints across species are to be identified and charted out (den Boer and Murray, 2000; Sablowski and Carnier Dornelas, 2014; Keçeli et al., 2017; Barrada et al., 2019; Daigle et al., 2019; Eekhout and De Veylder, 2019; Lora et al., 2019; Lorenzo-Orts et al., 2019; Min et al., 2019; Pinto et al., 2019; Zühl et al., 2019). For easy reference and as a starting point, a list of potential candidate genes (including miRNA and other non-coding RNAs)

regulating reproductive development in angiosperms is provided in Table 1. A brief comparison between the identified taxa with its nearest taxa with normal (3n) endosperm, at structural and functional levels for those genes might throw some light on the existing variability and its underlying genetic mechanism.

### Epigenetic regulation of genomic imprinting

In addition to genetic mechanisms, the importance of epigenetic mechanisms, especially during ontogeny of endosperm and seed development, is well documented with a role for non-coding small RNAs (Gehring and Satyaki, 2017; Yu et al., 2020). Methylation patterns of DNA and various modifications (acetylation and methylation) of histone protein (Song and Chen, 2015), in addition to small RNAs (Ng et al., 2012; Yakovlev et al., 2020; Yu et al., 2020), are the epigenetic regulators playing key roles in developmental processes. The coordinated act of the Polycomb group (PcG) of proteins and the DNA methylation essentially regulates the epigenetic systems for endosperm development, including the central cell stage of mega-gametogenesis. Gene products of *MEDEA* (*Mea*) *FERTILIZATION INDEPENDENT SEED DEVELOPMENT* (*FIS2*) and *FIS3* (a.k.a:



**Fig. 1.** An overview of the genetic mechanism regulating endosperm and embryonic variability.

*FIE, FERTILIZATION INDEPENDENT ENDOSPERM*) classified as PcG type are well-known regulators of the embryo and endosperm development (Köhler et al., 2003). PcG proteins are well-known transcriptional regulators through histone modifications mediated gene silencing mechanisms, with its homologues present across the animal and plant kingdoms in establishing cell identity and memory (Hsieh et al., 2003; Di Croce and Helin, 2013). The activity of most of the imprinted genes of the endosperm are regulated through epigenetic mechanisms and its complex nature poses difficulties in association with the corresponding phenotype (Pignatta et al., 2018). Besides, the complexities involved in understanding the embryo or endosperm development *per se*, crosstalk between embryo and endosperm also plays a crucial role in reproductive isolation and seed development. During wide hybridization, such crosstalk induced barriers for genetic exchange lead to hybrid incompatibilities (Fishman and Sweigart, 2018; Roth et al., 2019). Importantly, crosstalk between embryo and endosperm in mature seed during germination is also crucial for successful germination through the supply of seed reserves for the germination processes (Yan et al., 2014; Doll et al., 2020).

Although methylation patterns generally occur in heterochromatic (gene-poor) regions, in transposable element regions, the genic regions appear more methylated than the flanking regions. However, when compared between embryo and endosperm for genome-wide methylation patterns, it was reported that the *DEMETER* (*DME*) gene product demethylates the genes of transposable elements in the central cell nuclei of maternal origin before fertilization (Gehring et al., 2009). This highlights the role of epigenetic regulation in enforcing the parent-of-origin effect on the endosperm development. In maize endosperm, through genome-wide methylation pattern analysis, it has been reported that there is a reduction of 13–34% methylation when compared to embryo or leaf tissues (Lauria et al., 2004; Wang et al., 2015). DNA glycosylases (*DME*) and RNA-directed DNA

methylation (*RdDM*) through siRNA pathways are the two major epigenetic mechanisms regulating plant development with the former for imprinting female gametogenesis and the latter for vegetative tissues (Law and Jacobsen, 2010). The importance of Histone H1.2 in regulating *DME* for its downstream epigenetic regulation to impart an imprinting effect has been identified (Rea et al., 2012). Also, histones modified by PcG proteins are well-documented regulators at the transcriptional level involved in the normal embryo and endosperm development (Schubert et al., 2005; Moreno-Romero et al., 2019). Possibly, this corroborates the reports on endoreduplication in endosperm cells of the central region to have a reduced H1/DNA ratio and thereby enhancing the transcription and translation towards grain filling in endosperm cells (Zhao and Grafi, 2000; Larkins et al., 2001). This highlights the importance of epigenetic regulation through silencing as well as expression (through negative repression) for normal endosperm and embryo development.

There are very few reports that underscore the association between epigenetic regulation and ploidy level at the species level but none to the author's knowledge with reference to endosperm or seed development. Primarily, a non-linear relationship between the DNA methylation pattern and ploidy level was documented (Li et al., 2011), when compared between species with different ploidy (di-, tri- and tetra-). Altered ploidy might potentially alter the epigenetic silencing mechanism and thereby altering the normal developmental processes (Scheid et al., 1996).

### Role of genomics on understanding the endosperm variability

The candidate gene approach is the most traditional, long-standing and widely used methodologies in identifying genes associated with the trait of interest, particularly when the trait is of a quantitative or complex nature (Pflieger et al., 2001; Tabor et al., 2002; Zhu and Zhao, 2007). However, preliminary

knowledge of the trait is required for the identification of the genes underlying it. Also, contrasting genotypes for the trait of interest may be identified at first for generating a QTL map (Wayne and McIntyre, 2002; Zhu and Zhao, 2007). With the availability of computational facilities and robust sequencing technologies, the importance of genomics for the identification of genes and pathways is undisputed. Notably, genome-wide scanning approaches and digital candidate gene identification approaches, such as QTL, LD mapping, GWAS, GBS, and functional annotation, either alone or in combination with traditional candidate gene approaches are now frequently used (Mackay and Powell, 2007; Götz et al., 2008; McCarthy et al., 2008; Chen et al., 2009; Zhang et al., 2010, 2020; Bush and Moore, 2012; Glaubitz et al., 2014; Torkamaneh et al., 2020). The following three subsections provide an overview of the strength of genomics, with special reference towards understanding endosperm variability (both across species and within the seed).

### Differential gene expression studies

Significant progress has been made on the genetic regulation of endosperm formation that enhances our understanding of endosperm ontogeny (Table 1). Most of the studies involved the model plant *Arabidopsis thaliana* to understand the basic genetic mechanism underlying endosperm ontogeny (Table 1). However, genetic regulation underlying the genome-scale variability in endosperm ontogeny between closely related taxa (within *Piperales*) are yet to be uncovered. Differential gene expression (RNA-seq) in combination with single-cell genomics (scRNA-seq) might be of huge potential to address this. Additionally, for apomixis developmental processes, candidate genes like *SERK* and *APOSTART* have been identified (Albertini et al., 2005; Podio et al., 2014b). Since apomixis is reported to be tightly associated with the polyploidy mechanism (Ortiz et al., 2013), species within a genus (or variable cytotypes within a species) exhibiting variable ploidy level with apomictic behaviour are ideal models to identify the molecular mechanisms underlying the apomictic phenomena (Ortiz et al., 2013; Felitti et al., 2015; Ochogavía et al., 2018).

Comparison between nearest taxa variables for apomictic behaviour helped identify a set of genes including *AGO9* (Olmedo-Monfil et al., 2010), *CASEIN KINASE* (Depetris et al., 2018), *LORELEI* (Felitti et al., 2011), a MAP3K coding gene *QUI-GON-JINN* (Mancini et al., 2018), *THAUMATIN-LIKE*, *COPIA*, *CCD*, *LEISHMANOLYSIN-LIKE PEPTIDASE* and *FAR1-RELATED* (Ortiz et al., 2017), *ORC3* (Siena et al., 2016), *DORN1* and eATP pathway (Choi et al., 2014; Felitti et al., 2015), *AMP SYNTHASE*, *EF-1 $\alpha$* , *COP9 SIGNALOSOME* and *ACETOLACTATE SYNTHASE* (Cervigni et al., 2008), long non-coding RNAs like *NI3* (Ochogavía et al., 2018) and *QGF* (Mancini et al., 2018), and small RNAs like *ATHILA*, *LINE* and *ATLANTYS* (Olmedo-Monfil et al., 2010), involved in the regulatory processes for apomictic expression. In addition, some recent reports have identified apomixis-related genes in *Paspalum* (de Oliveira et al., 2020) and *Boehmeria* (Tang et al., 2019), which provide insight in genetic control of apomeiosis and polyploidy events leading to apomixis behaviour (Savidan, 2000; Hand and Koltunow, 2014). Also, regulatory mechanisms modulated at the transcriptional, translational and post-translational levels dissecting the apomictic trait, including epigenetic regulation, signal transduction and evolutionary aspects have been covered in detail (Ortiz et al., 2013; Brukhin, 2017; Schmidt, 2020). Notably, most

of the genetic or molecular mechanisms associated with apomictic traits have been reported for the genus *Paspalum* (Quarin, 1999; Labombarda et al., 2002; Laspina et al., 2008; Polegri et al., 2010; Ortiz et al., 2013, 2017; Podio et al., 2014a; Felitti et al., 2015; Siena et al., 2016; Bocchini et al., 2018; Depetris et al., 2018; Mancini et al., 2018; Ochogavía et al., 2018; de Oliveira et al., 2020). These details will be of significant help in mining the genes associated with apomictic behaviour in other species as well, based on homology, using sequences obtainable through next-generation sequencing (NGS) tools, followed by further functional validation.

Differential gene expression in combination with digital candidate gene identification approaches using functional annotation tools is a potential method to identify candidate genes underlying endosperm variability and apomixis. This is easily doable in qualitative traits and is feasible in quantitative traits when the quantum of expression results in a differential expression pattern associated with the contrasting phenotypic traits (Rangan et al., 2020a,b). On these advantages, differential gene expression or RNA-seq with functional annotation is a robust upcoming tool for candidate gene identification. While this approach could be used directly for endosperm ploidy variability across species, its utility for endosperm ploidy variability within a seed, and apomictic developmental processes will require additional single-cell sequencing methods (Wagner et al., 2016; Tanay and Regev, 2017).

### Single-cell sequencing for ploidy and epigenetic identification

Single-cell sequencing to understand ploidy level (aneuploidy) and its effect on development are well documented in animal and cell models, and upcoming in plant systems. The techniques in the whole genome, epigenetic and transcriptome studies have been applied to identify differences underlying cancer cells and regular ones (Ferrarini et al., 2018). Gene regulation and genome dosage imbalance due to aneuploidy in humans for trisomy dependent regulation, studied using RNA from skin fibroblast cells through single-cell RNA-seq (scRNA-seq) approaches have been reported.

Utilization of single-cell genomics in generating a cell-type atlas, cell-type specificity, resolving molecular relations and functional genomics, and cell differentiation for tissue specificity underlying biological or analytical problems are well documented (Efroni and Birnbaum, 2016; Tanay and Regev, 2017; Ryu et al., 2019; Rich-Griffin et al., 2020). Such studies might potentially be of help as starting material to plan for combining RNA-seq with single-cell genomics – scRNA-seq (Ryu et al., 2019), to study ploidy variability of individual cells within a seed or endosperm tissues.

For this, NGS and single-cell sequencing methodologies (Nawy, 2014), with the current understanding of the endosperm ontogeny in model plants (Table 1) might be of great help in understanding genome-scale variability of the endosperm. This may provide deeper insights in sporogenesis, gametogenesis, endosperm ontogeny and its variability across species from evolutionary, breeding and crop improvement perspectives. Gene regulatory network relationships and molecular interaction could very well be revealed using the scRNA-seq approaches. The peculiarities of seed resource allocation between maternal and paternal parents are not well-founded on hypotheses explaining the variability (Cailleau et al., 2010). Such studies will help underscore the genetic mechanisms underlying the biology of dosage compensation, genome imprinting and maternal/paternal effect with special reference to seed development.

Single-cell genomics in combination with RNA-seq and functional annotation approaches might be a robust tool to study and elucidate the underlying genetic events affecting the sporophytic and gametophytic mega-gametogenesis and, in turn, fertilization and seed developmental processes. This will also help in identifying the key candidate genes or molecular events responsible for the variability in the genome/nuclear content of ES from 1n to 14n. Additionally, this may also provide a better understanding of the genetic mechanism that favours seed resource allocation that begins with the occurrence of meiosis and completes before fertilization in cycads and ginkgoes whereas the same begins only after fertilization in angiosperms, with conifers in between (Cailleau et al., 2010). It will also be of use in endoreduplication events occurring in the central region of endosperm tissue leading to higher ploidy in those specific cell types as compared to the ones at the periphery (Duncan and Ross, 1950; Swift, 1950; Kowles and Phillips, 1985; Larkins et al., 2001).

### Merits and challenges of single-cell RNA-sequencing

Key merit is the availability of techniques and tools reported in model organisms or cells or tissue types, which can be applied with minor modifications to plants (Efroni and Birnbaum, 2016; Rich-Griffin et al., 2020). However, the presence of cell walls and various secondary metabolites in plants warrants for protocol standardization. Great advantage and strength are expected for studying cell-type or single-cell systems in plants, including the availability of laser-assisted microdissection tools to isolate what we actually see (Kehr, 2003; Nelson et al., 2006; Brandt et al., 2018; Sakai et al., 2018; Florez-Rueda et al., 2020). With these strategies, it is quite possible to achieve the isolation of genetic material from the cell lines or cell types of interest in plants, especially from endosperm during seed development. Once the genetic material is isolated; then, it is equal to any other system wherein common tools and techniques available for single-cell sequencing are directly applicable. Differential expression through RNA-seq might help identify the key candidate genes involved in endosperm ploidy variability.

In spite of these merits and potential possibilities of the applicability of scRNA-seq methods to understand endosperm ploidy variability, one should be cautious on the possible challenges that might be faced during the process: (1) epigenetic regulation including parent-of-origin and endoreduplication mechanisms are modulated through cell cycle regulators and isolating cell types for comparison might end up in different results; (2) sufficient quantity of homogenous cell types at the same cellular stage within the group is also an important criterion. It could potentially affect the results, and hence, uniformity of samples is imperative; (3) requirement of sufficient biological replicates of the same cellular stage and developmental level for higher confidence scores when the results are subjected to statistical analyses. Addressing these challenges might help save time and resources with proper planning towards understanding the genetic mechanism modulating endosperm variability within the seed, and across species, and their differences.

### Conclusion

Understanding the overall seed development is important for seeds that are not just the fulcrum for the survival of many life forms but also are the carriers of genetic imprints across

generations, which helps pass through the successful speciation events in the evolutionary timeline. Sporophytic (Mendelian segregation pattern) mutations and their molecular interactions affecting mega-gametogenesis are known to affect fertilization events and, at times, leads to ovule abortion yielding fewer seeds. Variability in endosperm ploidy within a seed (between the central region and the periphery) and at species level has been known for some 90 years. However, its underlying (epi)genetic mechanisms linking imprinting and parent-of-origin effects during its ontogeny are not completely understood. Present-day genomic tools like scRNA-seq might help gain a better picture of gametophyte development and fertilization in plants, especially of the ontogeny of embryo and endosperm, with special reference to variability within the seed and across species. Comparative studies between taxa exhibiting variability for ES structure and genome dosage level might shed light on the tolerance for altered genome dosage, imprinting and endosperm development under varied genetic environments, and its mechanism of action from an evolutionary perspective. The role of epigenetic mechanisms (both silencing and expression through negative repression) in combination with varied ploidy level (within the seed and species level) interactively regulating the embryo and endosperm development are yet to be elucidated. Additionally, genetic mechanisms underlying the other forms of variability like the disintegration of endosperm after a few divisions in certain taxa, cell identity for endoreduplication in specific cells from the central region of the endosperm, chlorophyllous endosperm and its importance in contributing to carbon fixation, apomictic endosperm dosage, composite endosperm and polyendospermy would also be revealed. This will potentially shed light on the variability in resource allocation during seed development processes and might help extend the duration of resource allocation in agricultural crop plants for enhanced productivity.

To exploit the triploid nature ( $3n$ ) of endosperm tissues, endosperm culture (*in vitro*) is mostly used for generating triploid plants with superior traits like disease resistance, high yield, larger fruits, etc., in species where the seed is not an economic product (including propagation mode), or felt undesirable as in case of banana, grapes or berries (Thomas and Chaturvedi, 2008; Miyashita et al., 2009). Direct somatic embryogenesis or callus mediated *in vitro* propagation, as reported in many species (Trolinder and Goodin, 1987; Novak et al., 1989; Li et al., 1998; Rangan et al., 2011), might be useful in deriving triploid plantlets with increased vigour using endosperm culture techniques (Sita et al., 1980; Tulecke et al., 1988; Gmitter et al., 1990; Sun et al., 2011; Antoniazzi et al., 2018; Van Thang et al., 2018). In addition, endosperm cells are targeted for improved grain quality and nutrition, like quality protein maize (Gibbon and Larkins, 2005; Vivek et al., 2008) and golden rice (Beyer et al., 2002; Paine et al., 2005). Generation of soft, friable and off-white to creamy yellow callus from endosperm tissues of cereal crops, in combination with processing (dried and powdered forms), would gear up agriculture towards lab farming that might lead to culture edible endosperm directly on Petri dishes. At the verge of a climate change scenario, transformation and utilization of the understanding of variability towards an entirely new form (petri cultures of endosperm) may also help feed humanity in the future, contributing towards fulfilling the dreams of the great academician N.I. Vavilov: a hunger-free world.

**Acknowledgement.** The author is thankful to ICAR-NBPGR, New Delhi, for institutional support.



## References

- Adams S, Vinkenoog R, Spielman M, Dickinson HG and Scott RJ (2000) Parent-of-origin effects on seed development in *Arabidopsis thaliana* require DNA methylation. *Development* **127**, 2493–2502.
- Albertini E, Marconi G, Reale L, Barcaccia G, Porceddu A, Ferranti F and Falcinelli M (2005) *SERK* and *APOSTART*. Candidate genes for apomixis in *Poa pratensis*. *Plant Physiology* **138**, 2185–2199.
- Alves ER, Carneiro VT and Araujo AC (2001) Direct evidence of pseudogamy in apomictic *Brachiaria brizantha* (Poaceae). *Sexual Plant Reproduction* **14**, 207–212.
- Antoniazzi CA, de Faria RB, de Carvalho PP, Mikovski AI, de Carvalho IF, de Matos EM, Reis AC, Viccini LF, Pinto DLP and Rocha DI (2018) In vitro regeneration of triploid plants from mature endosperm culture of commercial passionfruit (*Passiflora edulis* Sims). *Scientia Horticulturae* **238**, 408–415.
- Arendt N (1970) Variation in the apomictic seedlings of certain species of ficus, pp. 52–63 in Khokhlov S (Ed.) *Apomixis and breeding*. New Delhi, Amerind Publishing Co Pvt Ltd.
- Arias T and Williams JH (2008) Embryology of *Manekia naranjoana* (Piperaceae) and the origin of tetrasporic, 16-nucleate female gametophytes in Piperales. *American Journal of Botany* **95**, 272–285.
- Artlip T, Madison J and Setter T (1995) Water deficit in developing endosperm of maize: cell division and nuclear DNA endoreduplication. *Plant, Cell & Environment* **18**, 1034–1040.
- Asker S and Jerling L (1992) *Apomixis in plants*. Baton Roca, FL, CRC Press.
- Barke BH, Daubert M and Hörandl E (2018) Establishment of apomixis in diploid F2 hybrids and inheritance of apospory from F1 to F2 hybrids of the *Ranunculus auricomus* complex. *Frontiers in Plant Science* **9**, 1111.
- Baroux C, Spillane C and Grossniklaus U (2002) Evolutionary origins of the endosperm in flowering plants. *Genome Biology* **3**, 1021–1025.
- Barrada A, Djendli M, Desnos T, Mercier R, Robaglia C, Montané M-H and Menand B (2019) A TOR-YAK1 signaling axis controls cell cycle, meristem activity and plant growth in *Arabidopsis*. *Development* **146**, dev171298.
- Bashaw E and Hanna WW (1990) Apomictic reproduction, pp. 100–130 in Chapman GP (Ed.) *Reproductive versatility in the grasses*. Cambridge, Cambridge University Press.
- Batista RA, Figueiredo DD, Santos-González J and Köhler C (2019) Auxin regulates endosperm cellularization in *Arabidopsis*. *Genes & Development* **33**, 466–476.
- Battaglia E (1971) The embryo sac of Podostemaceae – an interpretation. *Caryologia* **24**, 403–420.
- Becraft PW (2001) Cell fate specification in the cereal endosperm. *Seminars in Cell & Developmental Biology* **12**, 387–394.
- Becraft PW and Gutierrez-Marcos J (2012) Endosperm development: dynamic processes and cellular innovations underlying sibling altruism. *Wiley Interdisciplinary Reviews: Developmental Biology* **1**, 579–593.
- Becraft PW and Yi G (2010) Regulation of aleurone development in cereal grains. *Journal of Experimental Botany* **62**, 1669–1675.
- Berger F, Grini PE and Schnittger A (2006) Endosperm: an integrator of seed growth and development. *Current Opinion in Plant Biology* **9**, 664–670.
- Beyer P, Al-Babili S, Ye X, Lucca P, Schaub P, Welsch R and Potrykus I (2002) Golden rice: introducing the  $\beta$ -carotene biosynthesis pathway into rice endosperm by genetic engineering to defeat vitamin A deficiency. *The Journal of Nutrition* **132**, 506S–510S.
- Bicknell RA and Koltunow AM (2004) Understanding apomixis: recent advances and remaining conundrums. *Plant Cell* **16**, S228–S245.
- Bocchini M, Galla G, Pupilli F, Bellucci M, Barcaccia G, Ortiz JPA, Pessino SC and Albertini E (2018) The vesicle trafficking regulator *PN\_SCD1* is demethylated and overexpressed in florets of apomictic *Paspalum notatum* genotypes. *Scientific Reports* **8**, 1–11.
- Brandt R, Mascher M and Thiel J (2018) Laser capture microdissection-based RNA-Seq of barley grain tissues, pp. 397–409 in Murray G (Ed.) *Laser capture microdissection. Methods in molecular biology*. New York, Springer.
- Brukhin V (2017) Molecular and genetic regulation of apomixis. *Russian Journal of Genetics* **53**, 943–964.
- Burton GW (1948) The method of reproduction in common bahia grass, *Paspalum notatum* L. *Agronomy Journal* **40**, 443–452.
- Bush WS and Moore JH (2012) Genome-wide association studies. *PLoS Computational Biology* **8**, e1002822.
- Cailleau A, Cheptou P-O and Lenormand T (2010) Ploidy and the evolution of endosperm of flowering plants. *Genetics* **184**, 439–453.
- Carman JG, Jamison M, Elliott E, Dwivedi KK and Naumova TN (2011) Apospory appears to accelerate onset of meiosis and sexual embryo sac formation in sorghum ovules. *BMC Plant Biology* **11**, 9.
- Cervigni GD, Paniego N, Pessino S, Selva JP, Díaz M, Spangenberg G and Echenique V (2008) Gene expression in diplosporous and sexual *Eragrostis curvula* genotypes with differing ploidy levels. *Plant Molecular Biology* **67**, 11–23.
- Chen J, Bardes EE, Aronow BJ and Jegga AG (2009) ToppGene Suite for gene list enrichment analysis and candidate gene prioritization. *Nucleic Acids Research* **37**, W305–W311.
- Choi J, Tanaka K, Cao Y, Qi Y, Qiu J, Liang Y, Lee SY and Stacey G (2014) Identification of a plant receptor for extracellular ATP. *Science* **343**, 290–294.
- Coelho CM, Dante RA, Sabelli PA, Sun Y, Dilkes BP, Gordon-Kamm WJ and Larkins BA (2005) Cyclin-dependent kinase inhibitors in maize endosperm and their potential role in endoreduplication. *Plant Physiology* **138**, 2323–2336.
- Daigle C, Mazin B and Matton DP (2019) The *Solanum chacoense* Fertilization-Related Kinase 3 (ScFRK3) is involved in male and female gametophyte development. *BMC Plant Biology* **19**, 202.
- den Boer BG and Murray JA (2000) Triggering the cell cycle in plants. *Trends in Cell Biology* **10**, 245–250.
- de Oliveira FA, Vigna BB, da Silva CC, Fávero AP, de Matta FP, Azevedo AL and de Souza AP (2020) Coexpression and transcriptome analyses identify active apomixis-related genes in *Paspalum notatum* leaves. *BMC Genomics* **21**, 78.
- Depetris MB, Acuña CA, Pozzi FI, Quarin C and Felitti SA (2018) Identification of genes related to endosperm balance number insensitivity in *Paspalum notatum*. *Crop Science* **58**, 813–822.
- De Veylder L, Larkin JC and Schnittger A (2011) Molecular control and function of endoreduplication in development and physiology. *Trends in Plant Science* **16**, 624–634.
- Di Croce L and Helin K (2013) Transcriptional regulation by Polycomb group proteins. *Nature Structural & Molecular Biology* **20**, 1147–1155.
- Doll N, Royek S, Fujita S, Okuda S, Chamot S, Stintzi A, Widiez T, Hothorn M, Schaller A and Geldner N (2020) A two-way molecular dialogue between embryo and endosperm is required for seed development. *Science* **367**, 431–435.
- Dujardin M and Hanna WW (1983) Apomictic and sexual pearl millet  $\times$  *Pennisetum squamulatum* hybrids. *Journal of Heredity* **74**, 277–279.
- Duncan RE and Ross JG (1950) The nucleus in differentiation and development: III. Nuclei of maize endosperm. *Journal of Heredity* **41**, 259–268.
- Eekhout T and De Veylder L (2019) Plant stress: hitting pause on the cell cycle. *eLife* **8**, e46781.
- Efroni I and Birnbaum KD (2016) The potential of single-cell profiling in plants. *Genome Biology* **17**, 65.
- Engelen-Eigles G, Jones RJ and Phillips R (2000) DNA endoreduplication in maize endosperm cells: the effect of exposure to short-term high temperature. *Plant, Cell & Environment* **23**, 657–663.
- Ermakova M, Danila FR, Furbank RT and Von Caemmerer S (2020) On the road to C4 rice: advances and perspectives. *Plant Journal* **101**, 940–950.
- Felitti SA, Seijo JG, González AM, Podio M, Laspina NV, Siena L, Ortiz JPA and Pessino SC (2011) Expression of *lorelei*-like genes in aposporous and sexual *Paspalum notatum* plants. *Plant Molecular Biology* **77**, 337–354.
- Felitti SA, Acuña CA, Ortiz JPA and Quarin CL (2015) Transcriptome analysis of seed development in apomictic *Paspalum notatum*. *Annals of Applied Biology* **167**, 36–54.
- Ferrarini A, Forcato C, Buson G, Tononi P, del Monaco V, Terracciano M, Bolognesi C, Fontana F, Medoro G, Neves R, Möhlendick B, Rihawi K, Ardizzoni A, Sumanasuriya S, Flohr P, Lambros M, de Bono J, Stoecklein NH and Manaresi N (2018) A streamlined workflow for single-cells genome-wide copy-number profiling by low-pass sequencing of LM-PCR whole-genome amplification products. *PLoS ONE* **13**, e0193689.

- Fishman L and Sweigart AL** (2018) When two rights make a wrong: the evolutionary genetics of plant hybrid incompatibilities. *Annual Review of Plant Biology* **69**, 707–731.
- Fiume E and Fletcher JC** (2012) Regulation of Arabidopsis embryo and endosperm development by the polypeptide signaling molecule CLE8. *Plant Cell* **24**, 1000–1012.
- Florez-Rueda AM, Waser L and Grossniklaus U** (2020) Laser-assisted microdissection of plant embryos for transcriptional profiling, pp. 127–139 in Bayer M (Ed.) *Plant embryogenesis. Methods in molecular biology*. New York, Springer.
- Friedman WE** (1992) Evidence of a pre-angiosperms origin of endosperm: implications for the evolution of flowering plants. *Science* **255**, 336–339.
- Friedman WE** (1998) The evolution of double fertilization and endosperm: a “historical” perspective. *Sexual Plant Reproduction* **11**, 6–16.
- Friedman CMR and Sumner MJ** (2009) Maturation of the embryo, endosperm, and fruit of the dwarf mistletoe *Arceuthobium americanum* (Viscaceae). *International Journal of Plant Sciences* **170**, 290–300.
- Friedman WE, Madrid EN and Williams JH** (2008) Origin of the fittest and survival of the fittest: relating female gametophyte development to endosperm genetics. *International Journal of Plant Sciences* **169**, 79–92.
- García-Aguilar M, Michaud C, Leblanc O and Grimanelli D** (2010) Inactivation of a DNA methylation pathway in maize reproductive organs results in apomixis-like phenotypes. *Plant Cell* **22**, 3249–3267.
- Gehring M and Satyaki P** (2017) Endosperm and imprinting, inextricably linked. *Plant Physiology* **173**, 143–154.
- Gehring M, Bubb KL and Henikoff S** (2009) Extensive demethylation of repetitive elements during seed development underlies gene imprinting. *Science* **324**, 1447–1451.
- Gibbon BC and Larkins BA** (2005) Molecular genetic approaches to developing quality protein maize. *Trends in Genetics* **21**, 227–233.
- Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q and Buckler ES** (2014) TASSEL-GBS: a high capacity genotyping by sequencing analysis pipeline. *PLoS ONE* **9**, e90346.
- Gmitter F, Ling X and Deng X** (1990) Induction of triploid *Citrus* plants from endosperm calli in vitro. *Theoretical and Applied Genetics* **80**, 785–790.
- Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M, Talón M, Dopazo J and Conesa A** (2008) High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Research* **36**, 3420–3435.
- Greilhuber J, Ebert I, Lorenz A and Vyskot B** (2000) Origin of facultative heterochromatin in the endosperm of *Gagea lutea* (Liliaceae). *Protoplasma* **212**, 217–226.
- Grimanelli D, Hernández M, Perotti E and Savidan Y** (1997) Dosage effects in the endosperm of diplosporous apomictic *Tripsacum* (Poaceae). *Sexual Plant Reproduction* **10**, 279–282.
- Guerin J, Rossel JB, Robert S, Tsuchiya T and Koltunow A** (2000) A DEFICIENS homologue is down-regulated during apomictic initiation in ovules of Hieracium. *Planta* **210**, 914–920.
- Haig D and Westoby M** (1991) Genomic imprinting in endosperm: its effect on seed development in crosses between species, and between different ploidies of the same species, and its implications for the evolution of apomixis. *Philosophical Transactions: Biological Sciences* **333**, 1–13.
- Hand ML and Koltunow AM** (2014) The genetic control of apomixis: asexual seed formation. *Genetics* **197**, 441–450.
- Henry RJ, Rangan P, Furtado A, Busch FA and Farquhar GD** (2017) Does C4 photosynthesis occur in wheat seeds? *Plant Physiology* **174**, 1992–1995.
- Hsieh T-F, Hakim O, Ohad N and Fischer RL** (2003) From flour to flower: how Polycomb group proteins influence multiple aspects of plant development. *Trends in Plant Science* **8**, 439–445.
- Hu G, Liang G and Wassom C** (1991) Chemical induction of apomictic seed formation in maize. *Euphytica* **56**, 97–105.
- Huang Y, Wang H, Huang X, Wang Q, Wang J, An D, Li J, Wang W and Wu Y** (2019) Maize VKS1 regulates mitosis and cytokinesis during early endosperm development. *Plant Cell* **31**, 1238–1256.
- Huh JH, Bauer MJ, Hsieh T-F and Fischer R** (2007) Endosperm gene imprinting and seed development. *Current Opinion in Genetics & Development* **17**, 480–485.
- Imai KK, Ohashi Y, Tsuge T, Yoshizumi T, Matsui M, Oka A and Aoyama T** (2006) The A-type cyclin CYCA2;3 is a key regulator of ploidy levels in Arabidopsis endoreduplication. *Plant Cell* **18**, 382–396.
- Jankowicz-Cieslak J and Till B** (2015) Forward and reverse genetics in crop breeding, pp. 215–240 in Al-Khayri J; Jain S and Johnson D (Eds) *Advances in plant breeding strategies: breeding, biotechnology and molecular tools*. Switzerland, Springer Nature.
- Johnston S, Den Nijs T, Peloquin S and Hanneman R** (1980) The significance of genic balance to endosperm development in interspecific crosses. *Theoretical and Applied Genetics* **57**, 5–9.
- Johri B, Ambegaokar K and Srivastava P** (1992) *Comparative embryology of angiosperms*. Berlin, Heidelberg, Springer-Verlag.
- Joubès J and Chevalier C** (2000) Endoreduplication in higher plants. *Plant Molecular Biology* **43**, 735–745.
- Kaushal P, Dwivedi KK, Radhakrishna A, Srivastava MK, Kumar V, Roy AK and Malaviya DR** (2019) Partitioning apomixis components to understand and utilize gametophytic apomixis. *Frontiers in Plant Science* **10**, 256.
- Keçeli BN, De Storme N and Geelen D** (2017) In vivo ploidy determination of *Arabidopsis thaliana* male and female gametophytes, pp. 77–85 in Schmidt A (Ed.) *Plant germline development methods and protocols*. New York, Springer.
- Kehr J** (2003) Single cell technology. *Current Opinion in Plant Biology* **6**, 617–621.
- Khush GS** (Ed.) (1994) *Apomixis: exploiting hybrid vigor in rice*. Philippines, Int. Rice Res. Inst. IRRI.
- Kirkbride RC, Lu J, Zhang C, Mosher RA, Baulcombe DC and Chen ZJ** (2019) Maternal small RNAs mediate spatial-temporal regulation of gene expression, imprinting, and seed development in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 2761–2766.
- Knox R** (1967) Apomixis: seasonal and population differences in a grass. *Science* **157**, 325–326.
- Köhler C, Hennig L, Spillane C, Pien S, Grissem W and Grossniklaus U** (2003) The Polycomb-group protein MEDEA regulates seed development by controlling expression of the MADS-box gene *PHERES*. *Genes & Development* **17**, 1540–1553.
- Kollmann J, Steinger T and Roy BA** (2000) Evidence of sexuality in European *Rubus* (Rosaceae) species based on AFLP and allozyme analysis. *American Journal of Botany* **87**, 1592–1598.
- Koltunow AM** (1993) Apomixis: embryo sacs and embryos formed without meiosis or fertilization in ovules. *Plant Cell* **5**, 1425–1437.
- Kowles RV and Phillips RL** (1985) DNA amplification patterns in maize endosperm nuclei during kernel development. *Proceedings of the National Academy of Sciences of the United States of America* **82**, 7010–7014.
- Kowles RV, Srien F and Phillips RL** (1990) Endoreduplication of nuclear DNA in the developing maize endosperm. *Developmental Genetics* **11**, 125–132.
- Kowles R, McMullen M, Yerk G, Phillips R, Kraemer S and Srien F** (1992a) Endosperm mitotic activity and endoreduplication in maize affected by defective kernel mutations. *Genome* **35**, 68–77.
- Kowles R, Yerk G, Srien F and Phillips R** (1992b) Maize endosperm tissue as an endoreduplication system, pp. 65–88 in Setlow JK (Ed.) *Genetic engineering: principles and methods*. Boston, MA, Springer.
- Kowles R, Yerk G, Haas K and Phillips R** (1997) Maternal effects influencing DNA endoreduplication in developing endosperm of *Zea mays*. *Genome* **40**, 798–805.
- Labombarda P, Busti A, Caceres ME, Pupilli F and Arcioni S** (2002) An AFLP marker tightly linked to apomixis reveals hemizyosity in a portion of the apomixis-controlling locus in *Paspalum simplex*. *Genome* **45**, 513–519.
- Lampe L** (1931) A microchemical and morphological study of the developing endosperm of maize. *Botanical Gazette* **91**, 337–376.
- Larkins BA, Dilkes BP, Dante RA, Coelho CM, Woo Ym and Liu Y** (2001) Investigating the hows and whys of DNA endoreduplication. *Journal of Experimental Botany* **52**, 183–192.
- Laspina NV, Vega T, Seijo JG, González AM, Martelotto LG, Stein J, Podio M, Ortiz JPA, Echenique VC and Quarín CL** (2008) Gene expression analysis at the onset of aposporous apomixis in *Paspalum notatum*. *Plant Molecular Biology* **67**, 615–628.

- Lauria M, Rupe M, Guo M, Kranz E, Pirona R, Viotti A and Lund G (2004) Extensive maternal DNA hypomethylation in the endosperm of *Zea mays*. *Plant Cell* **16**, 510–522.
- Law JA and Jacobsen SE (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature Reviews Genetics* **11**, 204–220.
- Leblanc O, Pointe C and Hernandez M (2002) Cell cycle progression during endosperm development in *Zea mays* depends on parental dosage effects. *Plant Journal* **32**, 1057–1066.
- Lee HO, Davidson JM and Duronio RJ (2009) Endoreplication: polyploidy with purpose. *Genes & Development* **23**, 2461–2477.
- Li C and Song R (2020) The regulation of zein biosynthesis in maize endosperm. *Theoretical and Applied Genetics*. doi:10.1007/s00122-019-03520-z.
- Li Z, Traore A, Maximova S and Guiltinan MJ (1998) Somatic embryogenesis and plant regeneration from floral explants of cacao (*Theobroma cacao* L.) using thidiazuron. *In Vitro Cellular & Developmental Biology – Plant* **34**, 293–299.
- Li A, Hu B-Q, Xue Z-Y, Chen L, Wang W-X, Song W-Q, Chen C-B and Wang C-G (2011) DNA methylation in genomes of several annual herbaceous and woody perennial plants of varying ploidy as detected by MSAP. *Plant Molecular Biology Reporter* **29**, 784–793.
- Lin B-Y (1977) Ploidy variation in maize endosperm. *Journal of Heredity* **68**, 143–149.
- Lopes MA and Larkins BA (1993) Endosperm origin, development, and function. *Plant Cell* **5**, 1383–1399.
- Lora J, Yang X and Tucker MR (2019) Establishing a framework for female germline initiation in the plant ovule. *Journal of Experimental Botany* **70**, 2937–2949.
- Lorenzo-Orts L, Witthoef J, Deforges J, Martinez J, Loubéry S, Placzek A, Poirier Y, Hothorn LA, Jaillais Y and Hothorn M (2019) Concerted expression of a cell cycle regulator and a metabolic enzyme from a bicistronic transcript in plants. *Nature Plants* **5**, 184–193.
- Mackay I and Powell W (2007) Methods for linkage disequilibrium mapping in crops. *Trends in Plant Science* **12**, 57–63.
- Maheshwari P (1950) *An introduction to the embryology of angiosperms* (1st edn). New York, McGraw-Hill.
- Mambelli S and Setter TL (1998) Inhibition of maize endosperm cell division and endoreduplication by exogenously applied abscisic acid. *Physiologia Plantarum* **104**, 266–272.
- Mancini M, Permingeat H, Colono C, Siena L, Pupilli F, Azzaro C, de Alencar Dusi DM, de Campos Carneiro VT, Podio M and Seijo JG (2018) The MAP3K-coding QUI-GON JINN (QGJ) gene is essential to the formation of unreduced embryo sacs in *Paspalum*. *Frontiers in Plant Science* **9**, 1547.
- McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JP and Hirschhorn JN (2008) Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nature Reviews Genetics* **9**, 356–369.
- Meerow AW and Snijman DA (2001) Phylogeny of Amaryllidaceae tribe Amaryllideae based on nrDNA ITS sequences and morphology. *American Journal of Botany* **88**, 2321–2330.
- Min Y, Frost JM and Choi Y (2019) Nuclear chaperone ASF1 is required for gametogenesis in *Arabidopsis thaliana*. *Scientific Reports* **9**, 13959.
- Miyashita T, Ohashi T, Shibata F, Araki H and Hoshino Y (2009) Plant regeneration with maintenance of the endosperm ploidy level by endosperm culture in *Lonicera caerulea* var. *emphylocalyx*. *Plant Cell, Tissue and Organ Culture* **98**, 291–301.
- Montezuma-de-Carvalho J (1967) The effect of N<sub>2</sub>O on pollen tube mitosis in styles and its potential significance for inducing haploidy in potato. *Euphytica* **16**, 190–198.
- Moreno-Romero J, Del Toro-De León G, Yadav VK, Santos-González J and Köhler C (2019) Epigenetic signatures associated with imprinted paternally expressed genes in the Arabidopsis endosperm. *Genome Biology* **20**, 41.
- Motamayor JC, Venzon D, Bajon C, Sauvanet A, Grandjean O, Marchand M, Bechtold N, Pelletier G and Horlow C (2000) Switch (swi1), an Arabidopsis thaliana mutant affected in the female meiotic switch. *Sexual Plant Reproduction* **12**, 209–218.
- Nawy T (2014) Single-cell sequencing. *Nature Methods* **11**, 18.
- Nelson T, Tausta SL, Gandotra N and Liu T (2006) Laser microdissection of plant tissue: what you see is what you get. *Annual Review of Plant Biology* **57**, 181–201.
- Ng DW, Lu J and Chen ZJ (2012) Big roles for small RNAs in polyploidy, hybrid vigor, and hybrid incompatibility. *Current Opinion in Plant Biology* **15**, 154–161.
- Normile D (2008) Reinventing rice to feed the world. *Science* **321**, 330–333.
- Novak F, Afza R, Van Duren M, Perea-Dallos M, Conger B and Xiaolang T (1989) Somatic embryogenesis and plant regeneration in suspension cultures of dessert (AA and AAA) and cooking (ABB) bananas (*Musa* spp.). *Bio/Technology* **7**, 154–159.
- Ochogavía A, Galla G, Seijo JG, González AM, Bellucci M, Pupilli F, Barcaccia G, Albertini E and Pessino S (2018) Structure, target-specificity and expression of PN\_LNC\_N13, a long non-coding RNA differentially expressed in apomictic and sexual *Paspalum notatum*. *Plant Molecular Biology* **96**, 53–67.
- Olmedo-Monfil V, Durán-Figueroa N, Arteaga-Vázquez M, Demesa-Arévalo E, Autran D, Grimanelli D, Slotkin RK, Martienssen RA and Vielle-Calzada J-P (2010) Control of female gamete formation by a small RNA pathway in Arabidopsis. *Nature* **464**, 628–632.
- Olsen O-A (2020) The modular control of cereal endosperm development. *Trends in Plant Science* **25**, 279–290.
- Ortiz JPA, Quarin CL, Pessino SC, Acuña C, Martínez EJ, Espinoza F, Hojsgaard DH, Sartor ME, Cáceres ME and Pupilli F (2013) Harnessing apomictic reproduction in grasses: what we have learned from *Paspalum*. *Annals of Botany* **112**, 767–787.
- Ortiz JPA, Revale S, Siena LA, Podio M, Delgado L, Stein J, Leblanc O and Pessino SC (2017) A reference floral transcriptome of sexual and apomictic *Paspalum notatum*. *BMC Genomics* **18**, 318.
- Paine JA, Shipton CA, Chaggar S, Howells RM, Kennedy MJ, Vernon G, Wright SY, Hinchliffe E, Adams JL and Silverstone AL (2005) Improving the nutritional value of Golden Rice through increased provitamin A content. *Nature Biotechnology* **23**, 482–487.
- Pflieger S, Lefebvre V and Causse M (2001) The candidate gene approach in plant genetics: a review. *Molecular Breeding* **7**, 275–291.
- Pignatta D, Novitzky K, Satyaki P and Gehring M (2018) A variably imprinted epiallele impacts seed development. *PLoS Genetics* **14**, e1007469.
- Pinto SC, Mendes MA, Coimbra S and Tucker MR (2019) Revisiting the female germline and its expanding toolbox. *Trends in Plant Science* **24**, 455–467.
- Podio M, Cáceres ME, Samoluk SS, Seijo JG, Pessino SC, Ortiz JPA and Pupilli F (2014a) A methylation status analysis of the apomixis-specific region in *Paspalum* spp. suggests an epigenetic control of parthenogenesis. *Journal of Experimental Botany* **65**, 6411–6424.
- Podio M, Felitti SA, Siena LA, Delgado L, Mancini M, Seijo JG, González AM, Pessino SC and Ortiz JPA (2014b) Characterization and expression analysis of SOMATIC EMBRYOGENESIS RECEPTOR KINASE (SERK) genes in sexual and apomictic *Paspalum notatum*. *Plant Molecular Biology* **84**, 479–495.
- Polegri L, Calderini O, Arcioni S and Pupilli F (2010) Specific expression of apomixis-linked alleles revealed by comparative transcriptomic analysis of sexual and apomictic *Paspalum simplex* Morong flowers. *Journal of Experimental Botany* **61**, 1869–1883.
- Quarin CL (1999) Effect of pollen source and pollen ploidy on endosperm formation and seed set in pseudogamous apomictic *Paspalum notatum*. *Sexual Plant Reproduction* **11**, 331–335.
- Quarin CL, Burson BL and Burton GW (1984) Cytology of intra- and inter-specific hybrids between two cytotypes of *Paspalum notatum* and *P. cymorrhizon*. *Botanical Gazette* **145**, 420–426.
- Raghavan V (2006) *Double fertilization: embryo and endosperm development in flowering plants*. Berlin, Heidelberg, Springer Verlag.
- Ramulu K, Sharma V, Naumova T, Dijkhuis P and van Lookeren Campagne M (1999) Apomixis for crop improvement. *Protoplasma* **208**, 196–205.
- Rangan P, Venugopalan A, Giridhar P and Ravishankar G (2011) Somatic embryogenesis and Agrobacterium-mediated transformation in *Bixa orellana* L. *Plant Cell, Tissue and Organ Culture* **105**, 317–328.
- Rangan P, Furtado A and Henry RJ (2016) New evidence for grain specific C<sub>4</sub> photosynthesis in wheat. *Scientific Reports* **6**, 31721.

- Rangan P, Furtado A and Henry R (2020a) Transcriptome profiling of wheat genotypes under heat stress during grain-filling. *Journal of Cereal Science* **91**, 102895.
- Rangan P, Furtado A, Henry R and Gaikwad A (2020b) Development of transcriptome analysis methods in Smithers G and Knoerzer K (Eds) *Reference module in food science*. Australia, Elsevier. In press. <https://doi.org/10.1016/B978-0-08-100596-5.22752-2>
- Rea M, Zheng W, Chen M, Braud C, Bhangu D, Rognan TN and Xiao W (2012) Histone H1 affects gene imprinting and DNA methylation in Arabidopsis. *Plant Journal* **71**, 776–786.
- Rich-Griffin C, Stechemesser A, Finch J, Lucas E, Ott S and Schäfer P (2020) Single-cell transcriptomics: a high-resolution avenue for plant functional genomics. *Trends in Plant Science* **25**, 186–197.
- Rodrigo JM, Zappacosta DC, Selva JP, Garbus I, Albertini E and Echenique V (2017) Apomixis frequency under stress conditions in weeping lovegrass (*Eragrostis curvula*). *PLoS ONE* **12**, e0175852.
- Roth M, Florez-Rueda AM and Städler T (2019) Differences in effective ploidy drive genome-wide endosperm expression polarization and seed failure in wild tomato hybrids. *Genetics* **212**, 141–152.
- Ryu KH, Huang L, Kang HM and Schiefelbein J (2019) Single-cell RNA sequencing resolves molecular relationships among individual plant cells. *Plant Physiology* **179**, 1444–1456.
- Sablowski R and Carnier Dornelas M (2014) Interplay between cell growth and cell cycle in plants. *Journal of Experimental Botany* **65**, 2703–2714.
- Sakai K, Tacconat L, Borrega N, Yansouni J, Brunaud V, Paysant-Le Roux C, Delannoy E, Magniette M-LM, Lepiniec L and Faure JD (2018) Combining laser-assisted microdissection (LAM) and RNA-Seq allows to perform a comprehensive transcriptomic analysis of epidermal cells of Arabidopsis embryo. *Plant Methods* **14**, 10.
- Santeramo D, Howell J, Ji Y, Yu W, Liu W and Kelliher T (2020) DNA content equivalence in haploid and diploid maize leaves. *Planta* **251**, 30.
- Šarhanová P, Vašut RJ, Dančák M, Bureš P and Trávníček B (2012) New insights into the variability of reproduction modes in European populations of *Rubus* subgen. *Rubus*: how sexual are polyploid brambles? *Sexual Plant Reproduction* **25**, 319–335.
- Satyaki PR and Gehring M (2019) Paternally acting canonical RNA-directed DNA methylation pathway genes sensitize Arabidopsis endosperm to paternal genome dosage. *Plant Cell* **31**, 1563–1578.
- Savidan Y (2000) Apomixis: genetics and breeding. *Plant Breeding Reviews* **18**, 13–86.
- Scheid OM, Jakovleva L, Afsar K, Maluszynska J and Paszkowski J (1996) A change of ploidy can modify epigenetic silencing. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 7114–7119.
- Schmidt A (2020) Controlling apomixis: shared features and distinct characteristics of gene regulation. *Genes* **11**, 329.
- Schubert D, Clarenz O and Goodrich J (2005) Epigenetic control of plant development by Polycomb-group proteins. *Current Opinion in Plant Biology* **8**, 553–561.
- Schweizer L, Yerk-Davis G, Phillips R, Srienc F and Jones RJ (1995) Dynamics of maize endosperm development and DNA endoreduplication. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 7070–7074.
- Scott RJ, Spielman M, Bailey J and Dickinson HG (1998) Parent-of-origin effects on seed development in *Arabidopsis thaliana*. *Development* **125**, 3329–3341.
- Selva JP, Siena L, Rodrigo JM, Garbus I, Zappacosta D, Romero JR, Ortiz JPA, Pessino SC, Leblanc O and Echenique V (2017) Temporal and spatial expression of genes involved in DNA methylation during reproductive development of sexual and apomictic *Eragrostis curvula*. *Scientific Reports* **7**, 1–11.
- Sharbel TF, Voigt ML, Corral JM, Thiel T, Varshney A, Kumlehn J, Vogel H and Rotter B (2009) Molecular signatures of apomictic and sexual ovules in the *Boechera holboellii* complex. *Plant Journal* **58**, 870–882.
- Sherwood R, Berg C and Young B (1994) Inheritance of apospory in buffelgrass. *Crop Science* **34**, 1490–1494.
- Siena LA, Ortiz JPA, Calderini O, Paolucci F, Cáceres ME, Kaushal P, Grisan S, Pessino SC and Pupilli F (2016) An apomixis-linked ORC3-like pseudogene is associated with silencing of its functional homologue in apomictic *Paspalum simplex*. *Journal of Experimental Botany* **67**, 1965–1978.
- Simkin AJ, Faralli M, Ramamoorthy S and Lawson T (2020) Photosynthesis in non-foliar tissues: implications for yield. *Plant Journal* **101**, 1001–1015.
- Sita GL, Ram NR and Vaidyanathan C (1980) Triploid plants from endosperm cultures of sandalwood by experimental embryogenesis. *Plant Science Letters* **20**, 63–69.
- Song Q and Chen ZJ (2015) Epigenetic and developmental regulation in plant polyploids. *Current Opinion in Plant Biology* **24**, 101–109.
- Spillane C, Steimer A and Grossniklaus U (2001) Apomixis in agriculture: the quest for clonal seeds. *Sexual Plant Reproduction* **14**, 179–187.
- Spillane C, Curtis MD and Grossniklaus U (2004) Apomixis technology development — virgin births in farmers' fields? *Nature Biotechnology* **22**, 687–691.
- Sugimoto-Shirasu K, Roberts GR, Stacey NJ, McCann MC, Maxwell A and Roberts K (2005) RHL1 is an essential component of the plant DNA topoisomerase VI complex and is required for ploidy-dependent cell growth. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 18736–18741.
- Sun D-Q, Lu X-H, Liang G-L, Guo Q-G, Mo Y-W and Xie J-H (2011) Production of triploid plants of papaya by endosperm culture. *Plant Cell, Tissue and Organ Culture* **104**, 23–29.
- Swift H (1950) The constancy of desoxyribose nucleic acid in plant nuclei. *Proceedings of the National Academy of Sciences of the United States of America* **36**, 643–654.
- Tabor HK, Risch NJ and Myers RM (2002) Candidate-gene approaches for studying complex genetic traits: practical considerations. *Nature Reviews Genetics* **3**, 391–397.
- Tanay A and Regev A (2017) Scaling single-cell genomics from phenomenology to mechanism. *Nature* **541**, 331–338.
- Tang C-Y (1977) *Apomixis in sorghum: mode of reproduction, inheritance, and improvement of apomictic frequency*. PhD dissertation, Texas A&M University, College Station.
- Tang Q, Zang G, Cheng C, Luan M, Dai Z, Xu Y, Yang Z, Zhao L and Su J (2017) Diplosporous development in *Boehmeria tricuspis*: insights from de novo transcriptome assembly and comprehensive expression profiling. *Scientific Reports* **7**, 46043.
- Tang Q, Xu Y, Deng C, Cheng C, Dai Z, Yang Z, Liu C and Su J (2019) A full-length reference floral transcriptome of *Boehmeria tricuspis* provides insights into apomeiosis and polyploidy. *International Journal of Genomics* **2019**, 4025747.
- Thomas TD and Chaturvedi R (2008) Endosperm culture: a novel method for triploid plant production. *Plant Cell, Tissue and Organ Culture* **93**, 1–14.
- Tomaszewska P and Kosina R (2018) Instability of endosperm development in amphiploids and their parental species in the genus *Avena* L. *Plant Cell Reports* **37**, 1145–1158.
- Torkamaneh D, Boyle B, St-Cyr J, Légaré G, Pomerleau S and Belzile F (2020) NanoGBS: a miniaturized procedure for GBS library preparation. *Frontiers in Genetics* **11**, 67.
- Trolinder NL and Goodin J (1987) Somatic embryogenesis and plant regeneration in cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* **6**, 231–234.
- Tulecke W, McGranahan G and Ahmadi H (1988) Regeneration by somatic embryogenesis of triploid plants from endosperm of walnut, *Juglans regia* L. cv Manregian. *Plant Cell Reports* **7**, 301–304.
- Valle Cd, Glienke C and Leguizamón G (1994) Inheritance of apomixis in *Brachiaria*, a tropical forage grass. *Apomixis Newsletter* **7**, 42–43.
- Van Thang B, Van Viet N, Nam VQ, Tung HT and Nhut DT (2018) Triploid plant regeneration from immature endosperms of *Melia azedarach*. *Plant Cell, Tissue and Organ Culture* **133**, 351–357.
- Vavilov NI (1922) The law of homologous series in variation. *Journal of Genetics* **12**, 47–89.
- Vivek B, Krivanek A, Palacios-Rojas N, Twumasi-Afryie S and Diallo A (2008) *Breeding quality protein maize (QPM): protocols for developing QPM cultivars*. Mexico, CIMMYT.
- Von Wangenheim K (1962) Zur Ursache der Abortion von Samenanlagen in Diploid-Polyploid-Kreuzungen. II. Unterschiedliche Differenzierung von

- Endospermen mit gleichem Genom. *Zeitschrift für Vererbungslehre* **93**, 319–334.
- Wagner A, Regev A and Yosef N** (2016) Revealing the vectors of cellular identity with single-cell genomics. *Nature Biotechnology* **34**, 1145–1160.
- Wang P, Xia H, Zhang Y, Zhao S, Zhao C, Hou L, Li C, Li A, Ma C and Wang X** (2015) Genome-wide high-resolution mapping of DNA methylation identifies epigenetic variation across embryo and endosperm in Maize (*Zea mays*). *BMC Genomics* **16**, 21.
- Wangenheim K-HFv** (1957) Untersuchungen Über den Zusammenhang zwischen Chromosomenzahl und Kreuzbarkeit bei Solanum-Arten. *Zeitschrift für Induktive Abstammungs-und Vererbungslehre* **88**, 21–37.
- Warmke H** (1954) Apomixis in *Panicum maximum*. *American Journal of Botany* **41**, 5–11.
- Wayne ML and McIntyre LM** (2002) Combining mapping and arraying: an approach to candidate gene identification. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 14903–14906.
- Williams JH and Friedman WE** (2002) Identification of diploid endosperm in an early angiosperms lineage. *Nature* **415**, 522–526.
- Yadegari R and Drews GN** (2004) Female gametophyte development. *Plant Cell* **16**, S133–S141.
- Yakovlev I, Viejo M and Fossdal CG** (2020) microRNAs in the formation of epigenetic memory in plants: the case of Norway spruce embryos, pp. 57–72 in Miguel C; Dalmay T and Chaves I (Eds) *Plant microRNAs. Concepts and strategies in plant sciences*. Switzerland, Springer.
- Yan D, Duermeyer L, Leoveanu C and Nambara E** (2014) The functions of the endosperm during seed germination. *Plant and Cell Physiology* **55**, 1521–1533.
- Young B, Sherwood R and Bashaw E** (1979) Cleared-pistil and thick-sectioning techniques for detecting aposporous apomixis in grasses. *Canadian Journal of Botany* **57**, 1668–1672.
- Yu J, Xu F, Wei Z, Zhang X, Chen T and Pu L** (2020) Epigenomic landscape and epigenetic regulation in maize. *Theoretical and Applied Genetics*. doi:10.1007/s00122-020-03549-5.
- Yudin B** (1966) *Experimental induction of marked polyploids in maize and problem in apomixis*. Saratovska, Gos.
- Zhang Z, Ersoz E, Lai C-Q, Todhunter RJ, Tiwari HK, Gore MA, Bradbury PJ, Yu J, Arnett DK and Ordovas JM** (2010) Mixed linear model approach adapted for genome-wide association studies. *Nature Genetics* **42**, 355–360.
- Zhang Y-W, Tamba CL, Wen Y-J, Li P, Ren W-L, Ni Y-L, Gao J and Zhang Y-M** (2020) mrMLM v4. 0: an R platform for multi-locus genome-wide association studies. bioRxiv, 976464.
- Zhao J and Grafi G** (2000) The high mobility group I/Y protein is hypophosphorylated in endoreduplicating maize endosperm cells and is involved in alleviating histone H1-mediated transcriptional repression. *Journal of Biological Chemistry* **275**, 27494–27499.
- Zhou S** (1980) Preliminary study on parthenogenesis in cotton. *Acta Genetica Sinica* **7**, 247–256.
- Zhu M and Zhao S** (2007) Candidate gene identification approach: progress and challenges. *International Journal of Biological Sciences* **3**, 420–427.
- Zühl L, Volkert C, Ibberson D and Schmidt A** (2019) Differential activity of F-box genes and E3 ligases distinguishes sexual versus apomictic germline specification in *Boechera*. *Journal of Experimental Botany* **70**, 5643–5657.