






Neuroimaging Highlight

Progressive Thrombosis of a Dolicho-Basilar Artery and Fusiform Aneurysm Resulting in Diffuse Ischemic and Hemorrhagic Complications

Vincent Brissette¹ , Marina Saad¹, Devavrat Nene¹, Christine Van Winssen¹ , Ange Boubacar Diouf²,
Marlise P. dos Santos³ , Robert Fahed^{1,2}  and Céline Ducroux¹ 

¹Department of Medicine (Neurology), The Ottawa Hospital and Ottawa Hospital Research Institute, Ottawa, ON, Canada, ²Department of Medical Imaging, The Ottawa Hospital and Ottawa Hospital Research Institute, Ottawa, ON, Canada and ³Department of Radiology, Radiation Oncology and Medical Physics, Department of Surgery, Section of Neuroradiology, Section of Interventional Neuroradiology, Division of Neurosurgery, University of Ottawa, Brain and Mind Research Institute, Ottawa Hospital Research Institute, Ottawa, ON, Canada

Keywords: interventional neuroradiology; stroke imaging; stroke

(Received 30 January 2024; date of acceptance 17 March 2024; First Published online 4 April 2024)




Figure 1. (A) Unenhanced CT: tortuous fusiform dilatation of the mid-basilar segment with hyperdense mural thrombus; (B) Computed Tomography Angiogram (CTA), axial plane: fusiform dolichoectatic basilar artery, red arrows on thrombus; (C) CTA, coronal plane: dolichoectatic basilar artery, red arrows on thrombus, and (D) patent basilar tip.

Corresponding author: Vincent Brissette; Email: vbrissette@toh.ca

Cite this article: Brissette V, Saad M, Nene D, Van Winssen C, Diouf AB, dos Santos MP, Fahed R, and Ducroux C. (2025) Progressive Thrombosis of a Dolicho-Basilar Artery and Fusiform Aneurysm Resulting in Diffuse Ischemic and Hemorrhagic Complications. *The Canadian Journal of Neurological Sciences* 52: 164–165, <https://doi.org/10.1017/cjn.2024.51>

© The Author(s), 2024. Published by Cambridge University Press on behalf of Canadian Neurological Sciences Federation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.




Figure 2. MRI (A) Longitudinal relaxation time (T1)-Volumetric interpolated breath-hold examination (VIBE) post contrast demonstrated interval progression of thrombus to distal basilar artery segment compared to CTA performed 2 days earlier, despite thrombolysis; (B) DWI: acute pontocerebellar infarcts; (C) Fluid-attenuated inversion recovery (FLAIR) post contrast: basilar artery wall circumferential mural enhancement (inflammation versus slow flow), and (D) acute subarachnoid hemorrhage.

A 54-year-old male presented acutely with dense right-sided hemiplegia and dysarthria (NIHSS 12). Initial CT angiogram revealed diffuse intracranial dolichoectasia with fusiform vertebrobasilar circulation and a partially occlusive basilar artery thrombus (Figure 1). He was treated with Tenecteplase but did not undergo endovascular treatment (EVT) given its lack of evidence in dolichoectasia. He aspirated vomitus and was intubated. MRI angiogram 24 hours later showed progression of the mural thrombosis of the distal basilar artery, infarction secondary to basilar artery perforators and right superior cerebellar artery occlusions,¹ cortical subarachnoid hemorrhage and basilar artery wall enhancement (Figure 2). This patient died secondary to brainstem damage. Outcomes associated with vertebrobasilar dolichoectasia and fusiform aneurysms are potentially catastrophic. Consensus surrounding acute management is lacking. There is limited evidence for thrombolysis, and EVT is technically challenging with high complication rates,² highlighting the need for further research to better understand effects on prognosis.

Author contributions. VB participated in the data collection and analysis and wrote the manuscript. MS, DN, CVW, ABD and MP dos S participated in the review of the manuscript. RF and CD participated in the data collection and analysis, and in the review of the manuscript.

Funding statement. No targeted fundings reported.

Competing interests. The authors report no disclosures relevant to the manuscript.

References

1. Dobrocky T, Piechowiak E, Goldberg J, et al. Absence of pontine perforators in vertebrobasilar dolichoectasia on ultra-high resolution cone-beam computed tomography. *J Neurointerv Surg.* 2021;13:580–584.
2. Wang Y, Yu J. Prospects and dilemmas of endovascular treatment for vertebrobasilar dolichoectasia. *Front Neurol.* 2022;13:895527.