

Protein arginine methyltransferases in protozoan parasites

Mario Alberto Rodriguez 

Departamento de Infectómica y Patogénesis Molecular, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Ciudad de México, México

Review Article

Cite this article: Rodriguez MA (2022). Protein arginine methyltransferases in protozoan parasites. *Parasitology* **149**, 427–435. <https://doi.org/10.1017/S0031182021002043>

Received: 20 September 2021
Revised: 15 November 2021
Accepted: 25 November 2021
First published online: 6 December 2021

Key words:

Arginine methylation; post-translational modification; protein arginine methyltransferases; protozoan parasites

Author for correspondence:

Mario Alberto Rodriguez,
E-mail: marodri@cinvestav.mx

Abstract

Arginine methylation is a post-translational modification involved in gene transcription, signalling pathways, DNA repair, RNA metabolism and splicing, among others, mechanisms that in protozoa parasites may be involved in pathogenicity-related events. This modification is performed by protein arginine methyltransferases (PRMTs), which according to their products are divided into three main types: type I yields monomethylarginine (MMA) and asymmetric dimethylarginine; type II produces MMA and symmetric dimethylarginine; whereas type III catalyses MMA only. Nine PRMTs (PRMT1 to PRMT9) have been characterized in humans, whereas in protozoa parasites, except for *Giardia intestinalis*, three to eight PRMTs have been identified, where in each group there are at least two enzymes belonging to type I, the majority with higher similarity to human PRMT1, and one of type II, related to human PRMT5. However, the information on the role of most of these enzymes in the parasites biology is limited so far. Here, current knowledge of PRMTs in protozoan parasites is reviewed; these enzymes participate in the cell growth, stress response, stage transitions and virulence of these microorganisms. Thus, PRMTs are attractive targets for developing new therapeutic strategies against these pathogens.

Introduction

Post-translational modifications (PTMs) are biochemical changes of amino acid residues that alter the charge state, hydrophobicity, conformation or stability of the proteins. These modifications act as molecular switches regulating the interaction of proteins with nucleic acids, cofactors, lipids and other proteins (Venne *et al.*, 2014). PTMs can be ‘read’ by specific effectors to regulate several cellular processes in response to extracellular and intracellular stimuli. To date, >450 unique protein modifications have been identified, including phosphorylation, acetylation, ubiquitination, SUMOylation and methylation, among others (Venne *et al.*, 2014).

Protein methylation occurs at different amino acids and it has been estimated that from 0.6 to 1.6% of the eukaryote genomes encode methyltransferases, the enzymes that perform this PTM (Katz *et al.*, 2003). Although initially the studies had been largely focused on lysine and arginine methylation of histones, which impact the transcription regulation (Jambhekar *et al.*, 2019), nowadays it is known that methylation on non-histone proteins regulates other molecular mechanisms, including RNA stability, splicing and translation (Lanouette *et al.*, 2014; Blanc and Richard, 2017; Al-Hamashi *et al.*, 2020).

Arginine contains in its side chain an aliphatic sequence of three carbon atoms, ending with a guanidino group, which is protonated and positively charged at physiological pH, with five potential hydrogen-bonds donors to interact with other molecules (Bedford and Clarke, 2009). The addition of one or two methyl groups to arginine creates a bulkier and more hydrophobic residue that does not alter the global cationic charge of the amino acid, but disperses it towards the methyl groups (Bedford and Richard, 2005; Bedford and Clarke, 2009). Arginine methylation increases the affinity for specific proteins such as those containing aromatic cages, including the Tudor domain (Tripsianes *et al.*, 2011), which interpret specific methylation marks to regulate several molecular events such as epigenetics, mRNA splicing, mRNA translation, DNA damage and stress response, among others (Blanc and Richard, 2017).

Infections with protozoan parasites are responsible for high morbidity and mortality in humans worldwide. The diseases produced by these microorganisms depend on the expression and activity of their virulence factors as well as the response of parasites to host immunity and the transition to different stages and hosts, events that require changes in gene expression and activation of specific signalling pathways that could be regulated by arginine methylation. Some advances have been made regarding the role of arginine methylation and the enzymes that perform this PTM, named as protein arginine methyltransferases (PRMTs), in the biology of certain protozoan parasites, mainly in *Trypanosoma brucei*; however, in others, such as *Trypanosoma cruzi*, many *Leishmania* and *Plasmodium* species as well as *Trichomonas vaginalis*, the arginine methylation and PRMTs have not yet been studied. Here, current knowledge on arginine methylation and PRMTs in protozoan parasites is reviewed.

Protein arginine methyltransferases (PRMTs)

The addition of methyl groups to the guanidino nitrogen atoms of arginine residues is catalysed by the PRMTs, which use S-adenosyl-L-Methionine (AdoMet or SAM) as the donor of the methyl groups (Blanc and Richard, 2017). According to their activity, there are three main types of PRMTs, type I catalyses monomethylarginine (MMA) and asymmetric dimethylarginine (ADMA); type II yields MMA and symmetric dimethylarginine (SDMA); and type III produces MMA only (Fig. 1); additionally, yeasts have a type IV PRMT, which monomethylates the delta (δ) nitrogen atom of arginine (Bedford and Clarke, 2009). Most PRMTs substrates contain glycine- and arginine-rich (GAR) sequences that include multiple arginines in RGG or RXR contexts (Najbauer *et al.*, 1993; Gary and Clarke, 1998; Feng *et al.*, 2013).

PRMTs contain a conserved core region of approximately 310 amino acids comprising the characteristic methyltransferase motif for the binding of the AdoMet cofactor, formed by a seven-strand twisted beta-sheet and the PRMT-specific 'double E' and 'THW' loops (Cheng *et al.*, 2005). In addition, different studies demonstrated that the dimerization of PRMTs facilitates their AdoMet-binding and their catalytic activity (Zhang *et al.*, 2000; Zhang and Cheng, 2003; Xu, 2004; Li *et al.*, 2015; Zhou *et al.*, 2015).

Nine PRMTs have been characterized in humans (PRMT1-9). PRMT1, 2, 3, 4 (also known as CARM1), 6 and 8 are type I, PRMT5 and 9 are type II, and PRMT7 is type III (Bedford and Clarke, 2009). Some of these proteins contain specific domains, PRMT2 has an SH3 domain, PRMT3 a Zn-finger domain (Frankel and Clarke, 2000) and PRMT8 is myristoylated at its N-terminus (Frankel *et al.*, 2002). It has been demonstrated that most of type I PRMTs are active as homodimers (Zhang *et al.*, 2000; Zhang and Cheng, 2003; Xu, 2004; Li *et al.*, 2015; Zhou *et al.*, 2015), while that PRMT5 forms a hetero-octameric complex (methylosome) with MEP50 (four subunits of each), in which MEP50 binds to the substrate and stimulates PRMT5 activity (Ho *et al.*, 2013). PRMT1 is responsible for most of the total monomethylation and asymmetric dimethylation of arginine, whereas PRMT5 has the major type II activity (Bedford and Clarke, 2009).

Most of these enzymes are ubiquitously expressed, with exception of PRMT8, which is localized only in the central nervous system (Lee *et al.*, 2005). In general, PRMTs are found in the nucleus and cytoplasm, but PRMT3 is located principally in the cytoplasm, PRMT6 is concentrated mainly in the nucleus and PRMT8 is linked to the plasma membrane (Frankel and Clarke, 2000; Frankel *et al.*, 2002; Lee *et al.*, 2005). Arginine methylation on histones is involved in the co-activation or co-repression of gene transcription and the effects on transcription of specific residues modified by different PRMTs have been determined (reviewed by Rakow *et al.*, 2020). PRMT1 and PRMT4/CARM1 are the main co-activators (Bedford and Clarke, 2009), PRMT2 and PRMT7 function as co-repressors (Ganesh *et al.*, 2006; Karkhanis *et al.*, 2012), whereas PRMT5 and PRMT6 have been reported to have both functions (Zhao *et al.*, 2009; Migliori *et al.*, 2012; Casadio *et al.*, 2013; Cheng *et al.*, 2020). In addition, PRMTs also methylate several non-histone proteins such as transcription factors and protein involved in RNA metabolism, DNA damage repair and many signalling pathways (reviewed by Al-Hamashi *et al.*, 2020).

PRMTs have also been identified in non-mammalian animals, filamentous fungi, yeasts, plants and protozoan parasites (McBride *et al.*, 2007; Ahmad *et al.*, 2011; Fisk and Read, 2011; Zhao *et al.*, 2016; Hadjikyriacou and Clarke, 2017; Plett *et al.*, 2017; Wang *et al.*, 2017; Bauer *et al.*, 2019). Enzymes similar to

mammalian PRMT1, 3 and 5 are present in different eukaryotic organisms, whereas other PRMTs, such as PRMT2 and 8, are less conserved and it has been suggested that they could have evolved in the multicellular organisms as a requirement for tissue-specific functions (Bachand, 2007).

Arginine methylation and PRMTs in protozoan parasites

Protozoan parasites, except *Giardia intestinalis*, have several PRMTs that were named according to their similarity with the human enzymes [reviewed by Fisk and Read (2011)], although they have structural differences with their mammalian homologues (see below). All protozoan parasites have two or more PRMTs of type I and at least one of type II; interestingly, trypanosomatids also contain a type III enzyme (Fisk and Read, 2011). They also possess the canonical histones, suggesting that the arginine methylation observed in histones of other eukaryotes, as well as their effects in gene expression, are preserved in these microorganisms. In addition, arginine methylation in non-histone proteins may also be important in the biology of protozoan parasites. However, most of the non-histone protein targets in these pathogens remain to be identified.

Phylogenetic analysis shows that there is no strong relationship of PRMT3, PRMT4/CARM1 and PRMT6 of the parasites with their respective human enzymes (Fig. 2). Furthermore, this analysis clearly separates the PRMTs of parasites into four main clades, one of them grouped PRMTs of type I, most of them with a higher relationship with the human PRMTs 1, 3 and 8; the other clade comprises PRMTs of type II similar to human PRMT5; the third clade is present only in trypanosomatids and corresponds to type III (PRMT7); and the last clade includes the PRMT3 of *T. brucei* and the atypical PRMT of *E. histolytica* (Fig. 2).

Arginine methylation and PRMTs in trypanosomatids

The causative agents of African trypanosomiasis (*T. brucei*), Leishmaniasis (*Leishmania* spp.) and Chagas disease (*T. cruzi*) belong to the kinetoplastids group (trypanosomatids). The main mechanisms implicated in the regulation of gene expression in these parasites are the post-transcriptional processes, including RNA turnover, translation and editing, involving multiple RNA-binding proteins (RBPs), which are common targets of arginine methylation (Campagnaro *et al.*, 2021). These microorganisms contain five PRMTs, which are homologues to the mammalian PRMT1, 3, 5, 6 and 7 (Fisk and Read, 2011); the studies related to these enzymes have been performed on the five enzymes of *T. brucei* and on the PRMT7 of *Leishmania major*, whereas there are no reports about the role of PRMTs in other *Leishmania* species or in *T. cruzi*.

Although the number of annual cases of African trypanosomiasis has been progressively decreasing, it has been estimated that approximately 1.8 million people live in high or very high-risk areas and 11.3 million people in moderate-risk zones (Simarro *et al.*, 2015). *T. brucei* life cycle consists of bloodstream trypomastigotes into the vertebrate host and procyclic trypomastigotes, epimastigotes and metacyclic trypomastigotes into the tsetse fly (*Glossina* spp.).

A recent global proteomic analysis identified 1332 methylarginines in 676 proteins (approximately 10% of the proteome) (Lott *et al.*, 2013), demonstrating that arginine methylation is a common PTM in this parasite. In addition, both type I and type II PRMT activities have been detected in cell extracts of *T. brucei* (Pelletier *et al.*, 2002). TbPRMT1 catalyses the majority of the ADMA modification *in vivo* (Pelletier *et al.*, 2005). It is expressed in all stages of the parasite life cycle and synthetic peptides




Fig. 1. Arginine methylation. The methylation of arginine residues is catalysed by the protein arginine methyltransferases (PRMTs), which transfer a methyl group from the S-adenosylmethionone (SAM or AdoMet), resulting in the arginine methylation and S-adenosylhomocysteine (SAH). PRMTs of type I, II and III produce monomethylarginine (MMA). In addition, PRMTs of type I add other methyl groups to the same nitrogen atom to yield asymmetric dimethylarginine (ADMA), while PRMTs of type II attach the second methyl group to the other N-terminal nitrogen of the arginine residue, forming symmetric dimethylarginine (SDMA).

containing GAR sequences are methylated by the recombinant protein (Pelletier *et al.*, 2005). The TbPRMT1 knockdown does not produce defects in the *in vitro* growth of the procyclic or bloodstream forms, but its knockout downregulates enzymes involved in glycolysis and upregulates enzymes participating in proline degradation, reduces the cytoplasmic mRNA granules in the insect stage under starvation and decreases the virulence of *T. brucei* in mice (Pelletier *et al.*, 2005; Kafková *et al.*, 2017; Kafková *et al.*, 2018). Interestingly, TbPRMT1 directly binds to RNA and to a lipin homologue that possesses activity of phosphatidic acid phosphatase (Pelletier *et al.*, 2013; Kafková *et al.*, 2018). All these results indicate that TbPRMT1 participates in carbohydrates metabolism, RNA biology, phospholipid biosynthesis and virulence (Table 1).

Based on the homology to human PRMT3, TbPRMT3 was predicted to be a type I enzyme (Fisk and Read, 2011); however, most of the conserved amino acids involved in PRMT activity are absent in TbPRMT3, where both glutamate residues in the double E loop are changed to aspartate and tryptophan is the only conserved residue in the THW loop; concordantly, its recombinant protein has no enzymatic activity (Kafková *et al.*, 2017). In fact, TbPRMT3 functions as a proenzyme that stabilizes the TbPRMT1 enzyme, producing an active heterotetramer formed by two enzyme-proenzyme pairs (TbPRMT1/TbPRMT3, also named TbPRMT1^{ENZ}/TbPRMT1^{PRO}) (Kafková *et al.*, 2017). Our phylogenetic analysis suggested that TbPRMT3 is more related to the atypical enzyme of *Entamoeba histolytica* EhPRMTA than to other PRMT members (Fig. 2).

TbPRMT5 is constitutively expressed in procyclic and bloodstream forms, it is found mainly in the cytoplasm forming high-molecular-weight complexes that do not include to MEP50, but that contain a DEAD box protein and two kinetoplast-specific proteins involved in RNA editing (Pasternack *et al.*, 2007). Interestingly, genes encoding MEP50

homologue have not been found in the genome database of *T. brucei* and recombinant TbPRMT5 displays enzymatic activity on broad substrates that includes the RNA-binding protein 16 (Pasternack *et al.*, 2007). Thus, this enzyme participates in the processing or translation of RNA (Table 1).

TbPRMT6 was named based on its higher homology to human PRMT6 than to another type I enzymes, but in the parasite, protein lacks the N-terminal nuclear localization signal detected in the human enzyme and its crystal structure revealed several features that are distinct from previously characterized type I PRMTs, including four stretches of insertion, the absence of the β 15 strand and a unique dimerization arm (Fisk *et al.*, 2010; Wang *et al.*, 2014a). It methylates bovine H4R3, components of the nuclear pore complex and flagellar proteins, but it does not methylate several parasite proteins that contain GAR sequences (Fisk *et al.*, 2010; Wang *et al.*, 2014a). Remarkably, the TbPRMT6 knockdown reduces the growth rate of procyclic and bloodstream stages and yields aberrant morphologies in both forms (Fisk *et al.*, 2010). Therefore, this enzyme participates in epigenetics, cell growth and cell morphology of *T. brucei* (Table 1).

TbPRMT7 is expressed in the bloodstream and procyclic stages, but is apparently dispensable (Fisk *et al.*, 2009). Unlike human PRMT7, the trypanosomatid homologue is localized only in the cytoplasm and lacks the second AdoMet binding-like domain, which is required for the activity of mammalian PRMT7; however, the parasite enzyme showed a robust monomethylation activity on several targets (Fisk *et al.*, 2009). Homodimerization of TbPRMT7 is due to the interaction between a dimerization arm of one subunit with the AdoMet-binding domain of the other and consequently, mutations on the dimerization arm remove dimerization and significantly reduce the methyltransferase activity (Wang *et al.*, 2014b; Debler *et al.*, 2016). In addition, the change of a glutamate residue to aspartate in the double E loop (E181D)




Fig. 2. Phylogenetic relationship of protozoan parasites PRMTs. The predicted amino acid sequences of PRMTs from human, protozoan parasites and *Saccharomyces cerevisiae* (types I and IV) were aligned by MUSCLE. Then, data were submitted to phylogenetic analysis by UPGMA using MEGA version 5.05. Accession numbers for the protein sequences of *T. vaginalis* are indicated. The other PRMTs used for this analysis were: Human: HsPRMT1 (NP_938074.2), Hs PRMT2 (NP_001526.2), HsPRMT3 (NP_005779.1), HsPRMT4/CARM1 (NP_954592.1), HsPRMT5 (NP_006100.2); HsPRMT6 (NP_060607.2), HsPRMT7 (NP_061896.1), HsPRMT8 (NP_062828.3), HsPRMT9 (NP_612373.2); *S. cerevisiae*: ScRMT2 (NP_010753.1), ScHMT1 (NP_009590.1); *T. brucei*: TbPRMT1 (Tb927.1.4690), TbPRMT3 (Tb927.10.3560), TbPRMT5 (Tb927.10.640), TbPRMT6 (Tb927.5.3960), TbPRMT7 (Tb927.7.5490); *P. falciparum*: PfPRMT1 (PF14_0242), PfPRMT4/CARM1 (PF14_0242), PfPRMT5 (PF13_0323); *T. gondii*: TgPRMT1 (GT1_030400), Tg PRMT3 (GT1_001320), TgPRMT4/CARM1 (GT1_074560), TgPRMT5 (GT1_126490); *E. histolytica*: EhPRMT1a (EHI_105780), EhPRMT1b (EHI_152460), EhPRMT1c (EHI_202470), EhPRMTA (EHI_159780), EhPRMT5 (EHI_158560). The numbers at the nodes of the branches indicate the confidence percentages of the tree topology from the bootstrap analysis of 1000 replicates. Proteins are grouped into type I, II, III, IV and atypical.

confers to TbPRMT7 the ability to perform ADMA, whereas an additional change of glutamine residue to alanine in the THW loop (E181D/Q329A double mutant) generated SDMA (Jain *et al.*, 2016), indicating that specific residues in the double E and THW loops are responsible for the product formed by different PRMTs.

An study analysing single and double TbPRMTs knockdowns (Lott *et al.*, 2014) showed that: (i) the reduction of TbPRMT1 induces a decrease in TbPRMT3 protein levels and similarly, the TbPRMT3 knockdown reduced the TbPRMT1 expression; (ii) the knockdown of TbPRMT1 or TbPRMT3 produced a decrease in ADMA with a concomitant increase in MMA, confirming that TbPRMT1 and TbPRMT3 form functional heterooligomers; (iii) in TbPRMT7 knockdown, a very slight reduction of ADMA and unchangeable levels of MMA were observed, but the double TbPRMT1/7 knockdown dramatically reduced ADMA and MMA, proposing that both enzymes can compete for the arginine monomethylation on the same substrates; (iv) in single TbPRMT6 and double TbPRMT6/7 knockdowns no

significant changes were observed in ADMA, confirming that TbPRMT6 methylates a relatively narrow group of substrates, but double knockdowns showed a more dramatic growth defect in the procyclic form that single repression of TbPRMT6, indicating that patterns of ADMA and MMA are critical to proliferation of procyclic forms (Lott *et al.*, 2014). All these results indicate that TbPRMTs display a functional interplay at multiple levels.

Leishmaniasis is a group of tropical diseases that affect 12 million people worldwide with 1.5 million new cases diagnosed each year (Alvar *et al.*, 2012). *Leishmania* spp. has two forms: promastigote in the sand fly vector and amastigote into the infected macrophages of the vertebrate host. The presence of arginine-methylated proteins in *Leishmania* sp. was demonstrated in 1986 and some of them have been identified (Paolantonacci *et al.*, 1986). *Leishmania* sp. has five PRMT homologues to those of *T. brucei*, but studies showed certain differences, for example, the *L. major* PRMT3 has an intact double E loop, suggesting that it could be an active enzyme (Campagnaro *et al.*, 2021). The expression of LmPRMT7 is high in the promastigote

Table 1. Functions of characterized PRMTs in protozoan parasites

Parasite	PRMT	Type	Molecular and/or cellular function	Reference
<i>Trypanosoma brucei</i>	TbPRMT1 (Tb927.1.4690)	Type I	Forms active heterotetramers with TbPRMT3, RNA and carbohydrates metabolism, phospholipid biosynthesis, stress response, virulence, <i>in vitro</i> growth of bloodstream forms	Pelletier <i>et al.</i> (2013), Kafková <i>et al.</i> (2017, 2018)
	TbPRMT3 (Tb927.10.3560)	Atypical	Inactive, but forms active heterotetramers with TbPRMT1	Kafková <i>et al.</i> (2017)
	TbPRMT5 (Tb927.10.640)	Type II	RNA metabolism	Pasternack <i>et al.</i> (2007)
	TbPRMT6 (Tb927.5.3960)	Type I	Cell growth, morphology, together with PRMT7 is critical for proliferation of procyclic forms	Fisk <i>et al.</i> (2010), Lott <i>et al.</i> (2013)
	TbPRMT7 (Tb927.7.5490)	Type III	Together with TbPRMT6 is critical for proliferation of procyclic forms	Lott <i>et al.</i> (2013)
<i>Leishmania major</i>	LmPRMT7 (LMJSD75_060015100)	Type III	RNA metabolism, life cycle progression, <i>in vitro</i> and <i>in vivo</i> virulence	Ferreira <i>et al.</i> (2014, 2020), Alcoforado Diniz <i>et al.</i> (2021)
<i>Plasmodium falciparum</i>	PfPRMT1 (PF14_0242)	Type I	RNA metabolism	Fan <i>et al.</i> (2009)
	PfPRMT5 (PF13_0323)	Type II	RNA splicing	Hossain <i>et al.</i> (2013)
<i>Toxoplasma gondii</i>	TgPRMT1 (GT1_030400)	Type I	Epigenetics, RNA metabolism and splicing, cell duplication	Musiyenko <i>et al.</i> (2012), El Bissati <i>et al.</i> (2016)
	TgPRMT4/TgCARM 1 (GT1_074560)	Type I	Epigenetics, stage conversion	Saksouk <i>et al.</i> (2005)
	TgPRMT5 (GT1_126490)	Type II	Epigenetics, stage conversion	Liu <i>et al.</i> (2019)
<i>Entamoeba histolytica</i>	EhPRMT1a (EHI_105780)	Type I	Epigenetics	Borbolla-Vázquez <i>et al.</i> (2015)
	EhPRMTA (EHI_159780)	Atypical	RNA metabolism, cell growth, migration, stress response	Cázares-Apátiga <i>et al.</i> (2017), Medina-Gómez <i>et al.</i> (2021)

logarithmic growth phase and intracellular amastigotes, but low in the promastigote stationary phase (Ferreira *et al.*, 2014). Interestingly, the LmPRMT7 knockout augmented infectivity *in vitro* and *in vivo*; and in concordance, the overexpression of this enzyme leads to a reduced progression of the lesion; moreover, the LmPRMT7 deletion rescues the pathogenic phenotype of an attenuated strain of *L. major* (Ferreira *et al.*, 2014; Alcoforado Diniz *et al.*, 2021). The absence of LmPRMT7 also impairs parasite development within the sand fly vector *Phlebotomus duboscqi* (Alcoforado Diniz *et al.*, 2021). The proteome comparison of wild-type and $\Delta prmt7$ parasites allowed the identification of 40 hypomethylated proteins in the mutant cells, including 17 RBPs (Ferreira *et al.*, 2020). Interestingly, the half-life of the RBP16 protein was reduced in the stationary phase of $\Delta prmt7$ cells compared to wild-type parasites, whereas the half-life of Alba3 was similar between both populations, although the absence of PRMT7 diminished the binding of Alba3 to the mRNA of the virulence-factor δ -*amastin*, affecting its stability (Ferreira *et al.*, 2020), suggesting that LmPRMT7 may regulate the protein stability or the affinity of RBPs for their mRNA targets, impacting in the life cycle progression and virulence of *L. major* (Table 1).

Arginine methylation and PRMTs in *Plasmodium falciparum*

Parasites of the genus *Plasmodium* are the causative agents of malaria. The World Health Organization estimated about 229 million malaria cases and 409 000 malaria-related deaths in 2019 (World Health Organization, 2020). The major species infecting humans are *P. falciparum*, *P. vivax*, *P. malariae*, *P. ovale* and *P. knowlesi*, although the majority of the malaria

cases are produced by *P. falciparum*, which is associated with the most severe morbidity and mortality in Africa, whereas *P. vivax* is the specie more widely distributed (Jong *et al.*, 2020). Parasites of the genus *Plasmodium* have several stages in the life cycle: gametes, zygotes, ookyetes, oocysts and sporozoites in the insect vector (females mosquitoes of the genus *Anopheles*) and schizonts, merozoites, ring, trophozoites and gametocytes in the vertebrate hosts. In addition, in *P. vivax* and *P. ovale* infections, some parasites become in dormant forms called 'hypnozoites'.

Immunoprecipitation assays in blood stages of *P. falciparum* using anti-methyl arginine antibodies showed the presence of 843 proteins with this PTM involved in several functions such as protein synthesis, chromatin organization and haemoglobin metabolic processes, among others, indicating that arginine methylation may play an important regulatory role in the parasite biology, including host-parasite interactions (Zeeshan *et al.*, 2017). *P. falciparum* encodes three PRMTs: PfPRMT1, PfPRMT4/CARM1 and PfPRMT5 (Fisk and Read, 2011). Recombinant PfPRMT1 methylates histones H4 and H2A and several conserved substrates involved in RNA metabolism, such as fibrillarin, poly(A)-binding protein II, ribosomal protein S2 and a splicing factor; in concordance, this enzyme is localized in both the cytoplasm and the nucleus (Fan *et al.*, 2009). In summary, these studies indicate that PfPRMT1 may play important roles in epigenetics and RNA metabolism (Table 1).

A study about the organization and assembly of the Sm core complex in the *P. falciparum* spliceosome suggested that its proteins form a heptameric ring, in which the methylation of PfSmD1, catalysed by PfPRMT5, allows the interaction of this Sm protein with two proteins containing the Tudor domain:

PfSMN (survival of motor neurons) and PfTSN (Tudor Staphylococcal nuclease) (Hossain *et al.*, 2013), indicating that PfPRMT5 has an important role in splicing (Table 1).

At the moment, neither the role of PfPRMT4/CARM1 in the biology of *P. falciparum* nor the function of PRMTs in the other *Plasmodium* species has been investigated.

Arginine methylation and PRMTs in *Toxoplasma gondii*

Toxoplasma gondii is an obligate intracellular parasite that produces a disease called toxoplasmosis. It has been estimated that this microorganism infects 16–40% of the population in the USA and up to 80% of the population in other countries (Hill and Dubey, 2002). Members of the cat family are its final hosts, whereas various warm-blooded animals including humans, are intermediate hosts (Montoya and Liesenfeld, 2004). The *T. gondii* life cycle consists of two stages, the replicating tachyzoite and the latent bradyzoite. Bradyzoites can remain in host tissues for its lifetime without health consequences, but when host immunity is attenuated, bradyzoites can differentiate into actively replicating tachyzoites, causing life-long chronic infection.

This parasite has four PRMTs similar to PRMT1, PRMT3, PRMT4/CARM1 and PRMT5 (Fisk and Read, 2011). Approximately 50% of the MMA-proteins found in wild-type parasites do not exhibit this modification in TgPRMT1 knockout cells, indicating that they are putative substrates for this enzyme (Yakubu *et al.*, 2017). TgPRMT1 substrates include the H4 histone, two ApiAP2 transcription factors, several RBPs, a lysine methyltransferase containing a SET domain, myosin E and SPM2 (a component of the subpellicular microtubules that regulate cell division) and, interestingly, most of them are also susceptible to phosphorylation, suggesting a relationship between both PTMs (Saksouk *et al.*, 2005; Yakubu *et al.*, 2017). The fact that myosin E and SPM2 are putative substrates of TgPRMT1 could explain why this enzyme is detected concentrated in the centrosome and in a pericentriolar material during the replication of tachyzoites (El Bissati *et al.*, 2016). Another target of TgPRMT1 is the Argonaute protein (TgAgo), a component of the RNA-induced silencing complex (RISC) that in *T. gondii* displays low RNA slicer activity, but its methylation by TgPRMT1 induces the recruitment of the Tudor Staphylococcal Nuclease (TgTSN), making it a second potent RNA slicer for RISC (Musiyenko *et al.*, 2012). Parasites lacking TgPRMT1 grow slower than wild-type ones and show abnormal daughter buds, perturbed centrosome stoichiometry and loss of synchronous replication (Musiyenko *et al.*, 2012; El Bissati *et al.*, 2016). Thus, TgPRMT1 participates in epigenetics, RNA metabolism and splicing, mechanisms that may control cell duplication (Table 1).

TgPRMT4/CARM1 catalyses the methylation of H3R17, an epigenetic mark for gene activation that is present in genes specifically expressed by tachyzoites; concordantly, inhibition of TgPRMT4/CARM1 induces conversion to bradyzoites, as well as a decrease of H3R17 methylation (Saksouk *et al.*, 2005). On the other hand, TgPRMT5 produces MMA and SMDA on H3R26 and H4R3 and is expressed in both tachyzoites and bradyzoites, but it has different localization in both stages, in tachyzoites, it is found in the cytoplasm, while in bradyzoites it is located mainly in the nucleus (Liu *et al.*, 2019). Therefore, TgPRMT4/CARM1 and TgPRMT5 participate in epigenetics, which in turn regulates the stage conversion (Table 1).

Arginine methylation and PRMTs in *Entamoeba histolytica*

Entamoeba histolytica is the microorganism that causes human amebiasis. This parasite infects up to 50 million people worldwide each year, producing intestinal and extra-intestinal diseases that

cause between 40 000 and 100 000 deaths annually (Walsh, 1986). Its life cycle consists of two stages, the infective cysts and the invasive trophozoites.

The *E. histolytica* chromatin is organized in nucleosomes, but the distance of the DNA linker is irregular (Torres-Guerrero *et al.*, 1991). Moreover, its histone H4 is longer than the consensus due to the insertion at the N-terminus of three amino acid segments, one of them adds three lysine residues that could be susceptible to being modified by acetylation and/or methylation (Binder *et al.*, 1995; Lozano-Amado *et al.*, 2016). Regarding the arginine methylation of histones, the antibodies against MMA- and ADMA-H4R3 recognized nuclear proteins of amoeba, indicating the presence of this epigenetic mark in *E. histolytica*, although the alignment of the amino acid sequence of its H4 histone with the eukaryotic consensus sequence, showed that the modified arginine corresponds to position 8 (H4R8) (Borbolla-Vázquez *et al.*, 2015; Lozano-Amado *et al.*, 2016).

This parasite contains five PRMTs, four of them show similarity to type I enzymes, whereas the other is related to the PRMT5 family (type II) (Fisk and Read, 2011) (Fig. 2). The amino acid sequences of the four type I PRMTs display higher similarity to human PRMT1 and do not contain the specific signatures for other type I PRMTs; therefore, they were initially classified as PRMT1 homologues (Fisk and Read, 2011). However, our phylogenetic analysis suggested that one of these EhPRMTs does not show significant homology with any PRMT family (Borbolla-Vázquez *et al.*, 2015) (Fig. 2), therefore, we named it as atypical PRMT (EhPRMTA), whereas the other three type I enzymes were termed EhPRMT1a, EhPRMT1b and EhPRMT1c (Borbolla-Vázquez *et al.*, 2015). An antibody against human PRMT1 recognized these EhPRMT1 proteins and they were localized in the nucleus and the cytoplasm, suggesting that they could methylate both histones and non-histone proteins (Borbolla-Vázquez *et al.*, 2015). In fact, the recombinant protein of EhPRMT1a catalysed the asymmetric dimethylation of H4R3 in commercial histones (Borbolla-Vázquez *et al.*, 2015), indicating that it may be involved in epigenetics (Table 1); however, until now it is unknown whether the three EhPRMT1 proteins have redundant or different functional roles.

EhPRMTA is the type I enzyme that shows the lowest similarity to PRMT1 (Fisk and Read, 2011). In concordance, this protein was not recognized by an antibody against human PRMT1 (Borbolla-Vázquez *et al.*, 2015). In addition, it does not contain several of the canonical amino acid residues of type I PRMTs (Medina-Gómez *et al.*, 2021) and a phylogenetic analysis grouped this protein with the inactive PRMT3 of *T. brucei* (Fig. 1), although the EhPRMTA recombinant protein displayed enzymatic activity on commercial histones (Medina-Gómez *et al.*, 2021). On the other hand, EhPRMTA interacts with the EhTSN protein, which is a transcription factor of *E. histolytica* (Cázares-Apátiga *et al.*, 2017) and its homologues in other protozoan parasites participate in RNA metabolism (Hossain *et al.*, 2008; Musiyenko *et al.*, 2012), suggesting that this EhPRMT is involved in transcription regulation and in RNA metabolism; moreover, the EhPRMTA-knockdown increased cell growth and phagocytosis, but decreased cell migration and survival of trophozoites submitted to heat shock (Medina-Gómez *et al.*, 2021), indicating that this PRMT participates in these events (Table 1), although the identification of its protein targets remains to be determined.

Discussion

Because there is a correlation between the expression level of some PRMTs and cancer (Yang and Bedford, 2013), investigation in both academic laboratories and the pharmaceutical industry has

been leading to the discovery of several small molecules targeting different PRMTs that at micromolar and submicromolar concentrations display anti-tumour effects in cellular assays (Hu *et al.*, 2016; Zhang and Zheng, 2016; Guccione and Richard, 2019). In fact, three specific inhibitors for PRMT5 (GSK3325695, JNJ-64619178 and EPZ015938) and one for type I PRMTs (GSK3368715) have been analysed in phase I clinical trials (Rakow *et al.*, 2020). As PRMTs of protozoan parasites play important roles in their cell growth, virulence, conversion stages and response to stress conditions (Table 1), anti-PRMT drugs could also be effective for the treatment of diseases produced by those pathogenic microorganisms. In fact, several small molecules that inhibit the activity of PfPRMT1 also restrict the *P. falciparum* growth (Fan *et al.*, 2009). Moreover, a study analysing the anti-plasmodial activity of several compounds that inhibit epigenetic modifiers in cancer cells, showed that sinefungin, a pan-inhibitor of AdoMet-dependent methyltransferases, affected the asexual stage of *P. falciparum*, whereas the ellagic acid, a PRMT inhibitor, was 10-fold more potent against the asexual stage and early gametocytes than against mammalian cells (Coetzee *et al.*, 2020). In addition, the structural differences shown by the PRMTs of the parasites with respect to their human homologues could be used to design specific anti-parasitic drugs.

Conclusions

Methylation of arginine residues is involved in the regulation of several molecular processes, such as transcription, RNA processing and signal transduction pathways, which in parasites may control pathogenesis-related events. Although most PRMTs from protozoan parasites have not been characterized, current research on some of these enzymes shows that they participate in cell growth, metabolism, stress response, stage transitions and virulence. Thus, the PRMTs of protozoan parasites are attractive targets to develop new therapeutic strategies that could help in the control of the public health problems generated by these pathogens. Actually, the considerable advances in the identification of clinically relevant PRMT inhibitors for cancer therapy could be extended to find effective drugs against the protozoan parasites.

Financial support. This work was supported by Consejo Nacional de Ciencia y Tecnología (CONACyT, Mexico) Grant FORDECYT-PRONACES/194163/2020.

Conflict of interest. The author declares there are no conflicts of interest.

Ethical standards. Not applicable.

References

- Ahmad A, Dong Y and Cao X (2011) Characterization of the PRMT gene family in rice reveals conservation of arginine methylation. *PLoS One* **6**, e22664.
- Al-Hamashi AA, Diaz K and Huang R (2020) Non-histone arginine methylation by protein arginine methyltransferases. *Current Protein & Peptide Science* **21**, 699–712.
- Alcoforado Diniz J, Chaves MM, Vaselek S, Miserani Magalhães RD, Ricci-Azevedo R, de Carvalho RVH, Lorenzon LB, Ferreira TR, Zamboni D, Walrad PB, Volf P, Sacks DL and Cruz AK (2021) Protein methyltransferase 7 deficiency in *Leishmania major* increases neutrophil associated pathology in murine model. *PLoS Neglected Tropical Diseases* **15**, e0009230.
- Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J and Boer Md (2012) *Leishmaniasis* worldwide and global estimates of its incidence. *PLoS One* **7**, e35671.
- Bachand F (2007) Protein arginine methyltransferases: from unicellular eukaryotes to humans. *Eukaryotic Cell* **6**, 889–898.
- Bauer I, Lechner L, Pidroni A, Petrone A-M, Merschak P, Lindner H, Kremser L, Graessle S, Golderer G, Allipour S and Brosch G (2019) Type I and II PRMTs regulate catabolic as well as detoxifying processes in *Aspergillus nidulans*. *Fungal Genetics and Biology* **129**, 86–100.
- Bedford MT and Clarke SG (2009) Protein arginine methylation in mammals: who, what, and why. *Molecular Cell* **33**, 1–13.
- Bedford MT and Richard S (2005) Arginine methylation. *Molecular Cell* **18**, 263–272.
- Binder M, Ortner S, Plaimauer B, Födinger M, Wiedermann G, Scheiner O and Duchêne M (1995) Sequence and organization of an unusual histone H4 gene in the human parasite *Entamoeba histolytica*. *Molecular and Biochemical Parasitology* **71**, 243–247.
- Blanc RS and Richard S (2017) Arginine methylation: the coming of age. *Molecular Cell* **65**, 8–24.
- Borbolla-Vázquez J, Orozco E, Betanzos A and Rodríguez MA (2015) *Entamoeba histolytica*: protein arginine transferase 1a methylates arginine residues and potentially modify the H4 histone. *Parasites & Vectors* **8**, 219.
- Campagnaro GD, Nay E, Plevin MJ, Cruz AK and Walrad PB (2021) Arginine methyltransferases as regulators of RNA-binding protein activities in pathogenic kinetoplastids. *Frontiers in Molecular Biosciences* **8**, 692668. doi: 10.3389/fmolb.2021.692668
- Casadio F, Lu X, Pollock SB, LeRoy G, Garcia BA, Muir TW, Roeder RG and Allis CD (2013) H3R42me2a is a histone modification with positive transcriptional effects. *Proceedings of the National Academy of Sciences* **110**, 14894–14899.
- Cázares-Apátiga J, Medina-Gómez C, Chávez-Munguía B, Calixto-Gálvez M, Orozco E, Vázquez-Calzada C, Martínez-Higuera A and Rodríguez MA (2017) The Tudor staphylococcal nuclease protein of *Entamoeba histolytica* participates in transcription regulation and stress response. *Frontiers in Cellular and Infection Microbiology* **7**, 52. doi: 10.3389/fcimb.2017.00052.
- Cheng X, Collins RE and Zhang X (2005) Structural and sequence motifs of protein (histone) methylation enzymes. *Annual Review of Biophysics and Biomolecular Structure* **34**, 267–294.
- Cheng D, Gao G, Di Lorenzo A, Jayne S, Hottiger MO, Richard S and Bedford MT (2020) Genetic evidence for partial redundancy between the arginine methyltransferases CARM1 and PRMT6. *Journal of Biological Chemistry* **295**, 17060–17070.
- Coetzee N, von Grüning H, Opperman D, van der Watt M, Reader J and Birkholtz L-M (2020) Epigenetic inhibitors target multiple stages of *Plasmodium falciparum* parasites. *Scientific Reports* **10**, 2355.
- Debler EW, Jain K, Warmack RA, Feng Y, Clarke SG, Blobel G and Stavropoulos P (2016) A glutamate/aspartate switch controls product specificity in a protein arginine methyltransferase. *Proceedings of the National Academy of Sciences* **113**, 2068–2073.
- El Bissati K, Suvorova ES, Xiao H, Lucas O, Upadhya R, Ma Y, Hogue Angeletti R, White MW, Weiss LM and Kim K (2016) *Toxoplasma gondii* arginine methyltransferase 1 (PRMT1) is necessary for centrosome dynamics during tachyzoite cell division. *mBio* **7**, e02094-15. doi: 10.1128/mBio.02094-15.
- Fan Q, Miao J, Cui L and Cui L (2009) Characterization of PRMT1 from *Plasmodium falciparum*. *Biochemical Journal* **421**, 107–118.
- Feng Y, Maity R, Whitelegge JP, Hadjikyriacou A, Li Z, Zurita-Lopez C, Al-Hadid Q, Clark AT, Bedford MT, Masson J-Y and Clarke SG (2013) Mammalian protein arginine methyltransferase 7 (PRMT7) specifically targets RXR sites in lysine- and arginine-rich regions. *Journal of Biological Chemistry* **288**, 37010–37025.
- Ferreira TR, Alves-Ferreira EVC, Defina TPA, Walrad P, Papadopoulou B and Cruz AK (2014) Altered expression of an RBP associated arginine methyltransferase 7 in *Leishmania major* affects parasite infection. *Molecular Microbiology* **94**, 1085–1102.
- Ferreira TR, Dowle AA, Parry E, Alves-Ferreira EVC, Hogg K, Kolokousi F, Larson TR, Plevin MJ, Cruz AK and Walrad PB (2020) PRMT7 regulates RNA-binding capacity and protein stability in *Leishmania* parasites. *Nucleic Acids Research* **48**, 5511–5526.
- Fisk JC and Read LK (2011) Protein arginine methylation in parasitic protozoa. *Eukaryotic Cell* **10**, 1013–1022.
- Fisk JC, Sayegh J, Zurita-Lopez C, Menon S, Presnyak V, Clarke SG and Read LK (2009) A type III protein arginine methyltransferase from the protozoan parasite *Trypanosoma brucei*. *Journal of Biological Chemistry* **284**, 11590–11600.
- Fisk JC, Zurita-Lopez C, Sayegh J, Tomasello DL, Clarke SG and Read LK (2010) TbPRMT6 is a type I protein arginine methyltransferase that

- contributes to cytokinesis in *Trypanosoma brucei*. *Eukaryotic Cell* **9**, 866–877.
- Frankel A and Clarke S** (2000) PRMT3 is a distinct member of the protein arginine N-methyltransferase family. *Journal of Biological Chemistry* **275**, 32974–32982.
- Frankel A, Yadav N, Lee J, Branscombe TL, Clarke S and Bedford MT** (2002) The novel human protein arginine N-methyltransferase PRMT6 is a nuclear enzyme displaying unique substrate specificity. *Journal of Biological Chemistry* **277**, 3537–3543.
- Ganesh L, Yoshimoto T, Moorthy NC, Akahata W, Boehm M, Nabel EG and Nabel GJ** (2006) Protein methyltransferase 2 inhibits NF- κ B function and promotes apoptosis. *Molecular and Cellular Biology* **26**, 3864–3874.
- Gary JD and Clarke S** (1998) RNA and protein interactions modulated by protein arginine methylation. *Progress in Nucleic Acid Research and Molecular Biology* **61**, 65–131.
- Guccione E and Richard S** (2019) The regulation, functions and clinical relevance of arginine methylation. *Nature Reviews Molecular Cell Biology* **20**, 642–657.
- Hadjikyriacou A and Clarke SG** (2017) *Caenorhabditis elegans* PRMT-7 and PRMT-9 are evolutionarily conserved protein arginine methyltransferases with distinct substrate specificities. *Biochemistry* **56**, 2612–2626.
- Hill D and Dubey JP** (2002) *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clinical Microbiology and Infection* **8**, 634–640.
- Ho M-C, Wilczek C, Bonanno JB, Xing L, Seznec J, Matsui T, Carter LG, Onikubo T, Kumar PR, Chan MK, Brenowitz M, Cheng RH, Reimer U, Almo SC and Shechter D** (2013) Structure of the arginine methyltransferase PRMT5-MEP50 reveals a mechanism for substrate specificity. *PLoS One* **8**, e57008.
- Hossain MJ, Korde R, Singh S, Mohammed A, Dasaradhi PVN, Chauhan VS and Malhotra P** (2008) Tudor domain proteins in protozoan parasites and characterization of *Plasmodium falciparum* Tudor staphylococcal nuclease. *International Journal for Parasitology* **38**, 513–526.
- Hossain M, Sharma S, Korde R, Kanodia S, Chugh M, Rawat K and Malhotra P** (2013) Organization of *Plasmodium falciparum* spliceosomal core complex and role of arginine methylation in its assembly. *Malaria Journal* **12**, 333.
- Hu H, Qian K, Ho M-C and Zheng YG** (2016) Small molecule inhibitors of protein arginine methyltransferases. *Expert Opinion on Investigational Drugs* **25**, 335–358.
- Jain K, Warmack RA, Debler EW, Hadjikyriacou A, Stavropoulos P and Clarke SG** (2016) Protein arginine methyltransferase product specificity is mediated by distinct active-site architectures. *Journal of Biological Chemistry* **291**, 18299–18308.
- Jambhekar A, Dhall A and Shi Y** (2019) Roles and regulation of histone methylation in animal development. *Nature Reviews Molecular Cell Biology* **20**, 625–641.
- Jong RM, Tebeje SK, Meerstein-Kessel L, Tadesse FG, Jore MM, Stone W and Bousema T** (2020) Immunity against sexual stage *Plasmodium falciparum* and *Plasmodium vivax* parasites. *Immunological Reviews* **293**, 190–215.
- Kafková L, Debler EW, Fisk JC, Jain K, Clarke SG and Read LK** (2017) The major protein arginine methyltransferase in *Trypanosoma brucei* functions as an enzyme-prozyme complex. *Journal of Biological Chemistry* **292**, 2089–2100.
- Kafková L, Tu C, Pazzo KL, Smith KP, Debler EW, Paul KS, Qu J and Read LK** (2018) *Trypanosoma brucei* PRMT1 is a nucleic acid-binding protein with a role in energy metabolism and the starvation stress response. *mBio* **9**. doi: 10.1128/mBio.02430-18.
- Karkhanis V, Wang L, Tae S, Hu Y-J, Imbalzano AN and Sif S** (2012) Protein arginine methyltransferase 7 regulates cellular response to DNA damage by methylating promoter histones H2A and H4 of the polymerase δ catalytic subunit gene, POLD1. *Journal of Biological Chemistry* **287**, 29801–29814.
- Katz JE, Dlakić M and Clarke S** (2003) Automated identification of putative methyltransferases from genomic open reading frames. *Molecular & Cellular Proteomics* **2**, 525–540.
- Lanouette S, Mongeon V, Figeys D and Couture J** (2014) The functional diversity of protein lysine methylation. *Molecular Systems Biology* **10**, 724.
- Lee J, Sayegh J, Daniel J, Clarke S and Bedford MT** (2005) PRMT8, a new membrane-bound tissue-specific member of the protein arginine methyltransferase family. *Journal of Biological Chemistry* **280**, 32890–32896.
- Li H-T, Gong T, Zhou Z, Liu Y-T, Cao X, He Y, Chen CD and Zhou J-Q** (2015) Yeast Hmt1 catalyses asymmetric dimethylation of histone H3 arginine 2 in vitro. *Biochemical Journal* **467**, 507–515.
- Liu M, Li F-X, Li C-Y, Li X-C, Chen L-F, Wu K, Yang P-L, Lai Z-F, Liu T, Sullivan WJ, Cui L and Chen X-G** (2019) Characterization of protein arginine methyltransferase of *TgPRMT5* in *Toxoplasma gondii*. *Parasites & Vectors* **12**, 221.
- Lott K, Li J, Fisk JC, Wang H, Aletta JM, Qu J and Read LK** (2013) Global proteomic analysis in trypanosomes reveals unique proteins and conserved cellular process impacted by arginine methylation. *Journal of Proteomics* **91**, 210–225.
- Lott K, Zhu L, Fisk JC, Tomasello DL and Read LK** (2014) Functional interplay between protein arginine methyltransferases in *Trypanosoma brucei*. *MicrobiologyOpen* **3**, 595–609.
- Lozano-Amado D, Herrera-Solorio AM, Valdés J, Alemán-Lazarini L, Almaraz-Barrera MdJ, Luna-Rivera E, Vargas M and Hernández-Rivas R** (2016) Identification of repressive and active epigenetic marks and nuclear bodies in *Entamoeba histolytica*. *Parasites & Vectors* **9**, 19.
- McBride AE, Zurita-Lopez C, Regis A, Blum E, Conboy A, Elf S and Clarke S** (2007) Protein arginine methylation in *Candida albicans*: role in nuclear transport. *Eukaryotic Cell* **6**, 1119–1129.
- Medina-Gómez C, Bolaños J, Borbolla-Vázquez J, Munguía-Robledo S, Orozco E and Rodríguez MA** (2021) The atypical protein arginine methyltransferase of *Entamoeba histolytica* (EhPRMTA) is involved in cell proliferation, heat shock response and in vitro virulence. *Experimental Parasitology* **222**, 108077.
- Migliori V, Müller J, Phalke S, Low D, Bezzi M, Mok WC, Sahu SK, Gunaratne J, Capasso P, Bassi C, Cecatiello V, De Marco A, Blackstock W, Kuznetsov V, Amati B, Mapelli M and Guccione E** (2012) Symmetric dimethylation of H3R2 is a newly identified histone mark that supports euchromatin maintenance. *Nature Structural & Molecular Biology* **19**, 136–144.
- Montoya J and Liesenfeld O** (2004) Toxoplasmosis. *The Lancet* **363**, 1965–1976.
- Musiyyenko A, Majumdar T, Andrews J, Adams B and Barik S** (2012) PRMT1 methylates the single Argonaute of *Toxoplasma gondii* and is important for the recruitment of Tudor nuclease for target RNA cleavage by antisense guide RNA. *Cellular Microbiology* **14**, 882–901.
- Najbauer J, Johnson BA, Young AL and Aswad DW** (1993) Peptides with sequences similar to glycine, arginine-rich motifs in proteins interacting with RNA are efficiently recognized by methyltransferase(s) modifying arginine in numerous proteins. *The Journal of Biological Chemistry* **268**, 10501–10509.
- Paolantonacci P, Lawrence F, Lederer F and Robert-Gero M** (1986) Protein methylation and protein methylases in *Leishmania donovani* and *Leishmania tropica* promastigotes. *Molecular and Biochemical Parasitology* **21**, 47–54.
- Pasternack DA, Sayegh J, Clarke S and Read LK** (2007) Evolutionarily divergent type II protein arginine methyltransferase in *Trypanosoma brucei*. *Eukaryotic Cell* **6**, 1665–1681.
- Pelletier M, Xu Y, Wang X, Zahariev S, Pongor S, Aletta J and Read LK** (2002) Arginine methylation of a mitochondrial guide RNA binding protein from *Trypanosoma brucei*. *Molecular and Biochemical Parasitology* **18**, 49–59.
- Pelletier M, Pasternack DA and Read LK** (2005) *In vitro* and *in vivo* analysis of the major type I arginine methyltransferase from *Trypanosoma brucei*. *Molecular and Biochemical Parasitology* **144**, 206–217.
- Pelletier M, Frainier AS, Munini DN, Wiemer JM, Karpie AR and Sattora JJ** (2013) Identification of a novel lipid homologue from the parasitic protozoan *Trypanosoma brucei*. *BMC Microbiology* **13**, 101.
- Plett KL, Raposo AE, Bullivant S, Anderson IC, Piller SC and Plett JM** (2017) Root morphogenic pathways in *Eucalyptus grandis* are modified by the activity of protein arginine methyltransferases. *BMC Plant Biology* **17**, 62.
- Rakow S, Pullamsetti SS, Bauer U-M and Bouchard C** (2020) Assaying epigenome functions of PRMTs and their substrates. *Methods (San Diego, Calif)* **175**, 53–65.
- Saksouk N, Bhatti MM, Kieffer S, Smith AT, Musset K, Garin J, Sullivan WJ, Cesbron-Delauw M-F and Hakimi M-A** (2005) Histone-modifying complexes regulate gene expression pertinent to the differentiation of the protozoan parasite *Toxoplasma gondii*. *Molecular and Cellular Biology* **25**, 10301–10314.
- Simarro PP, Cecchi G, Franco JR, Paone M, Diarra A, Priotto G, Mattioli RC and Jannin JG** (2015) Monitoring the progress towards the elimination of gambiense human African trypanosomiasis. *PLoS Neglected Tropical Diseases* **9**, e0003785.

- Torres-Guerrero H, Peattie DA and Meza I** (1991) Chromatin organization in *Entamoeba histolytica*. *Molecular and Biochemical Parasitology* **45**, 121–130.
- Tripsianes K, Madl T, Machyna M, Fessas D, Englbrecht C, Fischer U, Neugebauer KM and Sattler M** (2011) Structural basis for dimethylarginine recognition by the Tudor domains of human SMN and SPF30 proteins. *Nature Structural & Molecular Biology* **18**, 1414–1420.
- Venne AS, Kollipara L and Zahedi RP** (2014) The next level of complexity: crosstalk of posttranslational modifications. *Proteomics* **14**, 513–524.
- Walsh JA** (1986) Problems in recognition and diagnosis of amebiasis: estimation of the global magnitude of morbidity and mortality. *Clinical Infectious Diseases* **8**, 228–238.
- Wang C, Zhu Y, Chen J, Li X, Peng J, Chen J, Zou Y, Zhang Z, Jin H, Yang P, Wu J, Niu L, Gong Q, Teng M and Yunyu S** (2014a) Crystal structure of arginine methyltransferase 6 from *Trypanosoma brucei*. *PLoS One* **9**, e87267.
- Wang C, Zhu Y, Caceres TB, Liu L, Peng J, Wang J, Chen J, Chen X, Zhang Z, Zuo X, Gong Q, Teng M, Hevel JM, Wu J and Shi Y** (2014b) Structural determinants for the strict monomethylation activity by *Trypanosoma brucei* protein arginine methyltransferase 7. *Structure (London, England: 1993)* **22**, 756–768.
- Wang Y-C, Wang C-W, Lin W-C, Tsai Y-J, Chang C-P, Lee Y-J, Lin M-J and Li C** (2017) Identification, chromosomal arrangements and expression analyses of the evolutionarily conserved prmt1 gene in chicken in comparison with its vertebrate paralogue prmt8. *PLoS One* **12**, e0185042.
- World Health Organization** (2020) *World Malaria Report 2020: 20 Years of Global Progress and Challenges*. Geneva: World Health Organization.
- Xu W** (2004) A methylation-mediator complex in hormone signaling. *Genes & Development* **18**, 144–156.
- Yakubu RR, Simon de Moterri NC, Nieves E, Kim K and Weiss LM** (2017) Comparative monomethylarginine proteomics suggests that protein arginine methyltransferase 1 (PRMT1) is a significant contributor to arginine monomethylation in *Toxoplasma gondii*. *Molecular & Cellular Proteomics* **16**, 567–580.
- Yang Y and Bedford MT** (2013) Protein arginine methyltransferases and cancer. *Nature Reviews Cancer* **13**, 37–50.
- Zeeshan M, Kaur I, Joy J, Saini E, Paul G, Kaushik A, Dabral S, Mohammed A, Gupta D and Malhotra P** (2017) Proteomic identification and analysis of arginine-methylated proteins of *Plasmodium falciparum* at asexual blood stages. *Journal of Proteome Research* **16**, 368–383.
- Zhang X and Cheng X** (2003) Structure of the predominant protein arginine methyltransferase PRMT1 and analysis of its binding to substrate peptides. *Structure (London, England: 1993)* **11**, 509–520.
- Zhang J and Zheng YG** (2016) SAM/SAH analogs as versatile tools for SAM-dependent methyltransferases. *ACS Chemical Biology* **11**, 583–597.
- Zhang X, Zhou L and Cheng X** (2000) Crystal structure of the conserved core of protein arginine methyltransferase PRMT3. *The EMBO Journal* **19**, 3509–3519.
- Zhao Q, Rank G, Tan YT, Li H, Moritz RL, Simpson RJ, Cerruti L, Curtis DJ, Patel DJ, Allis CD, Cunningham JM and Jane SM** (2009) PRMT5-mediated methylation of histone H4R3 recruits DNMT3A, coupling histone and DNA methylation in gene silencing. *Nature Structural & Molecular Biology* **16**, 304–311.
- Zhao X-X, Zhang Y-B, Ni P-L, Wu Z-L, Yan Y-C and Li Y-P** (2016) Protein arginine methyltransferase 6 (Prmt6) is essential for early zebrafish development through the direct suppression of gadd45a stress sensor gene. *Journal of Biological Chemistry* **291**, 402–412.
- Zhou R, Xie Y, Hu H, Hu G, Patel VS, Zhang J, Yu K, Huang Y, Jiang H, Liang Z, Zheng YG and Luo C** (2015) Molecular mechanism underlying PRMT1 dimerization for SAM binding and methylase activity. *Journal of Chemical Information and Modeling* **55**, 2623–2632.