


## Regular Article

# Callous-unemotional traits and fearlessness: A cardiovascular psychophysiological perspective in two adolescent samples using virtual reality

Nicholas D. Thomson<sup>1</sup> , Steven M. Gillespie<sup>2</sup> and Luna C.M. Centifanti<sup>2</sup>

<sup>1</sup>Division of Acute Care Surgical Services, Department of Surgery, Virginia Commonwealth University Health & Department of Psychology, University of Durham, UK, and <sup>2</sup>Department of Psychological Sciences, Institute of Psychology, Health, and Society, University of Liverpool, Liverpool, UK.

### Abstract

There has been a longstanding debate about the link between callous-unemotional traits and fearlessness. However, biological evidence for a relationship in adolescents is lacking. Using two adolescent samples, we measured emotional reactivity and cardiac measures of sympathetic (pre-ejection period) and parasympathetic (respiratory sinus arrhythmia) reactivity during 3D TV and virtual reality fear induction. Study 1 included 62 community adolescents from a stratified sample. Study 2 included 60 adolescents from Emotional and Behavioral Difficulties schools. Results were consistent across both studies. Adolescents with high callous-unemotional traits showed coactivation of the sympathetic and parasympathetic nervous system. Consistent with these results, youths with callous-unemotional traits self-reported that they felt more in control after the fear induction. Thus, in both samples, youth with callous-unemotional traits displayed a physiological and emotional profile suggesting they maintained control during fear induction. Therefore, it is proposed here that a shift in thinking of youth with callous-unemotional traits as fearless to youth with callous-unemotional traits are better able to manage fearful situations, may be more appropriate.

**Keywords:** Callous-unemotional traits, fearlessness, pre-ejection period, respiratory sinus arrhythmia

(Received 31 January 2019; revised 23 April 2019; accepted 2 June 2019)

The presence of Callous-Unemotional (CU) traits denotes a particular subgroup of children who are characterized by more severe and frequent acts of aggression, greater harm to the victim, and greater use of instrumental or proactive forms of aggressions (see Frick, Ray, Thornton, & Kahn, 2014; Frick & Viding, 2009 for reviews). The importance of CU traits for identifying and understanding this high-risk subgroup of youth has led to its inclusion in the Diagnostic and Statistical Manual of Mental Disorders-5 (DSM-5; American Psychiatric Association, 2013) as a specifier for Conduct Disorder, named as Limited Prosocial Emotions (Fanti et al., 2019). These traits include a lack of remorse or guilt, callous lack of empathy, shallow or deficient affect, and a lack of concern about performance (APA, 2013; Blair, Leibenluft, & Pine, 2014). Theoretically, these characteristics are thought to reflect an absence of the conscious experience of fear, or reduced automatic reactivity to threatening or fear-inducing stimuli (Lykken, 1995). However, this position—commonly termed the “low fear hypothesis”—has been subject to some debate (Newman & Brinkley, 1997). The majority of

research on the low fear explanation has been focused on self-reported feelings of fear, the ability to recognize and understand fear in others (e.g., fearful facial expression recognition), or the capacity to form learned aversive associations between a neutral and threatening stimulus (Hoppenbrouwers, Bulten, & Brazil, 2016). However, few studies have examined the relationship of CU traits with both self-reported fear (i.e., fear that is consciously experienced), as well as physiological reactivity to a fear-inducing stimulus (i.e., automatic reactivity to threat). In this paper, we report the results of two studies that examined both self-reported experiential fear, and psychophysiological indices of threat reactivity, in adolescents with varying levels of CU traits.

Attempts to resolve the problem of whether CU traits are associated with low fear have been faced with the challenges of defining and measuring fear (Hoppenbrouwers et al., 2016). The term fear is most commonly used to refer to the aversive feeling of being afraid when one is in danger, that is, the conscious experience of fear. However, the term fear has also been used with reference to the activation of systems that detect and respond to threats in the environment, yet there is an absence of compelling evidence that this activation is necessarily tied to the conscious experience of fear (LeDoux, 2013). In support of this distinction, it has been reported that conditioned or unconditioned threats presented outside of conscious awareness elicit physiological responses without the person's awareness of the stimulus (Bornemann, Winkielman, & der Meer, 2012; Ohman & Soares,

**Author for Correspondence:** Nicholas D. Thomson, PhD, Division of Acute Care Surgical Services, Department of Surgery, Virginia Commonwealth University Health, Richmond, Virginia, 23298. E-mail: [Nicholas.Thomson@VCUHealth.org](mailto:Nicholas.Thomson@VCUHealth.org)

**Cite this article:** Thomson ND, Gillespie SM, Centifanti LCM (2020). Callous-unemotional traits and fearlessness: A cardiovascular psychophysiological perspective in two adolescent samples using virtual reality. *Development and Psychopathology* 32, 803–815. <https://doi.org/10.1017/S0954579419001196>

1998; Olsson & Phelps, 2004) and without their reporting any particular feeling (Bornemann et al., 2012). A potential solution to this measurement problem is to distinguish between indices of threat reactivity, including visceral responses to threatening stimuli (e.g., changes in autonomic nervous system [ANS] activity), and the conscious experience of fear (e.g., recognizing that one is feeling scared; Hoppenbrouwers et al., 2016).

Studies that have examined the relationships of psychopathic tendencies with participants' feelings of being scared have often used self-report measures. For example, CU traits in youths aged 10–17 years were associated with reductions in the subjective experience of fear, but not other emotions, while children recalled an emotionally evocative life event (Marsh et al., 2011). People with CU traits also seem unaware of the behaviors that make others afraid (Marsh & Cardinale, 2014). Therefore, youth with CU traits may experience low levels of fear as well poor interpretation of fear cues in others. Ratings of fearlessness also appear to vary with stability and change in levels of CU traits and conduct problems (CP) over time, with findings from a longitudinal study showing that teacher reports of fearlessness were highest for children with stable high CP and CU traits, and increases in CU traits were associated with increased fearlessness (Klingzell et al., 2016). Further, children with decreasing CP and CU traits were characterized by decreases in their levels of fearlessness (Klingzell et al., 2016).

To address the limitations associated with self-report measures, psychophysiological techniques have also been employed. These studies have revealed reduced autonomic reactivity among youth with CU traits using a variety of techniques. For example, CU traits were found to be associated with reduced heart rate reactivity to emotionally evocative films (Anastassiou-Hadjicharalambous & Warden, 2008; de Wied, van Boxtel, Matthys, & Meeus, 2012), and reduced skin conductance reactivity to provocation (Kimonis, Frick, Munoz, & Aucoin, 2008; Muñoz, Frick, Kimonis, & Aucoin, 2008) and during a pain procedure (Northover, Thapar, Langley, & van Goozen, 2015). While these findings are valuable for understanding biological markers of CU traits, they are less revealing about autonomic responses to fear in particular. This is, in part, because the stimuli used in these studies have tended to vary in content (e.g., violence, provocation), emotion (e.g., pain, anger), and valence.

More recently, a number of studies by Fanti and colleagues have examined the relationship of CU traits with fear-potentiated startle reflex, a well-established indicator of defensive motivation. These studies have revealed associations of CU traits with fearlessness, and reduced fear-potentiated startle to violent films (Fanti, Panayiotou, Kyranides, & Avraamides, 2016; Kyranides, Fanti, & Panayiotou, 2016) and during fearful mental imagery (Fanti, Panayiotou, Lazarou, Michael, & Georgiou, 2016). However, a study that used the full dimensional scale of psychopathic traits showed that when viewing violent scenes, CU traits were associated with reduced startle potentiation, but only the grandiose-manipulation facet was associated with reduced heart rate reactivity (Fanti et al., 2017). Importantly, CU traits are associated with a reduction in startle reflex even among young adults without CP (Fanti, Panayiotou, Kyranides, et al., 2016). This result indicates that the presence of fearlessness is specific to CU traits in particular, rather than antisociality more generally. Consistent with the findings reviewed here, brain imaging studies have shown that the CU dimension is inversely related to activation of the amygdala, a neuroanatomical region that typically responds to fear-related stimuli, while viewing others' fearful expressions (Dackis,

Rogosch, & Cicchetti, 2015; Jones, Laurens, Herba, Barker, & Viding, 2009; Lozier, Cardinale, VanMeter, & Marsh, 2014; Marsh et al., 2008).

Although there is overwhelming evidence that CU traits are associated with reduced neural and autonomic arousal to emotionally evocative stimuli (Fanti, 2018), the precise mechanisms underlying this pattern of hypoarousal remain unclear. For example, low startle potentiation may not be indicative of low fear, but instead may reflect greater attention to the stimuli (Anthony & Graham, 1985; Bradley, Cuthbert, & Lang, 1990; Patrick, Bradley, & Lang, 1993). The use of heart rate reactivity as a measure of fear is also limited by a need to account for the underlying systems that contribute to changes in the beat-to-beat interval of the heart. The physiological changes that typically accompany emotional responses are mediated by the relative actions of the two branches of the ANS: the sympathetic nervous system (SNS) and the parasympathetic nervous system (PNS). An increase in SNS activity is associated with increases in heart rate and greater expenditure of energy, whereas increases in PNS activity are associated with reductions in heart rate and increased conservation of resources. Because of the interacting effects of the SNS and PNS, failure to account for both SNS and the PNS could lead to inconsistent results, which may explain why some studies have failed to replicate the association of autonomic activity with CU traits (de Wied et al., 2012; Wagner & Abaied, 2015) and antisocial behavior (Calkins & Dedmon, 2000; Pine et al., 1998; Scarpa, Haden, & Tanaka, 2010).

Activation of the SNS and the PNS may be indexed using values of pre-ejection period (PEP) and respiratory sinus arrhythmia (RSA), respectively (Beauchaine, 2001). These measures of autonomic functioning have been used before to gain a better understanding of developmental psychopathology, but the interaction of PEP and RSA in relation to CU traits during fear induction remains poorly understood. PEP refers to a systolic time interval (composed of the electrical-mechanical delay occurring between the onset of depolarization and the beginning of ventricular contraction) and reflects  $\beta$ -adrenergic influences on the heart (Newlin & Levenson, 1979). As such, PEP can be used as an index of SNS activity. SNS reactivity appears to be most notably associated with approach-avoidance tendencies (Beauchaine, Gatzke-Kopp, & Mead, 2007; Brenner, Beauchaine, & Sylvers, 2005), and it has been used as a marker of sensitivity to reward during incentive conditions. Consistent with a role in reward sensitivity, longer PEP durations during reward processing—indicative of lower SNS activity—have been identified among children and adolescents with externalizing behavior disorders (Beauchaine et al., 2007; Beauchaine, Katkin, Strassberg, & Snarr, 2001; Crowell et al., 2006) and aggression (Beauchaine, Hong, & Marsh, 2008). Similar findings have also been observed during emotion induction and emotion regulation among children with low prosocial behavior—used as a proxy for CU traits (Musser, Galloway-Long, Frick, & Nigg, 2013). These findings contribute to a growing evidence base that suggests a reduced sympathetic arousal and lowered sensitivity to reward in those children with aggressive and antisocial behavior problems.

While PEP values index SNS activity, RSA is often used to index parasympathetic influences on the heart; this is mediated by the vagus (10th cranial) nerve (Beauchaine, 2001; Porges, 1995). Theoretically, RSA represents a marker of emotion regulation (Beauchaine, 2015; Thomson, Kiehl, & Bjork, 2018), and several reviews have shown that a reduction in vagally mediated influences on heart are associated with emotion dysregulation

and compromised functioning of emotion regulatory neural circuitry (Gillespie, Brzozowski, & Mitchell, 2018; Thayer, Åhs, Fredrikson, Sollers, & Wager, 2012; Thayer & Lane, 2009). Thus, it is perhaps not surprising that lower levels of RSA are associated with reactive aggression and symptoms of borderline personality disorder, characterized by emotional lability and emotion dysregulation (Thomson & Beauchaine, 2018). More proactive, instrumental types of aggression, on the other hand, appear to be associated with increases in vagally mediated influences on the heart (Brzozowski, Gillespie, Dixon, & Mitchell, 2018; Scarpa et al., 2010; Thomson, Kiehl, et al., 2018). The precise relationship of CU traits with RSA remains unclear, but some studies have reported a negative association between CU traits and RSA at rest (Fanti, 2018).

Physiological systems work dynamically (Porges, 2003, 2007) and exploring interactions between SNS and PNS reactivity may provide a more precise understanding of the relationship between CU traits and fearlessness. Reciprocal SNS activation occurs when both branches act to increase physiological arousal (i.e., an increase in SNS accompanied by PNS withdrawal) and is considered a normative physiological response to dealing with stressful or challenging situations (El-Sheikh et al., 2009). Reciprocal PNS activation, on the other hand, has the net effect of reducing physiological arousal (i.e., PNS activation accompanied by reduced SNS activity). In contrast to reciprocal modes, nonreciprocal patterns of ANS activity can also occur, where joint PNS and SNS activation can act concurrently on the same target organ (Berntson, Cacioppo, & Quigley, 1991). Depending on the relative dominance of the two branches, nonreciprocal modes can yield diametrically opposite responses in the target organ.

Both coinhibition and coactivation have been linked with greater levels of externalizing behaviors in young children (Boyce et al., 2001; Wagner & Abaied, 2015), and it is proposed in the “adaptive calibration model” that unemotional individuals may be characterized by coinhibition to unclear or ambiguous situations but show coactivation in response to immediate threat (Del Giudice, Ellis, & Shirtcliff, 2011; Del Giudice, Hinnant, Ellis, & El-Sheikh, 2012). This coactivation may represent an optimal response to facilitate behavioral and cognitive functioning in high-intensity situations, allowing the individual to be alert and attentive to potential danger, while nonetheless remaining calm and in control (Thomson, Aboutanos, et al., 2018). To date, only one study has examined interactions between the SNS and the PNS during fear induction in relation to psychopathic traits. In a sample of young adults, Thomson, Aboutanos, et al. (2018) found that participants with higher levels of interpersonal and affective psychopathic traits showed coinhibition of the SNS and PNS in response to virtual reality fear induction, coupled with increased self-reported feelings of happiness. In contrast, increasing behavioral and antisocial characteristics were associated with PNS reactivity and reduced feelings of control. The precise pattern of cardio-autonomic activation in relation to the CU dimension in particular remains relatively unknown, and these relationships are yet to be investigated in adolescent samples.

### The present study

The purpose of the present paper was to examine the relation of CU traits with SNS (PEP) and PNS (RSA) activation and consciously experienced fear during fear induction. In Study 1, we tested the relationship of CU traits with fear reactivity among

typically developing adolescents from community schools by using a stratified sampling technique to create two groups that were distinguishable on the presence of high versus low CU traits (see Dadds, El Masry, Wimalaweera, & Guastella, 2008). In Study 2, we examined the continuous relationships of CU traits with fear reactivity in youth with high levels of antisocial behavior by recruiting adolescents with behavioral problems from Emotional and Behavioral Difficulties (EBD) schools. Previous work on physiological reactivity among youth with CP and CU traits has typically used picture, imagery, and movie stimuli (Fanti, 2018) to induce a fearful response. For the first time in an adolescent sample, Study 1 included a 3D rollercoaster experience using a cinema screen, and Study 2 included a novel three-dimensional (3D) virtual reality (VR) roller coaster simulation, each providing a more ecologically valid experience of fear. Because this is the first study to date to test cardiac SNS and PNS reactivity to fear in adolescents with CU traits, we made two competing hypotheses. Hypothesis one was based on the predictions of Del Giudice et al. (2011, 2012) that unemotional individuals would show coactivation (i.e., SNS and PNS reactivity) under immediate threat. Thus, a positive relationship of CU traits with both SNS and PNS reactivity, and with self-reported feelings of being in control, would be expected under fear induction. Alternatively, affective and interpersonal psychopathic traits have been linked with coinhibition during a VR horror game in young adults (Thomson, Aboutanos, et al., 2018). Based on these results, we developed a competing hypothesis that CU traits, while under fear induction, would be inversely associated with both SNS and PNS reactivity and with feelings of being in control. The findings of these studies will help to clarify the association of CU traits with self-reported fearlessness and with psychophysiological indices of arousal during fear induction. Understanding these relationships will set the stage for future research to understand the causal relations between CU traits, fearlessness, and CP, and for developing interventions directed at reducing CU traits.

### Study 1

Study 1 was designed to assess the link between CU traits and fear reactivity in typically developing adolescents. The community sample was selected to represent adolescents without serious antisocial behavior problems. Consistent with prior research, a stratified sampling method was used to compare adolescents who displayed the highest (within the top 20%) and lowest (within the bottom 20%) levels of CU traits within the community (see Dadds, El Masry, Wimalaweera, & Guastella, 2008). The objective was to assess whether adolescents with high CU traits were fearless at the physiological and emotional level.

### Method

#### Procedure

Six hundred and ninety-six adolescents aged between 12 to 14 years from three large community schools in the North East of England were screened on the Inventory of Callous-Unemotional Traits (ICU). Participants completed the questionnaire at school within classrooms. A stratified sampling technique was used to recruit adolescents who were high (top 20%) and low (lowest 20%) in CU traits. Based on these scores, participants' parents/caregivers were invited to bring their child to complete the laboratory-based part of the study at the university 3D laboratory. From the 155

participants who were invited, 62 participants accepted the invitation. Both parent/caregiver and participants completed questionnaires before the experiment. This accommodated a stabilization period for the physiological measures. Next, participants were asked to sit and relax for a three-minute rest period (baseline condition). Participants wore 3D glasses for the 90 second roller coaster, and the six-minute space documentary (control condition). Participants completed each condition in the same order: (a) baseline (rest period), (b) fear induction (3D roller coaster), and (c) vanilla baseline (3D space documentary). After each condition, participants reported their emotional state using the SAM. Participants received a gift voucher for completing the study, and parents/caregivers were compensated for travel expenses.

### Participants

Sixty-two adolescents were included in the final experiment (low CU group  $n = 35$ , high CU group  $n = 27$ ). Participants were predominantly male ( $n = 53$ ), White British (89%), and aged between 12 and 14 years old ( $M = 12.54$ ,  $SD = .57$ ). Minority ethnicities included White other ( $n = 3$ ), mixed ( $n = 1$ ), African ( $n = 1$ ), and Bangladeshi ( $n = 1$ ). Seventy-nine percent of the sample was raised by both biological parents, 6.5% by biological mother and step father, 6.5% by biological mother alone, 3.2% biological father alone, and the remaining 4.8% included participants who were raised in a shared parental custody (3.2%) or by a guardian (1.6%).

### Measures

#### *Callous-unemotional traits*

The Inventory of Callous and Unemotional Traits (ICU; Frick, 2004) is a 24-item self-report scale designed to measure callous and unemotional traits in youth. Items are scored on a 4-point Likert scale from *Not at all true* (0) to *Definitely true* (3). The ICU is a valid measure of CU traits and has been widely used in community and incarcerated samples of youth (see Pihet, Etter, Schmid, & Kimonis, 2015). In the present study, the parent- ( $\alpha = .88$ ) and self-report ( $\alpha = .90$ ) yielded good internal consistency.

#### *Conduct problems and prosocial behavior*

Parent report of the Strengths and Difficulties Questionnaire (SDQ; Goodman, 1997) was administered to assess group differences in conduct problems and prosocial behavior. The SDQ items are scored from *Not true* (0) to *Certainly true* (2). The conduct problems and prosocial scales include five items. In the present sample, the internal consistency was poor for the conduct problems ( $\alpha = .45$ ) and acceptable for the prosocial scale ( $\alpha = .75$ ).

#### *Fear-inducing environment*

To safely measure emotional and physiological reactivity to fear, participants experienced a 90 second 3D roller coaster. The 3D roller coaster simulation video traverses mountains with steep drops and turns. One of the dips/valleys was determined by the computer scientists to be physiologically impossible for a human to withstand since the positive forces of gravity would be extreme at the lowest point. The control condition was the award-winning six-minute "Our Cosmic Origins" (Holliman, 2010) space documentary. All films were produced at the Durham Visualization Laboratory. The 3D videos were viewed on a 2.4 m rear projected PASCAD low-crosstalk screen (using a BARCO Gemini stereoscopic projection display). Participants wore lightweight glasses

during the videos. The BARCO display is linked to wireless devices, which allows 3D interaction and head tracking.

#### *ANS Fear reactivity and physiological data acquisition*

Two Ag-AgCl electrocardiogram electrodes in a modified Lead II configuration, and eight Ag-AgCl impedance cardiogram paired electrodes on the neck and torso (with at least a 3-cm distance between the paired electrodes as recommended; Sherwood et al., 1990) were placed on the participant. Respiration was recorded using RSPEC-R amplifier with a wireless respiration belt transducer. To ensure the belt was placed at maximum point of sensitivity, the participant was asked to exhale, at full exhalation the respiration belt was fastened around the abdomen of the participant. Data were recorded using Biopac MP150 with BioNomadix module transmitter (MP150-BIOPAC Systems Inc., Goleta, CA), and sampled at 1000 Hz. Data were reduced and analyzed offline, using the Biopac's Acknowledge 4.3 software. Data were visually inspected for motion artifacts and outliers. Electrocardiogram and impedance cardiography were reduced offline and the waves were coded using computer-aided event detection, but modified by visual inspection so that mid-beats were created if missing (<.001%) and errors in R-wave detection were adjusted. To compensate for fluctuations due to movement, the electrocardiogram was reduced at 250 Hz and respiration was passed through a .5 Hz digital band filter. Pre-ejection period was calculated from the time between the onset of the Q wave of the ECG to the B point of the dZ/dt waveform (i.e., beginning of ejection). RSA was computed using AcqKnowledge automated function for RSA analysis, which applies the validated peak-valley method (Grossman, van Beek, & Wientjes, 1990). RSA values reflect the millisecond difference between the minimum and maximum R-R intervals during each respiration cycle.

Participants completed the following conditions in the same order: Resting baseline, 3D roller coaster, vanilla baseline which was the 3D space documentary (which we will call our control condition). We used the vanilla baseline as a comparison with the 3D roller coaster, because several studies show that a vanilla baseline is a better comparator for calculating psychophysiological reactivity than a resting baseline (Hastrup, 1986; Jennings, Kamarck, Stewart, Eddy, & Johnson, 1992; Piferi, Kline, Younger, & Lawler, 2000). Also, due to requests by our Psychology Ethics Subcommittee, the vanilla baseline was included last to allow for a relaxing condition to directly follow the fear-induction condition.

The measures of physiological reactivity were change scores calculated using RSA and PEP levels during the control condition and the roller coaster condition. Our primary measure of physiological reactivity was calculated for RSA by subtracting control task levels from roller coaster task levels, so higher scores indicated *increases* in PNS. PEP was calculated by subtracting roller coaster task levels from control task levels, so higher scores indicated *increases* in SNS. Thus, higher scores indicated *increases* in PNS and SNS reactivity, while lower scores indicated *low* SNS and PNS reactivity. We elected to use delta change in cardiovascular reactivity scores for three primary reasons. First, they are easily interpreted. Second, they have been found to be reliable across time and have been found to be as reliable as residualized change scores. Third, they can be compared with reactivity reported in other studies (Boyce et al., 2001; Thomson, Aboutanos, et al., 2018).



### Arousal and emotional reactivity

To assess self-report of arousal and valence to the roller coaster participants were asked to report on a nine-point scale how they felt after each condition using the Self-Assessment Manikin (SAM; Bradley & Lang, 1994). The SAM is a nonverbal pictographic scale designed to assess feelings across emotional dimensions. The valence scale ranges from a manikin who is *smiley and happy* (1) to *frowning and unhappy* (9). The arousal scale ranges from a manikin who looks *excited and wide-eyed* (1) to *relaxed and sleepy* (9). The dominance scale ranges from feeling *small and out of control* (1) to *in control* (9). Because reactivity was of interest, scores were computed by subtracting control condition averages from roller coaster averages. On the arousal scale, positive numbers represented feeling more relaxed and negative values indicated feeling more excited. On the valence scale, a negative value was indicative of feeling *more happy* and a positive value was indicative of feeling *less happy and more sad*. A negative value on the dominance scale is indicative of feeling *less in control* and a positive value indicated feeling *more in control*.

## Results

### Data analysis plan

To test if the stratified groups, which were selected to be different on CU traits, were also different on behavioral and emotional symptoms, independent samples *t* tests were conducted on parent-report of conduct problems and prosocial behavior. Responses to the rollercoaster videos were assessed as a test of the validity of these tasks and this was done with paired-samples *t* tests. The *t* tests were examined to determine if the change in arousal ratings and autonomic reactivity to the rollercoasters differed by CU group. Because we were interested in the interaction term between SNS and PNS statistically predicting CU traits, we conducted logistic regression. Thus, we attempted to determine whether mapping the autonomic space (via an interaction between parasympathetic and sympathetic activity) would assist in classification of the high CU group. This was done on the standardized change scores of RSA and PEP, with higher values indicating greater parasympathetic activation (roller coaster RSA — control RSA) and greater sympathetic activation (control PEP—roller coaster PEP). The same was done for self-report of valence, arousal, and dominance change scores. Logistic regressions were chosen for this study because this cohort was stratified, so we predicted group membership of high or low CU grouping.

### Stratification on self-report: Examining parent-reports

To test if the stratified groups were different on behavioral and emotional symptoms, independent samples *t* tests were conducted on parent-report of CU traits, conduct problems, and prosocial behavior. Group differences on sex and age were assessed using chi-square and a *t* test, respectively. The high CU group ( $n = 27$ ) was significantly higher on parent-reports of CU traits,  $M = 24.46$ ,  $SD = 9.61$ ;  $t(36) = -4.75$ ,  $p < .001$ , Cohen  $d = -1.32$ , and conduct problems,  $M = 2.00$ ,  $SD = 1.39$ ;  $t(60) = -2.11$ ,  $p = .039$ , Cohen  $d = -0.53$ , and lower on prosocial behaviors,  $M = 7.00$ ,  $SD = 2.42$ ;  $t(34) = 3.76$ ,  $p = .001$ , Cohen  $d = 1.11$ , than the low CU group,  $n = 35$ ;  $M = 13.78$ ,  $SD = 5.63$ ;  $M = 1.29$ ,  $SD = 1.27$ ;  $M = 8.89$ ,  $SD = 1.11$ , respectively. The CU groups did not significantly differ on age,  $t(55) = -.18$ ,  $p = .859$ , Cohen  $d = 0.05$ , or sex,  $\chi^2(1) = 1.07$ ,  $p = .302$ . Overall, when compared with the low CU

group, members of the high CU group were perceived by their parents as being high in CU traits, having a greater level of conduct problems, and having a lower level of prosocial behaviors.

### ANS and emotional reactivity to fear

Paired samples *t* tests were conducted on raw score values of the levels of RSA and PEP to establish whether the 3D roller coaster induced an autonomic response when compared with the control video (space documentary). We also examined the difference in levels of RSA and PEP between the resting baseline and the 3D roller coaster and the control 3D video. Mean level of PEP,  $t(61) = 4.02$ ,  $p < .001$ , was lower during the roller coaster ( $M = .130$ ,  $SD = .02$ ) than during the control condition,  $M = .136$ ,  $SD = .02$ ; Cohen  $d = .32$ , and baseline condition,  $M = .133$ ,  $SD = .02$ ;  $t(61) = 2.72$ ,  $p = .009$ , Cohen  $d = .22$ . Mean level of RSA was lower during the roller coaster condition ( $M = 4.57$ ,  $SD = .55$ ) than the control condition,  $M = 4.69$ ,  $SD = .59$ ;  $t(60) = -3.25$ ,  $p = .002$ , Cohen  $d = .35$ , and the baseline condition,  $M = 4.76$ ,  $SD = .59$ ;  $t(61) = -5.28$ ,  $p < .001$ , Cohen  $d = .59$ . PEP was higher during the control condition than during the resting baseline condition,  $t(62) = -2.49$ ,  $p = .016$ , Cohen  $d = -.33$ , suggesting less sympathetic nervous system activity during the control condition (i.e., vanilla baseline) than the resting baseline at the start of the testing session. There was no significant difference in RSA between the control condition and the baseline condition,  $t(60) = -1.36$ ,  $p < .178$ , Cohen  $d = .03$ . Overall, compared with the control condition, the roller coaster was effective at inducing sympathetic activation (shortened PEP) and withdrawal in parasympathetic activity (RSA), suggesting that the roller coaster induced reciprocal sympathetic activation (high SNS and low PNS).

Paired samples *t* tests were conducted on the complete sample to assess whether the roller coaster was valid at influencing emotional feelings on the arousal (feeling excited or relaxed), valence (happy or unhappy), and dominance (in control or out of control) scale of the self-assessment manikin. Reactivity was measured in terms of self-assessment after the roller coaster compared with the control condition. Participants reported feeling more excited,  $t(64) = 7.79$ ,  $p < .001$ , Cohen  $d = .95$ , after the roller coaster ( $M = 4.20$ ,  $SD = 2.46$ ) than did participants in the control condition ( $M = 6.17$ ,  $SD = 2.36$ ). Levels of valence were similar,  $t(64) = .72$ ,  $p = .472$ , Cohen  $d = .09$ , after the roller coaster ( $M = 2.55$ ,  $SD = 1.33$ ) and after the control condition ( $M = 2.40$ ,  $SD = 1.51$ ). Participants reported feeling less in control,  $t(63) = 2.73$ ,  $p = .008$ , Cohen  $d = .32$ , after the roller coaster ( $M = 5.75$ ,  $SD = 1.86$ ) when compared with the control condition ( $M = 6.33$ ,  $SD = 1.62$ ). In sum, participants found the roller coaster to increase the feeling of excitement and a loss of control. However, the roller coaster did not have a significant effect on feeling happy or sad.

### CU groups and ANS reactivity to fear


A hierarchical logistic regression was performed to assess whether ANS reactivity predicted CU group membership. Odds ratios reflect the odds likelihood of being in one group over the other, based on the level of the independent variable. Because of differences in scaling of PEP and RSA, these scores were normalized by transforming values to *z*-scores.

Step 1 included age, sex, and conduct problems, Step 2 added RSA and PEP, and Step 3 included the interaction term between

**Table 1.** ANS indices predicting CU groups (1 = High) in community sample

|               | <i>B</i> | <i>SE</i> | <i>z</i> value | OR   |      | CI | 2LL    |
|---------------|----------|-----------|----------------|------|------|----|--------|
| <b>Step 1</b> |          |           |                |      |      |    | 38.70  |
| Age           | −0.01    | 0.49      | −0.03          | 0.99 | 0.36 | –  | 2.62   |
| Sex           | −1.21    | 0.88      | −1.37          | 0.29 | 0.04 | –  | 1.49   |
| CP            | 0.40     | 0.21      | 1.95           | 1.49 | 1.01 | –  | 2.29   |
| <b>Step 2</b> |          |           |                |      |      |    | 37.22  |
| Age           | −0.03    | 0.50      | 0.07           | 1.04 | 0.38 | –  | 2.79   |
| Sex           | −1.37    | 0.91      | −1.51          | 0.25 | 0.03 | –  | 1.32   |
| CP            | 0.29     | 0.22      | 1.35           | 1.34 | 0.88 | –  | 2.10   |
| ΔRSA          | −0.20    | 0.28      | −0.73          | 0.81 | 0.46 | –  | 1.40   |
| ΔPEP          | 0.46     | 0.45      | 1.03           | 1.59 | 0.80 | –  | 4.57   |
| <b>Step 3</b> |          |           |                |      |      |    | 34.56* |
| Age           | 0.25     | 0.54      | 0.46           | 1.28 | 0.44 | –  | 3.75   |
| Sex           | −1.77    | 0.92      | −1.92          | 0.17 | 0.02 | –  | 0.91   |
| CP            | 0.32     | 0.23      | 1.39           | 1.37 | 0.89 | –  | 2.19   |
| ΔRSA          | −0.02    | 0.29      | −0.07          | 0.98 | 0.54 | –  | 1.75   |
| ΔPEP          | 1.27*    | 0.58      | 2.19           | 3.57 | 1.29 | –  | 13.35  |
| ΔRSA x ΔPEP   | 1.21*    | 0.58      | 2.07           | 3.36 | 1.19 | –  | 12.49  |

Note: Sex (0 = male, 1 = female); CP = Conduct Problems; ΔRSA = respiratory sinus arrhythmia reactivity; ΔPEP = pre-ejection period reactivity; CI = 95% confidence interval. \* $p < .05$



**Figure 1.** Probability of being in the high CU group based on the interaction between SNS and PNS reactivity.

RSA and PEP. Results are displayed in Table 1. Steps 1 ( $-2LL = 38.70$ ;  $\chi^2(3) = 6.35$ ,  $p = .10$ ) and 2 ( $-2LL = 37.21$ ;  $\chi^2(5) = 8.15$ ,  $p = .15$ ) were not significantly better fitting than the null model. RSA ( $p = .46$ ) and PEP ( $p = .30$ ) were not significant predictors. Step 3 was significantly better than both the null model ( $-2LL = 34.56$ ;  $\chi^2(6) = 13.46$ ,  $p = .04$ ) and Step 2 ( $p = .02$ ). The interaction term between RSA and PEP was positive and significant (OR = 3.36, 95% CI = [1.19, 12.49],  $p = .038$ ). Figure 1 shows that high SNS reactivity (increased SNS to the roller coaster compared with the control video greater than or equal to +1 SD) and high PNS reactivity (increased PNS to the roller coaster compared with the control video greater than or equal to +1 SD) increased the probability of being in the high CU group. Thus, coactivation of the SNS and PNS during the roller coaster condition is

associated with a three times greater likelihood of being in the high CU group. To assess whether members of the high CU group were low in reactivity, post hoc  $t$  tests were conducted to test for differences in absolute change scores. The high CU group did not significantly differ in RSA,  $M = .10$ ,  $SD = .33$ ;  $t(59) = .82$ ,  $p = .41$ , Cohen  $d = 0.20$ , or PEP reactivity,  $M = .007$ ,  $SD = .01$ ;  $t(59) = -1.56$ ,  $p = .12$ , Cohen  $d = 0.21$ , from the low CU group,  $M = .17$ ,  $SD = .35$ ;  $M = .003$ ,  $SD = .01$ , respectively; both showed small effect sizes. Thus, members of the high CU group were not characterized by generally lowered reactivity.

### CU Groups and emotional fear reactivity

A series of logistic regressions were conducted to determine if self-report of emotional reactivity to the roller coaster increased the likelihood of being in the high CU group. In all analyses age, sex, and conduct problems were included. Arousal,  $\beta = .10$ ,  $SE = .13$ , OR = 1.10, 95% CI = [.85, 1.45],  $p = .469$ , and valence,  $\beta = -.23$ ,  $SE = .17$ , OR = .79, 95% CI = [.55, 1.09],  $p = .169$ , were nonsignificant predictors of group membership. However, dominance was significant,  $\beta = .37$ ,  $SE = .19$ , OR = 1.45, 95% CI = [1.02, 2.16],  $p = .048$ . These findings suggest adolescents who reported remaining in emotional control and dominant during the roller coaster were more likely to be higher in CU traits. However, the levels of arousal or valence did not differentiate the high and low CU groups.

### Discussion

The results of Study 1 showed that coactivation of the ANS distinguished the high CU group from the low CU group of community adolescents. Thus, high CU adolescents responded to fear with greater sympathetic and parasympathetic activity,

which may be indicative of maintaining alertness and control (Del Giudice et al., 2012). Conscious fear, measured by emotional reactivity, differentiated those in the high or low CU groups only for the dominance scale, not for arousal or valence. Thus, the high CU group felt more in control after the fear induction than the low CU group. In sum, community adolescents high in CU traits displayed a biological profile that is indicative of being able to remain calm and alert while experiencing fear. This may give them the appearance of being fearless, and it explains their feelings of remaining in control after fear induction.

## Study 2

The purpose of Study 2 was to assess the replicability of Study 1 in a sample of adolescents with high levels of emotional and behavioral difficulties. By selecting a high-risk and a community sample of adolescents we sought to understand whether the relation between CU traits and fear reactivity in a community sample was replicable in a sample with conduct problems and antisocial behavior.

## Method

### Participants

Sixty adolescents were recruited from Emotional and Behavioral Difficulties (EBD) schools. Participants were predominantly male ( $n = 50$ ), White British (96%), and aged between 11 and 16 ( $M = 13.95$ ,  $SD = 1.31$ ). Based on school records, 23% had lived in care, 34% had a history of abuse, 52% had a diagnosis of Attention Deficit Hyperactivity Disorder (ADHD), 5% with Autism Spectrum Disorder, 5% with depression and a history of self-harm, and 2% with Reactive Attachment Disorder. Participants' legal caregivers gave consent for the participant, and the participant assented to be involved in the study.

### Procedure

Four EBD schools in the North East of England were included in the recruitment process. Each school varied on recruitment success rate (71%, 87%, 60%, and 42%), this was due to availability of the pupils and the number of pupils on the role at each school (from eight pupils to 78). Information sheets and consent forms were sent home to caregivers, and only those who had returned a signed consent form were allowed to participate in the study. The experiment took place in a quiet room within the school. Self-report questionnaires were completed by the participant prior to the experiment to accommodate a stabilization period for the physiological measures. Participants completed each condition in the same order: (a) baseline (rest period without VR), (b) fear induction (VR roller coaster), and (c) vanilla baseline or "control" condition (VR garden). First, participants completed a three-minute rest period where they were asked to sit still and try to relax. After the rest period participants reported how they felt using the Self-Assessment Manikin (SAM; Bradley & Lang, 1994). The VR headset was then placed on the participant's head, at which point the child was asked to describe the VR surroundings. This allowed the participant to become familiar with the surroundings (e.g., sitting on a roller coaster) and confirm their ability to see the display. The roller coaster lasted for 90 seconds, at which point the participant was asked to report how they

felt using the SAM. Next, participants were introduced to a control resting condition, which was a sunny garden set in Tuscany, Italy. Participants were asked to sit still and relax for three minutes. Participants wore headphones during the roller coaster and control condition. After the control condition participants reported how they were feeling using the SAM. All participants received a chocolate bar for completing the study.

## Measures

### Callous-unemotional Traits

Consistent with Study 1, CU traits were measured using the Inventory of Callous and Unemotional Traits (ICU; Frick, 2004). In the present study, the ICU yielded good internal consistency ( $\alpha = .82$ ).

### Fear-inducing environment

To safely measure emotional and physiological reactivity to fear, participants experienced a 90 second VR roller coaster. A roller coaster was selected as it is age and ethically appropriate to administer to children and adolescents for inducing a fearful response. Furthermore, in support of a roller coaster being a marker for fear (or fearlessness), the most widely used self-report measures often include roller coaster items (see the Fear Survey Schedule: Geer, 1965; and the Situated Fear Questionnaire: Campbell et al., 2016). The virtual reality headset, the Oculus Rift, has an 18-cm 3D screen (allowing for 100 degrees of direct view) with low latency 360 degree head tracking capabilities. The headset is comfortable and lightweight, which makes the headset suitable for ages seven years and up. Participants wore noise cancelling headphones while wearing the headset. The roller coaster video (RiftCoaster; Oculus VR, 2013) lasted for 90 seconds, with steep drops, tunnels, turns, and jumps. The video was designed specifically for the use with the Oculus Rift. As a control condition (VR equivalent of a baseline) participants were "sat" in a VR garden based in Tuscany, Italy (Tuscany Demo; Oculus VR, 2013). Participants experienced the control condition for three minutes.

### Physiological data acquisition

The same method (and equipment) was employed from Study 1 for ANS data acquisition, reduction, and measure of ANS reactivity. Electrocardiogram and impedance cardiography were visually inspected so that mid-beats were created if missing (<.001%) and errors in R-wave detection were adjusted. PEP and RSA reactivity was computed so higher values indicated greater reactivity from the control condition. See Study 1 for full details.

### Arousal and emotional reactivity

As with Study 1, the Self-Assessment Manikin (SAM; Bradley & Lang, 1994) was used after each condition to measure valence, arousal, and dominance.

## Results

### Data analysis plan

Reactivity to the VR roller coaster was assessed as a test of validity. We conducted paired-samples *t* tests on ANS reactivity and self-reported emotional reactivity (i.e., valence, arousal, and dominance). Our main objective was to understand the physiological profiles of CU traits, which included the interaction term between

PNS and SNS activity. Therefore, we conducted hierarchical linear regressions to determine whether mapping the autonomic space (via an interaction between parasympathetic and sympathetic activity) would statistically predict levels of CU traits. This was done on the standardized change scores of RSA and PEP, with higher values indicating greater parasympathetic activation (roller coaster RSA–control RSA) and greater sympathetic activation (control PEP–roller coaster PEP). The same was done for self-report of valence, arousal, and dominance change scores.

### ANS and emotional reactivity to fear

To establish whether the VR roller coaster was valid at inducing an autonomic response, paired samples *t* tests were conducted comparing the raw score values of levels of RSA and PEP during the roller coaster ride with the span of time of the control condition (VR garden video). Again, we examined the difference in levels during the resting baseline and the roller coaster and control condition. For the entire sample, mean levels were significantly lower for RSA,  $t(59) = 2.25, p = .028$ , Cohen  $d = .26$ , during the roller coaster ( $M = 4.19, SD = .74$ ) when compared with RSA during the control condition ( $M = 4.34, SD = .63$ ) and resting baseline,  $M = 4.55, SD = .68; t(59) = 4.19, p < .001$ , Cohen  $d = .53$ . Mean levels of PEP were lower during the roller coaster ( $M = .137, SD = .02$ ) than during the control condition,  $M = .444, SD = .01; t(59) = 6.69, p < .001$ , Cohen  $d = .77$ , and resting baseline,  $M = .140, SD = .01; t(59) = 3.63, p = .001$ , Cohen  $d = .50$ . RSA was higher in the baseline condition than in the control condition,  $t(59) = 3.46, p = .001$ , Cohen  $d = .43$ , suggesting higher PNS activity at rest than during the control condition. PEP was lower in the baseline condition than in the control condition,  $t(59) = -6.23, p < .001$ , Cohen  $d = .83$ , suggesting higher SNS activity at rest than during the control condition.

Overall, compared with the control condition, the roller coaster induced reactivity on both PEP and RSA. Based on these results, the roller coaster produced sympathetic activation (lower PEP) and withdrawal in parasympathetic activity (lower RSA), suggesting that the roller coaster induced reciprocal sympathetic activation (high SNS and low PNS), which is considered the most common physiological response to dealing with challenging situations (El-Sheikh et al., 2009).

To establish whether the VR roller coaster was valid at inducing emotional feelings, such as arousal (feeling excited or relaxed), valence (happy or unhappy), or dominance (in control or out of control), paired samples *t* tests were conducted on the entire sample comparing SAM after the roller coaster with the control condition. Reported arousal levels were higher,  $t(49) = 8.69, p < .001$ , Cohen  $d = 1.15$ , after the roller coaster ( $M = 3.73, SD = 2.74$ ) than after the control condition ( $M = 7.46, SD = 2.28$ ), with participants reporting feeling more excited after the roller coaster. Levels of valence were similar,  $t(49) = 1.02, p = .315$ , Cohen  $d = .19$ , after the roller coaster ( $M = 1.75, SD = 1.13$ ) than after the control condition ( $M = 2.00, SD = 1.71$ ). Therefore, the roller coaster did not make the participant feel more or less happy. On the dominance scale, participants felt less in control,  $t(49) = 2.36, p = .022$ , Cohen  $d = .37$ , after the roller coaster ( $M = 6.67, SD = 2.34$ ), than after the control condition ( $M = 7.54, SD = 1.74$ ). Overall, the roller coaster did not affect adolescents' feeling happy or sad, but it did make them feel more excited and less in control.

**Table 2.** Summary of hierarchical regression for CU traits: The predictive effects of RSA and PEP

|                                    | $\beta$ | SE   | <i>t</i> | $\Delta R^2$ |
|------------------------------------|---------|------|----------|--------------|
| <b>Step 1</b>                      |         |      |          | .013         |
| Age                                | 0.09    | 0.94 | 0.68     |              |
| Sex                                | 0.07    | 3.28 | 0.49     |              |
| <b>Step 2</b>                      |         |      |          | .051         |
| Age                                | -0.02   | 1.04 | -0.12    |              |
| Sex                                | 0.08    | 3.27 | 0.60     |              |
| $\Delta$ RSA                       | 0.25    | 1.44 | 1.63     |              |
| $\Delta$ PEP                       | 0.00    | 1.34 | 0.03     |              |
| <b>Step 3</b>                      |         |      |          | .150*        |
| Age                                | 0.11    | 1.00 | 0.80     |              |
| Sex                                | 0.23    | 3.24 | 1.77     |              |
| $\Delta$ RSA                       | 0.01    | 1.51 | 0.04     |              |
| $\Delta$ PEP                       | 0.02    | 1.24 | 0.18     |              |
| $\Delta$ RSA $\times$ $\Delta$ PEP | 0.48*   | 0.91 | 3.21     |              |

Note: Sex (0 = male, 1 = female);  $\Delta$ RSA = respiratory sinus arrhythmia reactivity;  $\Delta$ PEP = pre-ejection period reactivity; \* $p < .01$ .


### CU Traits and ANS Reactivity to Fear

To test whether ANS reactivity predicted CU traits, a hierarchical multiple regression was performed with R (R Core Team, 2016). Post hoc testing of the significant interaction term was completed using simple slopes analysis in accordance to procedures described by Aiken and West (1991), using Pequod Package (Mirisola & Seta, 2011). Collinearity diagnostic tests were conducted to ensure regression analyses were not affected by multicollinearity (tolerance  $> .10$  and variance inflation factor  $< 10$ ; Tabachnick & Fidell, 2013). Tolerance values  $> 0.65$  and a variance inflation factor  $< 1.54$  were well within the acceptable range. The residuals scatter plots indicated the assumption of linearity and normality were not violated for all the regression models (Tabachnick & Fidell, 2013). Because of differences in scaling of PEP and RSA, these scores were normalized by transforming values to *z*-scores, which is consistent with prior research (Berntson, Norman, Hawley, & Cacioppo, 2008; Bylsma et al., 2015; Crowell et al., 2006).

Results of the hierarchical regression are displayed in Table 2. Step 1 included age and sex, but was not significant,  $F(2, 57) = .38, p = .683$ . Step 2 included age, sex, RSA, and PEP, but was not significant,  $F(4, 55) = .95, p = .445$ . Step 3, which included all the predictors from Step 2 and the interaction between RSA and PEP, was significant,  $F(5, 54) = 2.94, p = .020$ . The interaction between RSA and PEP reactivity was significant,  $\beta = .48, SE = .91, t = 3.21, p = .002$ . Examination of the interaction using simple slopes analysis revealed that CU traits was predicted by high (+1 SD) RSA reactivity and high (+1 SD) PEP reactivity,  $\beta = 2.96, p = .03$ . Lower levels (-1 SD) of RSA and lower levels (-1 SD) of PEP reactivity was not significant,  $\beta = -2.85, p = .18^1$ .

<sup>1</sup>To address the concern that ADHD or history of abuse may have influenced the findings, we have provided results including ADHD and history of abuse as covariates. Step 1 ( $F(4, 54) = .31, p = .871$ ; ADHD  $p = .676$ ; Abuse  $p = .839$ ) and step 2 ( $F(6, 52) = .73$ ,





**Figure 2.** Interaction between SNS and PNS reactivity prediction CU Traits.

Therefore, a coactivated parasympathetic and sympathetic nervous system during fear induction was associated with high levels of CU traits (see Figure 2).

### CU traits and reported fear reactivity

To test whether self-reported fear reactivity (self-report reactivity of valence, dominance, and arousal to fear induction) predicted higher levels of CU traits, a series of multiple regressions were conducted. Consistent with the ANS analyses, the regressions included change scores between the control condition and the roller coaster. The models including arousal,  $F(3, 49) = 2.03$ ,  $p = .123$ , and valence,  $F(3, 49) = .86$ ,  $p = .471$ , were not significant. However, the model including dominance was significant,  $F(3, 49) = 2.81$ ,  $p = .049$ . Dominance was the only predictor of CU traits,  $\beta = .48$ ,  $SE = .91$ ,  $t = 3.21$ ,  $p = .002$ . Thus, youth who felt more in control during fear induction showed the highest levels of CU traits.

### Discussion

The results show that adolescents with high levels of CU traits display coactivation of SNS and PNS during fear induction. Coactivation occurs in a minority of people and has been said to characterize those individuals who show an unemotional predisposition (Alkon et al., 2003; Del Giudice et al., 2011). The present findings support this assertion and show that coactivation characterizes adolescents with high levels of CU traits. Thus, adolescents who are high in CU traits appear to biologically respond to fear in a way that enables them to maintain “tight self-control” (Del Giudice et al., 2011 p.19). The only significant finding for conscious (or self-reported) fear was dominance, which suggests that adolescents who felt more in control after the roller coaster had higher levels of CU traits. The results demonstrate that adolescents with CU traits are biologically different in their autonomic and emotional response to fear.

$p = .622$ ) were not significant. Step 3 ( $F(7, 51) = 2.22$ ,  $p = .048$ ) and the interaction between RSA and PEP was significant ( $\beta = .49$ ,  $SE = .92$ ,  $t = 3.21$ ,  $p = .002$ ). Simple slopes analysis showed the interaction was significant for high PEP and high RSA ( $p = .036$ ), but not low PEP and low RSA ( $p = .200$ ). Therefore, the results were consistent with the primary analyses.

### General Discussion

Children with CU traits are described as being fearless (Fanti, Panayiotou, Lazarou, et al., 2016; Frick & Morris, 2004; Pardini, 2006), and to some degree the present findings support this assertion for adolescents. By employing cardiac measures of SNS and PNS reactivity, a consistent autonomic pattern of fear reactivity was established for adolescents high in CU traits from two different samples. That is, adolescents high in CU traits, regardless of their severity of emotional and behavioral problems, displayed coactivation of the sympathetic and parasympathetic divisions of the ANS while experiencing fear. Therefore, elevated CU traits were associated with physiological reactivity to fear, but not in a manner that is considered normative (e.g., reciprocal sympathetic activity; El-Sheikh et al., 2009). In accordance with the adaptive calibration model, coactivation of the PNS and SNS during frightening or high-risk situations may give the appearance of the individual being less afraid (see Del Giudice et al., 2011), and this may explain why youth with CU traits typically present as fearless (Fanti, Panayiotou, Lazarou, et al., 2016; Frick & Viding, 2009).

Coactivation is considered an optimal response during high-intensity situations, allowing for increased behavioral and cognitive functioning (Allison et al., 2012). For example, during high intensity situations the individual may be alert and attentive (facilitated by higher sympathetic activity) while being able to remain calm and in control of the situation (facilitated by increase in parasympathetic activity; Allison et al., 2012; Del Giudice et al., 2011, 2012). In other words, coactivation may help the individual to remain in control during frightening situations. If, however, the situation escalates and requires an immediate response, the parasympathetic “brake” on the heart is withdrawn, allowing for a fuller expression of sympathetic activity, resulting in an explosive response to deal with the situation (Carrive, 2006). Coactivation also has functional benefits (Paton, Boscan, Pickering, & Nalivaiko, 2005) such that the myocardial contractility increases without increasing heart rate, which allows for more efficient cardiac output (i.e., longer ventricular filling times and higher contractility; Koizumi, Terui, Kollai, & Brooks, 1982). Cardiac efficiency among youth with CU traits would suggest that these youth are “better” able to respond to fearful events. For example, during aggressive confrontation, parasympathetic upregulation enables the individual to remain calm, and an increase in sympathetic activity heightens vigilance and attentiveness (Del Giudice et al., 2011). Maintaining a physiological state of self-control in high intensity situations may explain why youth with CU traits are able to successfully manipulate, intimidate, and carry out goal-directed aggression (Fanti, Frick, & Georgiou, 2009). Further, coactivation is thought to augment the rush from high-risk activities (Allison et al., 2012), which may explain why youth with CU traits engage in risky behaviors (White & Frick, 2010). Thus, a coactivated physiological state during high intensity situations may differentiate youth with and without CU traits with conduct problems, providing a biological marker for the heterogeneity in conduct problems. However, this will need to be tested further.

It is important to highlight that this is the first study to assess fear induction while measuring cardiac PNS and SNS reactivity in adolescents with CU traits. While the main findings of this paper are novel, there may be an overlap with startle potentiation research in children and adults. Increased sympathetic activity, greater cardiac vagal control (deceleration of heart rate), and reduced startle response all indicate that the individual may be allocating and orienting attention to the fear stimuli (Öhman &

Wiens, 2003). Thus, it is interesting that prior research has found low fear startle in children (Fanti, Panayiotou, Lazarou, et al., 2016) and young adults with CU traits (Fanti, Panayiotou, Kyranides, et al., 2016) and in adults high in psychopathy (see Patrick et al., 1993; Rothmund et al., 2012). Therefore, youth with CU traits are not physiologically unresponsive, but rather, they display diminished reactivity (Fanti, Panayiotou, Kyranides, et al., 2016). Taken together, these findings suggest that youth with CU traits are paying greater attention to the fear stimuli (orienting response; Anthony & Graham, 1985; Bradley et al., 1990; Lang, Bradley, & Cuthbert, 1997; Patrick et al., 1993). Therefore, during fear inducing events, youth with CU traits may not typically respond to fear by losing physiological self-control, but rather they display a physiological profile that maintains calmness, vigilance, and attentiveness to the situation (Del Giudice et al., 2011).

In a recent study including young adults using the 2-factor model of psychopathy, interpersonal-affective psychopathic traits were related to coinhibition (low SNS and PNS) in response to a VR horror game (Thomson, Aboutanos, et al., 2018). There are possible explanations of the different findings that should be considered when making comparisons. In contrast to a horror game, a VR roller coaster may be more thrilling and exciting than fear inducing. Therefore, a coactivated autonomic state may reflect an attentiveness to the thrilling stimuli rather than a fear response. This may be supported by the self-report data from both samples in the present study, which showed participants found the roller coaster to increase excitement and feeling out of control but did not significantly induce feelings of unpleasantness. In contrast, the VR horror game employed by Thomson, Aboutanos, et al. (2018) induced feelings of unpleasantness as well as excitement and feelings of being less in control. Another methodological difference is that the present study focused on CU traits, whereas Thomson, Aboutanos, et al. (2018) included the 2-factor model of psychopathy (interpersonal-affective and impulsive-antisociality). Because CU traits reflect the affective facet and not the combined affective-interpersonal facet of adult psychopathy, our contrasting results may be reflective of different personality constructs being measured.

Prior research has found that children with CU traits report a reduced conscious experience of fear (Fanti, Panayiotou, Lazarou, et al., 2016). In the present study, we found that adolescents high in CU traits did not differ in their states of arousal or valence compared with youth low on CU traits. However, CU traits were related to feeling more in control after fear induction, which supports the idea that adolescents with CU traits manage fearful stimuli without losing control. This finding is complementary to the coactivated autonomic profiles associated with CU traits, and it supports the conclusion that youth with CU traits maintain a tight physiological state of control during fear induction.

In light of the findings, the present study was unable to assess sex differences because of the disproportionate number of boys in both samples. In one previous study, it was found that aggressive males tend to exhibit lower baseline PEP, whereas no significant differences were found between aggressive and nonaggressive females (Beauchaine et al., 2008). Thus, we would urge that future research should test for sex differences in autonomic profiles of adolescents with CU traits while experiencing fear. Further, it would have been beneficial to include supplementary physiological indices to measure valence during the rollercoaster. For example, the use of electromyography would have provided

information at the physiological level as to whether the participants experienced the roller coaster as appetitive or aversive. This would have allowed for a further test of the relationship between valence and CU traits. Nevertheless, the study also has many strengths. To date, this is the first study to assess PEP and RSA reactivity to fear induction in relation to adolescent levels of CU traits from both community and EBD schools. Including both community and EBD adolescent samples allowed for a direct test of the hypothesis that the autonomic profile of CU traits while experiencing fear may not be explained by high levels of conduct problems. Thus, based on these two studies, we would suggest that coactivation of PNS and SNS is specific to CU traits. This methodology has, for the first time, offered a more complete understanding of the relationship of adolescent CU traits with autonomic operations in response to fear and supported the replicability of this finding in two adolescent populations.

Prior research has suggested that youth with CU traits are characteristically unemotional and fearless (Fanti, Panayiotou, Lazarou, et al., 2016; Frick & Morris, 2004). At the biological level, the present findings may be interpreted to support this assertion. However, it is proposed here that a shift in thinking from being *fearless* to being *better able to manage fearful situations* may be more appropriate. Our results show that youth with high levels of CU traits are not unresponsive to fear, neither physiologically nor in terms of consciously experienced fear. Instead, CU traits in the present study were associated with increases in both SNS and PNS reactivity during fear induction, indicative of greater physiological control and feelings of dominance. Therefore, adolescents with CU traits show changes in their emotional experience during fear induction, but they are able to respond in a way that may be considered more optimal for dealing with and maintaining control of a fearful situation. This autonomic profile may give rise to the appearance of fearlessness that has been thought to characterize youth with CU traits. Having the psychophysiological disposition to better manage high intensity and fearful situations, coupled with an unemotional and callous lack of concern for others, it is perhaps unsurprising that adolescents with CU traits are able to predatorily aggress and commit more severe forms of violence.

## References

- Aiken, L. S., & West, S. G. (1991). *Multiple regression: Testing and interpreting interactions*. Thousand Oaks, CA: Sage Publications, Inc.
- Alkon, A., Goldstein, L. H., Smider, N., Essex, M. J., Kupfer, D. J., & Boyce, W. T. (2003). Developmental and contextual influences on autonomic reactivity in young children. *Developmental Psychobiology*, 42, 64–78. <https://doi.org/10.1002/dev.10082>
- Allison, A. L., Peres, J. C., Boettger, C., Leonbacher, U., Hastings, P. D., & Shirtcliff, E. A. (2012). Fight, flight, or fall: Autonomic nervous system reactivity during skydiving. *Personality and Individual Differences*, 53, 218–223. <https://doi.org/10.1016/j.paid.2012.03.019>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Anastassiou-Hadjicharalambous, X., & Warden, D. (2008). Physiologically-indexed and self-perceived affective empathy in conduct-disordered children high and low on callous-unemotional traits. *Child Psychiatry and Human Development*, 39, 503–17. <https://doi.org/10.1007/s10578-008-0104-y>
- Anthony, B. J., & Graham, F. K. (1985). Blink reflex modification by selective attention: evidence for the modulation of “automatic” processing. *Biological Psychology*, 21, 43–59.
- Beauchaine, T. P. (2001). Vagal tone, development, and Gray’s motivational theory: Toward an integrated model of autonomic nervous system

- functioning in psychopathology. *Development and Psychopathology*, 13, 183–214. <https://doi.org/https://doi.org/10.1017/S0954579401002012>
- Beauchaine, T. P. (2015). Respiratory Sinus Arrhythmia: A Transdiagnostic Biomarker of Emotion Dysregulation and Psychopathology. *Current Opinion in Psychology*, 3, 43–47. <https://doi.org/10.1016/j.copsyc.2015.01.017>
- Beauchaine, T. P., Gatzke-Kopp, L., & Mead, H. K. (2007). Polyvagal theory and developmental psychopathology: Emotion dysregulation and conduct problems from preschool to adolescence. *Biological Psychology*, 74, 174–84. <https://doi.org/10.1016/j.biopsycho.2005.08.008>
- Beauchaine, T. P., Hong, J., & Marsh, P. (2008). Sex differences in autonomic correlates of conduct problems and aggression. *Journal of the American Academy of Child and Adolescent Psychiatry*, 47, 788–96. <https://doi.org/10.1097/CHI.0b013e318172ef4b>
- Beauchaine, T. P., Katkin, E. S., Strassberg, Z., & Snarr, J. (2001). Disinhibitory psychopathology in male adolescents: Discriminating conduct disorder from attention-deficit/hyperactivity disorder through concurrent assessment of multiple autonomic states. *Journal of Abnormal Psychology*, 110, 610–24.
- Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1991). Autonomic determinism: The modes of autonomic control, the doctrine of autonomic space, and the laws of autonomic constraint. *Psychological Review*, 98, 459–87. <https://doi.org/https://doi.org/10.1037//0033-295x.98.4.459>
- Berntson, G. G., Norman, G. J., Hawley, L. C., & Cacioppo, J. T. (2008). Cardiac autonomic balance versus cardiac regulatory capacity. *Psychophysiology*, 45, 643–52. <https://doi.org/10.1111/j.1469-8986.2008.00652.x>
- Blair, R. J. R., Leibenluft, E., & Pine, D. S. (2014). Conduct disorder and callous-unemotional traits in youth. *New England Journal of Medicine*, 371, 2207–2216. <https://doi.org/10.1056/NEJMra1315612>
- Bornemann, B., Winkielman, P., & der Meer, E. van. (2012). Can you feel what you do not see? Using internal feedback to detect briefly presented emotional stimuli. *International Journal of Psychophysiology*, 85, 116–124. <https://doi.org/10.1016/j.ijpsycho.2011.04.007>
- Boyce, W., Quas, J., Alkon, A., Smider, N. A., Essex, M. J., & Kupfer, D. J. (2001). Autonomic reactivity and psychopathology in middle childhood. *The British Journal of Psychiatry: The Journal of Mental Science*, 179, 144–50. <https://doi.org/https://doi.org/10.1192/bjp.179.2.144>
- Bradley, M. M., Cuthbert, B. N., & Lang, P. J. (1990). Startle reflex modification: Emotion or attention? *Psychophysiology*, 27, 513–522. <https://doi.org/10.1111/j.1469-8986.1990.tb01966.x>
- Bradley, M. M., & Lang, P. J. (1994). Measuring emotion: The self-assessment manikin and the semantic differential. *Journal of Behavior Therapy and Experimental Psychiatry*, 25, 49–59. [https://doi.org/https://doi.org/10.1016/0005-7916\(94\)90063-9](https://doi.org/https://doi.org/10.1016/0005-7916(94)90063-9)
- Brenner, S. L., Beauchaine, T. P., & Sylvers, P. D. (2005). A comparison of psychophysiological and self-report measures of BAS and BIS activation. *Psychophysiology*, 42, 108–15. <https://doi.org/10.1111/j.1469-8986.2005.00261.x>
- Brzozowski, A., Gillespie, S. M., Dixon, L., & Mitchell, I. J. (2018). Cardiac autonomic function and psychological characteristics of heterosexual female perpetrators of intimate partner physical aggression. *Journal of Interpersonal Violence*, Article 088626051877574. Retrieved from <https://doi.org/10.1177/0886260518775748>
- Bylsma, L. M., Yaroslavsky, I., Rottenberg, J., Jennings, J. R., George, C. J., Kiss, E., ... Kovacs, M. (2015). Juvenile onset depression alters cardiac autonomic balance in response to psychological and physical challenges. *Biological Psychology*, 110, 167–74. <https://doi.org/10.1016/j.biopsycho.2015.07.003>
- Calkins, S. D., & Dedmon, S. E. (2000). Physiological and behavioral regulation in two-year-old children with aggressive/destructive behavior problems. *Journal of Abnormal Child Psychology*, 28, 103–118. <https://doi.org/10.1023/A:1005112912906>
- Campbell, A., Coombes, C., David, R., Opre, A., Grayson, L., & Muncer, S. (2016). Sex differences are not attenuated by a sex-invariant measure of fear: The situated fear questionnaire. *Personality and Individual Differences*, 97, 210–219. <https://doi.org/10.1016/j.paid.2016.03.049>
- Carrive, P. (2006). Dual activation of cardiac sympathetic and parasympathetic components during conditioned fear to context in the rat. *Clinical and Experimental Pharmacology and Physiology*, 33, 1251–1254. <https://doi.org/10.1111/j.1440-1681.2006.04519.x>
- Crowell, S. E., Beauchaine, T. P., Gatzke-Kopp, L., Sylvers, P., Mead, H., & Chipman-Chacon, J. (2006). Autonomic correlates of attention-deficit/hyperactivity disorder and oppositional defiant disorder in preschool children. *Journal of Abnormal Psychology*, 115, 174–8. <https://doi.org/10.1037/0021-843X.115.1.174>
- Dackis, M. N., Rogosch, F. A., & Cicchetti, D. (2015). Child maltreatment, callous-unemotional traits, and defensive responding in high-risk children: An investigation of emotion-modulated startle response. *Development and Psychopathology*, 27, 1527–45. <https://doi.org/10.1017/S0954579415000929>
- Dadds, M. R., El Masry, Y., Wimalaweera, S., & Guastella, A. J. (2008). Reduced eye gaze explains. *Journal of the American Academy of Child & Adolescent Psychiatry*, 47, 455–463.
- Del Giudice, M., Ellis, B. J., & Shirtcliff, E. A. (2011). The adaptive calibration model of stress responsivity. *Neuroscience and Biobehavioral Reviews*, 35, 1562–92. <https://doi.org/10.1016/j.neubiorev.2010.11.007>
- Del Giudice, M., Hinnant, J. B., Ellis, B. J., & El-Sheikh, M. (2012). Adaptive patterns of stress responsivity: A preliminary investigation. *Developmental Psychology*, 48, 775–90. <https://doi.org/10.1037/a0026519>
- de Wied, M., van Boxtel, A., Matthys, W., & Meeus, W. (2012). Verbal, facial and autonomic responses to empathy-eliciting film clips by disruptive male adolescents with high versus low callous-unemotional traits. *Journal of Abnormal Child Psychology*, 40, 211–23. <https://doi.org/10.1007/s10802-011-9557-8>
- El-Sheikh, M., Kouros, C. D., Erath, S., Cummings, E. M., Keller, P., & Staton, L. (2009). Marital conflict and children's externalizing behavior: interactions between parasympathetic and sympathetic nervous system activity. *Monographs of the Society for Research in Child Development*, 74, 1–79. <https://doi.org/10.1111/j.1540-5834.2009.00501.x>
- Fanti, K. A. (2018). Understanding heterogeneity in conduct disorder: A review of psychophysiological studies. *Neuroscience & Biobehavioral Reviews*, 91, 4–20.
- Fanti, K. A., Eisenbarth, H., Goble, P., Demetriou, C., Kyranides, M. N., Goodwin, D., ... Cortese, S. (2019). Psychophysiological activity and reactivity in children and adolescents with conduct problems: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, 100, 98–107. <https://doi.org/10.1016/J.NEUBIOREV.2019.02.016>
- Fanti, K. A., Frick, P. J., & Georgiou, S. (2009). Linking callous-unemotional traits to instrumental and non-instrumental forms of aggression. *Journal of Psychopathology and Behavioral Assessment*, 31, 285–298. <https://doi.org/10.1007/s10862-008-9111-3>
- Fanti, K. A., Kyranides, M. N., Georgiou, G., Petridou, M., Colins, O. F., Tuvblad, C., & Andershed, H. (2017). Callous-unemotional, impulsive-irresponsible, and grandiose-manipulative traits: Distinct associations with heart rate, skin conductance, and startle responses to violent and erotic scenes. *Psychophysiology*, 54, 663–672. <https://doi.org/10.1111/psyp.12837>
- Fanti, K. A., Panayiotou, G., Kyranides, M. N., & Avraamides, M. N. (2016). Startle modulation during violent films: Association with callous-unemotional traits and aggressive behavior. *Motivation and Emotion*, 40, 1–13. <https://doi.org/10.1007/s11031-015-9517-7>
- Fanti, K. A., Panayiotou, G., Lazarou, C., Michael, R., & Georgiou, G. (2016). The better of two evils? Evidence that children exhibiting continuous conduct problems high or low on callous-unemotional traits score on opposite directions on physiological and behavioral measures of fear. *Development and Psychopathology*, 28, 185–98. <https://doi.org/10.1017/S0954579415000371>
- Frick, P. J. (2004). *The Inventory of Callous-Unemotional Traits*. The University of New Orleans. Available at <http://labs.uno.edu/developmental-psychopathology/ICU.html>
- Frick, P. J., & Morris, A. S. (2004). Temperament and developmental pathways to conduct problems. *Journal of Clinical Child and Adolescent Psychology: The Official Journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*, 33, 54–68. [https://doi.org/10.1207/S15374424JCCP3301\\_6](https://doi.org/10.1207/S15374424JCCP3301_6)



- Frick, P. J., Ray, J. V., Thornton, L. C., & Kahn, R. E. (2014). Annual research review: A developmental psychopathology approach to understanding callous-unemotional traits in children and adolescents with serious conduct problems. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 55, 532–48. <https://doi.org/10.1111/jcpp.12152>
- Frick, P. J., & Viding, E. (2009). Antisocial behavior from a developmental psychopathology perspective. *Development and Psychopathology*, 21, 1111–31. <https://doi.org/10.1017/S0954579409990071>
- Geer, J. H. (1965). The development of a scale to measure fear. *Behaviour Research and Therapy*, 3, 45–53.
- Gillespie, S. M., Brzozowski, A., & Mitchell, I. J. (2018). Self-regulation and aggressive antisocial behaviour: Insights from amygdala-prefrontal and heart-brain interactions. *Psychology, Crime & Law*, 24, 243–257. <https://doi.org/10.1080/1068316X.2017.1414816>
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38, 581–586. <https://doi.org/10.1111/j.1469-7610.1997.tb01545.x>
- Grossman, P., van Beek, J., & Wientjes, C. (1990). A comparison of three quantification methods for estimation of respiratory sinus arrhythmia. *Psychophysiology*, 27, 702–14. <https://doi.org/https://doi.org/10.1111/j.1469-8986.1990.tb03198.x>
- Hastrup, J. L. (1986). Duration of initial heart rate assessment in psychophysiology: Current practices and implications. *Psychophysiology*, 23, 15–18.
- Holliman, N. (2010). Cosmic origins: Experiences making a stereoscopic 3D movie. *International Society for Optics and Photonics*, 7524.
- Hoppenbrouwers, S. S., Bulten, B. H., & Brazil, I. A. (2016). Parsing fear: A reassessment of the evidence for fear deficits in psychopathy. *Psychological Bulletin*, 142, 573–600. <https://doi.org/https://doi.org/10.1037/bul0000040>
- Jennings, J. R., Kamarck, T., Stewart, C., Eddy, M., & Johnson, P. (1992). Alternate cardiovascular baseline assessment techniques: vanilla or resting baseline. *Psychophysiology*, 29, 742–50.
- Jones, A. P., Laurens, K. R., Herba, C. M., Barker, G. J., & Viding, E. (2009). Amygdala hypoactivity to fearful faces in boys with conduct problems and callous-unemotional traits. *The American Journal of Psychiatry*, 166(9), 95–102. <https://doi.org/10.1176/appi.ajp.2008.07071050>
- Kimonis, E. R., Frick, P. J., Munoz, L. C., & Aucoin, K. J. (2008). Callous-unemotional traits and the emotional processing of distress cues in detained boys: Testing the moderating role of aggression, exposure to community violence, and histories of abuse. *Development and Psychopathology*, 20, 569–89. <https://doi.org/10.1017/S095457940800028X>
- Klingzell, I., Fanti, K. A., Colins, O. F., Frogner, L., Andershed, A.-K., & Andershed, H. (2016). Early childhood trajectories of conduct problems and callous-unemotional traits: The role of fearlessness and psychopathic personality dimensions. *Child Psychiatry & Human Development*, 47, 236–247. <https://doi.org/10.1007/s10578-015-0560-0>
- Koizumi, K., Terui, N., Kollai, M., & Brooks, C. M. (1982). Functional significance of coactivation of vagal and sympathetic cardiac nerves. *Proceedings of the National Academy of Sciences of the United States of America*, 79, 2116–20.
- Kyranides, M. N., Fanti, K. A., & Panayiotou, G. (2016). The disruptive adolescent as a grown-up: Predicting adult startle responses to violent and erotic films from adolescent conduct problems and callous-unemotional traits. *Journal of Psychopathology and Behavioral Assessment*, 38, 183–194. <https://doi.org/10.1007/s10862-015-9520-z>
- Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (1997). Motivated attention: Affect, activation, and action. In P. Lang, R. F. Simons, & M. T. Balaban (Eds.), *Attention and orienting: Sensory and motivational processes* (pp. 97–135). Lawrence Erlbaum Associates Publishers.
- LeDoux, J. E. (2013). The slippery slope of fear. *Trends in Cognitive Sciences*, 17, 155–6. <https://doi.org/10.1016/j.tics.2013.02.004>
- Lozier, L. M., Cardinale, E. M., VanMeter, J. W., & Marsh, A. A. (2014). Mediation of the relationship between callous-unemotional traits and proactive aggression by amygdala response to fear among children with conduct problems. *JAMA Psychiatry*, 71, 627–36. <https://doi.org/10.1001/jamapsychiatry.2013.4540>
- Lykken, D. (1995). *The antisocial personalities*. Hillsdale, NJ: Erlbaum.
- Marsh, A. A., & Cardinale, E. M. (2014). When psychopathy impairs moral judgments: neural responses during judgments about causing fear. *Social Cognitive and Affective Neuroscience*, 9, 3–11. <https://doi.org/10.1093/scan/nss097>
- Marsh, A. A., Finger, E. C., Mitchell, D. G. V. V., Reid, M. E., Sims, C., Kosson, D. S., ... Blair, R. J. R. R. (2008). Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *The American Journal of Psychiatry*, 165, 712–20. <https://doi.org/10.1176/appi.ajp.2007.07071145>
- Marsh, A. A., Finger, E. C., Schechter, J. C., Jurkowitz, I. T. N., Reid, M. E., & Blair, R. J. R. (2011). Adolescents with psychopathic traits report reductions in physiological responses to fear. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 52, 834–41. <https://doi.org/10.1111/j.1469-7610.2010.02353.x>
- Mirisola, A., & Seta, L. (2011). *Pequod package: Moderated regression with mean and residual centering and multiple slope analysis*. Perugia: University of Perugia.
- Muñoz, L. C., Frick, P. J., Kimonis, E. R., & Aucoin, K. J. (2008). Types of aggression, responsiveness to provocation, and callous-unemotional traits in detained adolescents. *Journal of Abnormal Child Psychology*, 36, 15–28. <https://doi.org/10.1007/s10802-007-9137-0>
- Musser, E. D., Galloway-Long, H. S., Frick, P. J., & Nigg, J. T. (2013). Emotion regulation and heterogeneity in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 52, 163–171.e2. <https://doi.org/10.1016/j.jaac.2012.11.009>
- Newlin, D. B., & Levenson, R. W. (1979). Pre-ejection period: Measuring beta-adrenergic influences upon the heart. *Psychophysiology*, 16, 546–53.
- Newman, J. P., & Brinkley, C. A. (1997). Reconsidering the low-fear explanation for primary psychopathy. *Psychological Inquiry*, 8, 236–244. [https://doi.org/10.1207/s15327965phi0803\\_14](https://doi.org/10.1207/s15327965phi0803_14)
- Northover, C., Thapar, A., Langley, K., & van Goozen, S. H. M. (2015). Pain sensitivity in adolescent males with attention-deficit/hyperactivity disorder: Testing for associations with conduct disorder and callous and unemotional traits. *PLoS One*, 10, e0134417. <https://doi.org/10.1371/journal.pone.0134417>
- Ohman, A., & Soares, J. J. (1998). Emotional conditioning to masked stimuli: Expectancies for aversive outcomes following nonrecognized fear-relevant stimuli. *Journal of Experimental Psychology: General*, 127, 69–82.
- Öhman, A., & Wiens, S. (2003). On the automaticity of autonomic responses in emotion: An evolutionary perspective. In R. J. Davidson, K. R. Scherer, & H. H. Goldsmith (Eds.), *Handbook of affective sciences. Series in affective science*. New York: Oxford University Press.
- Olsson, A., & Phelps, E. A. (2004). Learned fear of “unseen” faces after Pavlovian, observational, and instructed fear. *Psychological Science*, 15, 822–828. <https://doi.org/10.1111/j.0956-7976.2004.00762.x>
- Pardini, D. (2006). The callousness pathway to severe violent delinquency. *Aggressive Behavior*, 32, 590–598. <https://doi.org/10.1002/ab.20158>
- Paton, J. F. R., Boscan, P., Pickering, A. E., & Nalivaiko, E. (2005). The yin and yang of cardiac autonomic control: Vago-sympathetic interactions revisited. *Brain Research. Brain Research Reviews*, 49, 555–65. <https://doi.org/10.1016/j.brainresrev.2005.02.005>
- Patrick, C. J., Bradley, M. M., & Lang, P. J. (1993). Emotion in the criminal psychopath: Startle reflex modulation. *Journal of Abnormal Psychology*, 102, 82–92. <https://doi.org/10.1037//0021-843X.102.1.82>
- Piferi, R. L., Kline, K. A., Younger, J., & Lawler, K. A. (2000). An alternative approach for achieving cardiovascular baseline: Viewing an aquatic video. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, 37, 207–17.
- Pihet, S., Etter, S., Schmid, M., & Kimonis, E. R. (2015). Assessing Callous-unemotional traits in adolescents: Validity of the Inventory of Callous-Unemotional Traits across gender, age, and community/institutionalized status. *Journal of Psychopathology and Behavioral Assessment*, 37, 407–421. <https://doi.org/10.1007/s10862-014-9472-8>
- Pine, D. S., Wasserman, G. A., Miller, L., Coplan, J. D., Bagiella, E., Kovelencu, P., ... Sloan, R. P. (1998). Heart period variability and psychopathology in urban boys at risk for delinquency. *Psychophysiology*, 35, 521–9.
- Porges, S. W. (1995). Orienting in a defensive world: Mammalian modifications of our evolutionary heritage. A polyvagal theory. *Psychophysiology*, 32, 301–18. <https://doi.org/https://doi.org/10.1111/j.1469-8986.1995.tb01213.x>
- Porges, S. W. (2003). The polyvagal theory: Phylogenetic contributions to social behavior. *Physiology & Behavior*, 79, 503–513. [https://doi.org/10.1016/S0031-9384\(03\)00156-2](https://doi.org/10.1016/S0031-9384(03)00156-2)



- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, *74*, 116–143. <https://doi.org/10.1016/j.biopsycho.2006.06.009>
- R Core Team. (2016). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rothmund, Y., Ziegler, S., Hermann, C., Gruesser, S. M., Foell, J., Patrick, C. J., & Flor, H. (2012). Fear conditioning in psychopaths: Event-related potentials and peripheral measures. *Biological Psychology*, *90*, 50–9. <https://doi.org/10.1016/j.biopsycho.2012.02.011>
- Scarpa, A., Haden, S. C., & Tanaka, A. (2010). Being hot-tempered: Autonomic, emotional, and behavioral distinctions between childhood reactive and proactive aggression. *Biological Psychology*, *84*, 488–96. <https://doi.org/10.1016/j.biopsycho.2009.11.006>
- Sherwood, A., Allen, M. T., Fahrenberg, J., Kelsey, R. M., Lovallo, W. R., & van Doornen, L. J. (1990). Methodological guidelines for impedance cardiography. *Psychophysiology*, *27*, 1–23.
- Tabachnick, B. G., & Fidell, L. S. (2013). *Using multivariate statistics* (6th). Boston: Pearson.
- Thayer, J. F., Åhs, F., Fredrikson, M., Sollers, J. J., & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews*, *36*, 747–756. <https://doi.org/10.1016/j.NEUBIOREV.2011.11.009>
- Thayer, J. F., & Lane, R. D. (2009). Claude Bernard and the heart-brain connection: further elaboration of a model of neurovisceral integration. *Neuroscience and Biobehavioral Reviews*, *33*, 81–8. <https://doi.org/10.1016/j.neubiorev.2008.08.004>
- Thomson, N. D., Aboutanos, M. B., Kiehl, K. A., Neumann, C. S., Galusha, C., & Fanti, K. A. (2018). Physiological reactivity in response to a fear induced virtual reality experience: Associations with psychopathic traits. *Psychophysiology*.
- Thomson, N. D., & Beauchaine, T. P. (2018). Respiratory sinus arrhythmia mediates links between borderline personality disorder symptoms and both aggressive and violent behavior. *Journal of Personality Disorders*, *32*, 1–16. [https://doi.org/https://doi.org/10.1521/pedi\\_2018\\_358](https://doi.org/https://doi.org/10.1521/pedi_2018_358)
- Thomson, N. D., Kiehl, K. A., & Bjork, J. M. (2018). Predicting violence and aggression in young women: The importance of psychopathy and neurobiological function. *Physiology & Behavior*, *201*, 130–138. <https://doi.org/10.1016/J.PHYSBEH.2018.11.043>
- Wagner, C. R., & Abaied, J. L. (2015). Relational victimization and proactive versus reactive relational aggression: The moderating effects of respiratory sinus arrhythmia and skin conductance. *Aggressive Behavior*, *41*, 566–579. <https://doi.org/10.1002/ab.21596>
- White, S. F., & Frick, P. J. (2010). Callous-unemotional traits and their importance to causal models of severe antisocial behavior in youth. In R. Salekin & D. Lynam (Eds.), *Handbook of child and adolescent psychopathy* (pp. 135–155). New York: The Guilford Press.