

Hippocampal sclerosis induced in mice by a *Taenia crassiceps* metacestode factor

N. Zepeda¹, N. Copitin¹, J.L. Chávez², F. García³, F. Jaimes-Miranda³, R. Rincón-Heredia³, R. Paredes³, S. Solano⁴, A.M. Fernández⁴ and J.L. Molinari¹

Research Paper

Cite this article: Zepeda N *et al* (2019). Hippocampal sclerosis induced in mice by a *Taenia crassiceps* metacestode factor. *Journal of Helminthology* **93**, 690–696. <https://doi.org/10.1017/S0022149X18000755>

Received: 8 February 2018

Accepted: 26 July 2018

First published online: 23 August 2018

Key words:

apoptosis; Barnes maze; hippocampal sclerosis, metacestode factor (Fac); mouse model; neurocysticercosis; *Taenia crassiceps*; *Taenia solium*

Author for correspondence:

J.L. Molinari, E-mail: jmolinar@ifc.unam.mx

¹Departamento de Bioquímica y Biología Estructural, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, Ciudad de México, C.P. 04510 A.P. 70-242; ²Departamento de Neurociencia Cognitiva, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, Ciudad de México, C.P. 04510 A.P. 70-242; ³Unidad de Imagenología, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, Ciudad de México, C.P. 04510 A.P. 70-242 and ⁴Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México, 04510

Abstract

An experimental *Taenia crassiceps* mouse model was used to assess the role of *Taenia solium* metacestode factor (Fac) in human neurocysticercosis. Intraperitoneal infection with *T. crassiceps* metacestodes or subcutaneous inoculation with a *T. crassiceps* metacestode factor (Fac) produced significant impairment of performance (learning) in the Barnes maze and induced bilateral hippocampal sclerosis in mice. Several staining techniques revealed important cell dispersion, extensive apoptosis and cell loss in the dentate gyrus, hilus and CA1-CA3 regions of both hippocampi, as well as intense deterioration of the adjacent cortex. An outstanding disruption of its histoarchitecture in the surrounding tissue of all these regions and apoptosis of the endothelial cells were also observed.

Introduction

Human beings acquire neurocysticercosis (NCC) by ingesting foods contaminated with *Taenia solium* eggs. The acidic gastric juice breaks up the inter-embryophore cement of the eggs. Then, the sheath-covered embryos are carried to the duodenum, where the pancreatic juice and bile dissolve the external lipid layer of the sheath, allowing the entrance of water and nutrients, which activate the previously dormant embryos. The activated embryos rip their envelopes with the help of their six sharp hooks, and they promptly leave these broken structures, as described in an *in vitro* study (Molinari *et al.*, 1993). Released from their envelopes, activated embryos invade the intestine, as with *Hymenolepis nana* (Miyasato *et al.*, 1977), migrate through the bloodstream and implant in several tissues, such as the striated muscles and central nervous system, where they develop into the metacestode stage.

Neurocysticercosis is considered to be the most common parasitic infestation of the central nervous system and the single most common cause of epilepsy. Seizures, headaches and neurological deficits are described in human neurocysticercosis. Signs of psychiatric disease (65.8%), cognitive decline (87%), altered memory (25%) and attention deficits (100%) among others were reported in a heterogeneous sample of 38 patients (Forlenza *et al.*, 1997). Other features are hydrocephalous, chronic meningitis, and impairment in executive functions, verbal and non-verbal memory, praxis and verbal fluency (Varghese *et al.*, 2016).

Hippocampal atrophy without seizures in patients with calcified neurocysticercosis (cNCC) has been observed, suggesting inflammatory-mediated mechanisms provoked by cysticercosis, which may contribute or even lead to refractory mesial temporal lobe epilepsy with hippocampal sclerosis (MTLE-HS) by mechanisms that need to be clarified (Bianchin *et al.*, 2015; Griper and Welburn, 2017). To address the role that *T. solium* metacestode factor (Fac) (Molinari *et al.*, 1990) could play in the production of hippocampal sclerosis, whose cause in human neurocysticercosis is poorly understood, we developed an experimental *Taenia crassiceps*-mouse model to study and understand the effects of an infection of mice with *T. crassiceps* metacestodes and the subcutaneous inoculation of mice with a *T. crassiceps* factor on hippocampal cells; results showed an extensive and significant apoptosis of dentate gyrus, CA1, CA2, CA3 and hilus cells compared with intact control cells (Zepeda *et al.*, 2017).

In the current research, we extended the time of infection and the inoculation with Fac for 110 days to try to induce a more severe hippocampal injury than that reported recently (Zepeda *et al.*, 2017). Histopathologically, temporal lobe epilepsy (TLE) associated with HS has been characterized by cell death and neuronal cell loss, which is most severe in the CA1 and CA3 regions and in the hilus of the dentate gyrus (DG), volume loss, reactive gliosis, and dispersion of the dentate gyrus granule cells (Sloviter, 1994; Lurton *et al.*, 1998).

Materials and methods

Treatment of mice

BALB/c mice (28 days old, 20–22 g) were maintained under standard animal housing conditions with access to Purina chow and water *ad libitum*. *Taenia crassiceps* metacystodes (WFU strain) were maintained in our laboratory via peritoneal passage in female mice, as described (Everhard *et al.*, 2004). Peritoneal fluid from healthy female controls (CPF) and *T. crassiceps* metacystode-infected mice, as well as *T. crassiceps* metacystode factor (Fac), were obtained according to Zepeda *et al.* (2011). To determine the effects of infection with *T. crassiceps* metacystodes and Fac treatment on hippocampal cells and the surrounding tissue, 18 female mice were divided into three groups. The control (Con) group (n = 6) and Factor (Fac) group (n = 6) were inoculated subcutaneously with 0.1 ml CPF, or Fac containing 50 µg of carbohydrates, as determined by the anthrone method (Ashwell, 1957) every 4 days for 100 days. The infected (Inf) group (n = 6) was infected intraperitoneally with up to 40 *T. crassiceps* metacystodes, measuring 2–3 mm in diameter and suspended in 0.2 ml phosphate buffered saline (PBS).

Barnes maze

At 110 days post infection with intraperitoneal *T. crassiceps* metacystodes or subcutaneous inoculation with *T. crassiceps* Fac, the Barnes maze procedure for small rodents was followed to estimate cognitive impairment in learning and memory (Rosenfeld and Ferguson, 2014), with minor modifications. Briefly, a bright illumination overhead was used. The maze was centred below the lights. Mice were tested without the use of a wall or cues. The maze was 90 cm in diameter and 90 cm in height, with 20 escape holes that each had a diameter of 5 cm, and there was only one hole with an escape cage. No hole was plugged in this test. After 30 minutes of habituation, each mouse was gently removed from his home cage and placed onto the maze centre. Then, the overhead lights were turned on. Every mouse's performance was recorded using a Canon VIXIA HF R70 camera (Canon, Melville, NY, USA). If the mouse failed to enter the escape cage within 5 minutes, it was guided gently to the correct location and into the escape cage, where it remained for 2 minutes. Each mouse was tested for two trials/day with an inter-trial interval of approximately 30 minutes. All mice were tested for five consecutive days. Videos were analysed carefully. The Barnes maze test was used to study the behaviour of the control, experimental cysticercotic and Fac treated-mice.

Histology and cellular apoptosis

At 115 days post treatment and infection, all mice were anaesthetized with pentobarbital and perfused transcardially with saline followed by phosphate-buffered 10% formalin. Then, the brains were carefully removed, and intraperitoneal metacystodes were obtained and counted. Two coronal cuts were made in each brain: an anterior cut was made at Bregma –1.43 mm and the other at Bregma –5.19 mm (Paxinos and Franklin, 2013). The intermediate fragments of each brain were fixed with 2.5% paraformaldehyde in PBS for 12 h at room temperature. Next, the fragments of each brain were embedded in paraffin. Coronal sections that were 5 µm thick were prepared on a rotary microtome, placed on poly-L-lysine-coated microscopic slide, and subjected to haematoxylin/eosin (H/E) and DAPI staining to analyse the




Fig. 1. Mean escape latency in seconds as a measure of performance of mice in the Barnes maze. Values of the second trials for 5 days. Control group (○); Infected group (□); Factor group (▲). ** $P < 0.01$ Control vs Infected. * $P < 0.05$ Control vs Factor.




Fig. 2. Haematoxylin/eosin-stained brain section that contains part of the dentate gyrus and hilus from a control mouse (a & b), a mouse infected with *T. crassiceps* metacystodes (c & d), and a mouse inoculated with *T. crassiceps* Fac (e & f). Cell loss (*); bleached nuclei (arrows); tears (arrowheads); gyrus branches angle (vertical bar). Scale bar = 25 µm. 40×.

cellular morphology and histoarchitecture of the dentate gyrus and CA1-CA3 regions. Sections stained with H/E were analysed with an Olympus IX71 microscope and IMAGE PRO-PCUS 60 software. Sections stained with DAPI were analysed with a Zeiss LSM 800 confocal microscope and ZEN version 2.1 software. The presence of DNA strand breaks was evaluated using an *in situ* cell death detection kit (Roche). TUNEL assays were analysed with a confocal microscope Leica TCS-SPS (Carl Zeiss de México-División Microscopía, Ciudad de México).

Data analysis

The numbers of haematoxylin/eosin-stained and TUNEL-positive cells were measured by manual counting (area in µm²) with the aid of Image J software (National Institutes of Health, Bethesda, MD, USA). For statistical data, one-way analyses of variance and Tukey's test were performed using GraphPad software




Fig. 3. Haematoxylin/eosin-stained brain sections that contain portions of CA1, CA2 and CA3 regions from a control mouse (a–c), a mouse infected with *T. crassiceps* metacystodes (d–f) and a mouse inoculated with *T. crassiceps* Fac (g–i). Scale bar = 25 μm. 40×. (j) Number of apoptotic cells/100 μm². Bars represent the mean ± SD. * $P < 0.001$ Infected vs Control. ** $P < 0.001$ Factor vs Control.

(GraphPad Instat, San Diego, CA, USA). Data are presented as the mean ± SD. $P < 0.05$ was considered significant.

Results

Barnes maze

Both intraperitoneal infection with *Taenia crassiceps* metacystodes and subcutaneous inoculation with a *T. crassiceps* metacystode factor (Fac) produced significant impairment on performance in the Barnes maze and induced bilateral hippocampal sclerosis in mice. The performance in the Barnes maze was mainly assessed by the escape latency parameter (mean ± SD), measured in the

number of seconds it took to reach and completely hide in the escape hole.

During all time (second trial) to learn this task, both infected and factor mice were significantly slower in reaching and hiding in the escape hole, in comparison with control mice. On the fifth day, all control mice ($n = 6$) were able to hide in the escape hole in 28 ± 44 s after reaching it. Latency values of control mice were significantly less than those of experimental mice ($P < 0.05$ Con vs Fac, and $P < 0.01$ Con vs Inf.; fig. 1). Infected mice ($n = 4$) showed difficulty in moving in the maze; on the fourth day only one (25%) was able to hide at 151 s after reaching the escape hole. On the fifth day only one reached the escape hole in 73 s, and stayed there the rest of time without hiding in the hole. On




Fig. 4. DAPI-stained brain sections that contain portions of CA1, CA2 and CA3 from a control mouse (a–c), a mouse infected with *T. crassiceps* metacystodes (d–f) and a mouse inoculated with *T. crassiceps* Fac (g–i). Scale bar = 25 μ m. 40 \times .




Fig. 5. TUNEL-stained brain sections that contain a portion of the dentate gyrus from a control mouse (a), and a portion of the dentate gyrus, hilus, CA1, CA2 and CA3 from a mouse infected with *T. crassiceps* metacystodes (b–f). Scale bar = 50 μ m. 60 \times .




Fig. 6. TUNEL-stained brain sections that contain a portion of dentate gyrus from a control mouse (a), a portion of dentate gyrus and hilus from a mouse infected with *T. crassiceps* Fac (b and c), a portion of CD1, CD2 and CD3 from a mouse inoculated with *T. crassiceps* Fac (d–f), a portion of the adjacent cortex from a mouse inoculated with Fac (g), and the border of the adjacent cortex showing apoptotic endothelial cells within sections of the capillary (arrow) (h). Scale bar = 50 μm . 60 \times . (i) is an amplification of (h). Scale bar = 10 μm . 200 \times .

the fifth day Factor mice ($n = 6$) showed hyperactivity, reaching the escape hole at 67 ± 38 s. Only five mice had hidden at 73 ± 85 s after reaching the escape hole. The sixth mouse did reach the escape hole at 99 s, then left the hole four times to explore several holes and came back, but it never hid (fig. 1).

Histopathological studies

A heavy parasite load was found in the peritoneal cavity of mice infected with *T. crassiceps* metacystodes (485 ± 78 metacystodes per mouse). Haematoxylin/eosin staining showed that all sub-regions of hippocampi from experimental animals were affected in their tissue architecture either in the cellular morphology or in the surrounding tissue. The dentate gyri, as well as the hilus, CA1, CA2 and CA3 cells displayed dispersion, bleached nuclei, cell loss and many tears in the surrounding white matter, compared with the integrity of the cells and the white matter of the control hippocampi. Dentate gyrus branches from experimental mice (fig. 2c, e) formed a smaller angle compared to those of the control hippocampi (fig. 2a). The number of cells that had altered morphology and staining were statistically ($P < 0.001$) more than those of the control mice (fig. 3). DAPI staining displayed generalized pyknosis or hypercondensation of the chromatin of the CA1, CA2 and CA3 nuclei from *T. crassiceps* infected and Fac-treated mice. Apparently, all nuclei in these sub-regions were collapsed, and their number was

less than those of the control mice. Additionally, numerous empty spaces among the reduced number of pyknotic nuclei of these sub-regions were observed, which suggested cell loss (fig. 4). Pyknosis is the accepted sign of programmed cell death (Burgoyne, 1999). To evaluate DNA fragmentation, DNA strand breaks were labelled using fluorescein-dUTP. This staining showed generalized and significant TUNEL-positive cells in the three layers of the dentate gyrus, hilus and in cells from the CA1, CA2 and CA3 regions, including the adjacent cortex ($P < 0.01$ Inf and Fac vs. Con). In all these places, many fragmented nuclei were observed, compared to TUNEL-negative nuclei of the control hippocampi. The intensity of the fluorescence of the nuclei in the above-mentioned regions was not homogeneous, suggesting that apoptosis was induced at different times. Nuclei with a lower fluorescence intensity and of reduced size could be among the first in developing apoptosis. Dispersion and cell loss were also observed (figs 5 and 6). Numerous TUNEL-positive cells were also found in the neighbouring cortex. A remarkable finding was the observation of apoptosis of endothelial cells in several places in all the studied regions (figs 6 and 7).

Discussion

In the current research, the experimental *T. crassiceps* mouse model (Zepeda *et al.*, 2017) was used to investigate parasite mechanisms involved in the production of hippocampal sclerosis




Fig. 7. Statistical data from the sections stained with TUNEL. Bars represent the mean \pm SD of the dentate gyrus from control mice (a & b), dentate gyrus (DG), hilus (H), CA1, CA2 and CA3 from infected mice (a), and DG, H, CA1, CA2, CA3 and Cortex (Cx) from factor mice (b). * $P < 0.01$ DG, H, CA1, CA2, CA3 and Cx vs Con.

(HS), in which some patients suffer from neurocysticercosis. A deficit in learning and memory functions was observed using the Barnes maze. This deficit was more pronounced in mice infected with *T. crassiceps* metacystodes, perhaps due to a heavy parasitic load and an intense intraperitoneal inflammatory reaction as described (Zepeda *et al.*, 2010), as well as structural damage of the histoarchitecture of the hippocampi from experimental mice. This damage was characterized by significant cell death (apoptosis), cell dispersion and cell loss, in all sub-regions of the hippocampus either in the infected group or the Fac-treated animals, such as dentate gyrus, hilus, CA1, CA2, CA3, and the adjacent cortex, compared with the integrity of the cells in control mice. Structural damage was also observed in the neighbouring white matter.

The Barnes maze has been used to relate memory loss and long-term potentiation (LTP) dysfunction in the hippocampus, which are involved in Alzheimer's disease. Using the Barnes maze, Hadipour *et al.* (2018) found that administration of crocin in rats significantly improves spatial memory indicators such as latency time. The learning deficit in A β (beta-amyloid)-treated animals correlated with a reduction of LTP in hippocampal synapses. Crocin significantly reduced the number of TUNEL-positive cells in the CA1 sub-region compared to the A β -group. Barbe *et al.* (2016) utilized the Barnes maze to test for long-term memory deficits, with good success. Their results showed in the CA1-3 regions of the hippocampus of mice significantly reduced

numbers of neuronal cell bodies overall, and reduced numbers of Pur-alpha+ neuronal cell bodies and dendrites in heterozygous Pur-alpha (+/-) mice, compared to wild-type (+/+) littermates.

Here, in every sub-region studied, a very interesting finding was the observation of apoptotic endothelial cells dispersed in the white matter near the neuronal packages and in the border of the adjacent cortex. This finding suggests that the Fac apoptogenic activity on endothelial cells could impair the blood-brain barrier, allowing the entry of the Fac (that migrated via the bloodstream when the peritoneal cavity of the mouse was infected with *T. crassiceps* metacystodes or the abdominal skin of the mouse was inoculated with Fac) into the brain, so that it can reach the hippocampal cells and other regions of the brain, such as thalamic nuclei (Zepeda *et al.*, 2017). The apoptogenic activity of Fac seems to have been cumulative, as was observed with the TUNEL staining.

The pathological characteristic produced by *T. crassiceps* metacystode infection or by *T. crassiceps* Fac in all sub-regions of the hippocampus resembles human hippocampal sclerosis, such as volume loss, which indicates atrophy (Jeniffer *et al.*, 2015). The learning deficit and hyperactivity are neurological manifestations of structural damage in the hippocampus, which rely on other functions, such as attention, language and spatial memory. Another feature of hippocampal sclerosis described in the literature is the increase of astrocytes during the scarring process in places where there has been cell loss due to some injury (Sloviter, 1994). We looked for this feature using a rabbit antibody anti-GFAP fluorescein mouse, but we could not obtain positive results in this way. So, in the next study, we intend to test this hypothesis.

In humans, NCC damage to the hippocampus caused by the *T. solium* Fac must be a very chronic process, proportional to the number of living parasites; thus, when the neurological deficits are apparent, including epilepsy, the parasites are probably already calcified. Molinari *et al.* (1987) reported that in porcine cysticercosis, the suppression of T-cells was proportional to the number of cysticerci per 1 kg of parasitized meat. Human neurocysticercosis has been associated with hippocampal atrophy, and the results of a study by Del Brutto *et al.* (2017) suggested that NCC-related hippocampal atrophy takes a long time to develop.

Taenia crassiceps Fac are chromatographic Sephadex G-50 fractions containing substances of low molecular weight (<1250 Da) obtained by ultrafiltration of the peritoneal fluid from female mice infected intraperitoneally with *T. crassiceps* metacystodes (Zepeda *et al.*, 2011). The complete chemical characterization of these substances has not been possible. By preliminary NMR and infrared studies, these Sephadex G-50 fractions contain at least three aromatic components, carbohydrates such as glucose, and can induce a specific antibody response in rabbits and mice, which indicate that this factor is produced by the parasite (data not published). The non-specific way this Fac induces apoptosis in different types of cells, such as testis cells (Zepeda *et al.*, 2011), ovary cells (Solano *et al.*, 2015), spleen cells (Zepeda *et al.*, 2016), and hippocampal cells (Zepeda *et al.*, 2017) is unknown. The learning deficit and the extensive apoptosis of hippocampal cells of experimental mice indicate a close relationship between both structure and function. Thus, it may be reasonable to assume *T. solium* metacystode factor is the causative agent of brain injuries such as hippocampal sclerosis in human neurocysticercosis. It will be important to contrast these results with relevant studies in human neurocysticercosis, as has been noted by Del Brutto *et al.* (2016).

Using the Barnes maze, experimental mice showed a learning deficit closely related to hippocampal injury and very similar to that observed in patients with neurocysticercosis. Apoptosis of hippocampal endothelial cells suggested that *T. crassiceps* metacystode factor was able to break the BBB, enter the brain and produce hippocampal sclerosis.

Acknowledgments. We are grateful to Teresa Cortés and Daniela Rodríguez for performing histological procedures, and Claudia Rivera and Héctor Malagón for providing the healthy mice. Nadia Zepeda was supported for her graduate studies by the Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México (UNAM) and CONACyT (scholarship number 262633).

Financial support. This research was supported by Universidad Nacional Autónoma de México.

Conflict of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the laws of Mexico on the care and use of laboratory animals. Animal studies were performed under an institutional protocol similar to that of the U.S. Public Health Service's *Guide for the Care and Use of Laboratory Animals*. From the Secretariat of Agriculture (SAGARPA NOM-062-ZOO-1999).

References

- Ashwell G (1957) Colorimetric analysis of sugars. In Colowick SP and Kaplan NO (eds), *Methods in Enzymology*. III. New York, NY: Academic Press, pp. 73–84.
- Barbe MF et al. (2016) Memory deficits, gait ataxia and neuronal loss in the hippocampus and cerebellum in mice that are heterozygous for Pur-alpha. *Neuroscience* **337**, 177–190.
- Bianchin MM et al. (2015) Neuroimaging observations linking neurocysticercosis and mesial temporal lobe epilepsy with hippocampal sclerosis. *Epilepsy Research* **116**, 34–39.
- Burgoyne LA (1999) The mechanisms of pyknosis: hypercondensation and death. *Experimental Cell Research* **248**, 214–222.
- Del Brutto OH et al. (2016) Update on cysticercosis epileptogenic: the role of the hippocampus. *Current Neurology Neuroscience Reports* **16**, 1.
- Del Brutto OH et al. (2017) The association between neurocysticercosis and hippocampal atrophy is related to age. *American Tropical Medicine and Hygiene* **96**, 243–248.
- Everhard ME, Kuhn RE and Zelmer DA (2004) Infrapopulation dynamics of a wild strain of *Taenia crassiceps* (WFU) (Cestoda Taeniidae) in BALB/cJ mice. *Journal of Parasitology* **90**, 79–84.
- Forlenza OV et al. (1997) Psychiatric manifestations of neurocysticercosis: a study of 38 patients from a neurology clinic in Brazil. *Journal of Neurology, Neurosurgery and Psychiatry* **62**, 612–616.
- Griper LB and Welburn SC (2017) The causal relationship between neurocysticercosis infection and the development of epilepsy – a systematic review. *Infectious Diseases of Poverty* **6**, 31.
- Hadipour M et al. (2018) Crocin improved amyloid beta induced long-term potentiation and memory deficits in the hippocampal CA1 neurons in freely moving rats. *Synapse* **72**, e22026.
- Jeniffer VN, Udayakumar S and Pushpalatha K (2015) A clinical study to identify the possible etiology of complex partial seizures using magnetic resonance imaging brain findings and its implications on treatment. *Journal of Pediatric Neurosciences* **10**, 350–354.
- Lurton D et al. (1998) Granule cell dispersion is correlated with early epileptic events in human temporal lobe epilepsy. *Journal of Neurological Science* **154**, 133–136.
- Miyasato T et al. (1977) Electron microscopic observations on the penetration of *Hymenolepis nana* into the intestine of the mouse. *Acta Medica Kinki University* **2**, 1–18.
- Molinari JL, Tato P and Valles Y (1987) Immunosuppression T-lymphocytes in swine, modulated by *Cysticercus cellulosae*. *Revista Latinoamericana de Microbiología* **29**, 293–300.
- Molinari JL et al. (1990) Depressive effect of a *Taenia solium* cysticercus factor on cultured human lymphocytes stimulated with phytohaemagglutinin. *Annals of Tropical Medicine and Parasitology* **84**, 205–208.
- Molinari JL et al. (1993) Effects of serum from neurocysticercosis patients on the structure and viability of *Taenia solium* oncospheres. *Journal of Parasitology* **79**, 124–127.
- Paxinos G and Franklin KBJ (2013) *The Mouse Brain in Stereotaxic Coordinates*. 4th Edn. Oxford: Elsevier.
- Rosenfeld CS and Ferguson SA (2014) Barnes maze testing strategies with small and large rodent models. *Journal of Visualized Experiments* **84**, 51194–51219.
- Sloviter RS (1994) The functional organization of the hippocampal dentate gyrus and its relevance to the pathogenesis of temporal lobe epilepsy. *Annals of Neurology* **35**, 640–654.
- Solano S et al. (2015) A *Taenia crassiceps* metacystode factor enhance ovarian follicle atresia and oocyte degeneration in female mice. *Journal of Helminthology* **89**, 49–59.
- Varghese V et al. (2016) Cognitive dysfunction and its determinants in patients with neurocysticercosis. *Indian Journal of Psychological Medicine* **38**, 142–146.
- Zepeda N et al. (2010) Decrease of peritoneal inflammatory CD4+, CD8+, CD19+ lymphocytes and apoptosis of eosinophils in a murine *Taenia crassiceps* infection. *Parasitology Research* **127**, 1129–1135.
- Zepeda N et al. (2011) *Taenia crassiceps*: a secretion substance of low molecular weight leads to disruption and apoptosis of seminiferous epithelium cells in male mice. *Experimental Parasitology* **128**, 184–191.
- Zepeda N et al. (2016) A *Taenia crassiceps* factor induces apoptosis of spleen CD4+ T cells and TGF- β and Foxp3 gene expression in mice. *Journal of Helminthology* **90**, 223–231.
- Zepeda N et al. (2017) Apoptosis of mouse hippocampal cells induced by *Taenia crassiceps* metacystode factor. *Journal of Helminthology* **91**, 215–221.