

---

# Counting Intersecting and Pairs of Cross-Intersecting Families

---

PETER FRANKL and ANDREY KUPAVSKII<sup>†</sup>

<sup>1</sup>Department of Discrete Mathematics, Moscow Institute of Physics and Technology, Moscow, Russia  
and  
Ecole Polytechnique Fédérale de Lausanne, Switzerland  
(e-mail: kupavskii@yandex.ru)

Received 5 December 2016; revised 23 August 2017; first published online 16 October 2017

A family of subsets of  $\{1, \dots, n\}$  is called *intersecting* if any two of its sets intersect. A classical result in extremal combinatorics due to Erdős, Ko and Rado determines the maximum size of an intersecting family of  $k$ -subsets of  $\{1, \dots, n\}$ . In this paper we study the following problem: How many intersecting families of  $k$ -subsets of  $\{1, \dots, n\}$  are there? Improving a result of Balogh, Das, Delcourt, Liu and Sharifzadeh, we determine this quantity asymptotically for  $n \geq 2k + 2 + 2\sqrt{k \log k}$  and  $k \rightarrow \infty$ . Moreover, under the same assumptions we also determine asymptotically the number of *non-trivial* intersecting families, that is, intersecting families for which the intersection of all sets is empty. We obtain analogous results for pairs of cross-intersecting families.

2010 Mathematics subject classification: 05D05

## 1. Introduction

A *family* is a collection of subsets of an  $n$ -element set  $[n]$ . Collections  $\mathcal{F} \subset \binom{[n]}{k}$  are called  *$k$ -uniform families*. A family  $\mathcal{F}$  is called *intersecting* if  $F \cap F' \neq \emptyset$  holds for all  $F, F' \in \mathcal{F}$ . Similarly,  $\mathcal{F} \subset \binom{[n]}{k}$  and  $\mathcal{G} \subset \binom{[n]}{l}$  are called *cross-intersecting* if, for all  $F \in \mathcal{F}, G \in \mathcal{G}$ , one has  $F \cap G \neq \emptyset$ .

The research concerning intersecting families was initiated by Erdős, Ko and Rado, who determined the maximum size of intersecting families.

**Theorem 1.1 (Erdős, Ko and Rado [3]).** *Suppose that  $n \geq 2k > 0$  and  $\mathcal{F} \subset \binom{[n]}{k}$  is intersecting. Then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1}. \quad (1.1)$$

<sup>†</sup> Research by Andrey Kupavskii is supported by grant RNF 16-11-10014.

The family of all  $k$ -sets containing a fixed element shows that (1.1) is best possible. Hilton and Milner proved in a stronger form that for  $n > 2k$  these are the only families on which the equality is attained. We say that an intersecting family  $\mathcal{F}$  is *non-trivial* if  $\bigcap_{F \in \mathcal{F}} F = \emptyset$ , that is, if it cannot be pierced by a single point.

**Theorem 1.2 (Hilton and Milner [7]).** *Let  $n > 2k > 0$  and suppose that  $\mathcal{F} \subset \binom{[n]}{k}$  is a non-trivial intersecting family. Then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1. \tag{1.2}$$

For  $n = 2k + 1$  the difference between the upper bounds (1.1) and (1.2) is only  $k - 1$ . However, as  $n - 2k$  increases, this difference gets much larger. The number of subfamilies of  $\mathcal{F}$  is  $2^{|\mathcal{F}|}$ , and thus the ratio between the number of subfamilies of the Erdős–Ko–Rado family and that of the Hilton–Milner family is  $2^{k-1}$  for  $n = 2k + 1$  and grows very fast as  $n - 2k$  increases. This serves as an indication that most intersecting families are *trivial*, that is, they satisfy  $\bigcap_{F \in \mathcal{F}} F \neq \emptyset$ .

In an important recent paper Balogh, Das, Delcourt, Liu and Sharifzadeh [1] proved this in the following quantitative form. Let  $I(n, k)$  denote the *total* number of intersecting families  $\mathcal{F} \subset \binom{[n]}{k}$ .

**Theorem 1.3 (Balogh, Das, Delcourt, Liu and Sharifzadeh [1]).** *If  $n \geq 3k + 8 \log k$ , then*

$$I(n, k) = (n + o(1))2^{\binom{n-1}{k-1}}, \tag{1.3}$$

where  $o(1) \rightarrow 0$  as  $k \rightarrow \infty$ .

One of the main tools of the proof of (1.3) is a nice bound on the number of *maximal* (i.e. non-extendable) intersecting families (see Lemma 2.4 below). They obtain this bound using the following fundamental result of Bollobás.

**Theorem 1.4 (Bollobás [2]).** *Suppose that  $\mathcal{A} \subset \binom{[n]}{a}$ ,  $\mathcal{B} \subset \binom{[n]}{b}$ , with  $\mathcal{A} = \{A_1, \dots, A_m\}$ ,  $\mathcal{B} = \{B_1, \dots, B_m\}$ , satisfy  $A_i \cap B_j = \emptyset$  if and only if  $i = j$ . Then*

$$m \leq \binom{a+b}{a}. \tag{1.4}$$

Note that the bound (1.4) is independent of  $n$ . In [2] it is proved in a more general setting, not requiring uniformity, i.e. for  $\mathcal{A}, \mathcal{B} \subset 2^{[n]}$ . The uniform version (1.4) was rediscovered several years later by Jaeger and Payan [8] and Katona [9].

We are going to use (1.4) to obtain an upper bound on the number of maximal pairs of cross-intersecting families. Let  $CI(n, a, b, t)$  ( $CI(n, a, b, [t_1, t_2])$ ) denote the *number of pairs of cross-intersecting families*  $\mathcal{A} \subset \binom{[n]}{a}, \mathcal{B} \subset \binom{[n]}{b}$  with  $|\mathcal{A}| = t$  ( $t_1 \leq |\mathcal{A}| \leq t_2$ ). We also let

$$CI(n, a, b) := \sum_t CI(n, a, b, t).$$

We prove the following bound for the number of pairs of cross-intersecting families.

**Theorem 1.5.** Choose  $a, b, n \in \mathbb{N}$  and put

$$c := \max\{a, b\}, \quad T := \binom{n-a+b-1}{n-a}.$$

For  $n \geq a + b + 2\sqrt{c \log c} + 2 \max\{0, a - b\}$ ,  $a, b \rightarrow \infty$ , and  $b \gg \log a$  we have

$$CI(n, a, b) = (1 + \delta_{ab} + o(1))2^{\binom{n}{c}}, \tag{1.5}$$

$$CI(n, a, b, [1, T]) = (1 + o(1)) \binom{n}{a} 2^{\binom{n}{b} - \binom{n-a}{b}}, \tag{1.6}$$

where  $\delta_{ab} = 1$  if  $a = b$ , and 0 otherwise.

In fact, (1.6) easily implies (1.5) (see Section 2 for details). For a family  $\mathcal{F} \subset \binom{[n]}{k}$  we define the diversity  $\gamma(\mathcal{F})$  of  $\mathcal{F}$  to be  $|\mathcal{F}| - \Delta(\mathcal{F})$ , where

$$\Delta(\mathcal{F}) := \max_{i \in [n]} |\{F : i \in F \in \mathcal{F}\}|.$$

For an integer  $t$ , let  $I(n, k, t)$  ( $I(n, k, \geq t)$ ) denote the number of intersecting families with diversity  $t$  (at least  $t$ ). In particular,  $I(n, k, \geq 1)$  is the number of non-trivial intersecting families. With the help of (1.6) we obtain a refinement of Theorem 1.3.

**Theorem 1.6.** For  $n \geq 2k + 2 + 2\sqrt{k \log k}$  and  $k \rightarrow \infty$  we have

$$I(n, k) = (n + o(1))2^{\binom{n-1}{k-1}}, \tag{1.7}$$

$$I(n, k, \geq 1) = (1 + o(1))n \binom{n-1}{k} 2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}}. \tag{1.8}$$

Again, it is easy to see that (1.8) implies (1.7) (see Section 3 for details). In the next section we present the proof of Theorem 1.5, and in Section 3 we give the proof of Theorem 1.6.

### 2. Cross-intersecting families

Let us define the *lexicographic* order on the  $k$ -subsets of  $[n]$ . We have  $F \prec G$  in the lexicographic order if  $\min F \setminus G < \min G \setminus F$  holds. For example,  $\{1, 10\} \prec \{2, 3\}$ . For  $0 \leq m \leq \binom{[n]}{k}$ , let  $\mathcal{L}^{(k)}(m)$  denote the family of first  $m$   $k$ -sets in the lexicographic order. For example,

$$\mathcal{L}^{(k)}\left(\binom{n-1}{k-1}\right) = \left\{F \in \binom{[n]}{k} : 1 \in F\right\}.$$

Next we state the Kruskal–Katona theorem [11, 10], which is one of the most important results in extremal set theory.

**Theorem 2.1 (Kruskal [11], Katona [10]).** If  $\mathcal{A} \subset \binom{[n]}{a}$  and  $\mathcal{B} \subset \binom{[n]}{b}$  are cross-intersecting, then  $\mathcal{L}^{(a)}(|\mathcal{A}|)$  and  $\mathcal{L}^{(b)}(|\mathcal{B}|)$  are cross-intersecting as well.

Computationwise, the bounds arising from the Kruskal–Katona theorem are not easy to handle. Lovász [13] found a slightly weaker but very handy form, which may be stated as follows.

**Theorem 2.2 (Lovász [13]).** *Let  $n \geq a + b$ , and consider a pair of cross-intersecting families  $\mathcal{A} \subset \binom{[n]}{a}$ ,  $\mathcal{B} \subset \binom{[n]}{b}$ . If  $|\mathcal{A}| = \binom{x}{n-a}$  for a real number  $x \geq n - a$ , then we obtain*

$$|\mathcal{B}| \leq \binom{n}{b} - \binom{x}{b}. \tag{2.1}$$

Note that for  $x \geq k - 1$  the polynomial  $\binom{x}{k}$  is a monotone increasing function of  $x$ . Thus  $x$  is uniquely determined by  $|\mathcal{A}|$  and  $a$ .

We would also need the following result, whose proof is based on Theorem 2.1 and which is a combination of a result of Frankl and Tokushige [5] (Theorem 2 in [5]) and two results of Kupavskii and Zakharov [12] (part 1 of Theorem 1 and Corollary 1).

**Theorem 2.3 (Frankl and Tokushige [5], Kupavskii and Zakharov [12]).** *Let  $n > a + b$  and suppose that the families  $\mathcal{A} \subset \binom{[n]}{a}$ ,  $\mathcal{B} \subset \binom{[n]}{b}$  are cross-intersecting. If, for some real number  $\alpha \geq 1$ , we have*

$$\binom{n - \alpha}{n - a} \leq |\mathcal{A}| \leq \binom{n - a + b - 1}{n - a},$$

then

$$|\mathcal{A}| + |\mathcal{B}| \leq \binom{n}{b} + \binom{n - \alpha}{a - \alpha} - \binom{n - \alpha}{b}. \tag{2.2}$$

Note that the upper bound on  $|\mathcal{A}|$  in this theorem is exactly the same as in (1.6).

We now begin the proof of Theorem 1.5. First we show that (1.6) implies (1.5). Without loss of generality we may assume for this paragraph that  $c = b \geq a$ . For  $b > a$  we have  $T \geq \binom{n}{a}$  and thus  $CI(n, a, b, t) = 0$  for  $t > T$ . Therefore, we have  $CI(n, a, b) = CI(n, a, b, 0) + CI(n, a, b, [1, T])$ . If  $a = b$ , then  $T = \binom{n-1}{a-1}$ , and it follows from Theorem 2.1 that if  $\mathcal{A}, \mathcal{B} \subset \binom{[n]}{a}$  are cross-intersecting, then  $\min\{|\mathcal{A}|, |\mathcal{B}|\} \leq \binom{n-1}{a-1}$ . Therefore, in the case  $a = b$  we have

$$2CI(n, a, b, 0) - 1 \leq CI(n, a, b) \leq 2(CI(n, a, b, 0) + CI(n, a, b, [1, T])).$$

The ‘ $-1$ ’ in the first inequality stands for a pair of empty families, which is counted twice. At the same time, we have  $CI(n, a, b, 0) = 2\binom{n}{b}$ . Thus, in both cases  $b > a$  and  $b = a$  it is sufficient to show that the right-hand side of (1.6) is  $o(2\binom{n}{b})$ . We first note that  $n - a - b \geq \sqrt{n}$  for  $b \geq a$ ,  $n \geq a + b + 2\sqrt{b \log b}$ , since  $4b \log b \geq a + b + 2\sqrt{b \log b}$ . The rest is done by a simple calculation:

$$\frac{CI(n, a, b, [1, T])}{2\binom{n}{b}} = \binom{n}{a} 2^{-(n-a)} \leq 2^{n-(b+\sqrt{n})} = o(1).$$

Next, we discuss the proof of the lower bound in (1.6). To obtain that many pairs of intersecting families, take

$$\mathcal{A} := \{A\}, A \in \binom{[n]}{a}, \quad \mathcal{B}(A) := \{B \in \binom{[n]}{b} : B \cap A \neq \emptyset\}.$$

Next, choose an arbitrary subfamily  $\mathcal{B} \subset \mathcal{B}(A)$ . We only need to ensure that few of these pairs of subfamilies are counted twice. In fact we count a pair of families twice only in the case when  $a = b$  and both  $\mathcal{A}, \mathcal{B}$  consist of one set. The number of such pairs is  $\binom{[n]}{a}^2$ , and is negligible compared to the right-hand side of (1.6).

We pass to the proof of the upper bound.

(i)  $2 \leq |\mathcal{A}| \leq n - a$ . Applying Theorem 2.1, the size of the (unique) maximal family  $\mathcal{B}'$  that forms a cross-intersecting pair with  $\mathcal{A}$  is maximized if  $\mathcal{A}$  consists of two sets  $A_1, A_2$  that intersect in  $a - 1$  elements. Therefore,

$$|\mathcal{B}'| \leq \binom{n}{b} - \binom{n-a+1}{b} + \binom{n-a-1}{b-2}.$$

Any other family  $\mathcal{B}$  that forms a cross-intersecting pair with  $\mathcal{A}$  must be a subfamily of  $\mathcal{B}'$ .

So we can bound the number of pairs of cross-intersecting families  $\mathcal{A}, \mathcal{B}$  with  $2 \leq |\mathcal{A}| \leq n - a$  as follows:

$$\frac{\sum_{t=2}^{n-a} CI(n, a, b, t)}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq \sum_{t=2}^{n-a} \binom{\binom{n}{a}}{t} \frac{2^{\binom{n}{b} - \binom{n-a+1}{b} + \binom{n-a-1}{b-2}}}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq 2^{n^2} 2^{-(n-a-1)} = o(1).$$

(ii)  $n - a + 1 \leq |\mathcal{A}| \leq \binom{n-u}{n-a}$ , where  $u = \sqrt{c \log c} + \max\{0, a - b\}$ . Note that  $n - a + 1 = \binom{n-a+1}{n-a}$ . In this case the bound is similar, but we use Theorem 2.2 to bound the size of  $|\mathcal{B}|$ . For  $\mathcal{A}$  with

$$|\mathcal{A}| = t := \binom{n-u'}{n-a},$$

where  $u \leq u' \leq a - 1$ , we get

$$|\mathcal{B}| \leq 2^{\binom{n}{b} - \binom{n-u'}{b}},$$

and since  $\binom{\binom{n}{a}}{t} \leq 2^{nt}$ , we have the following bound:

$$\frac{CI(n, a, b, t)}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq \binom{\binom{n}{a}}{t} \frac{2^{\binom{n}{b} - \binom{n-u'}{b}}}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq 2^{n \binom{n-u'}{n-a}} 2^{-(n-u'-1)}.$$

At the same time we have  $n \geq a + b + 2u$  and

$$\begin{aligned} \frac{\binom{n-u'}{n-a}}{\binom{n-u'-1}{b-1}} &= \frac{n-u'}{b} \prod_{i=0}^{n-a-b-1} \frac{n-b-u'-i}{n-a-i} \\ &\leq \frac{n}{b} \prod_{i=0}^{n-a-b-1} \frac{n-a-\sqrt{c \log c}-i}{n-a-i} \\ &\leq \frac{n}{b} e^{-\sqrt{c \log c} \sum_{i=b+1}^{n-a} (1/i)} \leq \frac{1}{2n} \end{aligned} \tag{2.3}$$

for sufficiently large  $c$ . Indeed,

$$\sqrt{c \log c} \sum_{i=b+1}^{n-a} \frac{1}{i} \geq \sqrt{c \log c} \sum_{i=b+1}^{b+2\sqrt{c \log c}} \frac{1}{i} \geq (1+o(1)) \frac{2(\sqrt{c \log c})^2}{c} = (2+o(1)) \log c,$$

which justifies (2.3) for  $n \leq b^{3/2}$ . For  $n > b^{3/2}$  we have

$$\sqrt{c \log c} \sum_{i=b+1}^{n-a} \frac{1}{i} \geq (1 + o(1)) \sqrt{c \log c} \log \frac{n}{b} \geq (1 + o(1)) \sqrt{c \log c} \log n^{2/3} \gg \log n,$$

which justifies (2.3) for  $n > b^{3/2}$ .

We conclude that

$$\sum_{t=n-a+1}^{\binom{n-u}{n-a}} \frac{CI(n, a, b, t)}{2^{\binom{n}{b} - \binom{n-b}{b}}} \leq 2^{-\frac{1}{2} \binom{n-u-1}{b-1}} = o(1).$$

(iii)  $\binom{n-u}{n-a} < |\mathcal{A}| \leq T$ , where  $u = \sqrt{b \log b} + \max\{0, a - b\}$ . Using the Bollobás set-pair inequality, it is not difficult to obtain the following bound on the number of maximal pairs of cross-intersecting families.

**Lemma 2.4.** *The number of maximal cross-intersecting pairs  $\mathcal{A}' \subset \binom{[n]}{a}, \mathcal{B}' \subset \binom{[n]}{b}$  is at most*

$$\left[ \binom{n}{a} \binom{n}{b} \right]^{\binom{a+b}{a}}.$$

We note that the proof is very similar to the proof of an analogous statement for intersecting families from [1].

**Proof.** Find a minimal  $\mathcal{B}'$ -generating family  $\mathcal{M} \subset \mathcal{A}'$  such that

$$\mathcal{B}' = \left\{ B \in \binom{[n]}{b} : B \cap M \neq \emptyset \text{ for all } M \in \mathcal{M} \right\}.$$

We claim that  $|\mathcal{M}| \leq \binom{a+b}{a}$ . Indeed, due to minimality, for each set  $M' \in \mathcal{M}$  the family

$$\mathcal{B}'' := \left\{ B \in \binom{[n]}{b} : B \cap M \neq \emptyset \text{ for all } M \in \mathcal{M} - \{M'\} \right\}$$

strictly contains  $\mathcal{B}'$ . Therefore, there is a set  $B$  in  $\mathcal{B}'' \setminus \mathcal{B}'$  such that  $B \cap M' = \emptyset, B \cap M \neq \emptyset$  for all  $M \in \mathcal{M} - \{M'\}$ . Applying the inequality (1.4) to  $\mathcal{M}$  and the collection of such sets  $B$ , we get that  $|\mathcal{M}| \leq \binom{a+b}{b}$ .

Interchanging the roles of  $\mathcal{A}'$  and  $\mathcal{B}'$ , we get that a minimal  $\mathcal{A}'$ -generating family has size at most  $\binom{a+b}{b}$  as well. Now the bound stated in the lemma is just a crude upper bound on the number of ways one can choose these two generating families out of  $\binom{[n]}{a}$  and  $\binom{[n]}{b}$ , respectively.  $\square$

Combined with the bound (2.2) on the size of any maximal pair of families with such cardinalities, we get that

$$\frac{CI(n, a, b, [\binom{n-u}{n-a}, T])}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq \left[ \binom{n}{a} \binom{n}{b} \right]^{\binom{a+b}{a}} \frac{2^{\binom{n}{b} + \binom{n-u}{n-a} - \binom{n-u}{b}}}{2^{\binom{n}{b} - \binom{n-a}{b}}} \leq 2^{2n \binom{a+b}{b}} 2^{\binom{n-u}{n-a} - \binom{n-u-1}{b-1}}. \tag{2.4}$$

We also have

$$\frac{\binom{a+b}{b}}{\binom{n-u-1}{b-1}} = \frac{n-u}{b} \prod_{i=0}^{b-1} \frac{a+b-i}{n-u-i} \leq n \left(\frac{a+b}{n-u}\right)^b \leq \frac{1}{4n}.$$

Indeed, the last inequality is clearly valid for  $n \geq (a+b)^2, b \rightarrow \infty$ . If  $n < (a+b)^2$ , then the penultimate expression is at most

$$e^{\log n - (b(n-a-b-u))/(n-u)} \leq e^{2\log(a+b) - (b(u+\max\{0, a-b\}))/O(a+b)} \leq e^{2\log(a+b) - \Omega(\min\{b, u\})}.$$

Since by the assumption we have  $b \gg \log(a+b)$  and also, obviously,  $u \gg \log(a+b)$ , the last expression is at most  $e^{-4\log(a+b)} < 1/(4n)$ .

Taking into account (2.3), which is valid for  $u' = u$ , we conclude that the right-hand side of (2.4) is  $o(1)$ .

### 3. Intersecting families

We need a theorem due to Frankl [4], proved in the following, slightly stronger form in [12].

**Theorem 3.1 (Frankl [4], Kupavskii and Zakharov [12]).** *Let  $\mathcal{F} \subset \binom{[n]}{k}$  be an intersecting family, and  $n > 2k$ . Then, if  $\gamma(\mathcal{F}) \geq \binom{n-u-1}{k-u}$  for some real  $3 \leq u \leq k$ , then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1} + \binom{n-u-1}{k-u} - \binom{n-u-1}{k-1}. \tag{3.1}$$

We go on to the proof of Theorem 1.6. Let us first show that (1.8) implies (1.7). Indeed, using that  $n - 2k - 1 \geq \sqrt{n}$  and  $k \rightarrow \infty$  in the assumptions of Theorem 1.6, we get

$$\frac{I(n, k, \geq 1)}{2^{\binom{n-1}{k-1}}} \sim n \binom{n-1}{k} 2^{-\binom{n-k-1}{k-1}} \leq 2^{n+\log n - \binom{k+\sqrt{n}}{k-1}} = o(1).$$

Therefore,

$$I(n, k, \geq 1) = o(2^{\binom{n-1}{k-1}}).$$

On the other hand, it is easy to see that

$$I(n, k, 0) = (n + o(1))2^{\binom{n-1}{k-1}}$$

(for the proof see [1]).

Let us prove the lower bound in (1.8). For  $S \in \binom{[n]}{k}, i \in [n] \setminus S$ , define the family

$$\mathcal{H}(i, S) := \{S\} \cup \left\{ H \in \binom{[n]}{k} : i \in H, H \cap S \neq \emptyset \right\}.$$

Due to Theorem 1.2, these families are the largest non-trivial intersecting families. We have

$$|\mathcal{H}(i, S)| = \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1,$$

and each such family contains no fewer than

$$2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}} - k2^{\binom{n-2}{k-2}} = (1 + o(1))2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}} \tag{3.2}$$

non-trivial intersecting subfamilies, as  $k \rightarrow \infty$ . Indeed, a subfamily of  $\mathcal{H}(i, S)$  containing  $S$  is non-trivial unless all sets containing  $i$  contain also a fixed  $j \in S$ . In other words, they must be a subset of a family

$$\mathcal{I}(i, j, S) := \{S\} \cup \left\{ I \in \binom{[n]}{k} : i, j \in S \right\}.$$

The number of subfamilies of  $\mathcal{I}(i, j, S)$  containing  $S$  is  $2^{\binom{n-2}{k-2}}$ . Next, we have

$$\binom{n-1}{k-1} - \binom{n-k-1}{k-1} - \binom{n-2}{k-2} \geq \binom{n-3}{k-2},$$

and thus the last inequality in the displayed formula above holds since  $2^{\binom{n-3}{k-2}} \gg k$ . Denote the set of all non-trivial subfamilies of  $\mathcal{H}(i, S)$  by  $\tilde{\mathcal{H}}(i, S)$ .

Therefore,

$$\sum_{S \in \binom{[n]}{k}, i \notin S} |\tilde{\mathcal{H}}(i, S)| = (1 + o(1))n \binom{n-1}{k} 2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}}.$$

On the other hand, the pairwise intersections of these families are small: the families from  $\tilde{\mathcal{H}}(i, S) \cap \tilde{\mathcal{H}}(i, S')$  form the set  $I(n, k, 2)$ , and we do (somewhat implicitly) show in the proof that  $I(n, k, 2) = o(I(n, k, 1))$ . It could also be verified by a simple, direct, but somewhat tedious calculation. Therefore, the lower bound is justified.

Next we prove the upper bound. For  $i \in [n]$  and  $\mathcal{F} \subset \binom{[n]}{k}$  we use the standard notation

$$\begin{aligned} \mathcal{F}(i) &:= \{F - \{i\} : i \in F \in \mathcal{F}\} \subset \binom{[n] - \{i\}}{k-1}, \\ \mathcal{F}(\bar{i}) &:= \{F \in \mathcal{F} : i \notin F\} \subset \binom{[n] - \{i\}}{k}. \end{aligned}$$

Note that if  $\mathcal{F}$  is intersecting then  $\mathcal{F}(i)$  and  $\mathcal{F}(\bar{i})$  are cross-intersecting.

We count the number of families with different diversity separately. The number of families  $\mathcal{F}$  with  $i$  being the most popular element and  $\gamma(\mathcal{F}) \leq \binom{n-3}{k-2}$  is at most the number of cross-intersecting pairs  $\mathcal{F}(\bar{i}), \mathcal{F}(i)$ .

Therefore, we may apply (1.6) with  $n' := n - 1, a := k, b := k - 1$ , and get that the number of such families  $\mathcal{F}$  is at most

$$(1 + o(1)) \binom{n-1}{k} 2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}}.$$

Note that  $n' \geq a + b + 2\sqrt{a \log a} + 2$  and, in terms of Theorem 1.5, we have  $T = \binom{n-3}{k-2}$  for our case. Multiplying the number of such families by the number of choices of  $i$ , we get the claimed asymptotic.

It remains only to prove that there are few families with diversity larger than  $\binom{n-3}{k-2}$ . Using the upper bound  $\binom{n}{k} \binom{2k-1}{k-1}$  for the number of maximal intersecting families in  $\binom{[n]}{k}$  obtained in [1] (see Lemma 2.4 for the proof of a similar statement), combined with the bound (3.1) on the size of



any maximal family with such diversity, we get that

$$\frac{I(n, k, \geq \binom{n-3}{k-2})}{2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}}} \leq \binom{n}{k} \frac{\binom{2k-1}{k-1} 2^{\binom{n-1}{k-1} + \binom{n-4}{k-3} - \binom{n-4}{k-1}}}{2^{\binom{n-1}{k-1} - \binom{n-k-1}{k-1}}} \leq 2^{n \binom{2k-1}{k-1}} 2^{\binom{n-4}{k-3} - \binom{n-5}{k-2}}. \quad (3.3)$$

Putting  $n = 2k + x$ , we have

$$\frac{\binom{n-4}{k-3}}{\binom{n-5}{k-2}} = \frac{(n-4)(k-2)}{(n-k-1)(n-k-2)} \leq \frac{(2k+x)k}{(k+x-2)^2} \leq 1 - \frac{x^2}{(k+x)^2} \leq 1 - \frac{1}{k}.$$

On the other hand,

$$\frac{\binom{2k-1}{k-1}}{\binom{n-5}{k-2}} = \frac{n-4}{k-1} \prod_{i=1}^k \frac{2k-i}{n-3-i} \leq n \left( \frac{2k}{n-3} \right)^k \leq \frac{1}{2kn},$$

where the last inequality is clearly valid for  $n \geq 2k + 2 + 2\sqrt{k \log k}$  and sufficiently large  $k$ . We conclude that the right-hand side of (3.3) is at most  $2^{-\frac{1}{2k} \binom{n-5}{k-2}} = o(1)$ .

## References

- [1] Balogh, J., Das, S. A., Delcourt, M., Liu, H. and Sharifzadeh, M. (2015) Intersecting families of discrete structures are typically trivial. *J. Combin. Theory Ser. A* **132** 224–245.
- [2] Bollobás, B. (1965) On generalized graphs. *Acta Math. Acad. Sci. Hungar.* **16** 447–452.
- [3] Erdős, P., Ko, C. and Rado, R. (1961) Intersection theorems for systems of finite sets. *Quart. J. Math.* **12** 313–320.
- [4] Frankl, P. (1987) Erdős–Ko–Rado theorem with conditions on the maximal degree. *J. Combin. Theory Ser. A* **46** 252–263.
- [5] Frankl, P. and Tokushige, N. (1992) Some best possible inequalities concerning cross-intersecting families. *J. Combin. Theory Ser. A* **61** 87–97.
- [6] Hilton, A. J. W. (1976) The Erdős–Ko–Rado theorem with valency conditions. Unpublished manuscript.
- [7] Hilton, A. J. W. and Milner, E. C. (1967) Some intersection theorems for systems of finite sets. *Quart. J. Math. Oxford* **18** 369–384.
- [8] Jaeger, F. and Payan, C. (1971) Nombre maximal d’arêtes d’un hypergraphe critique de rang  $h$ . *CR Acad. Sci. Paris* **273** 221–223.
- [9] Katona, G. O. H. (1974) Solution of a problem of A. Ehrenfeucht and J. Mycielski. *J. Combin. Theory Ser. A* **17** 265–266.
- [10] Katona, G. O. H. (1987) A theorem of finite sets. In *Classic Papers in Combinatorics* (I. Gessel and G.-C. Rota, eds), Birkhäuser, pp. 381–401.
- [11] Kruskal, J. B. (1963) The number of simplices in a complex. In *Mathematical Optimization Techniques* (R. Bellman, ed.), University of California Press, pp. 251–278.
- [12] Kupavskii, A. and Zakharov, D. Regular bipartite graphs and intersecting families. Accepted at *J. Combin. Theory Ser. A* arXiv:1611.03129
- [13] Lovász, L. (1979) *Combinatorial Problems and Exercises*, North-Holland.