

EFFECTIVE CYCLES ON SOME LINEAR BLOWUPS OF PROJECTIVE SPACES

NORBERT PINTYE AND ARTIE PRENDERGAST-SMITH 

Abstract. We compute cones of effective cycles on some blowups of projective spaces in general sets of lines.

§1. Introduction

Cones of curves and divisors have played a central role in birational geometry since the groundbreaking work of Mori in the early 1980s. There are general results, such as the cone theorem, describing the structure of these cones, as well as numerous explicit calculations in cases of geometric interest.

More recently, there has been increased interest in cycles of intermediate dimensions. Debarre *et al.* [DELV] showed that, in general, these cycles do not share the good properties of divisors or curves: in particular, numerical positivity need not imply geometric positivity for such cycles. Nevertheless, there has been significant progress in extending the theoretical understanding of such cycles, due to Fulger and Lehmann [FL1, FL2], Ottem [Ott] and others. By contrast, the number of examples in which cones of effective cycles have been explicitly computed is relatively small. The most significant results to date were found by Coskun *et al.* [CLO], who computed cones of cycles on blowups of projective spaces at sets of points.

In this paper, we compute cones of effective cycles on some varieties obtained by blowing up general sets of lines in projective space. Blowups of this kind can be seen as the next natural examples to consider after those of Coskun–Lesieutre–Ottem; cones of divisors on these spaces were computed in some cases by Dumitrescu *et al.* [DPU].

Cones of cycles on these spaces are more complicated to compute than those of point blowups in two ways: first, a hyperplane in projective space cannot contain many general lines, and so inductive techniques tend to be less useful; second, the coefficients of the intersection form on the blowup vary with dimension, making uniform statements more difficult to find. In spite of these difficulties, we are able to compute cones in some interesting examples, which we now explain.

Blowing up a small number of lines in projective space gives a toric variety, so the cone of effective cycles is generated by torus-invariant subvarieties, hence linear subspaces. Our main results show that linear generation continues to hold when the number of lines is increased beyond the toric range: for example, the blowup of \mathbf{P}^4 in more than two lines is no longer toric, but we show in Theorem 4.2 that its cone of 2-cycles is still linearly generated when we blow up in three or four lines. Similarly, in Theorem 5.1, we show that the cones of 2-cycles is linearly generated when we blow up at most five lines in \mathbf{P}^5 , but the cone of 3-cycles fails to be linearly generated once we blow up four lines. Finally,

Received January 8, 2019. Revised November 13, 2019. Accepted November 13, 2019.
2010 Mathematics subject classification. 14C25, 14N20.

in Section 6, we complement these theorems with some results about linear generation of cones of curves and divisors.

Our results are summarized in the following tables. In each table, the entry in row k and column r shows whether the cone of effective k -cycles on the blowup of projective space of the relevant dimension in r general lines is linearly generated (or if the answer is not known). Note that once linear generation fails for a blowup, it fails for all further blowups, so any entry to the right of the symbol \times in a given row is also not linearly generated.

Dimension 4

$k \setminus r$	≤ 2	3	4	5	6	7	8	9	10
1	✓	✓	✓	✓	✓	✓ _{5.1}	?	?	× _{5.1}
2	✓	✓	✓ _{3.2}	× _{3.2}					
3	✓	✓	✓ _{5.3}	× _{5.3}					

Dimension 5

$k \setminus r$	≤ 3	4	5	6
1	✓	✓	✓ _{5.2}	× _{5.2}
2	✓	✓	✓ _{4.1}	?
3	✓ _{1.4}	× _{4.2}		
4	✓ _{1.4}	× _{5.4}		

The pattern we find agrees with Coskun–Lesieutre–Ottaviani’s results, namely that as we blow up more, cones of lower-dimensional cycles remain linearly generated for longer than do cones of higher-dimensional cycles. It would be interesting to find uniform bounds ensuring linear generation for blowups of projective space in general sets of linear subspaces of arbitrary dimension.

§2. Preliminaries

We work throughout over an algebraically closed field of characteristic zero. This assumption will be used in some computational proofs, where our calculations take place over \mathbf{Q} .

2.1 Intersection theory

Our goal in this paper is to compute cones of cycles. The natural contexts for these cones are the spaces of numerical classes of cycles, which we now introduce. In the examples we will consider, these spaces are just Chow groups with real coefficients, but we use the language of numerical classes for consistency with the general theory.

Let X be a smooth proper variety of dimension n . Let $Z_k(X)$ denote the group of algebraic cycles of dimension k on X . We define the vector space of numerical classes of k -cycles to be

$$N_k(X) := (Z_k(X)/\equiv) \otimes \mathbf{R},$$

where \equiv denotes numerical equivalence of cycles. For each k , this is a finite-dimensional real vector space, and intersection gives a perfect pairing $N_k(X) \times N_{n-k}(X) \rightarrow \mathbf{R}$. For convenience, we often write $N^k(X)$ instead of $N_{n-k}(X)$. For a k -dimensional subvariety Z in X , we write $[Z]$ to denote its class in $N_k(X)$. A fundamental feature of this product is positivity of proper intersections: if X is a smooth proper variety of dimension n , and V

and W are subvarieties of dimension k and $n - k$, respectively, such that $V \cap W$ is a finite set, then $[V] \cdot [W] \geq 0$.

A class $\alpha \in N_k(X)$ is *effective* if there are subvarieties Z_1, \dots, Z_m and nonnegative real numbers r_1, \dots, r_m such that $\alpha = \sum_{i=1}^m r_i [Z_i]$. A class $\alpha \in N^k(X)$ is called *nef* if $\alpha \cdot [Z] \geq 0$ for every k -dimensional subvariety Z in X or, equivalently, if $\alpha \cdot \beta \geq 0$ for every effective class $\beta \in N_k(X)$. We need some basic facts about the behavior of nef cycles under morphisms.

PROPOSITION 2.1. *Let $f : Y \rightarrow X$ be a morphism of smooth projective varieties.*

- (a) *If $\alpha \in N^k(X)$ is nef, then $f^*\alpha \in N^k(Y)$ is nef.*
 (b) *If f is surjective and $\alpha \in N_k(X)$ is a cycle such that $f^*\alpha$ is nef, then α is nef.*

Proof. (a): If $\beta \in N_k(Y)$ is effective, then $f_*\beta$ is also effective by definition of pushforward. So if $\alpha \in N^k(X)$ is nef, then using the projection formula for cycles, we get $f^*\alpha \cdot \beta = \alpha \cdot f_*\beta \geq 0$ for every effective cycle β in $N_k(Y)$.

(b): Let $\beta \in N^k(X)$ be an effective class. Since f is surjective, by a standard hyperplane section argument, there exists an effective class $\tilde{\beta} \in N^k(Y)$ such that $f_*\tilde{\beta} = \beta$. By the projection formula and nefness of $f^*\alpha$, we have $\alpha \cdot \beta = f^*\alpha \cdot \tilde{\beta} \geq 0$, showing that α is nef as required. \square

In general, the intersection of nef cycles need not be nef [DELV, Corollary 2.2], but for divisors, this is true.

LEMMA 2.2. *Let X be a smooth projective variety. If D and E are nef divisor classes on X , then DE is a nef class in $N^2(X)$.*

The statement is true more generally for the intersection of any number of nef divisor classes, but we will only prove the case that we need.

Proof. We need to prove that for any effective class $\alpha \in N_2(X)$, we have $DE \cdot \alpha \geq 0$. Since E is nef, we can find a sequence of ample divisor classes $\{E_i\}$ converging to E in $N^1(X)$. For each E_i , the intersection $E_i\alpha$ is effective, and so $D \cdot (E_i\alpha) \geq 0$. Taking the limit, we get $DE \cdot \alpha = \lim_i D \cdot (E_i\alpha) \geq 0$, as required. \square

2.2 Numerical classes on blowups

In the rest of the paper, we will write $X_{r,s}^n$ to denote the blowup of \mathbf{P}^n in a collection of r general lines L_1, \dots, L_r and s general points p_1, \dots, p_s . Our main examples have $s = 0$, and we denote these simply by X_r^n .

The ring $N^*(X_{r,s}^n) = CH(X_{r,s}^n) \otimes \mathbf{R}$ is generated by classes H, E_i for $i = 1, \dots, r$ and e_i for $i = 1, \dots, s$, which are, respectively, the pullback of the hyperplane class on \mathbf{P}^n , the exceptional divisors of the blowups of the L_i and the exceptional divisors of the blowups of the p_i . We will use the following intersection numbers among these classes [EH, Corollary 9.12]:

$$\begin{aligned} H^n &= 1, & E_i^n &= (-1)^n(n-1), & e_i^n &= (-1)^{n-1} \\ H \cdot E_i^{n-1} &= (-1)^n, & H^j \cdot E_i^{n-j} &= H^k \cdot e_i^{n-k} = 0 & \text{for } i \geq 1, \quad k \geq 0. \end{aligned}$$

We also need to know the numerical classes on $X_{r,s}^n$ of the proper transforms of certain subvarieties of \mathbf{P}^n . The blowup formula [Fu, Theorem 6.7] allows us to calculate these as long as we know the Segre classes of the blowup center inside the subvariety: in particular,

when we blow up lines and points, these are easy to compute. In particular, we note the following.

COROLLARY 2.3. *Let $X_{r,s}^n$ be the blowup of \mathbf{P}^n in r general lines and s general points. Let:*

- U be a linear space of codimension k intersecting L_i transversely;
- V be a linear space of codimension k containing L_i and
- Q be a quadric of codimension k containing L_i .

The numerical classes of the proper transforms of these spaces have the following coefficients:

	H^k	HE_i^{k-1}	E_i^k
$[\tilde{U}]$	1	$(-1)^{k-1}$	0
$[\tilde{V}]$	1	$(-1)^{k-1}k$	$(-1)^k$
$[\tilde{Q}]$	2	$(-1)^{k-1}(k+1)$	$(-1)^k$

If Z is any subvariety of codimension k containing p_i as a smooth point, then the coefficient of e_i^k in $[\tilde{Z}]$ equals $(-1)^k$.

2.3 Cones of cycles

For a smooth projective variety X , the *pseudoeffective cone* $\overline{\text{Eff}}_k(X)$ is the closed convex cone in $N_k(X)$ generated by numerical classes of k -dimensional subvarieties of X . The *nef cone* $\text{Nef}^k(X)$ is the cone spanned by all nef classes in $N^k(X)$: in other words, it is the dual cone of $\overline{\text{Eff}}_k(X)$.

Now we specialize the discussion to our examples $X_{r,s}^n$. A subvariety of $X_{r,s}^n$ is called *linear* if it is one of the following:

- (a) the proper transform on $X_{r,s}^n$ of a linear subspace of \mathbf{P}^n ;
- (b) the pullback to $E_i \cong \mathbf{P}^1 \times \mathbf{P}^{n-2}$ of a linear subspace in one of the factors or
- (c) a linear subspace in $e_i \cong \mathbf{P}^{n-1}$.

The *linear cone* $\text{Lin}_k(X_{r,s}^n)$ is the cone in $N_k(X_{r,s}^n)$ generated by the finitely many classes of k -dimensional linear subvarieties. We say that the pseudoeffective cone of k -cycles on $\overline{\text{Eff}}(X_{r,s}^n)$ is *linearly generated* if it equals the linear cone $\text{Lin}_k(X_{r,s}^n)$. Note that any blowup map $X_{r,s}^n \rightarrow X_{r-a,s-b}^n$ maps the effective cone onto the effective cone and the linear cone onto the linear cone, so if $\overline{\text{Eff}}(X_{r,s}^n)$ is linearly generated, then so too is $\overline{\text{Eff}}(X_{r-a,s-b}^n)$.

2.4 Toric varieties

Cones of cycles on toric varieties are well understood. For later use, let us record the facts we need.

PROPOSITION 2.4. *Let X be a normal proper toric variety. Then, $\overline{\text{Eff}}_k(X)$ is generated by the finitely many classes of k -dimensional torus-invariant subvarieties on X . Consequently, if the variety $X_{r,s}^n$ is toric, then $\overline{\text{Eff}}_k(X_{r,s}^n)$ is linearly generated for all k .*

Proof. The first statement is well known; a reference is [Li, Proposition 3.1].

For the second statement, note that the torus-invariant subvarieties of \mathbf{P}^n are exactly the coordinate subspaces, so if $X_{r,s}^n$ is a toric blowup of \mathbf{P}^n , any torus-invariant subvariety on $X_{r,s}^n$ that comes from \mathbf{P}^n is the proper transform of a coordinate subspace and hence is linear.

On the other hand, every exceptional divisor of $X \rightarrow \mathbf{P}^n$ is of the form $E_i \cong \mathbf{P}^1 \times \mathbf{P}^{n-2}$ or $e_i \cong \mathbf{P}^{n-1}$, so the torus-invariant subvarieties of the exceptional divisor are also linear. \square

2.5 Computations

In this paper, we will use computer algebra in several different contexts. In all cases, we use the computer algebra system Macaulay2. In particular, for all computations of dual numerical cones, we use the package `Normaliz` [Nor] for Macaulay2. Note that for compactness, we always list the generators of all cones “up to permutation”: that is, a full list of generators is obtained from our list by permuting indices in the appropriate way.

The full outputs of our computations are available in ancillary files provided with this paper [M2]. The name of each file in the repository indicates the result in the paper in which the output of the computation is used.

§3. Codimension-2 linear spaces

In this section, we prove that codimension-2 linear spaces incident to lines give nef classes in X_r^n for $r \leq n \leq 5$. The main idea of the proof is to verify by a dimension count that we can find such a linear space properly intersecting any given subvariety of complementary dimension. As mentioned in Section 1, proper intersections are nonnegative, so this is sufficient to prove our claim.

We begin with some preparatory results about intersections of Schubert cycles.

LEMMA 3.1. *Let l_1, \dots, l_4 be a set of four distinct lines in \mathbf{P}^3 and let $\Lambda \subset \mathbf{G}(1, 3)$ be the set of lines touching all four. Then, one of the following is true:*

- (a) *the set Λ has dimension 2, in which case one of the following is true:*
 - (i) *all four lines are concurrent (that is, they all meet in a single point) or*
 - (ii) *all four lines are coplanar;*
- (b) *the set Λ has dimension 1, in which case one of the following is true:*
 - (i) *the lines are all pairwise skew and lie on a smooth quadric surface $Q \in \mathbf{P}^3$;*
 - (ii) *there are exactly two pairs of intersecting lines, say l_1, l_2 and l_3, l_4 , and the intersection point of l_1, l_2 lies in the plane spanned by l_3, l_4 ;*
 - (iii) *there are three concurrent lines, say l_1, l_2, l_3 , and the line l_4 is skew to all others*
or
 - (iv) *there are three coplanar lines, say l_1, l_2, l_3 , and the line l_4 is skew to all the others;*
- (c) *the set Λ has dimension 0.*

Proof. For each case listed in (a) and (b) above, the given dimension count is straightforward to verify. It remains to check that in all other cases, the set Λ has dimension 0. In the case that all lines are pairwise skew, this is well known, so we must consider the cases in which some of the lines intersect. There are two possibilities not covered by the list above:

- two lines, say l_1 and l_2 , intersect, and all other pairs are skew;
- there are exactly two pairs l_1, l_2 and l_3, l_4 of intersecting lines, and neither of the intersection points of the two pairs lies in the plane spanned by the other pair.

In the first case, any line intersecting all four lines must either lie in the plane spanned by l_1 and l_2 or pass through the intersection point of l_1 and l_2 . In each case, however, there is such a unique line which also intersects l_3 and l_4 .

In the second case, no line contained in either of the planes spanned by two intersecting lines can intersect the other two lines. So the only line intersecting all four lines is the line joining the two intersection points of the pairs l_1, l_2 and l_3, l_4 . \square

LEMMA 3.2. *Let l_1, \dots, l_n be a set of n general lines in \mathbf{P}^n for $n = 4$ or 5 . Let $\Lambda \subset \mathbf{G}(n - 2, n)$ be the subset of the Grassmannian parametrizing codimension-2 linear spaces intersecting all the lines. Then:*

- (a) Λ is irreducible of dimension $n - 2$;
- (b) the intersection of all the linear spaces parametrized by points of Λ is empty.

The restriction on n can be removed at the cost of a more complicated proof, but the statement above is sufficient for our applications in later sections. The word “general” in the statement of the lemma means that the proof works for a Zariski-open subset of points in the space of sets of n lines; however, the proof does not produce such an open subset explicitly.

Proof. (a) Let $U_n \subset \mathbf{G}(1, n)^n$ be the open subset parametrizing sets of n distinct lines. Let $I \subset U_n \times \mathbf{G}(n - 2, n)$ be the incidence correspondence consisting of pairs $((L_1, \dots, L_n), L)$, where L is a codimension-2 linear space intersecting all of the L_i . Let $f : I \rightarrow U_n$ be the projection. We want to prove that a general fiber of f is irreducible of dimension $n - 2$.

Each fiber over a point $u \in U_n$ is cut out from $\mathbf{G}(n - 2, n)$ by n hyperplane sections, and hence has dimension at least

$$\dim \mathbf{G}(n - 2, n) - n = n - 2.$$

If we can find a single fiber $f^{-1}(u)$ of dimension $n - 2$, then by semicontinuity, there is an open set $V \subset U_n$ containing u over which all fibers have dimension $n - 2$. Moreover, the open set $f^{-1}(V)$ is local complete intersection, hence Cohen–Macaulay, so the morphism f is flat over V . By [EGA, Theorem 12.2.1(x)], the set of points $u \in V$ over which fibers of f are integral is an open set $W \subset V$. If we can further show that the fiber $f^{-1}(u)$ is integral, then W is nonempty, and so a general fiber of f is integral, in particular irreducible, as required. So it remains to find $u \in U_n$ over which the fiber of f is of dimension $n - 2$ and integral.

We do this using Macaulay 2. For the case $n = 4$, Outputs 1 and 2 from lemma.2.2-P4.m2 show that for a certain point $u_4 \in U_4$, the fiber $f^{-1}(u_4)$ is integral of dimension 2. Similarly, for the case $n = 5$, Outputs 1 and 2 from lemma.2.2-P5.m2 show that for a certain point $u_5 \in U_5$, the fiber $f^{-1}(u_5)$ is integral of dimension 3.

(b) Suppose there is a point $p \in \mathbf{P}^n$ such that every linear space parametrized by Λ passes through p . Let $\Sigma_p \subset \mathbf{G}(n - 2, n)$ be the Schubert cycle parametrizing linear spaces passing through p . In particular, we should have $\Lambda \subset \Sigma_p$. Let us show that this containment is impossible.

First, note that Λ has codimension n in $\mathbf{G}(n - 2, n)$, while for any p , the Schubert variety Σ_p has codimension 2. Considering the Plücker embedding of the Grassmannian $\mathbf{G}(n - 2, n)$ in projective space, we can view Λ as $\mathbf{G}(n - 2, n) \cap H_1 \cap \dots \cap H_n$ for certain

hyperplanes H_i . Therefore, if $\Lambda \subset \Sigma_p$, we must have $\Lambda \subset \Sigma_p \cap H_1 \cap \cdots \cap H_{n-2}$. Let us analyze the two possibilities for this intersection and show that both lead to a contradiction.

The first possibility is that the intersection $\Sigma_p \cap H_1 \cap \cdots \cap H_{n-2}$ is of the maximal codimension n . In this case, Λ must be an irreducible component of $\Sigma_p \cap H_1 \cap \cdots \cap H_{n-2}$. However, the degree of $\Sigma_p \cap H_1 \cap \cdots \cap H_{n-2}$ is the same as the degree of Σ_p , which by projection away from p is the same as the degree of $\mathbf{G}(n-3, n-1)$: for $n=4$, this equals 2, while for $n=5$, it equals 5. On the other hand, since Λ is a linear section of $\mathbf{G}(n-2, n)$, its degree equals the degree of $\mathbf{G}(n-2, n)$: for $n=4$, this equals 5, while for $n=5$, it equals 14. In both cases, the degree of Σ_p is strictly less than the degree of Λ , so the latter cannot be an irreducible component of the former. So this case is impossible.

The second possibility that $\Sigma_p \cap H_1 \cap \cdots \cap H_{n-2}$ is not of the maximal codimension n . We claim that we can move the hyperplanes H_i to new hyperplanes H'_i such that both of the following hold:

- $\Sigma_p \cap H'_1 \cap \cdots \cap H'_{n-2}$ is of codimension n ;
- $H'_1 \cap \cdots \cap H'_{n-2} = H_1 \cap \cdots \cap H_{n-2}$.

(Note that the new hyperplanes H'_i in general no longer correspond to Schubert varieties in the Grassmannian $\mathbf{G}(n-2, n)$, but that does not affect our proof.) Given the claim, we can then write Λ as $\mathbf{G}(n-2, n) \cap H'_1 \cap \cdots \cap H'_{n-2}$, and the argument from the first case applies again to complete the proof.

It remains to prove the claim. Write $Z = H_1 \cap \cdots \cap H_{n-2}$. For $i < n-2$, assume that we have chosen hyperplanes H'_1, \dots, H'_i such that each of them contains Z , and $\Sigma_p \cap H'_1 \cap \cdots \cap H'_i$ has codimension $i+2$. Since $i+2 < n$, we see that $\Sigma_p \cap H'_1 \cap \cdots \cap H'_i$ is not contained in Λ , and since $\Lambda = \mathbf{G}(n-2, n) \cap Z$, this proves that it is not contained in Z either. So we can find another hyperplane H'_{i+1} which contains Z but does not contain $\Sigma_p \cap H'_1 \cap \cdots \cap H'_i$. Hence, the intersection $\Sigma_p \cap H'_1 \cap \cdots \cap H'_i \cap H'_{i+1}$ has codimension $i+3$. Continuing in this way, we end up with hyperplanes H'_1, \dots, H'_{n-2} satisfying the two bulleted conditions above, as required. \square

Now we can prove our first main result about nefness of codimension-2 linear spaces in \mathbf{P}^4 . The idea is to project away from a point and use the information from the previous lemmas about configurations of 4 lines in \mathbf{P}^3 .

THEOREM 3.3. *Let $r \leq 4$. Let L_r^4 be the proper transform on X_r^4 of a codimension-2 linear space in \mathbf{P}^4 that intersects all the blown-up lines properly. Then, L_r^4 is nef.*

Proof. We first observe that if L_r^4 is nef on X_r^4 , then L_{r-1}^4 is nef on X_{r-1}^4 . To see this, note that the pullback of the class $[L_{r-1}^4]$ equals $[L_r^4] + [F]$, where F is a fiber of the blowup. If L_r^4 is nef, then any irreducible surface that has a negative intersection with the pullback of L_{r-1}^4 must have a negative intersection with F and so must be contained in E_r since F is a nef divisor in E_r . But surfaces contained in E_r are contracted by the blowup map, so they have zero intersection with the pullback of $[L_{r-1}^4]$ by the projection formula. So the pullback of $[L_{r-1}^4]$ is nef, and, therefore, L_{r-1}^4 is nef by Proposition 2.1. So it suffices to prove that L_4^4 is nef.

The restriction of L_4^4 to any of the divisors E_i is an effective curve class, hence nef, so if S is an irreducible surface contained inside one of the divisors E_i , then $L_4^4 \cdot \tilde{S} \geq 0$. We can therefore restrict our attention to irreducible surfaces \tilde{S} that are proper transforms of surfaces S in \mathbf{P}^4 . For such a surface, the intersection $\tilde{S} \cap E_i$ is one-dimensional, hence a

union of curves. We can write it in the form $\tilde{S} \cap E_i = C_1 \cup \dots \cup C_k \cup \Gamma_1 \cup \dots \cup \Gamma_j$, where the C_i are curves contained in fibers of the blowdown map $\pi : X_4^4 \rightarrow \mathbf{P}^4$, and the Γ_j intersect each fiber of π in finitely many points. By Lemma 3.2, we can choose a plane L_4^4 that is disjoint from any given finite set of fibers of π , and for such a plane, we get $L_4^4 \cap \tilde{S} \cap E_i = L_4^4 \cap (\bigcup_{k=1}^j \Gamma_j)$, a finite set of points. Therefore, if \tilde{S} is a surface intersecting every plane L_4^4 nonproperly, we see that $S \cap L$ has dimension at least 1 for every plane $L \subset \mathbf{P}^4$ intersecting all four lines.

So suppose that $S \subset \mathbf{P}^4$ is an irreducible surface such that $\dim(S \cap L) \geq 1$ for every plane $L \subset \mathbf{P}^4$ intersecting all four lines. We form the following incidence correspondence:

$$I = \{(L, p) \mid L \in \Lambda, p \in S \cap L\} \xrightarrow{\pi_1} S$$

$$\downarrow \pi_2$$

$$\Lambda$$

Here, $\Lambda \subset \mathbf{G}(2, 4)$ is the subset of the Grassmannian parametrizing planes intersecting all four lines. By Lemma 3.2, Λ is irreducible of dimension 2. Hence, by our assumption on the dimension of the fibers of π_2 , we see that I has dimension at least 3.

Every fiber $\pi_1^{-1}(p)$ is a subset of Λ , which is irreducible of dimension 2, so any fiber of dimension 2 must equal Λ . But if a fiber $\pi_1^{-1}(p)$ equals Λ , then all the planes parametrized by Λ pass through the point p , contradicting Lemma 3.2(b). Hence, no fiber $\pi_1^{-1}(p)$ has dimension 2.

Therefore, every fiber of π_1 has dimension 1, and so π_1 is surjective. That is, for every point $p \in S$, there are infinitely many planes L passing through p and intersecting all four lines. We will show that this is impossible.

By Lemma A.1, we may assume that S is not contained in any of the linear spaces $\text{Span}(L_i, L_j)$. By this assumption, if $p \in S$ is a general point, then when we project away from p , the images of our lines L_1, \dots, L_4 give four skew lines l_1, \dots, l_4 in \mathbf{P}^3 . Under this projection, planes $L \subset \mathbf{P}^4$ passing through p and intersecting all the lines L_i correspond to lines $l \subset \mathbf{P}^3$ intersecting all the lines l_i . So if there are infinitely many planes L passing through p and intersecting all four lines, then there must be infinitely many lines in \mathbf{P}^3 intersecting the four skew lines l_1, \dots, l_4 .

For the four skew lines l_1, \dots, l_4 in \mathbf{P}^3 , there are at most two lines intersecting them all unless the four lines all lie on a quadric $Q \subset \mathbf{P}^3$. So we must have that p is contained in the vertex of a quadric cone $Q' \subset \mathbf{P}^4$, which also contains the lines L_1, \dots, L_4 . Let us examine the possibilities for the rank of Q' :

- rank 1: in this case, all the lines L_i would be contained in a hyperplane, contradicting generality;
- rank 2: in this case, all the lines would be contained in a union of two hyperplanes whose intersection contains p . Each of the two hyperplanes would be spanned by two of the lines L_i , contradicting the assumption that p is not contained in the span of any two of the L_i ;
- rank 3: in this case, the vertex of Q' is a line L , and projecting from L , maps Q' to a smooth conic $Q'' \subset \mathbf{P}^2$. On the other hand, any line in Q' which is disjoint from L would map to a line in \mathbf{P}^2 contained in Q'' , which is impossible. So all lines in Q' , in particular all the L_i , must intersect a fixed line L . Again, by generality, this is impossible.

We conclude that any such quadric Q' must have rank 4; hence, its vertex has dimension 0.

The linear system V of quadrics containing the L_i has dimension 2. In order to complete the proof, we now analyze two possible cases.

If the general member of V is smooth, then the subset of singular quadrics has dimension at most 1. We just proved that, except for the three quadrics of rank 2 which are unions of hyperplanes $\text{Span}(L_i, L_j)$, the vertex of any such quadric has dimension 0. So we get a one-dimensional set of vertices of quadrics outside the subsets $\text{Span}(L_i, L_j)$. This one-dimensional set cannot contain any surface S , so there cannot exist a surface S outside the subspaces $\text{Span}(L_i, L_j)$ such that through each point of S , there pass infinitely many planes touching all the lines L_i .

If the general member of V is singular, then Bertini's theorem still guarantees that the set of singularities of a general member of V is contained in the base locus $\text{Bs}(V)$. Other than the three rank-2 quadrics from the last paragraph, the set of members of V whose singular set is not contained in $\text{Bs}(V)$ is at most one-dimensional, so the set of singular points of such quadrics again gives a one-dimensional set. On the other hand, $\text{Bs}(V)$ is also one-dimensional, as one sees, for example, by intersecting the three rank-2 quadrics, so we get a one-dimensional set of vertices altogether. Again, this set cannot contain a surface S . \square

Next, we prove the corresponding result for codimension-2 linear spaces in \mathbf{P}^5 . The idea of the proof in this case is to project away from a line, rather than a point, and then argue as before.

THEOREM 3.4. *Let $r \leq 5$. Let L_r^5 be the proper transform on X_r^5 of a codimension-2 linear space in \mathbf{P}^5 that intersects all the blown-up lines properly. Then, L_r^5 is nef.*

Proof. As in the previous theorem, it suffices to prove the result when $r = 5$. We suppose for contradiction that there is an irreducible surface $S \subset \mathbf{P}^5$ such that $\dim(S \cap L) \geq 1$ for every codimension-2 linear space L that intersects all five lines. Again, we form the incidence correspondence

$$I = \{(L, p) \mid L \in \Lambda, p \in S \cap L\} \xrightarrow{\pi_1} S$$

$$\downarrow \pi_2$$

$$\Lambda$$

where now $\Lambda \subset \mathbf{G}(3, 5)$ is the subset of the Grassmannian parametrizing linear spaces intersecting all five lines. Arguing exactly as before, we see that all fibers of π_1 must have dimension 2. We will show that the locus of points $p \in \mathbf{P}^5$ through which we have a two-dimensional family of linear spaces from Λ does not contain any irreducible surfaces except for those contained in subspaces $\text{Span}(L_i, L_j)$. As the proper transform of such a subspace is a toric variety, its cone of surfaces is linearly generated, and so L_5^5 has a nonnegative intersection product with the class of any such surface.

So assume $p \in \mathbf{P}^5$ is a point such that the set Λ_p of linear spaces in Λ that pass through p is two-dimensional. By Proposition 2.4, we can assume that the surface S above does not lie in one of the linear spaces $\text{Span}(L_i, L_j)$, so it is enough to consider points p not in any of these linear spaces.

Fix one of the lines, say L_1 . First, we claim that for any point $q \in L_1$, the subset $\Lambda_{pq} \subset \Lambda_p$ consisting of linear spaces through both p and q has dimension 1. If this were not the case, there would be a point $q \in L_1$ such that the family of linear spaces through p and q has dimension 2. Projecting away from the line joining p and q , the lines L_2, L_3, L_4, L_5 would

then map to lines l_2, l_3, l_4, l_5 in \mathbf{P}^3 with a two-dimensional family of lines intersecting all four. By Lemma 3.1, this can only happen in the following cases: first, two of the lines coincide; second, all four lines pass through a common point $p \in \mathbf{P}^3$; third, all four lines lie in a common plane $P \subset \mathbf{P}^3$. The first case only occurs if the center of projection \overline{pq} is contained in $\text{Span}(L_i, L_j)$ for some $i \neq j$, but since q is a point in L_1 , this means that L_1 intersects $\text{Span}(L_i, L_j)$, contradicting generality of the lines. The second case occurs only if there is a two-dimensional linear space in \mathbf{P}^5 (namely, the cone over the point p) intersecting all five lines L_i , and again, this contradicts generality. The third case only occurs if there is a hyperplane in \mathbf{P}^5 (namely, the cone over P) containing all five lines L_i , and again, this contradicts generality.

So we see that for any $q \in L_1$, the set of linear spaces through p and q and intersecting the lines L_2, L_3, L_4, L_5 has dimension 1. We may assume that the line \overline{pq} is not contained in any of the linear subspaces $\text{Span}(L_i, L_j)$, so projecting away from the line \overline{pq} , we obtain a set of four distinct lines in \mathbf{P}^3 such that the family of lines in \mathbf{P}^3 touching all four has dimension 1. Again using Lemma 3.1, this implies that either two of the lines intersect or else they are pairwise skew and lie on a smooth quadric in \mathbf{P}^3 .

Let us first deal with the case when two of the lines intersect. We will think of the projection away from the line \overline{pq} as the projection away from p first, followed by the projection away from the image of q in \mathbf{P}^4 . As explained above, we can assume that p does not lie in any of the linear spaces $\text{Span}(L_i, L_j)$, so first projecting away from p gives five skew lines l_1, \dots, l_5 in \mathbf{P}^4 . We next project away from a point \tilde{q} on l_1 . If l_1 is contained in any of the hyperplanes $\text{Span}(l_i, l_j)$, then in \mathbf{P}^5 , we would have three lines L_1, L_i, L_j contained in a hyperplane, contradicting generality. So l_1 meets each of the hyperplanes $\text{Span}(l_i, l_j)$ in a single point. Choosing \tilde{q} to be different from all of these points, the projection away from q then gives us four pairwise skew lines in \mathbf{P}^3 .

So we may suppose that the four lines are pairwise skew and lie on a smooth quadric surface in \mathbf{P}^3 . By taking the cone over this quadric, we get a quadric in \mathbf{P}^5 of corank 2 that contains L_2, L_3, L_4, L_5 and whose vertex is a line intersecting L_1 and passing through p . Moreover, for each $q \in L_1$, we get such a quadric, so there is a one-dimensional family of lines through p that are vertices of quadrics of this type. We will prove that the set of such points p either has dimension at most 1 or is a plane in \mathbf{P}^5 .

By Lemma 3.5, the family of quadrics in \mathbf{P}^5 of corank 2 that contain the lines L_2, \dots, L_5 and whose vertex intersects L_1 is of dimension 2. Call this two-dimensional family \mathcal{F} and consider the following incidence correspondence:

$$\begin{array}{ccc}
 J = \{(Q, p) \mid Q \in \mathcal{F} \text{ and } p \text{ lies on the vertex line of } Q\} & \xrightarrow{\pi_1} & \mathbf{P}^5 \\
 \downarrow \pi_2 & & \\
 \mathcal{F} & &
 \end{array}$$

All fibers of π_2 are lines, so every irreducible component of J has dimension 3. We may assume that J is irreducible: if not, we apply the same argument to each component of J in turn. We distinguish two possible cases. If π_1 is generically finite, then the points $p \in \mathbf{P}^5$ which lie on a one-dimensional family of vertex lines of members of \mathcal{F} are contained in a proper closed subset Z of $\pi_1(J)$. The preimage $\pi_1^{-1}(Z)$ is a proper closed subset of J , hence has dimension at most 2, and the fibers of π_1 over points of Z are one-dimensional by hypothesis. Hence, Z has dimension at most 1. If π_1 is not generically finite, then $\pi_1(J)$ is irreducible of dimension at most 2. For each point $p \in \pi_1(J)$, there is a one-dimensional

family of vertex lines touching L_1 and passing through p . Such a family sweeps out a plane Π inside \mathbf{P}^5 , and so $\pi_1(J)$ is a plane. \square

LEMMA 3.5. *For any $k \in \{0, \dots, n-1\}$ and any N , the set $\Lambda(k, N, n)$ of quadrics in \mathbf{P}^n of corank k and containing N general lines has the expected codimension*

$$e(k, N, n) := \max \left\{ 3N + \binom{k+1}{2}, \binom{n+2}{2} \right\}.$$

Moreover, for $k \geq 1$, the set $\Lambda_v(k, N, n)$ of those quadrics in $\Lambda(k, N, n)$ whose vertex intersects another general line has the expected codimension

$$\epsilon(k, N, n) := \max \left\{ e(k, N) + n - k - 1, \binom{n+2}{2} \right\}.$$

In particular, with $n = 5$, $N = 4$ and $k = 2$, we see that the locus $\Lambda_v(2, 4, 5)$ of quadrics in \mathbf{P}^5 of corank 2 containing four general lines and with vertex intersecting another general line has dimension

$$\binom{5+2}{2} - 1 - \epsilon(2, 4, 5) = 20 - 3 \cdot 4 - \binom{3}{2} - 3 = 2$$

as claimed in the proof of Theorem 3.4.

Proof. For $1 \leq i \leq N$, let $\Lambda(L_i)$ denote the set of quadrics in \mathbf{P}^n that contain the i th line L_i , and let $\lambda(L_i)$ denote the intersection of $\Lambda(L_i)$ with the set R_k of quadrics of corank k . Then $\lambda(L_i)$ has codimension 3 in R_k . To see this, one can, for example, fix the vertex l and project away $\pi_l: \mathbf{P}^n \dashrightarrow \mathbf{P}^{n-k}$: quadrics with vertex l and containing L_i then correspond to smooth quadrics in \mathbf{P}^{n-k} containing $\pi_l(L_i)$. If L_i is disjoint from l , this clearly gives a set of codimension 3. Varying l among all linear spaces disjoint from L_i , we then get a subset of codimension 3 in R_k . If L_i intersects l , then $\pi_l(L_i)$ is a point, so we get one condition on the smooth quadrics; however, for $n \geq 4$, the condition for l to intersect a fixed line imposes $n-2 \geq 2$ conditions, and so we get codimension at least 3 in this case too.

For any k , the group $PGL(n+1)$ acts transitively on R_k and maps $\lambda(L_i)$ to $\lambda(L'_i)$ for some other line L'_i in \mathbf{P}^5 . For each i , we can apply Kleiman's transversality theorem [KI] to each component of $\lambda(L_i)$ to find a Zariski-open subset of $PGL(n+1)$ that moves the component into proper position relative to $\bigcap_{1 \leq j < i} \lambda(L_j)$. Intersecting these open subsets, we get a nonempty subset of elements moving every component of $\lambda(L_i)$ into proper position relative to $\bigcap_{1 \leq j < i} \lambda(L_j)$, and, therefore, the intersection $\lambda(L'_i) \cap \bigcap_{j < i} \lambda(L'_j)$ has the expected codimension $\binom{k+1}{2} + 3i$. Putting $i = n$, we get the claimed codimension $e(k, N, n)$ of $\Lambda(k, N, n)$.

To prove the claimed codimension $\epsilon(k, N, n)$ of $\Lambda_v(k, N, n)$, for a line L , we write $\lambda_v(L)$ to denote the set of quadrics in R_k whose vertex intersects L . Then $\lambda_v(L)$ has codimension $n-k-1$ in R_k , as one sees again by the projection away from the vertex. Then the same argument as in the previous paragraph applies again to show that the codimension of $\Lambda_v(k, N, n)$ in $\Lambda(k, N, n)$ is $n-k-1$. \square

§4. 2-cycles on X_r^4 for $r \leq 4$

In the next two sections, we will prove our main results about linear generation of cones of cycles. We begin with the case of lines in \mathbf{P}^4 . In this case, $N_2(X_r^4) = N^2(X_r^4)$ has a basis

consisting of the classes

$$H^2, \quad F_i := HE_i, \quad G_i := -E_i^2 \quad (i = 1, \dots, r),$$

where we have chosen signs so that effective classes in the exceptional divisors have positive coefficients with respect to the basis.

The intersections among these classes are given by the following matrix:

	H^2	F_i	G_i
H^2	1	0	0
F_j	0	0	$-\delta_{ij}$
G_j	0	$-\delta_{ij}$	$3\delta_{ij}$

Using Corollary 2.3, we can write down all the classes of linear subvarieties in X_4^4 . The linear cone $\text{Lin}_2(X_4^4)$ is then generated by the following list of classes, in which (as explained in Section 2.5) we list generators up to permutations of indices:

	H^2	F_1	F_2	F_3	F_4	G_1	G_2	G_3	G_4
	0	1	0	0	0	0	0	0	0
	0	1	0	0	0	1	0	0	0
	1	-1	-1	-1	-1	0	0	0	0
	1	-2	-1	-1	0	-1	0	0	0

(The class $F_1 + G_1$ appearing here is the class of a linear subvariety $\mathbf{P}^1 \times \mathbf{P}^1$ inside the exceptional divisor $E_1 \cong \mathbf{P}^1 \times \mathbf{P}^2$.)

Before stating our first main result on linear generation, we record one fact that will save work when verifying that certain classes are nef.

LEMMA 4.1. *Let $\alpha \in N^k(X_r^n)$ be a nef class. Let β be any class of the form $\beta = \alpha + \sum_i [Z_i]$, where $\{Z_i\}$ are subvarieties of X_r^n contained in exceptional divisors. If β is contained in $\text{Lin}_k^*(X_r^n)$, then β is also nef.*

Proof. We must show that for every irreducible subvariety S of dimension k in X_r^n , we have $\beta \cdot [S] \geq 0$.

If $S \subset E_j$ for some j , then since E_j is toric, we have $[S] \in \text{Lin}_k(E_j) \subset \text{Lin}_k(X_r^n)$, and hence by hypothesis, $\beta \cdot [S] \geq 0$.

If S is not contained in any exceptional divisor E_j , then it intersects each E_j either in the empty set or in a set of dimension $k - 1$. If $S \cap E_j$ is nonempty and $Z_i \subset E_j$ is one of the subvarieties appearing in β , then we can compute $[S] \cdot [Z_i]$ as $([S \cap E_j] \cdot [Z_i])_{E_j}$, where the subscript indicates that the intersection is considered in the ambient space E_j . Since $E_j \cong \mathbf{P}^1 \times \mathbf{P}^{n-2}$, the intersection of any two effective cycles is again effective, so $[S] \cdot [Z_i] = ([S \cap E_j] \cdot [Z_i])_{E_j} \geq 0$. Since α is nef, we conclude that $\beta \cdot [S] \geq 0$, as required. □

Here is our first main result about linear generation.

THEOREM 4.2. *The effective cone of 2-cycles $\overline{\text{Eff}}_2(X_r^4)$ is linearly generated if and only if $r \leq 4$.*

Proof. As explained in Section 2.3, to prove linear generation, it is enough to consider the case $r = 4$. Our strategy is to use the list of linear classes above to compute generators

for the dual of the linear cone $\text{Lin}_2(X_4^4)^*$ and verify that the generators are indeed nef classes. According to the output of `theorem_3.2.m2`, the generators of $\text{Lin}_2(X_4^4)^*$ are as follows:

	H^2	F_1	F_2	F_3	F_4	G_1	G_2	G_3	G_4
α	1	0	0	0	0	0	0	0	0
β	1	-1	0	0	0	0	0	0	0
γ	1	-1	-1	0	0	0	0	0	0
δ	1	-1	-1	-1	0	0	0	0	0
ε	1	-1	-1	-1	-1	0	0	0	0
π	1	-2	0	0	0	-1	0	0	0
λ	3	-2	-2	-2	-1	-1	-1	-1	0
μ	4	-3	-3	-2	-2	-1	-1	-1	-1
ν	4	-3	-3	-3	-2	-1	-1	-1	-1
ξ	4	-3	-3	-3	-3	-1	-1	-1	-1

The class ε is represented by the proper transform of a two-dimensional linear subspace in \mathbf{P}^4 intersecting all four lines; hence, it is nef by Theorem 3.3. Lemma 4.1 then implies that the classes α to δ are also nef.

The class π is pulled back from a class π' on the toric variety X_1^4 . It is straightforward to check that π' is in the cone $\text{Lin}_2^*(X_1^4)$, so is nef by Proposition 2.4, and therefore by Proposition 2.1, the class π is nef too.

It remains to deal with the classes λ to ξ . Again, by Lemma 4.1, it is enough to show that λ and ξ are nef.

To show that λ and ξ are nef classes, we will decompose them into effective classes and analyze the summands geometrically. In each table below, the rows sum up to the class in the top-left corner. The symbol π_{ij} denotes the class of the proper transform of a plane containing L_i and intersecting L_j , while γ_k denotes the class of the proper transform of a plane containing L_i .

λ	H^2	F_1	F_2	F_3	F_4	G_1	G_2	G_3	G_4
π_{14}	1	-2	0	0	-1	-1	0	0	0
γ_2	1	0	-2	0	0	0	-1	0	0
γ_3	1	0	0	-2	0	0	0	-1	0
ξ	H^2	F_1	F_2	F_3	F_4	G_1	G_2	G_3	G_4
π_{12}	1	-2	-1	0	0	-1	0	0	0
π_{21}	1	-1	-2	0	0	0	-1	0	0
π_{34}	1	0	0	-2	-1	0	0	-1	0
π_{43}	1	0	0	-1	-2	0	0	0	-1

We have already noted that the classes γ_2 and γ_3 are nef. The classes π_{ij} are not nef, but we will show that any surface intersecting a class π_{ij} from the above tables negatively must nevertheless have nonnegative intersection with λ and ξ .

For convenience, let us consider π_{12} ; other cases are identical. Let $H = \text{Span}(L_1, L_2)$, and let \tilde{H} be the proper transform of H on X_4^4 . By generality, the lines L_3 and L_4 each intersect H in a point, and so $\tilde{H} \cong X_{2,2}^3$. Now, π_{12} is a divisor inside \tilde{H} , and by Lemma A.1, it is nef. Therefore, if $Z \subset X_4^4$ is an irreducible surface that is not contained in \tilde{H} , we have $Z \cdot \pi_{12} > 0$. On the other hand, if Z is contained in \tilde{H} , then we know that Z is linear by

Lemma A.1. Since λ and ξ are both in the dual of the linear cone $\text{Lin}_2(X_4^4)$, they must both have nonnegative intersection with Z .

Finally, to prove that linear generation does not hold for $r \geq 5$, it is enough to consider the case $r = 5$. Choose any linear subspace spanned by two of the lines, say $H = \text{Span}(L_1, L_2)$. The other three lines intersect H in three points p_3, p_4, p_5 . Counting dimensions, there is a quadric surface Q inside H containing the lines L_1 and L_2 and the points p_3, p_4, p_5 . Blowing up, Corollary 2.3 tells us that the class of the proper transform of Q on X_5^4 is

$$[\tilde{Q}] = 2H^2 - 3F_1 - 3F_2 - F_3 - F_4 - F_5 - G_1 - G_2$$

and it is straightforward to check that this is not in the linear cone $\text{Lin}_2(X_5^4)$. □

§5. 2-cycles on X_r^5 for $r \leq 5$

The space $N^2(X_r^5)$ has a basis consisting of the classes

$$H^2, \quad F_i := HE_i \quad (i = 1, \dots, r), \quad G_i := -E_i^2 \quad (i = 1, \dots, r)$$

and the space $N_2(X_r^5)$ has a basis consisting of the classes

$$H^3, \quad f_i := -HE_i^2 \quad (i = 1, \dots, r), \quad g_i := E_i^3 \quad (i = 1, \dots, r)$$

where, again, signs are chosen so that effective cycles in exceptional divisors have positive coefficients in the basis.

The intersections among these are as follows:

	H^2	F_i	G_i
H^3	1	0	0
f_j	0	0	$-\delta_{ij}$
g_j	0	$-\delta_{ij}$	$4\delta_{ij}$

The linear cone $\text{Lin}_2(X_5^5)$ is then generated by the following classes:

	H^3	f_1	f_2	f_3	f_4	f_5	g_1	g_2	g_3	g_4	g_5
	0	1	0	0	0	0	0	0	0	0	0
	0	2	0	0	0	0	1	0	0	0	0
	1	-1	-1	-1	-1	0	0	0	0	0	0
	1	-3	-1	0	0	0	-1	0	0	0	0

We can now prove our second main result.

THEOREM 5.1. *The cone of effective 2-cycles $\overline{\text{Eff}}_2(X_r^5)$ is linearly generated for $r \leq 5$.*

Proof. As before, we compute the classes generating $\text{Lin}_2(X_5^5)^*$. To avoid an extremely long list, let us say that a subset $\{v_1, \dots, v_n\}$ of the full set of generators of $\text{Lin}_2(X_5^5)^*$ is *maximally incident* if every generator can be written in the form $v = v_i + \sum_j a_j F_j + \sum_k b_k (F_k + G_k)$ for some positive integers a_j and b_k . Using Lemma 4.1, it is sufficient to show that all generators in a maximally incident set are nef. A maximally incident set of generators for $\text{Lin}_2(X_5^5)^*$ is as follows:

	H^2	F_1	F_2	F_3	F_4	F_5	G_1	G_2	G_3	G_4	G_5
α	1	-2	0	0	0	0	-1	0	0	0	0
β	1	-1	-1	-1	-1	-1	0	0	0	0	0
γ	2	-2	-2	-1	-1	-1	-1	-1	0	0	0
δ	3	-3	-3	-3	-2	-2	-1	-1	-1	0	0
ε	4	-4	-4	-4	-4	-4	-1	-1	-1	-1	-1

Let us prove that each of these classes is nef:

- α : this class is pulled back from a class $\tilde{\alpha}$ on the toric variety X_1^5 . Since the effective cones of toric varieties are linearly generated, $\tilde{\alpha}$ is nef, and hence so too is α .
- β : this is the class of a codimension-2 linear space touching all five lines. We proved that this class is nef in Theorem 3.4.
- γ : let H denote the proper transform of a four-dimensional linear space containing L_1 and L_2 . We can write the class γ as $q + F_1 + F_2$, where q is the pushforward of a class in $H \cong X_{2,3}^4$. By Lemma 4.1, any subvariety intersecting γ negatively must intersect q negatively, but by Lemma A.3, we can see that q is nef in H , so any such subvariety must be contained in H . However, again by Lemma A.3, the cone of 2-cycles on H is linearly generated, so γ has positive degree on any subvariety contained in H .
- δ : we can prove this is nef by considering the following decomposition into classes of lower degrees:

δ	H^2	F_1	F_2	F_3	F_4	F_5	G_1	G_2	G_3	G_4	G_5
λ	1	-2	0	0	-1	-1	-1	0	0	0	0
q	2	-1	-3	-3	-1	-1	0	-1	-1	0	0

Let H_{23} denote a four-dimensional linear subspace containing the lines L_2 and L_3 . Then, q is the class of the proper transform of a quadric threefold in H_{23} containing L_2 and L_3 and the three points of intersection of the other lines with H_{23} . As for our proof above for γ , the proper transform of H_{23} is the fourfold $X_{2,3}^4$, and the class of a quadric containing all three points and two lines is nef on this space. So any surface class intersecting q negatively must be contained in $X_{2,3}^4$ and hence must be linearly generated.

We must now show the same for λ . This class is represented by a codimension-2 linear space containing the line L_1 and intersecting the lines L_4 and L_5 . Let H_{14} denote a four-dimensional linear space containing the lines L_1 and L_4 : then, λ is represented by any hyperplane inside H_{14} that contains L_1 and the point p_5 of intersection of L_5 with H_{14} . Now let S be an irreducible surface in X_5^5 . If S is contained in some linear space H_{14} , then again by Lemma A.3, S is linearly generated. If not, then for each choice of H_{14} , we have that the intersection $S \cap H_{14}$ is a curve C . If $S \cdot \lambda < 0$, then C must be contained in the base locus of the family of hyperplanes containing L_1 and p_5 , which is exactly the plane P spanned by L_1 and p_5 . As we vary the hyperplane H_{14} , the corresponding curves C will sweep out the whole surface S , and, therefore, S is contained in the union of all the planes P , which is exactly the span of L_1 and L_5 . Again, this shows that S is linearly generated.

- ε : this class can be written as D^2 , where $D = 2H - \sum_{i=1}^5 E_i$ is the class of the proper transform of a quadric containing all the lines. Since the intersection of nef divisors is nef by Lemma 2.2, it is enough to prove that D is nef. We claim that it is sufficient to show that there exists a set of five lines L_1, \dots, L_5 such that on the corresponding blowup, the class D contains exactly six divisors D_1, \dots, D_6 whose common intersection is empty. To prove the claim, write U to denote the subset of $\mathbf{G}(1, 5)^5$ consisting of ordered sets of five distinct lines. There are five tautological sections $U \rightarrow \mathbf{P}^5 \times U$, and blowing up along the image gives a family $\mathcal{X}_5^5 \rightarrow U$ whose fibers are varieties of type X_5^5 . The divisor class D extends to a class \mathcal{D} on X_5^5 . A simple dimension count shows that on any fiber, the line bundle D has at least six sections, so the divisors D_i extend to divisors \mathcal{D}_i on the

family. If we can show that, on a chosen fiber, the divisors D_i have empty intersection, by semicontinuity, the same is true for fibers in some open set $U' \subset U$, and therefore D is basepoint-free, in particular nef, on a general fiber.

To conclude, we choose five lines L_1, \dots, L_5 in \mathbf{P}^5 as specified in `theorem_4.1-b.m2` and use Macaulay 2 to find a basis of the linear system L of quadrics containing all five. Output 2 of `theorem_4.1-b.m2` shows that this linear system has six generators $\Delta_1, \dots, \Delta_6$ such that, as a scheme, the intersection $\bigcap_i \Delta_i$ is exactly the union of the L_i . In particular, the divisors in this linear system have no common normal directions along any of the lines L_i . Blowing up, the strict transforms D_i of the hypersurfaces Δ_i give divisors in the class D with no common point, and, therefore, D is basepoint-free as required. \square

5.1 3-cycles on X_r^5

As a complement to the previous result, we next show that for 3-cycles on blowups of \mathbf{P}^5 , linear generation fails as soon as we blow up four lines. This is in keeping with the results of [CLO] which show that as we blow up more, linear generation fails sooner for cones of higher-dimensional cycles.

For this result, recall that the Segre cubic threefold is a copy of $\mathbf{P}^1 \times \mathbf{P}^2$ embedded in \mathbf{P}^5 by sections of $O(1, 1)$.

PROPOSITION 5.2. *The cone of effective 3-cycles $\overline{\text{Eff}}_3(X_r^5)$ is not linearly generated for $r \geq 4$.*

Proof. It suffices to prove the claim for $r = 4$. For four general lines L_i in \mathbf{P}^5 , there is a Segre cubic S containing the lines as rulings $\mathbf{P}^1 \times \{\text{point}\}$. The normal bundle of L_i in S is easily shown to be $O \oplus O$. Fulton’s blowup formula [Fu, Theorem 6.7] then shows that the proper transform of S on X_4^5 has class

$$[\tilde{S}] = 3H^2 - \sum_{i=1}^4 (4F_i + G_i).$$

It is straightforward to check that $[\tilde{S}]$ is not in the linear cone $\text{Lin}_3(X_4^5)$. \square

§6. Curves and divisors on X_r^n

In this section, we round out the picture for cycles on the varieties X_r^n by considering linear generation of cones of curves and divisors. We write l to denote the pullback of the class of a line in \mathbf{P}^n and l_i for the class of a line in an exceptional divisor which is contracted by blowing down.

PROPOSITION 6.1. *For $r \leq 7$ lines in \mathbf{P}^4 , the cone of curves $\overline{\text{Eff}}_1(X_r^4)$ is linearly generated. For $r \geq 10$ lines in \mathbf{P}^4 , this cone is not linearly generated.*

Proof. For any three lines in \mathbf{P}^4 , there is a line intersecting all three. Therefore, the linear cone $\text{Lin}_1(X_r^4)$ is generated by classes l_i for $i = 1, \dots, r$ and classes $l - l_i - l_j - l_k$ for distinct $1 \leq i, j, k \leq r$. The dual cone $\text{Lin}_1(X_r^4)^*$ is spanned by H , classes $H - E_i$ for $i = 1, \dots, r$ and the class $3H - E_1 - \dots - E_r$.

We claim that the last class is nef for any $r \leq 7$. It suffices to prove this for $r = 7$. Exactly as in the proof of Theorem 5.1, it suffices to prove this class is basepoint-free for any chosen set of seven disjoint lines in \mathbf{P}^4 . The output of `proposition_5.1.m2` shows that, for a set of seven randomly chosen lines, the scheme-theoretic base locus of the linear system of cubics

containing the seven lines is exactly the union of the lines, and so blowing up the class $3H - E_1 - \cdots - E_7$ is basepoint-free as required.

In the other direction, using the intersection numbers in Section 2.1, we compute that the top self-intersection number of the divisor $3H - \sum_{i=1}^r E_i$ on X_r^4 is $81 - 9r$. For any $r \geq 10$, this is negative, so the class is not nef, and, therefore, $\text{Lin}_1(X_r^4)$ does not equal $\overline{\text{Eff}}_1(X_r^5)$. \square

For eight lines in \mathbf{P}^4 , the base locus of the corresponding class $3H - \sum_{i=1}^8 E_i$ has a component that comes from a curve C of degree 19 in \mathbf{P}^4 . Computation shows that C intersects each of the blown-up lines transversely in six points; if C were irreducible, we would be able to conclude that $3H - \sum_{i=1}^8 E_i$ is nef and hence that the cone of curves is again linearly generated in this case. Unfortunately, it seems to be out of reach of computation to decide whether C is irreducible.

PROPOSITION 6.2. *The cone of curves $\overline{\text{Eff}}_1(X_r^5)$ is linearly generated if and only if $r \leq 5$.*

Proof. In this case, the linear cone $\text{Lin}_1(X_r^5)$ is generated by the l_i together with classes $l - l_i - l_j$. The dual cone $\text{Lin}_1(X_r^5)^*$ is then spanned by H , classes $H - E_i$ and the class $2H - E_1 - \cdots - E_r$.

In the proof of Theorem 5.1, we showed that $2H - E_1 - \cdots - E_5$ is a nef divisor class on X_5^5 , and, therefore, $2H - E_1 - \cdots - E_r$ is nef on X_r^5 for any $r \leq 5$.

In the other direction, the top self-intersection number of the divisor $2H - \sum_{i=1}^r E_i$ on X_r^5 is $32 - 6r$. For any $r \geq 6$, this is negative, so $2H - \sum_{i=1}^r E_i$ is not nef. Hence, $\text{Lin}_1(X_r^5)$ does not equal $\overline{\text{Eff}}_1(X_r^5)$. \square

PROPOSITION 6.3. *The cone of divisors $\overline{\text{Eff}}^1(X_r^4)$ is linearly generated if and only if $r \leq 4$.*

Proof. It suffices to prove the linear generation claim for $r = 4$. The linear cone $\text{Lin}^1(X_4^4)$ is spanned by classes E_i and $H - E_i - E_j$, so as in Proposition 6.2, the dual cone $\text{Lin}^1(X_4^4)^*$ is spanned by curve classes l , $l - l_i$ and $2l - \sum_{i=1}^4 l_i$. Curves in the classes l and $l - l_i$ evidently sweep out dense open subsets of \mathbf{P}^4 , and consequently, they are nef. For the last class, we argue as follows. For any point $p \in \mathbf{P}^4$, Schubert calculus shows that there is a plane Π touching our four blown-up lines L_i and passing through p . There is a conic in Π passing through the points $L_i \cap \Pi$ and p . The proper transform of this conic on X_4^4 then has class $2l - \sum_{i=1}^4 l_i$. Since these conics sweep out a dense open subset of X_4^4 , the class is nef as required.

Now we will prove that the cone is not linearly generated for $r = 5$; again, this implies the claim for $r \geq 5$. In this case, the dual of the cone of linear divisors has an extremal ray spanned by the effective class $\gamma = 2l - \sum_{i=1}^5 l_i$. We claim that γ is not nef. To see this, it suffices to find a big divisor D on X_5^4 with $D \cdot \gamma = 0$; applying Kodaira's lemma, we can write $D \equiv A + E$ with A ample and E effective, so we must have $E \cdot \gamma < 0$. Choose $D = -K = 5H - 2 \sum_{i=1}^5 E_i$; we claim that D is big. A straightforward dimension count shows that the dimension of the space of sections of mD is bounded below by a polynomial with leading term km^4 , where

$$k = \frac{5^4}{4!} - 5 \cdot \frac{14}{3} > 0,$$

and, therefore, D is big as required. \square

PROPOSITION 6.4. *The cone of divisors $\overline{\text{Eff}}^1(X_r^5)$ is linearly generated if and only if $r \leq 3$.*

Proof. For $r \leq 3$, the variety $\overline{\text{Eff}}^1(X_r^5)$ is toric, so the claim follows from Proposition 2.4.

For the converse, as above, it suffices to prove the claim when $r = 4$. The divisor class $3H - 2E_1 - 2E_2 - 2E_3 - E_4$ is not in the linear cone. This class is represented by the proper transform of a cubic fourfold double along L_1, L_2, L_3 and containing L_4 . A straightforward dimension count shows that such fourfolds exist for any 4-tuple of lines in \mathbf{P}^5 , and, therefore, $\overline{\text{Eff}}^1(X_r^5)$ is not linearly generated. \square

Acknowledgments. Thanks to Izzet Coskun and Elisa Postinghel for helpful conversations.

Supplementary materials

Supplementary materials are available at <https://doi.org/10.1017/nmj.2019.41>.

Appendix A. 2-cycles on $X_{2,2}^3$ and $X_{3,2}^4$

In this section, we prove linear generation for the cones of effective 2-cycles on the spaces $X_{2,2}^3$ and $X_{3,2}^4$. These linear generation results were used in the proofs of Theorems 4.2 and 5.1.

LEMMA A.1. *The cone of effective 2-cycles $\overline{\text{Eff}}_2(X_{2,2}^3)$ is linearly generated.*

Proof. Writing down all linear classes on $X_{2,2}^3$ and computing the dual, the output of lemma_6.1.m2 shows that $\text{Lin}_2(X_{2,2}^3)^*$ is spanned by the classes

	H^2	HE_1	HE_2	$-E_3^2$	$-E_4^2$
α	1	0	0	0	0
β	1	-1	0	0	0
γ	1	-1	-1	0	0
δ	1	0	0	-1	0
ε	2	-1	0	-1	-1
κ	2	-1	-1	-1	-1

In each case, irreducible curves representing the class cover a dense open set in $X_{2,2}^3$. For example, the class κ is represented by proper transforms of conics touching L_1 and L_2 and passing through p_3 and p_4 . Choosing a general point $p \in \mathbf{P}^3$, there is a plane Π containing p, p_3 and p_4 ; this plane intersects L_1 and L_2 in points q_1 and q_2 , and there is an irreducible conic in Π through the five points q_1, q_2, p_3, p_4 and p . \square

LEMMA A.2. *The cone of effective 2-cycles $\overline{\text{Eff}}(X_{3,1}^3)$ is linearly generated.*

Proof. Similar to the previous lemma, the output of lemma_6.2.m2 shows that the dual $\text{Lin}_2(X_{3,1}^3)^*$ of the linear cone of 2-cycles is spanned by the classes

	H^2	HE_1	$-E_2^2$	$-E_3^2$	$-E_4^2$
α	1	0	0	0	0
β	1	-1	0	0	0
γ	1	0	-1	0	0
δ	2	-1	-1	-1	0
ε	3	-2	-1	-1	-1

Curves representing the first three classes evidently cover X , and hence are nef. For the class δ , picking any point p on L_1 , the plane spanned by p_2 , p_3 and p is covered by irreducible conics with class δ ; varying p along L_1 , these conics cover X , and so ε is nef. Finally, we can write ε as $\delta + (H^2 - HE_1 + E_4^2)$; since δ is nef, any divisor which is negative on ε must be negative on $H^2 - HE_1 + E_4^2$, which is the class of a line passing through p_4 and intersecting L_1 . If π is the plane spanned by p_4 and L_1 , these lines sweep out π , and therefore $H^2 - HE_1 + E_4^2$ and hence ε can be negative only on the proper transform of π . Note, however, that π is a linear class, and ε is in the dual of the linear cone; so ε is in fact nef. \square

LEMMA A.3. *The cone of effective 2-cycles $\overline{\text{Eff}}(X_{2,3}^4)$ is linearly generated.*

Proof. The strategy of proof is again very similar to the previous cases. The output of lemma_6.3.m2 shows that the dual $\text{Lin}_2(X_{2,3}^4)^*$ of the linear cone of 2-cycles is spanned by the classes

	H^2	HE_1	HE_2	$-E_1^2$	$-E_2^2$	$-E_3^2$	$-E_4^2$	$-E_5^2$
α_1	1	0	0	0	0	0	0	0
α_2	1	-1	0	0	0	0	0	0
α_3	1	-1	-1	0	0	0	0	0
α_4	1	0	0	-1	0	0	0	0
α_5	1	-2	0	-1	0	0	0	0
α_6	2	-2	-1	-1	-1	0	0	0
α_7	2	-1	-1	0	0	-1	-1	0
α_8	3	-3	-2	-1	0	-1	-1	-1
α_9	3	-3	-1	-3	-1	-1	0	0
α_{10}	4	-4	-4	-1	-1	-1	-1	-1

The first six classes are pulled back from classes on toric varieties that are easily checked to be nef. Similarly, α_7 is pulled back from a nef class on $X_{2,2}^4$. The last class α_{10} can be written as D^2 , where D is the divisor class $2H - \sum_i E_i$. This is the pullback of the class $\tilde{D} = 2H - \sum_i E_i$ on X_5^5 , which was shown to be nef in Theorem 5.1, so D is a nef divisor, and hence by Lemma 2.2, we know that $\alpha_{10} = D^2$ is nef too.

It remains to treat α_8 and α_9 , which we do by decomposition. We start with α_8 , which can be decomposed as follows:

α_8	H^2	HE_1	HE_2	$-E_1^2$	$-E_2^2$	$-E_3^2$	$-E_4^2$	$-E_5^2$
β_1	1	0	-1	0	0	0	0	-1
β_2	2	-3	-1	-1	0	-1	-1	0

The class β_1 is pulled back from a nef class on the toric variety $X_{1,1}^4$ and so is nef. The class β_2 is represented by the proper transform of a quadric containing L_1 , intersecting L_2 and passing through p_3 and p_4 . Let Π be a three-dimensional linear space containing L_1 and passing through p_3 and p_4 ; then Π intersects L_2 in a point, call it p_2 . Let π be the plane spanned by L_1 and p_2 and let π' be any plane containing p_3 and p_4 : then $Q = \pi \cup \pi'$ is a quadric with class β . Swapping the roles of p_2 and p_3 , say, we see that the base locus of the linear system $|Q|$ consists of a union of lines in Π . Writing down the classes of linear 1-cycles on $X_{3,1}^3$, we check that Q has a positive degree on any such class. So Q is nef inside the proper transform of Π . It follows that any 2-cycle which intersects α_8 and hence β_2 negatively must be contained in the proper transform of Π . On the other hand,

by Lemma A.2, we know that 2-cycles in $X_{3,1}^3$ are linearly generated, and α_8 is in the dual of the linear cone. Hence, α_8 is nef, as required.

For α_9 , we consider the following decomposition:

α_9	H^2	HE_1	HE_2	$-E_1^2$	$-E_2^2$	$-E_3^2$	$-E_4^2$	$-E_5^2$
γ_1	1	0	0	0	0	-1	0	0
γ_2	1	-2	-1	-1	0	0	0	0
γ_3	1	-1	-2	0	-1	0	0	0

Again, the first class is pulled back from a nef class on a toric variety, and hence is nef. For γ_2 (and similarly for γ_3), we argue as follows: γ_2 is represented by the proper transform of a plane π containing L_1 and intersecting L_2 in a point. Let Π be the three-dimensional space spanned by L_1 and L_2 . The proper transform $\tilde{\Pi}$ of Π is a toric variety X_2^3 . One checks that the proper transform of π is a nef divisor in X_2^3 , and hence any surface intersecting γ_2 negatively must lie in $\tilde{\Pi}$. On the other hand, as X_2^3 is toric, its cone of effective divisors is linearly generated, and so γ_2 has nonnegative degree on surfaces contained in X_2^3 . \square

REFERENCES

- [CLO] I. Coskun, J. Lesieutre and J. C. Ottem, *Effective cones of cycles on blowups of projective spaces*, Algebra Number Theory **10**(9) (2016), 1983–2014.
- [DELV] O. Debarre, L. Ein, R. Lazarsfeld and C. Voisin, *Pseudoeffective and nef classes on abelian varieties*, Compos. Math. **147**(6) (2011), 1793–1818.
- [DPU] O. Dumitrescu, E. Postingshel and S. Urbinati, *Cones of effective divisors on the blown-up \mathbf{P}^3 in general lines*, Rend. Circ. Mat. Palermo (2) **66**(2) (2017), 205–216.
- [EH] D. Eisenbud and J. Harris, *3264 And All That*, Cambridge University Press, Cambridge, 2016.
- [FL1] M. Fulger and B. Lehmann, *Zariski decompositions of numerical cycle classes*, J. Algebraic Geom. **26**(1) (2017), 43–106.
- [FL2] M. Fulger and B. Lehmann, *Positive cones of dual cycle classes*, Algebraic Geom. **4**(1) (2017), 1–28.
- [Fu] W. Fulton, *Intersection Theory*, 2nd ed. Springer, 1998.
- [EGA] A. Grothendieck, *Éléments de géométrie algébrique. IV. Étude locale des schémas et de morphismes des schémas. III*, Publ. Math. Inst. Hautes Études Sci. **28** (1966), Available at http://www.numdam.org/item/PMIHES_1966__28__5_0.
- [Kl] S. Kleiman, *The transversality of a general translate*, Compos. Math. **28**(3) (1974), 287–297.
- [Li] Q. Li, *Pseudo-effective and nef cones on spherical varieties*, Math. Z. **280**(3–4) (2015), 945–979.
- [M2] Macaulay2 computations. Ancillary files available at <https://arxiv.org/abs/1812.08476>.
- [Nor] W. Bruns, B. Ichim, T. Römer, R. Sieg and C. Söger, *Normaliz. Algorithms for rational cones and affine monoids*. Available at <https://www.normaliz.uni-osnabrueck.de>.
- [Ott] J. C. Ottem, *Ample subvarieties and q -ample divisors*, Adv. Math. **229**(5) 2868–2887.

Norbert Pintye

*Department of Mathematical Sciences, Loughborough University, LE11 3TU,
United Kingdom*

`norbert.pintye@gmail.com`

Artie Prendergast-Smith

*Department of Mathematical Sciences, Loughborough University, LE11 3TU,
United Kingdom*

`a.prendergast-smith@lboro.ac.uk`