

Video modeling for the development of personal hygiene skills in youth with autism spectrum disorder

S. Piccin¹†, A. Crippa², M. Nobile², A. Y. Hardan³ and P. Brambilla^{4,5*}

¹ IRCCS “Eugenio Medea” Scientific Institute, San Vito al Tagliamento, Pordenone, Italy

² IRCCS “Eugenio Medea” Scientific Institute, Bosisio Parini, Lecco, Italy

³ Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA

⁴ Department of Neurosciences and Mental Health, Fondazione IRCCS Ca’ Granda Ospedale Maggiore Policlinico, University of Milan, Milan, Italy

⁵ Department of Psychiatry and Behavioural Neurosciences, University of Texas at Houston, Houston, Texas.

Received 2 September 2017; Accepted 24 September 2017; First published online 10 November 2017

Key words: Behaviour, daily living activities, video modelling point of view, video self-modelling.

Autism spectrum disorder (ASD) is a neurodevelopmental disorder mainly characterised by deficits in social communication as well as by narrow patterns of behaviour and interests (American Psychiatric Association, 2013), often accompanied by language, intellectual and sensory impairments. The severity of these impairments may lead to deficits in the development of daily living activities such as simple meal preparation and feeding, community skills (e.g. buying groceries), personal care (e.g. dressing) and personal hygiene skills (bathing, toileting, hand washing, teeth brushing) needed for independence. Among others, the lack of independence in personal hygiene skills increases the burden of the caregiver and makes children with ASD more dependent (Flynn & Healy, 2012). Therefore, it is important to develop tools for helping individuals with ASD in increasing their ability to perform these basic life activities which will lead to savings that can be invested in other critical areas of needs.

Video Modeling (VM) is a successful strategy for developing various skills in individuals with ASD, such as social, communication and daily living skills (Rosenberg *et al.* 2010). The VM consists in showing a video in which a model performs a target behaviour or a specific task, then the participant has to perform the skills presented in the video (Shukla-Mehta *et al.* 2010). Any person familiar or unknown to the participant can act as a model, adult or peer. Two main

versions of the VM have been developed: video Self Modeling (VSM), where the video shows the participant itself as a model, and Video Modeling Point of View (VPOV), where the target skill is filmed from the participant’s perspective, at him/her eye level.

The advantages of these strategies include: (i) the use of visual information, which are better processed by individuals with ASD (Cottini, 2016); (ii) media support which attract their attention; (iii) procedures that eliminate visual and auditory interfering stimuli; (iv) no face-to-face interaction that may be stressful for individuals with ASD (Cottini, 2016). Furthermore, in ASD rehabilitation research, the high variability among the autism spectrum favours the use of single-subject designs, which have been preferred in many studies on VM interventions (Cottini, 2016).

Although VM is a widely used intervention in the field of ASD (Shukla-Mehta *et al.* 2010), a small body of research has been reported regarding the development of personal hygiene skills in children and adolescents with ASD. An overview of these studies can be found in Table 1. The studies were selected by a systematic search in PubMed and Google Scholar for peer-reviewed articles in the English language; search words included video modeling, video self modeling, autism spectrum disorder, hand washing, teeth brushing, toileting, hygiene skills, daily living skills, functional skills and self-help skills. This body of work focused on toileting, hand washing, teeth brushing and took place mostly in educational settings.

Investigators have examined the benefit of VM in targeting toileting skills. Keen *et al.* (2007) examined the effectiveness of a VM in combination with operant conditioning strategies in comparison to the latter alone in teaching urinary control to five children with ASD. Results showed gains for the children who received VM. More recent investigations used a

† Scientific Institute IRCCS ‘E. Medea’, Polo FVG, San Vito al Tagliamento, Pordenone, Italy.

* Address for correspondence: P. Brambilla, Department of Neurosciences and Mental Health, Fondazione IRCCS Ca’ Granda Ospedale Maggiore Policlinico, University of Milan, Milan, via F. Sforza, 35, 20122 Milan, Italy.

(Email: paolo.brambilla1@unimi.it)

Table 1. Studies using VM for the development of personal hygiene skills in children and adolescent with ASD

Author(s)	Targeted skill(s)	Participants	Setting	Study design	Inter Observer Agreement (I.O.A.)	Intervention Dependent variable	Type Model Custom-made/ available video	Outcome
Charlop-Christy, Le & Freeman (2000)	Brushing teeth (among others social skills: conversational speech, expressive labelling of emotions, independent play, spontaneous greetings, oral comprehension, cooperative play, social play)	One boy aged 7 (among other four children)	School	Multiple baseline design within child	I.O.A. = 90–100%	Compare VM (brushing teeth) and <i>in vivo</i> modeling (washing face) giving prompts exclusively for on-task behaviour. Dependent variable: correct performance with the target behaviour.	VM Adult (familiar) Custom-made video	With VM, the participant acquired the skill (brushing teeth) faster and generalised it.
Rayner (2010)	Brushing teeth (among other skill, i.e. unpacking bag)	One boy aged 12	School	Non-concurrent multiple baseline design	I.O.A. = 97%	VM with verbal and signal prompt provided when the child did not start independently. Prompting procedure and live modelling were then used only for teeth brushing. Dependent variable: number of steps correctly performed.	VM Symbols embedded in the video Adult (unfamiliar) Custom-made video	Limited gains with VM for brushing teeth (enhanced from baseline levels to a mean of 55% during the intervention); however, VM led to faster enhancements in the other skill. Generalisation was not assessed for brushing teeth.
Rosenberget al. (2010)	Hand washing	Three children aged 3–5	School	Concurrent multiple baseline across participants design	I.O.A. = 89%–100%.	Commercial VM and then customised VM were used. No prompts occurred. Dependent variable: number of correctly completed steps of the task.	VM Peer (familiar for custom video, unfamiliar for commercial video) Available video Custom-made video	One participant learnt 80% of the steps but two participants did not acquire from the commercial model. Then they received a customised VM, that lead some acquisition for the two participants who did not acquire from the commercial model. Gains were maintained for two participants during follow-up and generalised to a second setting.

Campbell <i>et al.</i> (2015)	Hand washing	Three students aged 17–19*	School	Multiple baseline across participants design	I.O.A. = 100%	VM on a portable handheld device. Basic prompts were used if it was necessary and scored as 0. Dependent variable: number of steps of the task performed independently.	VM n.a n.a	Substantial improvement skill level for two participants and moderate improvement for one participant.
Popple <i>et al.</i> (2016)	Brushing teeth	18 children aged 5–14**	n.a.	Randomized control trial	Not necessary	Randomised control trial – the VM intervention was delivered to the participants via the internet. Participants were assigned to an intervention or control video condition. Dependent variable: a plaque index was used to determine oral hygiene.	VM With narration and closed captioning Peer (unfamiliar) Available video	Oral hygiene enhancements in both groups, with greater improvement within the intervention condition.
Keen <i>et al.</i> (2007)	Toilet use	Five children aged 4–6	Home and educational setting	Multiple baseline design between and across groups	I.O.A. = n.a.	VM and operant conditioning strategies compared with operant conditioning strategies only. Minimum reinforcements were used. Dependent variable: frequencies of in-toilet urinations.	VM Animated video embedded colour, sound and music, accompanied by picture card Available video	Improvement in children who received VM. Three children showed maintenance during follow-up. Two children showed generalisation to a new setting
Drysdale <i>et al.</i> (2015)	Toilet use	Two boys aged 4–5***	Home	Multiple baseline across behaviour design	I.O.A. = 98%; 100%.	VM using both a ‘chaining procedure’ and prompts. Dependent variable: number of prompts required to complete the steps.	VSM/VPOV The video contained both VSM and VPOV elements Real model (self) and animated models Custom-made video	After the intervention, the children required less prompts to perform the toileting steps independently. Children showed maintenance and generalisation to another setting.

Continued

Table 1. *Continued*

Author(s)	Targeted skill(s)	Participants	Setting	Study design	Inter Observer Agreement (I.O.A.)	Intervention Dependent variable	Type Model Custom-made/ available video	Outcome
Lee <i>et al.</i> (2014)	Toilet train	One boy aged 4***	Home	Changing criterion design	I.O.A. = 100%.	VM intervention with reinforcement and picture prompts. Dependent variable: number of unprompted completion of the steps of the task.	VSM/VPOV The video contained both VSM and VPOV elements Custom-made video	Increase in most of the skills of the entire task, which generalised to a second setting. The child did not learn in-toilet voiding.
McLay <i>et al.</i> (2015)	Toileting skills	Two boys aged 7–8***	Home	Non-concurrent multiple baseline design across participants	I.O.A. = 99%; 93.5%	VM intervention that uses animation for in-toilet voiding associated with reinforcement procedures and prompts. Dependent variable: number of steps independently performed; frequency of urination and defecation.	VSM/VPOV The video contained both VSM and VPOV elements Urination and defecation were animated models Custom-made video	Children increased the percentage of steps independently performed in the toileting chain and successfully did in-toilet voiding. Participants showed maintenance and generalisation to a second setting for the steps achieved
Bainbridge & Myles (1999)	Initiate toilet use	One boy aged 3	Home	ABA design	I.O.A. = 100%	The child was exposed to a commercially VM with immediate verbal prompt Dependent variable: frequency in the toilet use initiation and the number of wet and dry diapers.	VM Peer (unfamiliar) Available video	Enhancement in the beginning of the toilet use and dry diapers.

VM, video Modelling; VSM, video self-modelling; VPOV, video modelling point of view; n.a., not available; I.O.A., Inter Observer Agreement (two independent raters scored the study and then an I.O.A. was obtained based on the following formula: $100 \times \text{agreement} / (\text{agreement} + \text{disagreement})$).

Note: the mental age of the participants was between 2 and 7 years old, *one participant with borderline to normal cognitive range, **unknown mental age, ***unknown mental age, presumably lower mental age.

custom-made video with VSM plus VPOV elements associated with reinforcements and prompts (Lee *et al.* 2014; Drysdale *et al.* 2015). One video incorporated animation model for urination and defecations. These studies obtained positive outcomes using VM to toilet train participants with ASD. However, the participants did not independently acquire the behaviours of the entire toileting chain (e.g. undressing, sitting on the toilet, in-toilet voiding, redressing and flushing) (Lee *et al.* 2014). McLay *et al.* (2015) reported on two participants who acquired the entire chain as well as in-toilet urination. In this study, the authors used a custom-made video with VSM plus VPOV elements associated with reinforcements and prompting procedure and the video included animation model for urination and defecations. Investigators examining the development of toilet skills in individuals with ASD largely used reinforcements and prompts procedures as well as elements in the videos. Interestingly, studies that used commercial available videos showed poorer acquisitions in contrast to investigations that used custom-made videos (McLay *et al.* 2015). With respect to hand washing, Rosenberg *et al.* (2010) explored the benefits from watching a commercially VM with three children with ASD and reported that one child learned 80% of the steps but the other two children did not profit. The three children then received a customised VM with a familiar peer as model, which led to some skill achievements for the two participants who did not learn from the commercially available model. Two children who acquired from commercial VM and one child who only achieved some skills with the customised VM showed maintenance and generalisation to a second setting. Campbell *et al.* (2015) implemented a VM on a handheld device to teach hand washing to three pre-adolescents with ASD. Two participants exhibited substantial improvement and one participant showed moderate improvement (up to 84.3% of task steps performed independently).

Three studies investigated the use of VM in oral hygiene in children with ASD. Charlop-Christy *et al.* (2000) used a VM to teach a child to brush his teeth and compared it to an *in vivo* modeling for teaching the same individual to wash his face (prompts were provided exclusively for on-task behaviour). The model in the video was a familiar adult. In this case, VM allowed faster achievement and generalisation of teeth brushing. However, limited gains with VM for brushing teeth were found in the report by Rayner (2010). Although video prompting and live modeling were introduced only for teeth brushing, the child showed limited gains in this skill while exhibiting rapid increases in the steps performed in the other skill trained (i.e. unpacking bag). Difficulties with motor skills related to the target behaviour could

explain this result. Furthermore, the oral reinforcements that the child received during the intervention may have competed with some of the actions required for the tooth brushing task. Finally, there is only one study (Popple *et al.* 2016) that randomly assigned children with ASD to a control group or an experimental condition. In this study, 18 participants were assigned to an intervention condition and received a 3-week VM via the internet compared to a control video condition. Oral hygiene enhancements were found in both groups, with greater improvement within the intervention condition.

The studies reviewed here have found positive results with VM or its variations in enhancing personal hygiene skills, with at least some benefits in selected domains. No studies found negative effects with VM. Interestingly, VM interventions often promoted generalisation and maintenance of the behaviours achieved in participants with ASD. Custom-made interventions and interventions using video with peer or familiar as model appear to be more effective. Furthermore, VM interventions are often effective in teaching some steps but not the entire chain of the task, especially in the toilet use interventions. Therefore, VM focusing on the task steps the participants performed poorly should be interesting to realise. Our review contributes to the small body of literature indicating the effectiveness of VM in helping children and adolescents with ASD to develop personal hygiene skills. In this mini-review, only one study performed a randomized control trial (RCT), while the others used single-subject designs. This is because rehabilitation programmes often involve one or few children. Moreover, the high variability within the autism spectrum, where different IQ levels can be present with jeopardised level of social and motor deficits, makes it difficult to recruit comparable ASD subjects in terms of the level of impairment. Nonetheless, it is crucial that future research includes large RCT on homogeneous ASD individuals with professionally developed videos on personal hygiene skills. The videos should be implemented on handheld technology to allow wide-broad dissemination to promote autonomy for children and adolescents with ASD.

Financial Support

Professor Brambilla (RF-2011-02352308) and Dr Crippa and Dr Piccin (GR-2011-02348929) were partly supported by grants from the Italian Ministry of Health.

Conflict of Interest

None.

References

- American Psychiatric Association** (2013). *Diagnostic and Statistical Manual of Mental Disorder*. DSM 5. APA Publishing: Washington.
- Bainbridge N, Myles BS** (1999). The use of priming to introduce toilet training to a child with autism. *Focus on Autism & Other Developmental Disabilities* **14**, 106–109.
- Campbell JE, Morgan M, Barnett V, Spreat S** (2015). Handheld devices and video modeling to enhance the learning of self-help skills in adolescents with autism spectrum disorder. *OTJR: Occupation, Participation and Health* **32**, 95–100.
- Charlop-Christy MH, Le L, Freeman KA** (2000). A comparison of video modeling with *in vivo* modeling for teaching children with autism. *Journal of Autism and Developmental Disorders*, **30**, 537–552.
- Cottini L** (2016). *Fare Ricerca con Singoli Soggetti. Principi Metodologici E Applicazioni in Educazione Speciale E in Psicologia Clinica*. IRFID: Ottaviano, Neaple.
- Drysdale B, Lee CYQ, Anderson A, Moore DW** (2015). Using video modeling incorporation animation to teach toileting to two children with autism spectrum disorder. *Journal of Developmental and Physical Disabilities* **27**, 149–165.
- Flynn L, Healy O** (2012). A review of treatments for deficits in social skills and self-help skills in autism spectrum disorder. *Research in Autism Spectrum Disorders* **6**, 431–441.
- Keen D, Brannigan KL, Cuskelly M** (2007). Toilet training for children with autism: the effects of video modeling. *Journal of Developmental and Physical Disabilities* **19**, 291–303.
- Lee CYQ, Anderson A, Moore DW** (2014). Using video modeling to toilet train a child with autism. *Journal of Developmental and Physical Disabilities* **26**, 123–134.
- McLay L, Carnett A, van der Meer L, Lang R** (2015). Using a video modeling-based intervention package to toilet train Two children with Autism. *Journal of Developmental and Physical Disabilities* **27**, 431–451.
- Popple B, Wall C, Flink L, Powell K, Discepolo K, Douglas K, Mademtz M, Volkmar F, Shic F** (2016). Brief report: remotely delivered video modeling for improving oral hygiene in children with ASD: a pilot study. *Journal of Autism and Developmental Disorders* **46**, 2791–2796.
- Rayner CS** (2010). Video-modelling to improve task completion in a child with autism. *Developmental Neurorehabilitation* **13**, 225–230.
- Rosenberg NE, Schwartz IS, Davis CA** (2010). Evaluating the utility of commercial videotapes for teaching hand washing to children with autism. *Education and Treatment of Children* **33**, 443–455.
- Shukla-Mehta S, Miller T, Callahan KJ** (2010). Evaluating the effectiveness of video instruction on social and communication skills training for children with autism spectrum disorders: a review of the literature. *Focus on Autism and Other Developmental Disabilities* **25**, 23–36.