

Research Article

Cite this article: Marsden SJ, Junaid AR, Kaprawi F, Muladi F, Aprianto GC, van Balen S(B), Saryanthi R, Collar NJ, Devenish C (2023). Distribution and abundance of threatened and heavily traded birds in the mountains of western Java. *Bird Conservation International*, **33**, e63, 1–9
<https://doi.org/10.1017/S095927092300014X>

Received: 26 October 2022
Revised: 22 January 2023
Accepted: 13 February 2023





Keywords:

Abundance; Asian Songbird Crisis; baseline; conservation; endemic; trapping

Corresponding author:

Stuart J. Marsden;
Email: s.marsden@mmu.ac.uk

Distribution and abundance of threatened and heavily traded birds in the mountains of western Java

Stuart J. Marsden¹ , Achmad Ridha Junaid², Fajar Kaprawi², Faris Muladi², Ganjar Cahyo Aprianto², S. (Bas) van Balen³ , Ria Saryanthi², Nigel J. Collar⁴  and Christian Devenish^{1,5} 

¹Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK; ²Burung Indonesia, Bogor, Indonesia; ³Basilornis Consults, Muntendampad 15, 6835 BE Arnhem, The Netherlands; ⁴BirdLife International, David Attenborough Building, Cambridge, UK and ⁵NatureMetrics, Surrey Research Park, Guildford, UK

Summary

There is serious concern for the future of a wide range of birds in Java and elsewhere in Indonesia due to both loss of habitat and trapping for the cagebird trade (the so-called “Asian Songbird Crisis”). Despite this concern, few data on presence and abundance of key species exist. We provide such data on 184 bird species from over two years of biodiversity surveys from 37 sites on 12 mountains in West and Central Java. Many of these species are heavily traded, endemic, and globally threatened. Several of the threatened endemics, notably Javan Trogon and Javan Cochoa, were often recorded, in terms of both geographical spread and numerical abundance. Rufous-fronted Laughingthrush, Spotted Crocias, and Orange-spotted Bulbul, believed to be threatened by trapping for the songbird trade, appear to remain fairly widespread. By contrast, Brown-cheeked Bulbul, Chestnut-backed (Javan) Scimitar-babbler, Javan Oriole, and especially Javan Blue-flycatcher, recorded on just a single occasion, and Javan Green Magpie, which we failed to record with certainty, now appear to be extremely rare. Our encounter rates, while not pinned to specific mountains for security reasons, represent an important baseline against which future changes in abundance can be gauged.

Introduction

For the world’s most populous island, with around 149 million inhabitants, and despite a long occupation by Europeans with a strong tradition of natural history, Java in Indonesia is remarkably poorly known ornithologically. Unlike its larger island neighbours Sumatra and Borneo, it has no modern checklist of birds, and the only recent field guide (Eaton *et al.* 2021) and bird atlas (Winnans *et al.* 2020) also serve much of the rest of Indonesia. However, the biodiversity of Java is of considerable importance: although the island forms part of the Greater Sunda biogeographical region (“Sundaland”) and shares many species with the Thai-Malay Peninsula, Sumatra, and Borneo, it is also a centre of endemism in its own right (Stattersfield *et al.* 1998); indeed, greater taxonomic scrutiny in the twenty-first century has shown this endemism to be far more pronounced than was previously apparent (del Hoyo and Collar 2014, 2016, Eaton *et al.* 2021). Much of Java is montane, concentrated in the tropical forests flanking the island’s many volcanoes; owing to the declining west–east rainfall gradient across the island however the highest endemism and overall biodiversity are concentrated in the west (Whitten *et al.* 1997).

Researchers wishing to study – and birdwatchers wishing simply to see – the birds endemic to montane western Java almost invariably visit the twin peaks of Mounts Gede and Pangrango which, being only 25 km south-east of Bogor, form easily the most accessible and much the best-known site (Andrew 1985). A consequence of this is that knowledge of the avifaunas of other forested volcanoes in the region has remained rudimentary. For example, the Rufous-fronted Laughingthrush *Garrulax rufifrons* is known from 15 volcanoes but, as documented in Collar and van Balen (2013), only Gede–Pangrango held records from the present century, while half of the other 14 involved records made in or before 1930. Similarly, the Javan Green Magpie *Cissa thalassina* has only been recorded in the twenty-first century at four of its 18 known sites (van Balen *et al.* 2013). The absence of recent information on these two species at so many sites, and indeed on the extent and condition of their habitat there, has rendered it problematic to assess their IUCN Red List status or to identify the most appropriate conservation measures; the same difficulty affects all other species occupying a similar range.

Prior to fieldwork, an analysis of satellite images of 19 volcanoes in West Java attempted to assess, as best as possible, the extent of remaining montane forest on their slopes (Higginbottom

© The Author(s), 2023. Published by Cambridge University Press on behalf of BirdLife International. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

et al. 2019). This indicated that much of the most accessible lower-altitude montane forest has already disappeared and only some 5,200 km² of montane forest remains, often as fragmented isolates, although official protection has slowed deforestation rates in recent decades (Higginbottom *et al.* 2019). However, a further problem in assessing the conservation status and needs of the bird species in these forests is the intense pressure on Java's songbirds exerted by the cagebird and song competition industries (Marshall *et al.* 2020). So great is the concern over the fate of the Rufous-fronted Laughingthrush and Javan Green Magpie that they have become the precautionary subject of intensive (and expensive) captive-breeding initiatives (Collar *et al.* 2012, Owen *et al.* 2014), despite the possibility that populations might survive in some of the forests where no surveys have been undertaken in 50 years or more. Equally, if such populations survive but are in poor condition or simply remain unknown, the opportunity may be lost to put in place measures to secure them for the long term. Moreover, a further value in a modern inventory of these forests is their potential for reintroductions of captive-bred birds, if (1) the sites prove to be in good condition but "empty", having lost the species in question to trapping, and (2) they can be better protected under new management systems.

There are, however, also concerns for the loss of numbers in once extremely common species, e.g. white-eyes, leafbirds, shrikes, bushlarks, and even sunbirds and weavers, and the ecosystem services, such as seed dispersal and pollination, that they provide (e.g. Barros *et al.* 2019). While evidence of declines due to excessive trapping is clear in species on the brink of extinction (e.g. van Balen and Collar 2021), in Java, as elsewhere, much less is known about the scale of declines in commoner species, largely due to a lack of baseline historical data (e.g. Hughes 2017). This knowledge gap is

slowly being filled in Java's lowlands by initiatives such as the *BigMonth2020* citizen science event and the Indonesian bird atlas (Squires *et al.* 2021), and targeted repeat surveys of individual species (van Balen *et al.* 2022). For Java's montane birds, knowledge is far more rudimentary and restricted to notes of visiting bird-watchers or records from consultants.

We undertook a two-year bird survey across 11 West (plus one western Central) Javan mountains aimed at identifying areas for enhanced biodiversity protection; mammals and certain amphibians were also surveyed (see, e.g., Devenish *et al.* 2022), but will be reported elsewhere. Here we present occurrence data for bird species from 622 km of transects from 37 sites on the 12 mountains, and encounter rate data (groups encountered per hour) aggregated across sites. We do not identify bird records with particular sites or mountains for reasons of security, but these data are available to *bona fide* individuals on request.

Methods

We chose mountain sites based on an evaluation of current knowledge of the fauna and forest status of 20 montane areas in West Java (and Mount Slamet in Central Java; Marsden and Collar 2018 unpublished report). Twelve mountains (Figure 1) were chosen for surveys based on their large extents of remaining forest (Higginbottom *et al.* 2019), with the potential to provide habitat for species of conservation interest such as Rufous-fronted Laughingthrush and Javan Green Magpie. At each mountain, we chose sites in consultation with local villagers, in areas accessible to a field team along mountain trails as far into the forest area as possible, but also in proximity to water for the camp sites (Figure 1; map of sites). Table 1 shows information on survey effort across the




Figure 1. Survey sites (circles) in 12 montane areas, 2018–2021, (differing shades of grey denote clusters of sites corresponding to labelled mountains) located in West and Central Java, Indonesia; forest cover shown in green shading.

Table 1. The twelve mountain regions visited with dates, altitudes worked and survey effort.

Mountain	No. sites	Dates	Alt. range (m)	Transects	km	hrs
Slamet	5	14/9–16/10/2018	808–2751	52	103	219
Cikuray	2	19/11–5/12/2018	1593–2806	13	23	45
Patuha	2	24/2–6/3/2019	1793–2354	9	16	28
Tilu	3	8/4–6/5/2019	1295–2116	29	72	105
Malabar	2	17/7–29/7/2019	754–2322	15	43	51
Wayang-Windu	1	7/10–9/10/2019	1808–2160	9	16	32
Limbang	3	11/12/19–3/2/2020	994–1782	22	44	73
Masigit	6	13/3–12/10/2020	1100–2047	43	116	175
Kencana	5	28/8–22/9/2020	1091–2116	37	82	132
Simpang	2	8/8–20/8/2020	1044–1594	16	36	54
Papandayan	3	5/12–24/12/2020	1982–2321	22	37	54
Guntur	3	10/2–6/3/2021	1377–1933	28	35	65

12 mountains. To support our analyses, we took habitat recordings at 8–21 10-m radius plots positioned every 200 m along transect lines at each site. At each plot we measured/estimated a range of habitat features, but in this paper, we include assessments of forest type, counts of cut stems to indicate forest disturbance, along with cut trails and signs of trapping (see Appendix S2).

At each site, we walked transects of variable length and duration along trails emanating from our camp, noting bird species, number of individuals per group, and time of day. Transects were generally walked during the period 06h00–09h00, at an approximate speed of 1 km per hour. In all, 295 transects totalling 622 km and 1,031 hours' effort were walked across the 37 sites over 127 days between 14 September 2018 and 06 March 2021. The mean length of each transect was 2.1 km, with a mean number of transects per site of 8 (min = 4, max = 16), and a mean length per site of 16.8 km. Transects were walked by one or more of 13 experienced recorders, but with three recorders (ARJ: 206 km; GCA: 189 km; FM: 63 km) contributing nearly 75% of all transect length.

We expressed bird occurrence as the number of transects, sites, and mountains in which the species was recorded. Encounter rates were expressed as mean number of encounters with groups or individuals per hour of each transect. These were then aggregated to site level (including transects on which the species was not recorded) and averaged across all sites, but only where the species was recorded at least once. We present the final figures as mean encounter rate \pm standard deviation and a minimum and maximum site-level encounter rate (site absences excluded).

Although comparisons with similarly collected historical data from the mountains of West and Central Java are understandably rare, we do make some broad comparisons of our encounter rates for selected species with those made by BvB in 1981 on Mounts Gede–Pangrango and Puncak, just to the west of our surveys, and from 1995 at two sites on Mount Slamet (van Balen 1984, van Balen unpublished data).

In our analyses, we consider predominantly submontane species with IUCN threatened or Near Threatened classifications (BirdLife International 2021), and non-threatened but traded submontane species, including those regularly recorded in market or household surveys (e.g. Marshall *et al.* 2020), and those identified in the priority species list by the IUCN Asian Songbird Trade

Specialist Group (ASTSG; www.asiansongbirdtradesg.com). It should be remembered when reviewing the results of these analyses that none of these species is restricted to the mountains covered by this study, being found on at least one other mountain in Java. For security reasons, we do not name any specific mountains or sites in the Results section. Taxonomy follows del Hoyo and Collar (2014, 2016) plus Lim *et al.* (2018) for Sangkar White-eye *Zosterops melanurus* and Gwee *et al.* (2019) for Javan Blue-flycatcher *Cyornis banyumas*.

Results

Altogether, 234 bird species were recorded at any time during the surveys, with 184 of these recorded on the transects themselves. Appendix S1 provides a full list of occurrence and encounter rate data for all species recorded on transects. Encounter rates were positively skewed, with the majority of species occurring on few transects and at low rates (Figure 2a). In fact, only 14 species were recorded at rates above 0.5 encounters per hour, just 0.1 encounters per hour greater than the median value. Only two species (Javan Tesia *Tesia supercilialis* and Pygmy Cupwing *Pnoepyga pusilla*) had rates of >1 encounters per hour. Rates decreased with decreasing site/transect occupancy (Figure 2a), but showed little difference across categories of extinction risk (Figure 2b).

A total of 32 species of elevated conservation concern (2 Critically Endangered, 5 Endangered, 9 Vulnerable, and 16 Near Threatened) were recorded, either on transects (26 species) or incidentally (Table 2). Orange-spotted Bulbul *Pycnonotus bimaculatus* was the only species of elevated conservation concern to be recorded on transects on every mountain, being also found on most transects and generally at high encounter rates. The transects were clearly more appropriate for recording some species than certain others such as the nocturnal species; for example, Salvadori's Nightjar *Caprimulgus pulchellus* was encountered incidentally on all 12 mountains but only on five mountains during transect surveys. Surprisingly, the Javan Cochoa *Cochoa azurea*, a retiring, unobtrusive species, was recorded at nearly every site and on all but one mountain, while Javan Trogon *Apalharpactes reinwardtii*, previously known from only three of the mountains sampled prior to our surveys (Collar and van Balen 2002), was recorded at nine of them




Figure 2. Mean encounter rates (groups per hour) for 184 bird species across 37 sites in 12 western Javan mountain regions. Shown are (a) the relationship between transect occupancy (number of transects with species presence) and encounter rates (grey bars show 1 standard deviation); (b) median and variability of encounter rates grouped by global Red List categories (2021 assessment). CR = Critically Endangered; EN = Endangered; VU = Vulnerable; NT = Near Threatened; LC = Least Concern.

and in around two-thirds of sites. These two species also had reasonably high and quite consistent (low standard deviation) encounter rates across transects. Two heavily trapped threatened species – Rufous-fronted Laughingthrush (around half of mountains and sites) and Spotted Crocias *Laniellus albonotatus* (around two-thirds of sites/mountains) – proved to occur quite widely. Incidental records of the “Critically Endangered” Javan Blue-banded Kingfisher *Alcedo euryzona* at single sites on four mountains are notable as there are just a handful of records of the species since the 1930s (Chan and Setiawan 2019).

Of the 26 non-threatened but trapped species we considered (Table 3), four stand out as present at few sites, rarely encountered, or both, namely Javan Oriole *Oriolus cruentus* (seven sites on three mountains; IUCN Red List category Data Deficient), Chestnut-backed Scimitar-babbler *Pomatorhinus montanus* (six sites on four mountains), and Mountain Serin *Chrysocorythus estherae* and Javan Blue-flycatcher (both single records only). In contrast, Chestnut-bellied Partridge *Arborophila javanica*, Sunda Minivet *Pericrocotus miniatus*, Rufous-tailed Fantail *Rhipidura phoenicura*, Chestnut-fronted Shrike-babbler *Pteruthius aenobarbus*, Javan Grey-throated White-eye *Heleia javanica*, Snowy-browed Flycatcher *Ficedula hyperythra*, Indigo Flycatcher *Eumyias indigo*, Little Pied Flycatcher *Ficedula westermanni*, and White-flanked Sunbird *Aethopyga eximia* all occurred on over half of transects, at the great majority of sites on nearly all mountains. While these were fairly consistently recorded across transects at sites, two species, Mountain White-eye *Zosterops japonicus* (16 of 37 sites but only 32 of 295 transects), and White-bibbed Babbler *Stachyris thoracica* (20 sites, 35 transects), were found at a reasonable number of sites but only on very few transects, suggesting their local rarity.

Table 4 shows comparisons of mean encounter rates (groups per hour) across the surveys. We were able to make reasonable comparisons for 11 species. Of these, we posit that Javan Green Magpie, Chestnut-backed Scimitar-babbler, Javan Fulvetta *Alcippe pyrrhoptera*, and Javan Grey-throated White-eye appear to have encounter rates from our study markedly lower than those presented previously.

Discussion

Java holds high levels of bird endemism, and yet our study represents a rare attempt – another such being van Balen *et al.* (1999) – to gauge abundance systematically in the island’s key birds. It also represents the first documented ornithological surveys of many of the mountains in decades. This was a data gap that needed to be filled, given the rates of environmental change on the island and especially the breadth and volume of bird trapping to supply demand for songbirds (Eaton *et al.* 2015, Marshall *et al.* 2020). During over two years of biodiversity surveys, we recorded 234 species, including 32 threatened or “Near Threatened” taxa. Some species suspected to be scarce were in fact widespread and reasonably often encountered at sites. Species such as Javan Trogon and Javan Cochoa, and, to a certain extent, the traded Rufous-fronted Laughingthrush, were encouragingly well-recorded. There was, however, a larger suite of species that were rarer than anticipated: Crested Jay *Platylophus galericulatus*, White-breasted Babbler, Sangkar White-eye, Javan Oriole, Brown-cheeked Bulbul, White-bellied Fantail *Rhipidura euryura*, Chestnut-backed Scimitar-babbler, Mountain Serin, and Javan Blue-flycatcher were all either restricted to a few sites, uncommonly recorded within sites, or both. Crested Jay, recorded

Table 2. Occurrence and encounter rates per site for 32 bird species of conservation concern.

	Transect (n = 295)	Sites (n = 37)	Mts (n = 12)	ER ± SD (min-max)
Asian Woollyneck <i>Ciconia episcopus</i> NT	1	1 (1)	1 (1)	0.06
Javan Hawk-eagle <i>Nisaetus bartelsi</i> EN * ^s	25	11 (21)	7 (9)	0.09 ± 0.06 (0.02–0.20)
Rufous-bellied Eagle <i>Lophotriorchis kienerii</i> NT	0	(3)	(3)	
Javan Woodcock <i>Scolopax saturata</i> NT ^m	5	4 (6)	3 (3)	0.07 ± 0.04 (0.03–0.12)
Sumatran Green-pigeon <i>Treeron oxyurus</i> NT ^m	1	1(2)	1 (2)	0.03
Yellow-throated Hanging-parrot <i>Loriculus pusillus</i> NT * ^s	36	18 (22)	7 (9)	0.11 ± 0.08 (0.03–0.30)
Javan Scops-owl <i>Otus angelinae</i> VU * ^m	5	3 (18)	2 (10)	0.07 ± 0.10 (0.01–0.18)
Salvadori's Nightjar <i>Caprimulgus pulchellus</i> NT † ^m	16	8 (29)	5 (12)	0.09 ± 0.08 (0.02–0.25)
Waterfall Swift <i>Hydrochous gigas</i> NT ^m	2	1 (5)	1 (4)	0.11
Volcano Swiftlet <i>Aerodramus vulcanorum</i> NT * ^m	0	(5)	(3)	
Javan Trogon <i>Apalharpactes reinwardtii</i> VU * ^m	59	24 (27)	9 (10)	0.11 ± 0.07 (0.02–0.25)
Javan Blue-banded Kingfisher <i>Alcedo euryzona</i> CR *	1	1 (4)	1 (4)	0.02
Rhinoceros Hornbill <i>Buceros rhinoceros</i> VU †	0	(1)	(1)	
Wreathed Hornbill <i>Rhyticeros undulatus</i> VU	9	3 (4)	3 (3)	0.13 ± 0.10 (0.02–0.22)
Black-banded Barbet <i>Psilopogon javensis</i> NT *	5	4 (4)	3 (4)	0.05 ± 0.02 (0.03–0.07)
Javan Yellownape <i>Chrysophlegma mentale</i> NT * ^s	41	20 (22)	8 (10)	0.10 ± 0.07 (0.02–0.24)
Javan Flameback <i>Chrysocolaptes strictus</i> VU * ^s	4	3 (5)	3 (4)	0.07 ± 0.02 (0.05–0.09)
White-rumped Woodpecker <i>Meiglyptes tristis</i> EN *	2	2 (2)	2 (2)	0.04 ± 0.01 (0.03–0.05)
Javan Broadbill <i>Eurylaimus javanicus</i> NT * ^s	93	23 (25)	9 (9)	0.32 ± 0.24 (0.02–0.80)
Crested Jay <i>Platylophus galericulatus</i> NT †	11	5 (9)	4 (5)	0.11 ± 0.07 (0.04–0.22)
Bar-winged Prinia <i>Prinia familiaris</i> NT	1	1 (7)	1 (4)	0.05
Ruby-throated Bulbul <i>Rubigula dispar</i> VU	1	1 (2)	1 (1)	0.03
Orange-spotted Bulbul <i>Pycnonotus bimaculatus</i> NT ^m	100	31(32)	12 (12)	0.23 ± 0.30 (0.03–1.50)
Brown-cheeked Bulbul <i>Alphoixus bres</i> EN *	2	1 (1)	1 (1)	0.23
White-breasted Babbler <i>Stachyris grammiceps</i> NT * ^l	3	3 (3)	3 (3)	0.09 ± 0.10 (0.02–0.21)
Rufous-fronted Laughingthrush <i>Garrulax rufifrons</i> CR * ^m	39	14 (14)	6 (6)	0.16 ± 0.20 (0.02–0.60)
Spotted Crocias <i>Laniellus albonotatus</i> NT * ^m	77	21 (22)	8 (9)	0.27 ± 0.34 (0.03–1.38)
Sangkar White-eye <i>Zosterops melanurus</i> VU ^s	37	15 (20)	8 (8)	0.13 ± 0.09 (0.03–0.32)
Javan Myna <i>Acridotheres javanicus</i> VU	0	(4)	(3)	
Javan Cochoa <i>Cochoa azurea</i> VU * ^m	100	31 (31)	11 (11)	0.18 ± 0.15 (0.03–0.51)
Greater Green Leafbird <i>Chloropsis sonnerati</i> EN † ^l	0	(1)	(1)	
Javan Leafbird <i>Chloropsis cochinchinensis</i> EN * ^s	0	(5)	(4)	

Summary data are given for observations on transects (26 species), including encounter rate ± standard deviation, number of transects, sites and mountain regions. Incidental records away from transects are included in the totals inside parenthesis; six species were only observed off transects. Species which are currently heavily trapped are shown in **bold**. IUCN Red List categories are CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened. * Species endemic to Java and Bali; † species with subspecies endemic to Java and Bali. Elevational ranges (see Hoogerwerf 1948, Eaton et al. 2021) are m = strictly montane, mainly 1000–3000m, but many start at a lower altitude, s = found at 0–1500m but preferring the higher parts of this zone, l = found at 0–1500m, more inland, higher than sea level, but usually not higher than 800m; all other species are less restricted, but below 1500m.

at just five sites on four mountains, and White-breasted Babbler *Stachyris grammiceps*, on just three transects on three mountains, were likely rare in our surveys as most effort was above the elevational range of the species (survey effort, at just three and seven sites respectively, was within the core elevational range of the species: Eaton et al. 2021). Javan Oriole is so poorly known that it may never have been that common in Java's mountains (BirdLife International 2021), but trapping for the cagebird trade must surely be a concern for several taxa. For both the traded or

threatened species and the common Least Concern birds, our occurrence and encounter rate data represent a first baseline against which future trends in bird abundance can be gauged.

A general frustration in conservation biology is the lack of comparable historical data against which to gauge current population densities, thus preventing population trends from being accurately assessed (e.g. Annorbah et al. 2016). In our case, a literature review revealed no published papers that had used similar encounter rates along transects to survey montane birds in Indonesia, but

Table 3. Occurrence and encounter rates for 26 heavily trapped but non-threatened birds across the twelve Javan mountains surveyed.

	Transect (n = 295)	Sites (n = 37)	Mts (n = 12)	ER ± SD (min–max)
Chestnut-bellied Partridge <i>Arborophila javanica</i> * ^m	205	37	12	0.47 ± 0.23 (0.09–1.02)
Pink-headed Fruit-dove <i>Ptilinopus porphyreus</i> ^m	76	24	9	0.21 ± 0.20 (0.02–0.68)
Dark-backed Imperial-pigeon <i>Ducula lacernulata</i> † ^m	32	11	6	0.11 ± 0.11 (0.02–0.35)
Sunda Minivet <i>Pericrocotus miniatus</i> ^m	173	37	12	0.33 ± 0.22 (0.03–0.92)
Javan Oriole <i>Oriolus cruentus</i> * ^m	11	7	3	0.12 ± 0.08 (0.04–0.28)
Rufous-tailed Fantail <i>Rhipidura phoenicura</i> * ^m	172	36	12	0.33 ± 0.20 (0.03–0.78)
White-bellied Fantail <i>Rhipidura euryura</i> * ^m	23	10	6	0.13 ± 0.10 (0.02–0.35)
Javan Bulbul <i>Ixos virescens</i> * ^m	109	25	9	0.38 ± 0.27 (0.03–0.86)
White-bibbed Babbler <i>Stachyris thoracica</i> * ^m	35	20	9	0.11 ± 0.18 (0.01–0.87)
Chestnut-fronted Shrike-babbler <i>Pteruthius aenobarbus</i> † ^m	156	31	12	0.42 ± 0.29 (0.04–1.07)
Chestnut-backed Scimitar-babbler <i>Pomatorhinus montanus</i> † ^m	10	7	4	0.06 ± 0.04 (0.02–0.13)
Mountain White-eye <i>Zosterops japonicus</i> ^m	32	16	10	0.08 ± 0.06 (0.03–0.23)
Javan Grey-throated White-eye <i>Heleia javanica</i> † ^m	149	33	12	0.43 ± 0.39 (0.04–2.00)
Velvet-fronted Nuthatch <i>Sitta frontalis</i> †	32	14	8	0.14 ± 0.13 (0.02–0.54)
Blue Nuthatch <i>Sitta azurea</i> † ^m	68	26	11	0.12 ± 0.08 (0.02–0.28)
Javan Shortwing <i>Brachypteryx montana</i> * ^m	44	18	10	0.13 ± 0.10 (0.03–0.30)
Sunda [Javan] Blue Robin <i>Myiomela diana</i> † ^m	47	22	9	0.12 ± 0.11 (0.02–0.45)
Snowy-browed Flycatcher <i>Ficedula hyperythra</i> ^m	174	34	12	0.46 ± 0.28 (0.03–1.07)
Little Pied Flycatcher <i>Ficedula westermanni</i> ^m	232	36	12	0.85 ± 0.47 (0.08–1.99)
Indigo Flycatcher <i>Eumyias indigo</i> * ^m	156	33	12	0.34 ± 0.21 (0.04–0.85)
Javan Blue-flycatcher <i>Cyornis banyumas</i> †	1	1	1	0.04
White-flanked Sunbird <i>Aethopyga eximia</i> * ^m	162	36	12	0.37 ± 0.29 (0.02–1.39)
Javan Sunbird <i>Aethopyga mystacalis</i> * ^s	40	15	8	0.24 ± 0.28 (0.01–1.00)
Tawny-breasted Parrotfinch <i>Erythrura hyperythra</i> † ^m	20	10	6	0.09 ± 0.06 (0.02–0.21)
Pin-tailed Parrotfinch <i>Erythrura prasina</i> ^s	4	4	3	0.03 ± 0.02 (0.01–0.05)
Mountain Serin <i>Chrysocorythus estherae</i> ^m	1	1	1	0.03

* Species endemic to Java and Bali; † species with subspecies endemic to Java and Bali. See Table 2 for elevational ranges.

we did have reasonably comparable counts made in the 1980s and 1990s on the same or nearby mountains. We acknowledge that we must interpret these encounter rates with great caution, for several reasons including survey effort and seasonal differences, but most importantly because we are not comparing the same sites. This said, we do suggest that some potentially interesting patterns emerge. Several species in the current study appear to occur at encounter rates fairly like those from the 1980s and 1990s, i.e. the fantails, White-bibbed and Crescent-chested Babblers, Indigo Flycatcher, and notably Rufous-fronted Laughingthrush among them. There is some support for the notion that Orange-spotted Bulbul, Javan Fulvetta (see Appendix S1), and Javan Grey-throated White-eye may have declined, but this is not strong, given the necessary caveats. In contrast, Chestnut-backed Scimitar-babbler does seem to have become scarcer.

Our work produced a number of new localities for species of conservation interest. We found Javan Scops Owl *Otus angelinae* and Brown (or Sunda) Wood Owl *Strix (leptogrammica) bartelsi* on Mount Slamet for the first time, both formerly known from only a few sites. The relatively large and colourful Javan Trogon was found on Slamet, Cikuray, Limbung, Patuha, Masigit, Tilu,

Kencana, Simpang, and Papandayan. The more cryptic Javan Cochoa was recorded at all the above plus Malabar and Guntur. Sunda Grasshopper Warbler *Locustella montis* was found on Slamet and Tilu. White-breasted Babbler, a species known to be present on the foothills of Patuha, Cikuray, and Slamet (van Balen *et al.* 2005), was not recorded at these sites but compensated by turning up at three new sites (Masigit, Kencana, Papandayan) during our surveys. Mountain White-eye was previously recorded only as far west as Papandayan (Mees 1996), but we recorded it at several mountains (Patuha, Masigit, Tilu, Malabar, Kencana and Wayang-Windu) up to 50 km further west. None of the new localities can be considered to reflect recent colonisations; rather they far more likely represent lack of contact in earlier surveys. However, our failure to find White-breasted Babblers at three known sites for the species should be treated as a warning signal: the species may simply have been missed, perhaps because most of our survey efforts were above the elevations where it usually occurs, but it is equally possible that it has steeply declined or disappeared entirely. This is a species that joins understorey mixed flocks in numbers (van Balen *et al.* 2005) and, as such, might be easily caught in mist-nets. We encourage

Table 4. Comparisons of mean encounter rates (bird groups per hour) for selected songbirds between surveys done in the 1980s and 1990s and our study (numbers in parentheses are maximum and minimum at occupied sites). Also shown are dates, altitudes and survey effort (hours of morning fieldwork).

Attribute/Species	Gede-Pangrango 1981	Puncak 1981	Slamet 1995	This study
Dates	2/4–20/7	31/3–22/6	28/6–undated	14/09/18–06/03/21
Hours/days surveying	35.5 / 6	27.7 / 7	10.4 / 3	1031/127
Altitudinal range (m)	1,450–1,700	~1,600	600–2,500	754–2,806
Rufous-tailed Fantail	0.25	0.14	0.96	0.33 (0.03–0.78)
White-bellied Fantail	0.11	0	0.19	0.13 (0.02–0.35)
Javan Green Magpie	0.06	0.07	0	0
Orange-spotted Bulbul	0.17	0.61	0.96	0.23 (0.03–1.50)
White-bibbed Babbler	0.08	0.11	0	0.11 (0.01–0.87)
Chestnut-backed Scimitar-babbler	0.82	0.90	1.05	0.06 (0.02–0.13)
Crescent-chested Babbler	0.23	0.61	2.01	0.90 (0.12–1.84)
Javan Fulvetta	1.49	0.98	1.73	0.76 (0.25–2.90)
Rufous-fronted Laughingthrush	0.42	0.22	0	0.16 (0.02–0.60)
Javan Grey-throated White-eye	0.90	1.48	2.11	0.43 (0.04–2.00)
Indigo Flycatcher	0.20	0.29	0.29	0.34 (0.04–0.85)

future visiting birdwatchers to determine which of these scenarios is correct.

As with most status assessments of species in tropical forests, the lack of a historical baseline against which to compare current bird abundance (e.g. Hughes 2017) is frustrating. This is especially true of most of the mountains included in our survey, some of which have not been visited by biologists and naturalists for decades (as inferred from the absence of their names in online search engines considering both academic and popular postings). Without such a baseline, we can at least report on current occurrence and likely abundance, as a core portion of the montane avifauna is both widespread across mountains and readily recorded within sites. This includes Sunda Minivet, Rufous-tailed Fantail, Chestnut-fronted Shrike-babbler, several flycatchers, and White-flanked Sunbird. The list even includes some Red List species, like Javan Cochoa and Orange-spotted Bulbul, classified as Vulnerable and Near Threatened on account of habitat loss and trapping respectively (BirdLife International 2021). The abundance of Chestnut-bellied Partridge is encouraging, given the concern for other ground-dwelling galliforms in Java and elsewhere in Indonesia (Boakes *et al.* 2019). It seems likely that, in this part of Java at least, the partridge is no longer targeted for food in numbers by trappers. That components of Java's montane avifauna remain largely intact bodes well for both their populations in coming years, and for ecosystem functioning (e.g. Loreau *et al.* 2001).

In stark contrast, however, we had only a single and unconfirmed record, from one of the team's local guides, of the Critically Endangered Javan Green Magpie, and we must assume that excessive trade has pushed this once reasonably widespread but perhaps never common species (MacKinnon 1988, van Balen *et al.* 2013) to the very brink of extinction. Javan Blue-flycatcher, also suffering from trade pressure (Eaton *et al.* 2015), was recorded just once, although most of our survey effort was above its usual elevational range. Hoogerwerf (1969–1971) described it as “one of the commonest flycatchers in Java, perhaps more common at moderate elevations than in the lowlands or highlands”, while in the 1980s the species was described as “one of the commoner flycatchers at moderate to

high elevations” (MacKinnon 1988). It is now extremely rare in the lowlands (Eaton *et al.* 2021) and was recorded only three times in over 20,000 bird lists in a month-long citizen science event in Java and Bali (Squires *et al.* 2021). This species, about to be recognised as Critically Endangered (BirdLife International pers. comm.), clearly warrants urgent searches in forests not covered in our survey. While several babbler species appear to be relatively widespread, the current rarity of Chestnut-backed Scimitar-babbler is a major concern, given its frequency in bird markets (Chng and Eaton 2016, S. Marsden pers. obs.), and especially given that the taxon is likely soon to be treated as a Javan endemic by BirdLife International. This species was, in the 1980s and 1990s, relatively easily recorded in Java's mountains and was described as “a not uncommon bird, found in loose parties” (MacKinnon 1988). Similarly, the abundance of White-bellied Fantail seems to have declined: around a century ago it was collected in numbers similar to those of the related Rufous-tailed Fantail (M. Bartels 1895–1936 unpublished data) and both species were considered fairly common by MacKinnon (1988), yet White-bellied was recorded on nearly 10 times fewer transects than Rufous-tailed in our fieldwork. It is likely that the latter's higher elevational preference has served it well in terms of protection against forest alteration, excessive trapping, or both. However, these suggestions of abundance declines must be interpreted cautiously, although they certainly are backed up by the perceptions of ornithologists with experience on the island for decades (BvB pers. obs.).

While some of Java's montane areas, such as Mounts Halimun–Salak and Gede–Pangrango, have been formally protected as national parks since the last century (www.protectedplanet.net), the majority of the forested highlands in West and Central Java are under either weaker management or no protection at all (Higginbottom *et al.* 2019). Indeed, the objective of our fieldwork was either to support moves towards gazetted further areas as formal reserves, or to enhance protection in alternative ways. Several mountains have stood out as particularly warranting protection, including Mount Slamet, the furthest east of our sites, along with Masigit, Kencana, and Tilu (Devenish *et al.* 2022). Key taxa

driving these judgements included the Endangered Javan Hawk-eagle *Nisaetus bartelsi*, Javan Leopard *Panthera pardus melas*, and Javan Gibbon *Hylobates moloch*, which are among the Indonesian governments “Priority” species for recovery (Mardiastuti *et al.* 2008), but others such as Critically Endangered Rufous-fronted Laughingthrush should also guide decisions. Arguably, however, there are key birds on all mountains surveyed.

How individual sites are best protected is open to debate, but the Indonesian authorities have recently moved away from the idea of further “national parks” towards a more integrated form of land management. This, largely but not wholly in partnership with private sector entities, involves land and forest protection combined with income generation in an “Essential Ecosystem Areas” (EEAs) framework (Sahide *et al.* 2020, Devenish *et al.* 2022). Some of our key species will of course benefit from forest protection and restoration, particularly at the lower sections of mountains which have lost most forest in recent decades (Higginbottom *et al.* 2019). We encountered evidence of bird trapping at all but six of the 38 sites we visited, and all but one site had cut trails that may well have been used for bird trapping (Appendix S2). The survival of a suite of species including Javan Green Magpie, Crested Jay, Javan Blue-flycatcher, and Chestnut-backed Scimitar-babbler will depend on efforts over the next decade to: (1) reduce demand for songbirds; (2) enforce restrictions on trapping and trading of key species; (3) work with local communities at individual sites either to protect remaining populations or to create socio-ecological conditions suitable for re-introductions. This last action appears the most feasible at present, and indeed initiatives centred on species such as Javan Green Magpie are underway. A cornerstone of such initiatives must be to identify and create alternative livelihoods for those who currently gain at least part of their income from bird trapping.

Acknowledgements. Our surveys were supported by the Ministry of Environment and Forestry (Kementerian Lingkungan Hidup dan Kehutanan/KLHK). Permission to work in protected areas was granted by the West Java Agency of Conservation of Natural Resources (Balai Besar Konservasi Sumber Daya Alam Jawa Barat) through permit numbers SI.745/K.1/BIDTEK.1/KSA/03/2019 (Gunung Tilu Nature Reserve and Cimanggu Nature Tourism Park), SI.2733/K.1/BIDTEK.1/KSA/08/2020 (Gunung Simpang Nature Reserve), SI.16/SKW.5/11/2020 (Mount Papandayan Nature Reserve), and SI.03/SKW.5/02/2021 (Kawah Kamojang Nature Reserve). Sites outside the protected areas were surveyed with the permission of Perum Perhutani, through their Regional Division of West Java and Banten (Perum Perhutani Divisi Regional Jawa Barat dan Banten) number 230/016.5/SDM/Divre Janten. We thank our primary sponsor, Rainforest Trust, along with the Silent Forest campaign of the European Association of Zoos and Aquaria (EAZA) and Toledo Zoo. Chester Zoo provided funding for our surveys on Mount Slamet. We are very grateful for the help and hospitality of our guides, porters, and other members of the local communities who kept us safe and comfortable in the field. For assistance with fieldwork on Mount Slamet, we thank Rick Stanley and Gabby Salazar. The following volunteer students joined the project at various stages: Andhika Prasetya, Jehuda Chris Wahyu, Joshua Muhammad Ridwan Firdaus, Wilterza Nababan (Bandung Institute of Technology); Ahmad Nabil Faturahman, Feril Fizar Roziki, Mausul Rohman, Muhammad Lutfhi Ashiddiqi, Umar Fadhli Kennedy, Vanny Lesmana (IPB University); Arya Adi Pratama, Hafiz Aulia Khairy Rakananda (Jenderal Soedirman University); Ahmad Yasin Chumaidi (Nusa Bangsa University); Arya Teguh Prakasa, Ghani Komarudin, Giancoro Juan Alifi S., Iqbal Abi Yaghshyah, Marwan Agung Nugraha, Raihan Anshari (Padjadjaran University); Fathurrohman (State University of Jakarta).

Supplementary material The supplementary material for this article can be found at <http://doi.org/10.1017/S095927092300014X>.

References

- Andrew, P. (1985) An annotated checklist of the birds of the Cibodas–Gunung Gede Nature Reserve. *Kukila* 2: 10–28.
- Annorbah, N. N. D., Collar, N. J. and Marsden, S. J. (2016) Trade and habitat change virtually eliminate the Grey Parrot *Psittacus erithacus* from Ghana. *Ibis* 158: 82–91.
- van Balen, S. (1984) Comparison of Bird Counts and Bird Observations in the Neighbourhood of Bogor (Indonesia). MSc Student Report, State University of Utrecht.
- van Balen, S. and Collar N. J. (2021) The vanishing act: a history and natural history of the Javan Pied Starling *Gracupica jalla*. *Ardea* 109: 41–54.
- van Balen, S., Collar, N. J., Liley, D. and Rudyanto (2005) The White-breasted Babbler *Stachyris grammiceps* of Java: natural history and conservation status, especially on Gunung Halimun. *Forktail* 21: 139–146.
- van Balen, S., Eaton, J. A. and Rheindt, F. E. (2013) Biology, taxonomy and conservation status of the Short-tailed Green Magpie *Cissa [t.] thalassina* from Java. *Bird Conserv. Internatn.* 23: 91–109.
- van Balen, S., Nijman, V. and Sözer, R. (1999) Distribution and conservation of the Javan Hawk-eagle *Spizaetus bartelsi*. *Bird Conserv. Internatn.* 9: 333–349.
- van Balen, S., Saryanthi, R. and Marsden, S. J. (2022) Evidence of steep declines in the heavily-traded Javan White-eye from repeated standardised surveys. *Bird Conserv. Internatn* <https://doi.org/10.1017/S0959270922000144>.
- Barros, F. M., Peres, C. A., Pizo, M. A. and Ribeiro, M. C. (2019) Divergent flows of avian-mediated ecosystem services across forest-matrix interfaces in human-modified landscapes. *Landsc. Ecol.* 34: 879–894.
- BirdLife International (2021) IUCN Red List for birds. Accessed online 25 May 2021 from <http://www.birdlife.org>.
- Boakes, E. H., Fuller, R. A. and McGowan, P. J. K. (2019) The extirpation of species outside protected areas. *Conserv. Lett.* 12: e12608.
- Chan, B. P. L. and Setiawan, A. (2019) New record of the Critically Endangered Javan Blue-banded Kingfisher *Alcedo euryzona* in Central Java, Indonesia. *BirdingASIA* 31: 24–27.
- Chng, S. C. L. and Eaton, J. A. (2016) *In the market for extinction: eastern and central Java*. Petaling Jaya, Selangor, Malaysia: TRAFFIC.
- Collar, N. J. and van Balen, S. (2002) The Blue-tailed Trogon *Harpactes (Apalharpactes) reinwardtii*: species limits and conservation status. *Forktail* 18: 121–125.
- Collar, N. J. and van Balen, S. (2013) Notes for the conservation of the Rufous-fronted Laughingthrush *Garrulax rufifrons*. *Forktail* 29: 15–18.
- Collar, N. J., Gardner, L., Jeggo, D. F., Marcordes, B., Owen, A., Pagel, T., Pes, T., *et al.* (2012) Captive breeding and the most threatened birds in Asia. *BirdingASIA* 18: 50–57.
- Devenish, C., Junaid, A. R., Andriansyah, Saryanthi, R., van Balen, S., Kaprawi, F., Aprianto, G. C., *et al.* (2022) Biological richness of Gunung Slamet, Central Java, and the need for its protection. *Oryx* 56: 429–438.
- Eaton, J. A., van Balen, B., Brickle, N. W. and Rheindt, F. E. (2021) *Birds of the Indonesian Archipelago: Greater Sundas and Wallacea*. Second edition. Barcelona, Spain: Lynx Edicions.
- Eaton J. A., Shepherd C. R., Rheindt F. E., Harris J. B. C., van Balen S. (B.), Wilcove D. S. and Collar N. J. (2015) Trade-driven extinctions and near-extinctions of avian taxa in Sundaic Indonesia. *Forktail* 31: 1–12.
- Gwee, C. Y., Eaton, J. A., Garg, K. M., Alström, P., van Balen, S. (B.), Hutchinson, R. O., Prawiradilaga, D. M., *et al.* (2019) Cryptic diversity in *Cyornis* (Aves: Muscicapidae) jungle-flycatchers flagged by simple bioacoustic approaches. *Zool. J. Linn. Soc.* 20: 1–17.
- Higginbottom, T. P., Collar, N. J., Symeonakis, E. and Marsden, S. J. (2019) Deforestation dynamics in an endemic-rich mountain system: conservation successes and challenges in West Java 1990–2015. *Biol. Conserv.* 229: 152–159.
- Hoogerwerf, A. (1948) Contribution to the knowledge of the distribution of birds on the island of Java, with remarks on some new birds. *Treubia* 19: 83–137.
- Hoogerwerf, A. (1969–1971) On the ornithology of the Rhino Sanctuary Ujung Kulon in West Java (Indonesia). *Nat. Hist. Bull. Siam Soc.* 23: 9–65, 447–500; 24: 79–135.

- del Hoyo, J. and Collar, N. J. (2014) *The HBW–BirdLife International illustrated checklist of the birds of the world, 1: non-passerines*. Barcelona, Spain: Lynx Edicions.
- del Hoyo, J. and Collar, N. J. (2016) *The HBW–BirdLife International illustrated checklist of the birds of the world, 2: passerines*. Barcelona, Spain: Lynx Edicions.
- Hughes, A. C. (2017) Mapping priorities for conservation in Southeast Asia. *Biol. Conserv.* **209**: 395–405.
- Lim, B. T. M., Sadanandan, K. R., Dingle, C., Leung, Y. Y., Prawiradilaga, D. M., Irham, M., Ashari, H., et al. (2018) Molecular evidence suggests radical revision of species limits in the great speciator white-eye genus *Zosterops*. *J. Ornithol.* **160**: 1–16.
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., Hooper, D. U., et al. (2001) Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science* **294**: 804–808.
- MacKinnon, J. (1988) *Field guide to the birds of Java and Bali*. Yogyakarta, Indonesia: Gadjah Mada University Press.
- Mardiastuti, A., Kusriani, M. D., Mulyani, Y. A., Manullang, S. and Soehartono, T. (2008) *Arahan strategis konservasi spesies nasional 2008–2018*. Jakarta, Indonesia: Direktorat Jenderal Perlindungan Hutan dan Konservasi Alam-Departemen Kehutanan RI.
- Marshall, H., Collar, N. J., Lees, A. C., Moss, A., Yuda, P. and Marsden, S. J. (2020) Spatio-temporal dynamics of consumer demand driving the Asian Songbird Crisis. *Biol. Conserv.* **241**: 108237.
- Mees, G. F. (1996) *Geographical variation in birds of Java*. Issue 26. Cambridge, MA, USA: Nuttall Ornithological Club.
- Owen, A., Wilkinson, R. and Sözer, R. (2014) *Ex situ* conservation breeding and the role of zoological institutions and private breeders in the recovery of highly endangered Indonesian passerine birds. *Internatn. Zoo Yearb.* **48**: 199–211.
- Sahide, M. A. K., Fisher, M., Nasri, N., Dharmiasih, W., Verheijen, B. and Maryudi, A. (2020) Anticipating a new conservation bureaucracy? Land and power in Indonesia's Essential Ecosystem Area policy. *Land Use Policy* **97**: 104789.
- Squires, T., Yuda, P., Akbar, P., Collar, N. J., Devenish, C., Taufiqurrahman, I., Wibowo, W., et al. (2021) Citizen science rapidly delivers extensive distribution data for birds in a key tropical biodiversity area. *Glob. Ecol. Conserv.* **28**: e01680.
- Stattersfield, A. J., Crosby, M. J., Long, A. J. and Wege, D. C. (1998) *Endemic bird areas of the world: priorities for biodiversity conservation*. Conservation Series 7. Cambridge, UK: BirdLife International.
- Whitten, T., Soeriaatmadja, R. E. and Afiff, S. A. (1997) *The ecology of Java and Bali*. Oxford, UK: Oxford University Press.
- Winnasis, S., Yuda, P., Imron, M. A., Iqbal, M., Rudyanto and Wahyudi, H. A. eds. (2020) *Atlas Burung Indonesia: wujud karya peneliti amatir dalam memetakan burung nusantara*. Batu, Indonesia: Yayasan Atlas Burung Indonesia.