



# Left rostrolateral prefrontal cortex lesions reduce suicidal ideation in penetrating traumatic brain injury

Matteo Pardini,<sup>1</sup>  Jordan Grafman,<sup>2\*</sup> Vanessa Raymont,<sup>3,4</sup> Mario Amore,<sup>1</sup>  
Gianluca Serafini,<sup>1</sup> Michael Koenigs,<sup>5</sup> and Frank Krueger<sup>6\*</sup> 

<sup>1</sup>Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genoa

<sup>2</sup>Ospedale Policlinico S. Martino-IRCCS, Genoa

<sup>3</sup>Department of Psychiatry, University of Oxford, Oxford, United Kingdom

<sup>4</sup>Centre for Dementia Prevention, University of Edinburgh, United Kingdom

<sup>5</sup>Department of Psychiatry, University of Wisconsin-Madison, Madison, WI, USA

<sup>6</sup>School of Systems Biology, George Mason University, Fairfax, VA, USA

**Objectives.** The objective of this study is to evaluate the relationship between suicidal ideation (SI), structural brain damage, and cognitive deficits in patients with penetrating traumatic brain injury (pTBI).

**Methods.** Vietnam War veterans ( $n = 142$ ) with pTBI to the prefrontal cortex (PFC) underwent combination of neuropsychological and psychiatric examinations and non-contrast CT brain scan. Patients were divided into SI positive (SI+) and SI negative (SI-) groups according to the SI item of the Beck Depression Inventory.

**Results.** Lesions to the left rostrolateral PFC (rlPFC) were associated with a *lower risk* of SI independent of depression and global functioning. Left rlPFC lesion also reduced abstract reasoning skills, which mediated the lesion effects on suicide ideation.

**Conclusions.** The left rlPFC plays a crucial role in SI independently of depression and global functioning.

Received 7 May 2018; Accepted 30 November 2018

**Key words:** Suicidality, traumatic brain injury, cognition, frontal lobe.

## Introduction

Suicidal ideation (SI), defined as the occurrence of suicidal thoughts or wishes, is one of the most relevant predictors of suicidal behavior.<sup>1</sup> Suicidal behavior is

among the most prominent public health problems worldwide, being responsible for approximately 800,000 deaths per year.<sup>2</sup>

To date, a relative wealth of data exists regarding brain abnormalities in individuals with a history of previous suicidal attempts, showing a dysregulation of connectivity between deep gray matter and different regions of the prefrontal cortex (PFC) such as medial and orbital PFC regions.<sup>3</sup> However, not all people with SI go on to attempt suicide, and in some who attempted suicide the SI phase can be very brief. People with SI without any previous suicide attempt are somewhat less studied than those with previous suicide attempters. However, the available evidence *suggests* that the PFC could also play a role in SI development and maintenance.<sup>1</sup> The PFC is involved in emotional regulation,<sup>4</sup> abstract reasoning, and self-monitoring,<sup>5</sup> which are key cognitive and emotional correlates of SI.<sup>1</sup> The PFC, moreover, plays a role in the modulation<sup>6</sup> of the hypothalamic–pituitary–adrenal axis and the autonomic nervous

\*Address correspondence to: Frank Krueger, PhD, School of Systems Biology, George Mason University, 4400 University Drive, Fairfax, MS 2A1, VA 22030, USA. (Email: [frueger@gmu.edu](mailto:frueger@gmu.edu)). For general information regarding the Vietnam Head Injury Study (VHIS) registry please contact: Jordan Grafman, PhD, Cognitive Neuroscience Laboratory, Shirley Ryan AbilityLab, Chicago, IL, USA. (Email: [jgrafman@northwestern.edu](mailto:jgrafman@northwestern.edu))

This work was supported by the National Institute of Neurological Disorders and Stroke intramural research program and a project grant from the United States Army Medical Research and Materiel Command administered by the Henry M. Jackson Foundation (Vietnam Head Injury Study Phase III: A 30 Year Post-Injury Follow-Up Study, Grant number DAMD17-01-1-0675). We are grateful to S. Bonifant, B. Cheon, C. Ngo, A. Greathouse, K. Reding, and G. Tasick for their invaluable help with the testing of participants and organization of this study. We thank Sergio Paradiso, MD, for his insights on the possible effects of pTBI on SI.

system, which often are dysregulated in people with SI.<sup>7</sup> In line with these observations, people with SI and major depressive disorder have been shown to present functional alterations in a vast network, including parietal and prefrontal brain regions (i.e., rostralateral PFC, dorsal PFC, and frontal pole).<sup>8</sup>

In this study, we decided to investigate anatomical underpinnings of SI in a population of Vietnam War veterans who suffered a penetrating traumatic brain injury (pTBI) to the PFC. SI and suicidal behavior represent a significant problem in both war veterans and pTBI patients,<sup>9</sup> and this is an active focus of research in military medicine, neuropsychiatry, and neurorehabilitation. A quantitative analysis of focal lesion distribution (as we utilized in the current study with pTBI patients) enables identification of causal brain-behavior relationships by highlighting brain regions that when damaged impair specific behaviors, thereby indicating its necessary role in subserving those behaviors.<sup>10,11</sup> The pTBI approach has been shown to represent a suitable model to disentangle the effect of lesions in key brain regions from those of psychosocial factors in conditions such as pathological aggressiveness,<sup>12,13</sup> major depression,<sup>14</sup> and post-traumatic stress disorder (PTSD).<sup>14,15</sup> Our pTBI population presents with some unique characteristics, including minimal possible confounding factors such as availability of a pre-injury intelligence measure, demographic homogeneity, lack of pre-injury comorbidities, and shared circumstances causing pTBI.<sup>16</sup> Here, we evaluated whether damage in PFC regions was associated with increased SI and whether this association was mediated by changes in cognitive performance.

## Methods

### Recruitment

Patients were drawn from the Vietnam Head Injury Study (VHIS) (Phase III).<sup>16</sup> Phase III (2003–2006) was conducted at the Bethesda National Naval Medical Center (36–39 years post-injury). They underwent neurologic and psychiatric examinations and a non-contrast CT brain scan. The patients' pre-injury characteristics and clinical follow-up data were available from military and Veterans Administration records. Based on the goal of the study, only veterans with pTBI to PFC regions were included, whereas veterans with a history of anxiety or psychotic disorders, or personality disorders were excluded. Further, none of the eligible veterans had symptoms compatible with the clinical onset of a neurodegenerative disease. Based on those inclusion criteria, 142 male veterans (years of age at evaluation,  $58.3 \pm 0.3$ ; years of education,  $14.7 \pm 0.2$ ; handedness, 120 subjects right-handed, 22 non-right-handed) out of 197 veterans of VHIS were included. All patients gave informed consent to the study, which was approved by the

Institutional Review Board at the Bethesda Naval Medical Center.


### Psychopathological and cognitive evaluations

Depression was evaluated with the Beck Depression Inventory, II Edition (BDI-II), a 21-item self-report instrument for measuring the current severity of specific symptoms of depressive disorder in the previous 2 weeks.<sup>17</sup> Patients rated each item on a scale ranging from 0 to 3, with higher scores indicating greater symptom severity. SI was evaluated with the ninth item of the BDI-II (9i-BDI-II). Patients with a 9<sup>th</sup>-BDI-II score of 0 were assigned to the SI negative (SI-) group, while participants with higher scores to the SI positive (SI+) group.

To quantify childhood traumatic experiences, patients completed the Early Trauma Inventory (ETI) scale.<sup>18</sup> The presence of PTSD symptoms was assessed with the Clinician-Administered PTSD Scale (CAPS) – a structured interview previously used for the assessment of PTSD in individuals with pTBI.<sup>19</sup> Global functioning was evaluated with the Global Assessment of Functioning (GAF) scale,<sup>20</sup> providing a measure for the overall mental health and day-to-day functioning. The presence of current and/or lifetime alcohol dependence or mood disorder diagnoses was evaluated with the Structured Clinical Interview for DSM-IV-TR Axis I Disorders. Moreover, different facets of executive abilities were assessed with the Delis-Kaplan Executive Function System (DK) battery<sup>21</sup>: Trail Making Test (task switching), Verbal Fluency Test (phonemic verbal fluency), Sorting Test (abstract reasoning and conceptualization), Twenty Question Test (category formation and feedback acquisition), and Tower Test (spatial planning). Lastly, memory abilities were evaluated with the Wechsler Memory scale (III edition), focusing on the general memory and working memory scores.

### Pre- and post-injury intelligence evaluation

Pre-injury intelligence was evaluated with the Armed Forces Qualification Test (AFQT-7A), which was administered upon entry into the military. The AFQT-7A is a standardized multiple choice test of cognitive aptitude, measuring verbal ability, visual-spatial organization, and arithmetic and functional associations via multiple choice questions that has been extensively standardized within the US military.<sup>22</sup> The total score ranges from 0 to 100. Scores are reported as percentiles (1–99). Post-injury intelligence was evaluated with the Wechsler Adult Intelligence Scale (WAIS-3, 3rd edition) – a widely used intelligence test which comprises 14 subtests grouped in four first-order factors: verbal comprehension, perceptual organization, working memory, and processing speed. The AFQT-7A and the WAIS-3 are highly correlated.<sup>22</sup>



**FIGURE 1.** VLSM results. Voxel-wise binomial Liebermeister comparison of lesion distribution between the SI+ and SI- groups ( $p(\text{FWE-permutations}) < 0.05$ ), showing greater damage in the rIPFC for the SI- group compared to the SI+ group. Values represent FWE-corrected z scores.

### Lesion identification and voxel-based lesion-symptom mapping

Axial non-contrast CT scans were acquired on a GE Medical Systems Light Speed Plus CT scanner in helical mode. Images were reconstructed with an in-plane voxel size of  $0.4\text{mm} \times 0.4\text{mm}$ , overlapping slice thickness of  $2.5\text{mm}$ , and a  $1\text{-mm}$  slice interval. Lesion location and volume were determined from CT images by manual tracing using the Analysis of Brain Lesion (ABLe) software implemented in MEDx v3.44 (Medical Numerics) with enhancements to support the Automated Anatomical Labeling (AAL) atlas.<sup>23</sup> A trained neuropsychiatrist (VR) performed the tracings which were then reviewed by an experienced observer (JG), who was blind to the results of the clinical evaluations. The skull and scalp components of the CT volume were then removed; each volume was spatially normalized to a de-skulled CT scan, which was previously spatially normalized to match the shape of the T1 Montreal Neurological Institute (MNI) brain (standard of the International Consortium for Brain Mapping). The ABLe program was used to exclude the manually delineated lesion from the spatial normalization process to improve registration accuracy. Spatial normalization was performed using an automated image registration algorithm using a 12-parameter affine model on de-skulled CT scans. Lesion distribution is represented in Supplementary Figure 1.

MRICron ([www.mccauslandcenter.sc.edu/mricron/mricron](http://www.mccauslandcenter.sc.edu/mricron/mricron)) was used to evaluate the difference in the pattern of brain damage between SI+ and SI- subjects using a voxel-wise, permutation-based analysis of the

normalized lesion maps as reported below and extract regional values of volume loss as previously described.<sup>14</sup> Lesion distribution was compared between the SI+ and SI- groups using a voxel-wise binomial Liebermeister test<sup>24</sup> with a threshold of  $p < 0.05$  permutation-family wise error (FWE) corrected ( $p(\text{permutation-FWE})$ ) for multiple comparisons (3000 permutations). The Liebermeister test is an alternative to the Fisher Chi-square test and is thought to better account for unconstrained marginals usually encountered in lesion studies.<sup>24</sup> To reduce false-positive findings, the analysis was limited to those voxels in the PFC territories which were damaged in at least five patients. The same threshold was also used to correlate the presence of structural damage in any given voxel with performance on the Delis-Kaplan Executive Function System (D-KEFS) Sorting Test using the voxel-wise, permutation based Brunel-Munzel test.<sup>24</sup> The Brunel-Munzel test is an alternative to the more widely used  $t$  test to perform voxel-based lesion-symptom mapping (VLSM) analysis, since it does not require a normal distribution, similar variance between groups, and an interval of target measures.

### Behavioral statistical analysis

Statistical behavioral analyses were performed with SPSS 21 ([www.ibm.com/software/analytics/spss/](http://www.ibm.com/software/analytics/spss/)) and Stata 13.1 ([www.stata.com](http://www.stata.com)). A  $p$  value lower than 0.05 (two-tailed) was considered as significant.

Differences between SI+ and SI- groups in cognitive and psychopathological tests and percentage of volume loss were evaluated with independent samples  $t$  tests. An analysis of covariance (ANCOVA) on volume loss was

**TABLE 1. Sociodemographic, neurocognitive and lesion measures for SI+ and SI– groups**

	SI– (121 subjects)	SI+ (21 subjects)	t test <sup>a</sup>
Sociodemographic and neurocognitive variables			
Age	58.4 (0.3)	57.7 (0.5)	p = 0.401
Education (years)	14.7 (0.2)	14.5 (0.4)	p = 0.677
Pre-injury IQ (AFQT)	59.2 (2.2)	64.2 (5.6)	p = 0.393
Post-injury IQ (WAISS total score)	101.1 (1.4)	106.7 (2.0)	p = 0.124
WMS general memory primary index sum of standard scores	48.8 (0.9)	47.5 (2.1)	p = 0.561
WMS working memory primary index sum of standard scores	19.7 (0.4)	20.4 (0.8)	p = 0.465
DK trail making test number-letter switching scaled score ( <i>task switching</i> )	8.7 (0.4)	9.8 (0.8)	p = 0.241
DK letter fluency total scaled score ( <i>phonemic verbal fluency</i> )	8.7 (0.3)	9.1 (0.8)	p = 0.634
DK free sorting description score ( <i>abstract reasoning and conceptualization</i> )	<b>10.4 (0.3)</b>	<b>12.5 (0.5)</b>	<b>t = 2.8, p &lt; 0.005</b>
DK twenty Questions, total question asked score ( <i>category formation and feedback acquisition</i> )	9.4 (0.3)	10.1 (0.5)	p = 0.369
DK tower test total achievement score ( <i>spatial planning</i> )	10.5 (0.3)	11.9 (0.6)	p = 0.135
Psychometric instruments			
Early trauma inventory total score	4.5 (0.4)	5.7 (0.8)	p = 0.235
Beck depression inventory total score	<b>6.8 (0.6)</b>	<b>22.2 (2.5)</b>	<b>t = 9.0, p &lt; 0.001</b>
Global assessment of functioning	<b>78.1 (1.0)</b>	<b>67.5 (2.3)</b>	<b>t = 4.1, p &lt; 0.001</b>
Clinician-administered PTSD scale: no. of symptoms	<b>3.8 (0.3)</b>	<b>7.5 (1.0)</b>	<b>t = 3.5, p &lt; 0.002</b>
Alcohol abuse disorder diagnosis (lifetime)	3 patients (2.5%)	1 patient (4.8%)	p = 0.453 (chi-square)
Major depressive disorder diagnosis (lifetime)	15 patients (12.4%)	6 patients (28.6%)	p = 0.08 (chi-square)
Bipolar disorder diagnosis (lifetime)	1 patient (0.8%)	0 patients (0%)	
Percentage of damage			
Left rPFC	<b>11.1 (1.9)</b>	<b>0.8 (0.5)</b>	<b>t = 5.4, p &lt; 0.001</b>
Right rPFC	10.1 (1.8)	7.0 (3.3)	p = 0.491
Whole brain	3.6 (0.3)	3.1 (0.8)	p = 0.569
dIPFC	6.1 (0.8)	4.4 (1.6)	p = 0.389
dmPFC	2.2 (0.5)	1.8 (0.9)	p = 0.682
vlPFC	8.4 (1.2)	5 (2.1)	p = 0.285
vmPFC	6.9 (1.2)	4.4 (2.7)	p = 0.452
Posterior lobes ( <i>mean BA percentage of damage</i> )	1.6 (0.2)	2.0 (0.6)	p = 0.490

AFQT, armed forces qualification test; WMS, Weshler Memory scale III edition; DK, Delis–Kaplan; rl, rostralateral; dl, dorsolateral; dm, dorsomedial; vl, ventrolateral; vm, ventromedial; PFC, prefrontal cortex; BA, Brodmann area. Data reported as means (standard errors). Significant results reported in bold font.

<sup>a</sup>All analyses adjusted for heteroskedasticity. Results confirmed using bootstrap (3000 permutations).

performed with group (SI+, SI–) as a between-subjects factor and PTSD symptoms, childhood psychological trauma, global functioning, and whole brain damage as covariates. Differences in the frequency of alcohol abuse or mood disorders diagnoses between the two groups were assessed with a Chi-Square test. Normal distributions of targets measures were confirmed using a non-parametric bias-corrected and accelerated bootstrap with 3000 replicates. Effect sizes were calculated using Cohen’s *d* for those measures with significant differences between groups.

To link patients’ cognitive profile with brain damage, a bivariate correlation between left rPFC damage and DK free sorting description score was performed, since those were the only measures that showed significant differences between the two groups (SI+, SI–). As a follow-up, a mediation analysis was performed to verify whether the influence of left rPFC damage (independent variable) on SI


(dependent variable) was mediated through free sorting description score (mediator variable). The model was adjusted for depressive symptoms, PTSD symptoms, childhood psychological trauma, and global functioning because of their impact on SI. Since our prediction was directional (i.e., pTBI impacts sorting performance and SI), 90% bootstrap confidence intervals (CIs) were constructed by resampling the data 10,000 times with replacement.<sup>25</sup> A significant evidence for mediation existed if the 90% CI for the indirect effect did not include zero.

## Results

### Sociodemographic, neurocognitive, and psychopathological measures

The sociodemographic, cognitive, and psychopathological measures for the SI+ group (*n* = 21) and SP– group (*n* = 121 veterans) are reported in Table 1.





**FIGURE 2.** (Color online) Overlap (yellow area) between areas found to be associated with SI (green area, voxel-wise binomial Liebermeister test) and with abstract reasoning (red area, voxel-wise Brunel–Munzel test). Statistical threshold set at  $p < 0.05$  FWE-corrected for multiple comparisons after 3000 permutations. Image in neurological convention.

### Abstract reasoning performance differences between SI+ and –SI– groups

The SI+ group demonstrated impaired abstract reasoning skills (as assessed with the DK free sorting description score) compared to the SI– group ( $d = 0.73$ , medium effect size), while performing similarly on other executive functions (Table 1).

### Lesion distribution differences between SI+ and SI– groups

The rostralateral prefrontal cortex (rPFC) was significantly more damaged in the SI– group compared to the SI+ group (peak at MNI coordinates:  $-26, 54, -4$ ) (Figure 1). No brain regions were more damaged in the SI+ than in the SI– group. Confirmatory analyses on regional volume loss showed that the SI– group had more damage to the left rPFC than the SI+ group ( $11.1 \pm 1.9$  vs.  $0.8 \pm 0.5$ ,  $t = 5.4$ ,  $p < 0.001$ ), but both groups showed similar damage across the entire brain and other brain regions (e.g., right rPFC and other PFC regions) (Table 1). The difference in left rPFC damage between the SI+ and SI– groups remained significant not only after controlling for differences in depressive symptoms, PTSD symptoms, childhood psychological trauma, and whole brain damage, but also after excluding participants with a history of mood disorders or alcohol-dependence disorder (see Supplementary Table 1).

### Mediation of rPFC damage on SI via abstract reasoning skills


A significant negative correlation was observed between left rPFC damage and abstract reasoning (as assessed

with the DK free sorting description score) ( $r = -0.247$ ,  $p < 0.003$ ), demonstrating that more damage in the left rPFC leads to lower performance in abstract reasoning. A VLSM analysis confirmed a role of left rPFC in abstract reasoning ( $p(\text{FWE}) < 0.05$ ) (Figure 2). The impact of left rPFC damage on SI was significantly mediated by changes in abstract reasoning abilities: 39% of the total impact of left PFC damage on SI was mediated by the role played by this region on abstract reasoning (Figure 3).

### Discussion

The goal of the study was to evaluate the role of structural PFC damage on SI and its cognitive *underpinnings* in patients with pTBI. The results showed that left rPFC damage was associated with reduced SI presence; this association was independent of depressive symptoms and global functioning and partly mediated by the role of left rPFC for abstract reasoning.

An association between rPFC and SI is in line with previous evidence. The rPFC plays a relevant role in different psychological processes associated with SI, including guilt,<sup>26</sup> ruminations,<sup>27</sup> and performance monitoring.<sup>28</sup> Moreover, in patients with mood disorders, SI and suicidal behavior have been associated with long-range disconnections of rPFC territories with other frontal and temporal regions<sup>8</sup> and with changes in rPFC functional architecture.<sup>29</sup> Further, a recent study in patients with major depressive disorder has shown that the rPFC is associated with an abnormal frontotemporal pattern of connectivity in patients with SI compared to those without SI. In particular, results from graph-theory analysis revealed that the severity of SI was associated



**FIGURE 3.** Mediation analysis results. All effects are significant as shown by the CIs. Upper panel: total effect (*c*) of rIPFC damage on SI. Lower panel: direct (*c'*) and mediated (*ab*; via free sorting performance) effects of rIPFC damage on SI. Model adjusted for depressive symptoms, post-traumatic stress disorder symptoms, childhood psychological trauma and global functioning. s.e., standard errors; SI, suicidal ideation; DK, Delis–Kaplan; rIPFC, rostralateral PFC.

with the strength of information crossing the rIPFC.<sup>8</sup> Finally, a diffuse increase in prefrontal activity (including rIPFC and medial/dorsal PFC territories) was observed during a functional MRI (fMRI) error-monitoring task in war veterans with SI compared to veterans without SI.<sup>28</sup>

A protective effect of structural damage on SI – the association between rIPFC damage and reduced SI – seems unexpected at a first glance. However, the crucial role of rIPFC in abstract reasoning as reported in our study and in the literature<sup>30</sup> could help to interpret this unexpected finding. Abstract reasoning abilities, such as deliberating the irreversibility of death, underlie some of the assumptions needed to develop SI.<sup>31</sup> Indeed, SI is uncommon in children<sup>31</sup> and adolescents with learning disabilities,<sup>32</sup> that is, before the maturation of abstract reasoning abilities. Moreover, SI has been reported to be associated with better cognitive abilities in subjects with schizophrenia, confirming the need for relatively preserved cognition to allow for the emergence of SI.<sup>33</sup> Thus, the reduction in abstract reasoning skills due to rIPFC damage could moderate SI. Although the mediation analysis was significant, only around a third of the observed impact of rIPFC damage on SI was mediated by abstract reasoning. This is not surprising given the multifaceted role played by rIPFC on executive functioning. Since a key role for the left hemisphere in SI has been reported in other mood disorder studies,<sup>8,34</sup> we argue that this lateralization could be due to the role played by the left hemisphere in some cognitive functions composing SI such as abstract reasoning<sup>30</sup> and emotion recognition.<sup>35,36</sup>

We acknowledge that the relationship between cognitive functioning and SI is complex. In elderly subjects, for example, worse executive functions have been associated with an increased prevalence of SI<sup>37</sup>; while in young subjects with major depression, SI has been associated with increased cognitive inflexibility and reduced executive performances.<sup>38</sup> These differences are probably due to

the heterogeneity in both the enrolled populations and the extent of cognitive deficits. Future studies using the same psychometric and cognitive measures across different age groups, as well as across neuropsychiatric diagnoses, are needed to better clarify the role of cognition on SI development in subjects with and without structural brain damage.

Regarding the relationship between SI and TBI, most of published studies report an increase in SI frequency after TBI. As in the general population, depressed mood, hopelessness, and low social support are thought to represent the more relevant risk factors for SI after TBI.<sup>39,40</sup> Conversely, TBI severity has not been firmly associated with increased risk of SI<sup>40</sup>; however, published studies have been focused on overall severity rather than the role of regional damage.

This work expands on a previous research study using the VHIS database focused on the relationship between emotional competence and SI, showing a key role of emotional competence in reducing the risk of SI.<sup>41</sup> Here we decided instead to focus on more basic cognitive functions (i.e., the different facets of executive abilities) to evaluate the cascade leading from prefrontal damage to cognitive functions to SI. These two points of view (i.e., a top-down method starting from higher-order cognitive constructs that influence SI and a bottom-up approach focused on regional damage associated with mental states) complement each other and help portray the complexity of the SI experience.

The present study needs to be considered in the light of the following limitations. First, a homogeneous population of male war veterans with pTBI was investigated, which reduces the generalizability of our findings. Second, no information was available on the nature of SI (e.g., degree of planning, possible lethality, and access to means) in our population. Therefore, future studies are needed to validate our results that focus on other clinical populations with discrete brain lesions (e.g.,

stroke and multiple sclerosis) and to associate psychopathological characteristics of SI with cognitive and lesion patterns in pTBI patients. Another limitation is represented by the use of a single item of the BDI to evaluate SI. While not available at the time of data collection, to date a number of new psychometric instruments have been developed to better probe the SI construct, such as the Columbia-suicide severity rating scale.<sup>42</sup> A positive replication of the present findings using these scales would strengthen the results reported in this work.

Overall, this study is not the first to report a paradoxical mitigating role of structural brain damage on a psychopathological construct. Using the same population, our group has shown previously that damage to the amygdala and the medial PFC is associated with reduced PTSD<sup>14</sup> and risk for depression<sup>14</sup>, respectively. Indeed, taking the results of these previous study into account, we have demonstrated an association between rIPFC and SI independently of depressive symptoms (total BDI scores), PTSD symptoms, and global functioning.

In summary, the results indicate that SI depends on the integrity of the left rIPFC and is at least partly dependent upon the cognitive and emotional processes required for complex ideational thinking. It may be that interventions designed to change and shape such deliberative thinking would be especially beneficial for people with SI.

## Disclosures

Matteo Pardini receives research support from Novartis and fees from Merck for participation in advisory board activities. Jordan Grafman, Vanessa Raymont, Mario Amore, Gianluca Serafini, Michael Koenigs, and Frank Krueger do not have anything to disclose.

## Supplementary materials

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1092852918001694>.

## REFERENCES:

- Mann JJ, Waternaux C, Haas GL, *et al.* Toward a clinical model of suicidal behavior in psychiatric patients. *Am J Psychiatry*. 1999; **156**(2): 181-189.
- Vijayakumar L, Phillips MR, Silverman MM, *et al.*, eds. *Mental, Neurological, and Substance Use Disorders: Disease Control Priorities* Vol. 4. 3rd ed. Washington, DC; World Bank; 2016.
- Ding Y, Lawrence N, Olie E, *et al.* Prefrontal cortex markers of suicidal vulnerability in mood disorders: a model-based structural neuroimaging study with a translational perspective. *Transl Psychiatry*. 2015; **5**: e516.
- Soares JC, Mann JJ. The anatomy of mood disorders - review of structural neuroimaging studies. *Biol Psychiatry*. 1997; **41**(1): 86-106.
- Serafini G, Pardini M, Pompili M, *et al.* Understanding suicidal behavior: the contribution of recent resting-state fMRI techniques. *Front Psychiatry*. 2016; **7**: 69.
- Sullivan RM, Gratton A. Prefrontal cortical regulation of hypothalamic-pituitary-adrenal function in the rat and implications for psychopathology: side matters. *Psychoneuroendocrinology*. 2002; **27**(1-2): 99-114.
- Braquehais MD, Picouto MD, Casas M, *et al.* Hypothalamic-pituitary-adrenal axis dysfunction as a neurobiological correlate of emotion dysregulation in adolescent suicide. *World J Pediatr*. 2012; **8**(3): 197-206.
- Myung W, Han CE, Fava M, *et al.* Reduced frontal-subcortical white matter connectivity in association with suicidal ideation in major depressive disorder. *Transl Psychiatry*. 2016; **6**(6): e835.
- Fonda JR, Fredman L, Brogly SB, *et al.* Traumatic brain injury and attempted suicide among veterans of the wars in Iraq and Afghanistan. *Am J Epidemiol*. 2017; **186**(2): 220-226.
- Kwok SC. Where neuroimaging and lesion studies meet. *J Neuroimaging*. 2013; **23**(1): 1-4.
- Adolphs R. Human lesion studies in the 21<sup>st</sup> century. *Neuron*. 2016; **90**(6): 1151-1153.
- Pardini M, Krueger F, Hodgkinson C, *et al.* Prefrontal cortex lesions and MAO-A modulate aggression in penetrating traumatic brain injury. *Neurology*. 2011; **76**(12): 1038-1045.
- Pardini M, Krueger F, Hodgkinson CA, *et al.* Aggression, DRD1 polymorphism, and lesion location in penetrating traumatic brain injury. *CNS Spectr*. 2014; **19**(5): 382-390.
- Koenigs M, Huey ED, Calamia M, *et al.* Distinct regions of prefrontal cortex mediate resistance and vulnerability to depression. *J Neurosci*. 2008; **28**(47): 12341-12348.
- Pardini M, Krueger F, Koenigs M, *et al.* Fatty-acid amide hydrolase polymorphisms and post-traumatic stress disorder after penetrating brain injury. *Transl Psychiatry*. 2012; **2**: e75.
- Raymont V, Salazar AM, Krueger F, *et al.* "Studying injured minds" - the Vietnam head injury study and 40 years of brain injury research. *Front Neurol*. 2011; **2**: 15.
- Beck AT, Steer RA, Ball R, *et al.* Comparison of beck depression inventories-IA and -II in psychiatric outpatients. *J Pers Assess*. 1996; **67**(3): 588-597.
- Bremner JD, Vermetten E, Mazure CM. Development and preliminary psychometric properties of an instrument for the measurement of childhood trauma: the Early Trauma Inventory. *Depress Anxiety*. 2000; **12**(1): 1-12.
- Blake DD, Weathers FW, Nagy LM, *et al.* The development of a clinician-administered PTSD Scale. *J Trauma Stress*. 1995; **8**(1): 75-90.
- Hall RC. Global assessment of functioning. A modified scale. *Psychosomatics*. 1995; **36**(3): 267-275.
- Keifer E, Tranel D. A neuropsychological investigation of the Delis-Kaplan Executive Function System. *J Clin Exp Neuropsychol*. 2013; **35**(10): 1048-1059.
- Grafman J, Jonas BS, Martin A, *et al.* Intellectual function following penetrating head injury in Vietnam veterans. *Brain*. 1988; **111**(Pt 1): 169-184.
- Solomon J, Raymont V, Braun A, *et al.* User-friendly software for the analysis of brain lesions (ABLE). *Comput Methods Programs Biomed*. 2007; **86**(3): 245-254.
- Rorden C, Karnath HO, Bonilha L. Improving lesion-symptom mapping. *J Cogn Neurosci*. 2007; **19**(7): 1081-1088.
- Coussons-Read ME, Lobel M, Carey JC, *et al.* The occurrence of preterm delivery is linked to pregnancy-specific distress and elevated inflammatory markers across gestation. *Brain Behav Immun*. 2012; **26**(4): 650-659.
- Pulec E, Zahn R, Elliott R. The role of self-blaming moral emotions in major depression and their impact on social-economical decision making. *Front Psychol*. 2013; **4**: 310.
- Jacobs RH, Jenkins LM, Gabriel LB, *et al.* Increased coupling of intrinsic networks in remitted depressed youth predicts rumination and cognitive control. *PLoS One*. 2014; **9**(8): e104366.

28. Matthews S, Spadoni A, Knox K, *et al.* Combat-exposed war veterans at risk for suicide show hyperactivation of prefrontal cortex and anterior cingulate during error processing. *Psychosom Med.* 2012; **74**(5): 471–475.
29. Pu S, Nakagome K, Yamada T, *et al.* Suicidal ideation is associated with reduced prefrontal activation during a verbal fluency task in patients with major depressive disorder. *J Affect Disord.* 2015; **181**: 9–17.
30. Fine EM, Delis DC, Dean D, *et al.* Left frontal lobe contributions to concept formation: a quantitative MRI study of performance on the Delis-Kaplan Executive Function System Sorting Test. *J Clin Exp Neuropsychol.* 2009; **31**(5): 624–631.
31. Garfinkel BD, Froese A, Hood J. Suicide attempts in children and adolescents. *Am J Psychiatry.* 1982; **139**(10): 1257–1261.
32. Carlson GA, Asarnow JR, Orbach I. Developmental aspects of suicidal behavior in children and developmentally delayed adolescents. *New Dir Child Dev.* 1994; **(64)**: 93–107.
33. Delaney C, McGrane J, Cummings E, *et al.* Preserved cognitive function is associated with suicidal ideation and single suicide attempts in schizophrenia. *Schizophr Res.* 2012; **140**(1–3): 232–236.
34. Marchand WR, Lee JN, Johnson S, *et al.* Striatal and cortical midline circuits in major depression: implications for suicide and symptom expression. *Prog Neuropsychopharmacol Biol Psychiatry.* 2012; **36**(2): 290–299.
35. Dal Monte O, Schintu S, Pardini M, *et al.* The left inferior frontal gyrus is crucial for reading the mind in the eyes: brain lesion evidence. *Cortex.* 2014; **58**: 9–17.
36. Leopold A, Krueger F, dal Monte O, *et al.* Damage to the left ventromedial prefrontal cortex impacts affective theory of mind. *Soc Cogn Affect Neurosci.* 2012; **7**(8): 871–880.
37. Gujral S, Dombrowski AY, Butters M, *et al.* Impaired executive function in contemplated and attempted suicide in late life. *Am J Geriatr Psychiatry.* 2014; **22**(8): 811–819.
38. Marzuk PM, Hartwell N, Leon AC, *et al.* Executive functioning in depressed patients with suicidal ideation. *Acta Psychiatr Scand.* 2005; **112**(4): 294–301.
39. Tsaousides T, Cantor JB, Gordon WA. Suicidal ideation following traumatic brain injury: prevalence rates and correlates in adults living in the community. *J Head Trauma Rehabil.* 2011; **26**(4): 265–275.
40. Simpson G, Tate R. Suicidality after traumatic brain injury: demographic, injury and clinical correlates. *Psychol Med.* 2002; **32**(4): 687–697.
41. Paradiso S, Beadle JN, Raymond V, *et al.* Suicidal thoughts and emotion competence. *J Clin Exp Neuropsychol.* 2016; **38**(8): 887–899.
42. Posner K, Brown GK, Stanley B, *et al.* The Columbia-Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am J Psychiatry.* 2011; **168**(12): 1266–1277.