

Original Article

Cite this article: Radford BN, Han VKM. (2019) Offspring from maternal nutrient restriction in mice show variations in adult glucose metabolism similar to human fetal growth restriction. *Journal of Developmental Origins of Health and Disease* 10: 469–478. doi: 10.1017/S2040174418000983

Received: 9 July 2018
Revised: 5 September 2018
Accepted: 31 October 2018
First published online: 3 December 2018

Key words:

fetal growth restriction; glucose metabolism; insulin sensitivity; liver

Address for correspondence:

V.K.M. Han, MD, Children's Health Research Institute, Victoria Hospital, 800 Commissioners Rd E, London, ON, Canada N6A 5W9.
E-mail: Victor.Han@lhsc.on.ca

Offspring from maternal nutrient restriction in mice show variations in adult glucose metabolism similar to human fetal growth restriction

B. N. Radford^{1,2} and V. K. M. Han^{1,2,3}

¹Department of Biochemistry, Schulich School of Medicine & Dentistry, Western University, London, ON, Canada, ²Children's Health Research Institute, London, ON, Canada and ³Department of Pediatrics, Schulich School of Medicine & Dentistry, Western University, London, ON, Canada

Abstract

Fetal growth restriction (FGR) is a pregnancy condition in which fetal growth is suboptimal for gestation, and this population is at increased risk for type 2 diabetes as adults. In humans, maternal malnutrition and placental insufficiency are the most common causes of FGR, and both result in fetal undernutrition. We hypothesized that maternal nutrient restriction (MNR) in mice will cause FGR and alter glucose metabolism in adult offspring. Pregnant CD-1 mice were subjected to MNR (70% of average *ad libitum*) or control (*ad libitum*) from E6.5 to birth. Following birth, mice were fostered by mothers on *ad libitum* feeds. Weight, blood glucose, glucose tolerance and tissue-specific insulin sensitivity were assessed in male offspring. MNR resulted in reduced fetal sizes but caught up to controls by 3 days postnatal age. As adults, glucose intolerance was detected in 19% of male MNR offspring. At 6 months, liver size was reduced ($P=0.01$), but pAkt-to-Akt ratios in response to insulin were increased 2.5-fold relative to controls ($P=0.004$). These data suggest that MNR causes FGR and long-term glucose intolerance in a population of male offspring similar to human populations. This mouse model can be used to investigate the impacts of FGR on tissues of importance in glucose metabolism.

Introduction

Fetal growth restriction (FGR) or intrauterine growth restriction (IUGR) is a pregnancy condition in which the fetus fails to achieve optimal growth expected for gestational age resulting in a small for gestational age (SGA) newborn with birth weight of <10th percentile. The most common causes include maternal malnutrition in developing countries and placental insufficiency in developed countries.¹ Irrespective of the cause, poor macro- and micro-nutrient and substrate concentrations in the fetus are common pathophysiological findings. Growth-restricted infants are at increased risk for morbidity and mortality in the perinatal period,² and for many chronic diseases, including cardiovascular diseases,³ metabolic disorders such as type 2 diabetes and glucose intolerance,³ and obesity and abnormal lipid metabolism.⁴

Associations between poor fetal growth and adult health and disease have led to the concept of *Developmental Origins of Health and Disease* (DOHaD), which proposes that environmental exposures such as nutrient restriction during development can result in metabolic adaptations,⁵ which may enhance the survival of the fetus but increase their risk for metabolic disorders as children and adults. The concept of a mismatched environment underlying the association between fetal nutrient environment and predisposition to adult diseases is defined in the 'thrifty phenotype hypothesis'.⁵

FGR has been studied using maternal nutrient restriction (MNR) in animals such as rats^{6–8} and guinea pigs,⁹ or larger mammals like pigs,^{10,11} providing a short lifespan or similar morphology and diet to humans, respectively. These models have a limited capacity for genetic manipulation in offspring to identify genes and epigenetic changes moderating impacts of fetal nutrition. Such limitations can be addressed by using MNR in mice.

Models of growth restriction in rodents with variation in the duration of nutrient restriction provide insight on certain types of fetal undernutrition, either due to the onset of placental insufficiency or food availability. For example, nutrient restriction before and throughout pregnancy^{9,12} in rodents model mothers that experience extended nutritional deficits, however, does not mimic fetal undernutrition due to placental insufficiency. Conversely,

nutrient restriction beginning later in gestation may represent placental insufficiency.⁶ Nutrient restriction just after implantation in the mouse at E6.5 has not been studied and would be more similar to early- to mid-gestation-onset placental insufficiency.

Evidence suggests that metabolic adaptations to fetal under-nutrition are tissue-specific and serum peptides for obesity and diabetes can aid in identification of impacted tissues. Incretin hormones, such as glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic peptide (GIP), are released by the gut upon ingestion of food and stimulate release of insulin.^{13,14} Pancreatic hormones, such as insulin and glucagon, regulate tissue glucose uptake and energy utilization. In addition to hormone release, c-peptide, the cleaved portion of proinsulin, can indicate changes in insulin maturation.¹⁵ Leptin, resistin¹⁶ and plasminogen activator inhibitor (PAI-1)¹⁷ are secreted primarily from adipose tissue in mice. Leptin along with ghrelin, which is released from the gut, regulate appetite.^{18,19} Changes in these markers in addition to other metabolic tests can provide insight into tissues with metabolic adaptations.

The objective of this study was to establish a moderate total calorie nutrient restriction in mice to mimic human FGR populations and examine the long-term metabolic complications. We hypothesized that a 30% reduction in maternal food consumption by weight during early and late pregnancy would result in reduced fetal and newborn weight and tissue-specific alterations in adult glucose metabolism similar to humans.

Methods

Animals

The animal experiments were conducted as approved by the Council on Animal Care at the University of Western Ontario. All mice received free access to water in a standard 12 h light/dark cycle. Eight-week-old virgin female CD-1 mice, obtained from Charles River Laboratories (Montreal, PQ, Canada), were mated and the presence of mucous plugs indicated E0.5. After mating mice were housed individually. To avoid reduction in the number of conceptuses of each litter, mated females received *ad libitum* until after embryo implantation at about E5. At E6.5, the mice were randomly assigned into two groups, the control group was fed *ad libitum* and the MNR group was fed 70% average *ad libitum* total calorie intake (both groups receiving #F0173, Bio-Serv, Flemington, NJ, USA). At E18.5, 3 control and 3 MNR pregnant females were euthanized by CO₂ narcosis. No differences between MNR and control litters (Supplementary Table 1). Litter size ranged from 6 to 22 pups. To control for variations in litter size, only litters with 11–15 pups were analyzed in fetal (control litters = 3, MNR litters = 3) and long-term studies (control litters = 5, MNR litters = 10). Because many studies suggest that male offspring are more sensitive to maternal undernutrition^{20–22} and that postreproductive changes complicate long-term studies in females,²³ only males were examined in this study. SRY polymerase chain reaction (forward primer: TGGACTGGTGA-CAARGCTA, reverse primer: TGGAAGTACAGGTGTGCACT CT) was used in fetal studies to include males only (control: $n = 21$, MNR: $n = 15$). Tissues were snap frozen in liquid nitrogen.

For postnatal studies, pups' paws were tattooed at birth in order to track weights for each pup with the Armis Microtattoo kit (Braintree Scientific, Inc., Braintree, MA, USA). All litters were cross-fostered to females that were fed *ad libitum* throughout

pregnancy and the litters were culled or fostered to 13 pups each. *Ad libitum* standard chow (Teklad LM-485 Mouse/Rat Sterilizable Diet, Mississauga, ON, Canada) was used for the remainder of the study in both MNR and control offspring.

At weaning (P21), only male offspring were kept for further study (control litters: $n = 5$, MNR litters: $n = 10$) and were housed with 3–5 siblings per cage. Weight was measured once a month. With the exception of littermates used for the intrahepatic portal vein insulin challenge, offspring at 1 month (control: $n = 8$, MNR: $n = 16$) and 6 months (control: $n = 25$, MNR: $n = 36$) were euthanized with CO₂ narcosis. Weight was measured, and serum was collected through intracardiac puncture. Individual liver lobes, perigonadal adipose tissue and quadriceps femoris were collected, snap frozen in liquid nitrogen and stored at -80°C until analysis. Animals used for the intrahepatic portal vein insulin challenge ($n = 8$ per group) were euthanized at 7 months after tissue collections.

Glucose tolerance and hepatic glucose production

Glucose tolerance was assessed in 36 MNR (10 litters) and 25 controls (5 litters, 3–8 pups/litter) one week before tissue collections at 6 months by intraperitoneal glucose tolerance testing (IP-GTT). Briefly, fasted mice (4 h) (at 8:00 am) were injected with glucose 2 g/kg body weight and blood glucose was measured via tail vein every 30 min for 2 h using a Freestyle Lite Glucometer (Abbott, Alameda, CA, USA). Hepatic glucose output was measured with a pyruvate challenge test (5 pups per group, 1 pup/litter). Pyruvate was injected (IP) into mice fasted for 4 h (at 8:00 am) at 2 g/kg body weight and blood glucose was measured via tail vein every 30 min for 2 h.

Random blood glucose

Once a month from 1 to 6 months, random blood glucose (nonfasted) was measured. Blood samples were collected at 10:30 am via tail vein prick and measured using a Freestyle Lite Glucometer.

Intrahepatic portal vein insulin challenge test

At 7 months old, mice were fasted for 4 h and anaesthetized with isoflurane. Using a mid-line incision, intestines were moved to expose the hepatic portal vein and 2 U/kg of insulin (Humalin R, Eli Lilly Canada Inc., Toronto, ON, Canada) (control and MNR $n = 5$, 1 pup/litter) or saline (control and MNR $n = 3$, 1 pup/litter) was injected. After 1 min and 15 s, a piece of right medial liver lobe, a quadriceps femoris muscle and perigonadal fat were frozen in liquid nitrogen. Samples were stored in -80°C until western blot analysis.²⁴

Protein isolations

The experiment was conducted with 50 mg of frozen liver and muscle and 100 mg of adipose tissue that were pulverized (Medium Bessman Tissue Pulverizer, Spectrum Laboratories, Rancho Dominguez, CA, USA) and then homogenized for 30 s with the Brinkmann Homogenizer Polytron[®] 3000 in 1 × cell lysis buffer (9803S, Cell Signaling Technology[®] Inc., Danvers, MA, USA) and protease inhibitor cocktail 1, 2 and 3 (P1860, P5726 and P0044, Sigma-Aldrich, St Louis, MO, USA). The homogenate was then sonicated (F550 Sonic Dismemberator, Fisher Scientific, Markham, ON, Canada) and rocked for 40 min

at 4°C. Samples were spun at 12,000 *g* for 10 min at 4°C and the supernatant was collected with care, not to collect the lipid layer, and stored at -20°C until analysis. Protein quantification was done with the Bio-Rad protein assay dye according to the manufacturer's protocol using bovine serum albumin (BSA) standards (Cat #500-0006, Bio-Rad Laboratories Inc., Des Plaines, IL, USA).

Western blotting

Protein samples were boiled in sodium dodecyl sulfate (SDS) sample loading buffer with dithiothreitol at 75°C for 5 min and then incubated on ice for 5 min. A 10 µg of protein was loaded and run on a 10% SDS polyacrylamide gel. Proteins were transferred to a polyvinylidene difluoride membrane using the Trans-Blot® Turbo™ Transfer System and Trans-Blot® Turbo™ RTA Midi Transfer Kit (170-4273, Bio-Rad Laboratories Inc.) according to the manufacturer's protocol.

Once the transfer was completed, membranes were dipped in methanol and blocked with 5% milk in TBST [tris buffered saline (TBS) + 1% tween]. Blocking buffer was rinsed from the membrane for 5 min (3 ×) with TBST at room temperature (RT). Primary antibody for pAkt (#4000, rabbit mAb, Cell Signaling Technology® Inc.) was diluted 1: 1000 into 5% BSA in TBST and incubated with blots shaking at overnight at 4°C. Membranes were rinsed with TBST 3 × (5 min) at RT and then incubated with secondary rabbit IgG (#170-6515, Bio-Rad Laboratories Inc.) diluted 1: 10 000 in 5% milk in TBST for 1 h at RT. After rinsing the secondary antibody, Clarity™ Western ECL substrate (#170-5061, Bio-Rad Laboratories Inc.) and a VersaDoc Imaging System (Bio-Rad Laboratories Inc.) were used to capture an image of the gel. Membranes were then rinsed and stripped with 0.5 M NaOH, rinsed in TBS (3 × 5 min at RT) and reprobed with Akt (#9272, Bio-Rad Laboratories Inc.), according to the protocol mentioned earlier.

Quantification of blots was done using Imagemag software® version 5.1 BETA (Bio-Rad Laboratories Inc.) and quantities were compared using pAkt to total Akt ratios.

Serum peptide markers for obesity and diabetes

Blood samples collected by intracardiac puncture at 1 month (control: *n* = 7, MNR: *n* = 18, 4 control and 10 MNR litters, 1–2 pups/litter) and 6 months (5 control and 10 MNR litters, 3–7 pups/litter, MNR: *n* = 52, control *n* = 30) in nonfasted mice (8:30–9:30 am) were used in this multiplex assay. At the time of collection, protease inhibitors, 10 mM DPP-IV (EMD Millipore, DPP4-010) and 1.3% aprotinin inhibitor (EMD Millipore, 616399), both in 0.9% sterile saline, were added to the blood samples (10 µl/ml). Samples were allowed to clot for 45 min and spun at twice at 12,000 *g* for 10 min to obtain serum. All serum samples were stored at -80°C. Ghrelin, leptin, insulin, glucagon, PAI-1, GLP, GIP and resistin were assayed on the Bio-Rad Multiplex at Lawson's Multiplex Facility, London, ON, using the Bio-Plex Pro Mouse Diabetes Assay (171F7001M, Bio-Rad Laboratories Inc.). Serum C-peptide was also quantified according to the manufacturer's protocol (Mouse C-Peptide ELISA, 80-CPTMA-E1, Alpco, Salem, NH).

Lipid quantification in serum and liver

An aliquot of nonfasted serum as described earlier was used for cholesterol (Cholesterol Quantification Kit MAK043, Sigma-Aldrich®, St Louis, MO, USA) and triglyceride quantification

(Triglyceride Quantification Assay Kit ab65336, Abcam®, Cambridge, MA, USA). Liver cholesterol was also isolated from nonfasted liver tissue and quantified according to the manufacturer's protocol. Measurements were taken from 5 control and 10 MNR litters, 3–6 pups/litter.

Statistical analysis

Data are presented as mean ± SEM. For glucose tolerance testing, pyruvate challenge tests and random blood glucose measurements, a repeated measure ANOVA and Bonferroni *post-hoc* correction were used. For all other analyses, Mann–Whitney test was used to compare means where samples sizes were >8, and an unpaired *t*-test with ≤8 samples. Means are considered significantly different if *P* < 0.05. All statistics and graphs were produced in Prism software (GraphPad®, version 5.2), or the R console (version 3.3.2) with ggplot2® (version 2.2.1) and Sciplot® (version 1.1–0) software packages.

Results

Maternal total-calorie nutrient restriction (MNR) resulted in a 13% reduction (*P* = 0.001) of the fetal weight at E18.5 in males (Fig. 1a), and 14% reduction (*P* = 1.5 × 10⁻⁸) in birth weight of males (Fig. 1b), relative to controls. Body weights are not significantly different in MNR offspring between day 3 of life and 6 months (Fig. 1c), except at 1 month (*P* = 0.005) when controls (23.5 ± 0.4) were heavier than MNR (22.3 ± 0.3). This protocol demonstrated a reproducible FGR from moderate MNR.

In addition to body weight, organ weights were taken at each dissection. Fetal liver-to-body weight ratio was 1.4-fold reduced in MNR offspring (0.05 ± 0.003) relative to controls (0.07 ± 0.002, *P* < 0.0001), indicating a disproportionate effect of maternal nutrition on the liver (Fig. 2a). All other organs were proportionally reduced with body weight. At 1 month, liver weight relative to body weight was similar in MNR offspring (*P* = 0.09) (Fig. 2b), but decreased again by 6 months (*P* = 0.03). These data suggest that the disproportionate impact on the liver growth is impacted by fetal nutrition and the effects extend into adulthood (Fig. 2c).

Whole body glucose metabolism and hepatic glucose output was assessed with IP-GTT and pyruvate challenge tests, respectively. Differences in glucose tolerance were detected with the area under the curve (AUC) in 6-month-old MNR offspring (1553 ± 57) relative to controls (1323 ± 68, *P* = 0.045) (Fig. 3a and b). Fasting blood glucose at the time of GTT was also relatively higher in MNR offspring (*P* = 0.3) (Fig. 3a); however, blood glucose at all time points and the AUC (Supplementary Fig. 1S) for pyruvate challenge testing indicated similar hepatic glucose output in MNR offspring relative to controls. Therefore, moderate differences in glucose metabolism were detected in the MNR offspring.

Random blood glucose was measured once a month until 6 months of age. MNR offspring blood glucose level was similar to controls each month (*P*-value = 0.4) (Supplementary Fig. 2S), and the AUC did not differ (*P*-value = 0.3) (Supplementary Fig. 2S).

Serum peptide markers for diabetes and obesity were also assayed at 1 and 6 months of age. At 1 month, PAI-1 was 1.9-fold higher in MNR offspring relative to controls (*P* = 0.04) [Fig. 4a(i)]. PAI-1 remained 1.5-fold higher in 6-month-old MNR offspring (*P* = 0.04) [Fig. 4b (i)]. Increased resistin was also detected at this time in MNR offspring (189.8 ± 9.8 ng/ml) relative to controls (171.2 ± 14.5 ng/ml, *P* = 0.04) [Fig. 4b(v)]. C-peptide,




Fig. 1. (a) Body weights of offspring of MNR and control from E18.5 until 6 months. At E18.5, fetal weights of MNR ($n = 15$) were 13% smaller than controls ($n = 20$) ($P = 0.001$) (3 Control and 3 MNR litters, 3–8 pups/litter). (b) Birth weight of MNR offspring remained 14% smaller ($P = 1.5 \times 10^{-8}$) (control: $n = 39$ from 5 litters, MNR: $n = 58$ from 10 litters, 4–9 pups/litter). Except at 1 month when controls were heavier than MNR ($P = 0.005$), no differences were observed in weight from day 3 to 6 months. (c) From birth to 1 month (control: $n = 37$, MNR: $n = 52$, 5 control and 10 MNR litters, 3–11 pups/litter), and from 2 to 6 months (control: $n = 27$, MNR: $n = 36$) (5 control and 10 MNR litters, 3–9 pups/litter). Growth curve data is plotted as mean \pm SEM. Asterisk represents significance ($P < 0.05$) evaluated by Mann-Whitney test.




Fig. 2. Organ weights of male offspring at E18.5, 1 month and 6 months. (a) At E18.5, liver weight was reduced in MNR offspring relative to their body weight ($P < 0.0001$) (control: $n = 20$, MNR: $n = 15$, 3 control and 3 MNR litters, 3–8 pups/litter). (b) Relative to body weight, the liver of MNR offspring was similar to controls at 1 month ($P = 0.09$) (control: $n = 12$ from 5 litters, MNR: $n = 19$ from 9 litters, 1–3 pups/litter), and significantly lower at 6 months ($P = 0.03$) (control: $n = 24$, MNR: $n = 30$, 5 control and 10 MNR litters, 3–8 pups/litter). (c) Fetal data are mean \pm SEM (a). Asterisk represents $P < 0.05$ evaluated by Mann-Whitney test.

insulin-to-glucagon ratio, GIP, GLP, ghrelin and leptin did not differ based on maternal nutrition at 1 and 6 months (Fig. 4).

In addition to peptide markers, lipids were measured in serum and the right lateral liver lobe at 6 months. No differences were observed between control and nutrient-restricted offspring in serum triglycerides ($P = 0.6$) (Supplementary Fig. 3S Ai) or total

cholesterol ($P = 0.7$) (Supplementary Fig. 3S Aii). Moreover, control and MNR offspring had similar liver cholesterol ($P = 0.07$) (Supplementary Fig. 3S B).

Although some differences were detected in whole-body glucose metabolism, a hepatic portal vein insulin challenge was used to examine tissue-specific insulin sensitivity. Relative to controls, the




Fig. 3. IP-GTT of male offspring at 6 months. (a) Blood glucose at 90 and 120 min was significantly higher in MNR offspring ($n = 36$, 10 litters) relative to controls ($n = 25$, 5 litters, 3–8 pups/litter). The AUC was also significantly higher ($P = 0.045$) (b). Glucose tolerance data (a) are mean \pm SEM. Asterisk represents $P < 0.05$ evaluated by Bonferroni *post hoc* (a) or Mann–Whitney test (b).




Fig. 4. Serum peptide markers for obesity and diabetes at (a) 1 month (control: $n = 7$, MNR: $n = 18$; 4 control and 10 MNR litters, 1–2 pups/litter), and (b) 6 months of age (5 control and 10 MNR litters, 3–7 pups/litter; MNR: $n = 52$, control: $n = 30$). PAI-1 was increased in MNR offspring at 1 month ($P = 0.04$) and 6 months ($P = 0.04$), (i) and resistin at 6 months ($P = 0.04$) (v). Ghrelin (ii), GLP-1 (iii), leptin (iv), GIP (vi), insulin-to-glucagon ratios (vii), and C-peptide (viii) levels were similar between control and MNR offspring. Asterisk represents $P < 0.05$ evaluated by Mann–Whitney test.

ratio of pAkt-to-Akt was 2.5-fold increased in the liver of MNR offspring in response to the insulin injection ($P = 0.004$) (Fig. 5a). Although not significant, insulin-stimulated phosphorylation of

Akt in adipose tissue (Fig. 5b) and skeletal muscle (Fig. 5c) was relatively increased by 1.5- and 1.6-fold in the nutrient-restricted offspring, respectively ($P = 0.08$ and 0.05 , respectively).




Fig. 5. Insulin signaling as determined by phosphorylation of Akt in response to an insulin bolus in male MNR and control offspring at 7 months of age. pAkt relative to Akt after an insulin dose was increased in MNR offspring relative to controls in the right medial liver lobe ($P=0.004$) (a). pAkt-to-Akt ratios were relatively increased in MNR adipose and (c) skeletal muscle of insulin injected mice, but this change was not significant ($P=0.08$ and 0.05 , respectively) (b). Saline injected $n=3$ and insulin injected $n=5$ for both maternal nutrition groups, 5 control and 8 MNR litters, 1 pup/litter. Means with asterisks are significantly different ($P<0.05$, unpaired t -test).

Although changes in glucose metabolism at the tissue-level occurred to some extent in all MNR offspring, there was a variation in the degree of response to fetal nutrient restriction on

adult glucose metabolism. At 6 months, 19% of the offspring were less tolerant to glucose (Fig. 6a), with a higher AUC than any controls (Fig. 6b). The remaining MNR have AUCs that were




Fig. 6. Glucose tolerance tests for all offspring of MNR and control at 6 months. (a) Individual glucose tolerance tests for controls demonstrate the range of responses. (b) 7 MNR (black) had AUC greater than other MNR or controls (grey), which are referred to as intolerant MNR (5 control and 10 MNR litters, 3–8 pups/litter).

similar to those of controls. Although differences in blood glucose regulation were evident after the glucose bolus, random blood glucose measurements from 1 to 6 months were not different in less-tolerant MNR (data not shown).

Serum peptide analysis in the less-tolerant mice showed increased insulin ($P=0.006$), leptin ($P=0.001$) (Fig. 7g) and resistin ($P=0.005$) (Fig. 7f) levels, with no changes in other assayed peptide markers (Fig. 7h).

Discussion

In humans, FGR can be symmetric or asymmetric. Symmetric FGR, meaning all organs including the brain are reduced proportionately to body weight, is often associated with an insult of early onset.²⁵ Conversely, asymmetric FGR is more common in late-onset growth restriction, and is where less essential organs are reduced but vital organs such as the brain and the heart are spared.²⁵ Similar to humans, maternal undernutrition in mice resulted in an asymmetric growth restriction, with the most significant impact on the liver.²⁶ The thymus and spleen are also smaller relative to body weight in SGA autopsies,²⁶ but were not measured in this study. Although reduced again at 6 months, 1-month relative liver weights were similar to controls, which might indicate differences in growth rate of the liver in MNR offspring. MNR in guinea pigs do not result in reduced liver-to-fetal weight ratios, but the fractional growth rate is increased near term.⁹ These data suggest that liver growth may be more sensitive than other organs to fetal undernutrition.

The variation in response to fetal nutrient restriction in our model was similar to the variations observed in the human FGR populations. Some individuals with extremely low birth weights (ELBW) although having three times greater risk will not develop glucose intolerance and type 2 diabetes.³ In adults, genetic variation and differential expression contribute to obesity and metabolic disorder resistant and/or susceptible populations of nonhuman primates²⁷ and rodent models^{28,29} in response to nutritional stress, such as a high fat diet. Offspring in models exposed to maternal high fat diet or nutrient restriction have metabolic,^{7,30} gut microbe³¹ and epigenomic changes,³² but to our knowledge none of these models focus on changes specific to susceptible and resistant populations in these offspring. Having glucose tolerant and intolerant offspring resulting from MNR

allow future studies to examine differences in glucose metabolism within the MNR populations, as well as adaptations that occur in all offspring exposed to nutrient restriction *in utero*.

In this study, normal glucose tolerance observed in 81% of the offspring might have been due to increased insulin sensitivity as adults, despite having smaller livers. Increased insulin sensitivity in peripheral tissues in response to growth restriction is thought to occur at birth and has been shown to persist a few weeks postnatally in lambs exposed to hypoxia-induced placental insufficiency.³³ To our knowledge, evidence of insulin desensitization^{6,7} has been observed rather than persistence of the increased sensitivity into adulthood. Intolerant MNR offspring, which had higher serum insulin and relatively higher insulin-to-glucagon ratios, might have insulin resistance that was not detectable during insulin sensitivity test due to technical limitations in sample size. Studies comparing liver responses in intolerant and tolerant MNR mice may provide insight into why some offspring are able to adapt to the postnatal parameters assessed in this study.

Despite having similar body weight at 6 months, PAI-1 and resistin were increased in MNR offspring. Both are produced primarily in adipose tissue in mice^{16,17} and are associated with obesity and meta-inflammation.^{16,17,34} Individuals born SGA with the highest body mass index (BMI) have reduced insulin sensitivity,³⁵ however, total fat pads are increased in female adult rats that experienced 50% nutrient restriction throughout gestation with similar adult body weights to controls.³⁶ ELBW babies also have similar BMI with increased body fat mass,³ suggesting that an increased markers for obesity and meta-inflammation may indicate reduced lean mass in response to fetal undernutrition in male offspring, despite no differences in adult body weight. MNR offspring with glucose intolerance had higher serum resistin, suggesting increased fat mass may contribute to the development of glucose intolerance.

Leptin was also increased in offspring with less glucose tolerance. Serum leptin is correlated with body fat mass in humans³⁷ and mice,³⁸ supporting the notion of differences in body composition. This marker could also indicate a decreased appetite in MNR offspring, particularly in those with glucose intolerance. Alternatively, leptin resistance occurs in diet-induced obesity, in part, via $A\beta_{1-42}$ -induced basal STAT3 phosphorylation in the hypothalamus,³⁹ indicating that the increased leptin in offspring with impaired glucose homeostasis may be a sign of leptin




Fig. 7. Serum peptide markers for obesity and diabetes in male MNR and control offspring at 6 months old according to glucose tolerance. Resistin ($P=0.005$) (d) and leptin ($P=0.001$) (e) were significantly increased in MNR offspring that are glucose intolerant. PAI-1 (a), Ghrelin (b), GLP-1(c), GIP(f), insulin-to-glucagon (g) and c-peptide (h) were not significantly different in less tolerant MNR. Control $n=30$ (5 litters, 3–8 pups/litter), tolerant MNR $n=43$ (10 litters, 3–8 pups/litter), intolerant MNR $n=7$ (5 litters, 1–2 pups/litter). Asterisk represents $P < 0.05$ by Mann-Whitney test.

resistance. Leptin may also play a role in regulation of energy expenditure³⁹ and behavior changes, such as food preference and total activity, which are also altered in adulthood following IUGR or undernourishment during development.^{40,41} Increased serum leptin levels suggest that behaviors and body mass composition may be altered in all or a portion of the MNR offspring and require further investigation.

Moderate nutrient restriction throughout gestation resulted in changes to the liver and adipose tissue, including decreased fetal liver weight, increased adult insulin sensitivity in the liver, and

increased serum markers for body mass composition and meta-inflammation. Liver is well known to be important in regulation of lipid and glucose metabolism and a key tissue in the development of type 2 diabetes. Although it is possible that some adaptations occur in all offspring exposed to MNR, additional physiologic and biochemical changes in the liver and other organs may be specific to those that develop glucose intolerance. This model provides an opportunity to investigate the tissue-specific molecular mechanism underlying changes between control and MNR offspring, as well as between more affected MNR and those similar to the controls, and

represents the diversity in human FGR populations and their susceptibilities toward type 2 diabetes.^{3,4}

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/S2040174418000983>

Acknowledgements. Serum peptide analysis was run by Ms. Shannon Seney at the Screening Lab for Immune Disorders, at the Lawson Health Research Institute, London, Ontario. Dr. Edith Arany for providing guidance for the method of insulin sensitivity testing. I would also like to thank members of my Graduate Advisory Committee, Dr. Greg Gloor, Dr. Nathalie Berube, Dr. Melissa Mann, and Dr. Daniel Hardy, for their guidance and support throughout the study.

Financial support. Grant support for these studies were provided by the Canadian Institutes of Health Research (V.K.M.H., Grant 111024), Douglas and Vivian Bocking Chair in Fetal and Newborn Growth from St Joseph's Health Care Foundation (V.K.M.H. and B.R.) and the Children's Health Foundation (V.K.M.H.).

Conflicts of interest. The authors have no conflicts of interest to disclose.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guides on the care and use of mice and has been approved by the Council on Animal Care at the University of Western Ontario (2017-033).

References

- Han VK, Seferovic MD, Albion CD, Gupta MB. Intrauterine growth restriction: intervention strategies. In *Neonatology* (eds. Buonocore G, Bracci R, Weindling M), 2012; pp. 89–93. Springer: Milano.
- Garite TJ, Clark R, Thorp JA. Intrauterine growth restriction increases morbidity and mortality among premature neonates. *Am J Obstet Gynecol*. 2004; 191, 481–487.
- Morrison KM, Ramsingh L, Gunn E, *et al.* Cardiometabolic health in adults born premature with extremely low birth weight. *Pediatrics*. 2016; 138, e20160515.
- Würtz P, Wang Q, Niironen M, *et al.* Metabolic signatures of birthweight in 18288 adolescents and adults. *Int J Epidemiol*. 2016; 45, 1539–1550.
- Hales C, Barker D. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia*. 1992; 35, 595–601.
- Oliveira D, Cezar J, Gomes RM, *et al.* Protein restriction during the last third of pregnancy malprograms the neuroendocrine axes to induce metabolic syndrome in adult male rat offspring. *Endocrinology*. 2016; 157, 1799–1812.
- Xiao D, Kou H, Zhang L, Guo Y, Wang H. Prenatal food restriction with postweaning high-fat diet alters glucose metabolic function in adult rat offspring. *Arch Med Res*. 2017; 48, 35–45.
- Lee S, You Y-A, Kwon EJ, *et al.* Maternal food restriction during pregnancy and lactation adversely affect hepatic growth and lipid metabolism in three-week-old rat offspring. *Int J Mol Sci*. 2016; 17, E2115.
- Nevin CL, Formosa E, Maki Y, *et al.* Maternal nutrient restriction in guinea pigs as an animal model for studying growth restricted offspring with post-natal catch-up growth. *Am J Physiol Regul Integr Comp Physiol*. 2018; 314, R647–R654.
- Wang J, Cao M, Zhuo Y, *et al.* Catch-up growth following food restriction exacerbates adulthood glucose intolerance in pigs exposed to intrauterine undernutrition. *Nutrition*. 2016; 32, 1275–1284.
- Liu Y, Ma C, Li H, *et al.* Effects of intrauterine growth restriction during late pregnancy on the cell apoptosis and related gene expression in ovine fetal liver. *Theriogenology*. 2017; 90, 204–209.
- Vomhof-DeKrey E, Darland D, Ghribi O, *et al.* Maternal low protein diet leads to placental angiogenic compensation via dysregulated M1/M2 macrophages and TNF α expression in Sprague-Dawley rats. *J Reprod Immunol*. 2016; 118, 9–17.
- Palha AM, Pereira SS, Costa MM, *et al.* Differential GIP/GLP-1 intestinal cell distribution in diabetics' yields distinctive rearrangements depending on Roux-en-Y biliopancreatic limb length. *J Cell Biochem*. 2018; 119, 7506–7514.
- Nauck MA, Meier JJ. Incretin hormones: their role in health and disease. *Diabetes Obes. Metab*. 2018; 20(Suppl 1), 5–21.
- Leighton E, Sainsbury CA, Jones GC. A practical review of c-peptide testing in diabetes. *Diabetes Ther*. 2017; 8, 475–487.
- Qatanani M, Szwegold NR, Greaves DR, Ahima RS, Lazar MA. Macrophage-derived human resistin exacerbates adipose tissue inflammation and insulin resistance in mice. *J Clin Invest*. 2009; 119, 531–539.
- Kaji H. Adipose tissue-derived plasminogen activator inhibitor-1 function and regulation. *Compr Physiol*. 2016; 6, 1873–1896.
- Bewick GA, Kent A, Campbell D, *et al.* Mice with hyperghrelinemia are hyperphagic and glucose intolerant and have reduced leptin sensitivity. *Diabetes*. 2009; 58, 840–846.
- Kalra SP, Ueno N, Kalra PS. Stimulation of appetite by ghrelin is regulated by leptin restraint: peripheral and central sites of action. *J Nutr*. 2005; 135, 1331–1335.
- Lecoutre S, Marouze L, Drougard A, *et al.* Maternal undernutrition programs the apelinergic system of adipose tissue in adult male rat offspring. *J Dev Orig Health Dis*. 2017; 8, 3–7.
- Simchen MJ, Weisz B, Zilberberg E, *et al.* Male disadvantage for neonatal complications of term infants, especially in small-for-gestational age neonates. *J Matern Fetal Neonatal Med*. 2014; 27, 839–843.
- Delahaye F, Wijetunga NA, Heo HJ, *et al.* Sexual dimorphism in epigenomic responses of stem cells to extreme fetal growth. *Nat Commun*. 2014; 5, 5187.
- Pae M, Baek Y, Lee S, Wu D. Loss of ovarian function in association with a high-fat diet promotes insulin resistance and disturbs adipose tissue immune homeostasis. *J Nutr Biochem*. 2018; 57, 93–102.
- Chamson-Reig A, Thyssen SM, Hill DJ, Arany E. Exposure of the pregnant rat to low protein diet causes impaired glucose homeostasis in the young adult offspring by different mechanisms in males and females. *Exp Biol Med*. 2009; 234, 1425–1436.
- Nardoza LMM, Araujo Júnior E, Barbosa MM, *et al.* Fetal growth restriction: current knowledge to the general Obs/Gyn. *Arch Gynecol Obstet*. 2012; 286, 1–13.
- Man J, Hutchinson JC, Ashworth M, *et al.* Organ weights and ratios for postmortem identification of fetal growth restriction: utility and confounding factors. *Ultrasound Obstet Gynecol Off J Int Soc*. 2016; 48, 585–590.
- Harris RA, Alcott CE, Sullivan EL, *et al.* Genomic variants associated with resistance to high fat diet induced obesity in a primate model. *Sci Rep*. 2016; 6, 36123.
- Weingarten A, Turchetti L, Krohn K, *et al.* Novel genes on rat chromosome 10 are linked to body fat mass, preadipocyte number and adipocyte size. *Int J Obes*. 2016; 40, 1832–1840.
- Chen K, Jih A, Osborn O, *et al.* Distinct gene signatures predict insulin resistance in young mice with high fat diet-induced obesity. *Physiol Genomics*. 2018; 50, 144–157.
- Zhu W-F, Zhu J-F, Liang L, Shen Z, Wang Y-M. Maternal undernutrition leads to elevated hepatic triglycerides in male rat offspring due to increased expression of lipoprotein lipase. *Mol Med Rep*. 2016; 13, 4487–4493.
- Ma J, Prince AL, Bader D, *et al.* High-fat maternal diet during pregnancy persistently alters the offspring microbiome in a primate model. *Nat Commun*. 2014; 5, 3889.
- Heo HJ, Tozour JN, Delahaye F, *et al.* Advanced aging phenotype is revealed by epigenetic modifications in rat liver after in utero malnutrition. *Aging Cell*. 2016; 15, 964–972.
- Camacho LE, Chen X, Hay WW, Limesand SW. Enhanced insulin secretion and insulin sensitivity in young lambs with placental insufficiency-induced intrauterine growth restriction. *Am J Physiol Regul Integr Comp Physiol*. 2017; 313, R101–R109.
- Kim S-H, Park H-S, Hong MJ, *et al.* Tongqiaohuoxue decoction ameliorates obesity-induced inflammation and the prothrombotic state by regulating adiponectin and plasminogen activator inhibitor-1. *J Ethnopharmacol*. 2016; 192, 201–209.
- Veening MA, Weissenbruch V, Waal MM, D de AH. Glucose tolerance, insulin sensitivity, and insulin secretion in children born small for gestational age. *J Clin Endocrinol Metab*. 2002; 87, 4657–4661.

36. de Souza AP, Pedroso AP, Watanabe RLH, *et al.* Gender-specific effects of intrauterine growth restriction on the adipose tissue of adult rats: a proteomic approach. *Proteome Sci.* 2015; 13, 32.
37. Soininen S, Sidoroff V, Lindi V, *et al.* Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6–8years of age – The Physical Activity and Nutrition in Children (PANIC) study. *Bone.* 2018; 108, 106–114.
38. Xiao X, Sun Q, Kim Y, *et al.* Exposure to permethrin promotes high fat diet-induced weight gain and insulin resistance in male C57BL/6J mice. *Food Chem Toxicol.* 2018; 111, 405–416.
39. Meakin PJ, Jalicy SM, Montagut G, *et al.* Bace1-dependent amyloid processing regulates hypothalamic leptin sensitivity in obese mice. *Sci Rep.* 2018; 8, 55.
40. Silveira PP, Pokhvisneva I, Gaudreau H, *et al.* Fetal growth interacts with multilocus genetic score reflecting dopamine signaling capacity to predict spontaneous sugar intake in children. *Appetite.* 2018; 120, 596–601.
41. Lira LA, Almeida LCA, da Silva AAM, *et al.* Perinatal undernutrition increases meal size and neuronal activation of the nucleus of the solitary tract in response to feeding stimulation in adult rats. *Int J Dev Neurosci.* 2014; 38, 23–29.