

Single-time Markovianized spectral closure in fluid turbulence

Takuya Kitamura[†]

Graduate School of Engineering, Nagasaki University, Nagasaki 852-8521, Japan

(Received 8 March 2020; revised 23 April 2020; accepted 18 May 2020)

Kaneda's (*J. Fluid Mech.*, vol. 107, 1981, pp. 131–145) Lagrangian renormalized approximation was extended to single-time spectral closure under two assumptions: (i) Markovianization and (ii) the Lagrangian velocity response function is expressed by $G(k, \tau) = \exp(-C_1(k)\tau - C_2(k)\tau^2/2)$. The unknown functions $C_1(k)$ and $C_2(k)$ are theoretically derived to be consistent with the exact short-time behaviour of $G(k, \tau)$ and the asymptotic short-time behaviour of assumed exponential form of $G(k, \tau)$, i.e. the present closure is derived from the Navier–Stokes equation without introduction of any adjustable parameters and it can calculate the statistical quantities by theory. The results show that the present closure has good agreement with direct numerical simulation for single- and two-point statistics.

Key words: turbulence theory, isotropic turbulence, homogeneous turbulence

1. Introduction

Turbulence has a stochastic nature, so that the randomness and complexity of turbulence make theoretical analysis difficult. However, such randomness may permit the statistical description of turbulence, and it is believed that there is a certain kind of universality at sufficiently small scales in fully developed turbulence away from boundaries (Kolmogorov 1941*a,b*). A striking feature of such universality is the existence of exact statistical laws for homogeneous isotropic turbulence at sufficiently high Reynolds numbers. Starting with Kraichnan (1959), much effort has been dedicated to deriving the statistical theory of turbulence using various field-theoretic approaches from first principles such as the Navier–Stokes equation (Leslie 1973; Monin & Yaglom 1975; McComb 1989, 2014). The goal is to accurately describe and predict statistical quantities of turbulence while maintaining a strong connection to the underlying dynamics of the Navier–Stokes equation. However, it has been long known that the equations for any statistical quantities are never closed because of the nonlinearity of the Navier–Stokes equation. The so-called closure problem has remained as a fundamental unsolved problem of fluid mechanics despite much effort by many researchers (Leslie 1973; Monin & Yaglom 1975; Davidson 2004; McComb 1989, 2014). The closure related to the present study belongs to the so-called spectral closure theory that postulates a dynamic equation for the energy transfer function and

[†] Email address for correspondence: t.kitamura@nagasaki-u.ac.jp

its spirit is to address in theory the characteristics of different scales that constitute a turbulent field. A major concern of the spectral closure theory of turbulence is a derivation of the second-order moments such as spectrum from the Navier–Stokes equation without any adjustable parameters.

As one of the great achievements in the statistical theory of turbulence, the direct interaction approximation (DIA) will be highlighted. In DIA, the two-point two-time correlation and the response function are introduced, and the nonlinear interactions are involved in the response function. This yields the non-negativity of the energy spectrum. Although DIA fails in the derivation of Kolmogorov's $-5/3$ law, DIA is attractive in the sense that there are several ways to derive the DIA equations, e.g. the weak dependence principle (Kraichnan 1959), self-consistent method (Herring 1965) and renormalized perturbation method (Kraichnan 1977). As a reason for the failure of DIA, the Eulerian coordinate system is unsuited to distinguishing between two possible effects such as uniform convection and distortion. To maintain Galilean invariance, the Lagrangian coordinate system is suited to distinguishing between the above effects, and Lagrangian two-point two-time spectral closures have succeeded in the derivation of Kolmogorov's $-5/3$ law (Kraichnan 1965; Kraichnan & Herring 1978; Kaneda 1981). Kolmogorov's $-5/3$ law has been observed in various turbulent motions, the interstellar medium (Armstrong, Rickett & Spangler 1995), solar wind (Matthaeus & Goldstein 1982) and atmosphere (Nastrom & Gage 1985; Tsuji 2004). Among various closures, abridged Lagrangian history DIA (ALHDIA), strain-based ALHDIA (SBALHDIA) and Lagrangian renormalized approximation (LRA) are representative closures which can provide Kolmogorov's $-5/3$ law through systematic ways without any adjustable parameters. As an exception to closures in the sense of the Eulerian viewpoint, local energy transfer is highlighted (McComb & Shanmugasundaram 1984). Local energy transfer is derived from a purely Eulerian viewpoint using a mapping function, whereas the result is compatible with Kolmogorov's $-5/3$ law. The Kolmogorov constant K_0 for an infinite Reynolds number is 1.77 and 1.72 for ALHDIA and LRA, respectively. On the other hand, K_0 for a finite Reynolds number is 1.78, 2.0, 1.67, 1.69 and 2.3 for ALHDIA, SBALHDIA, LRA, Markovianized LRA and local energy transfer, respectively (Herring & Kraichnan 1979; Gotoh, Kaneda & Bekki 1988; McComb & Shanmugasundaram 1984). Taking into account the facts that $K_0 = 1.62 \pm 0.17$ (Sreenivasan 1995) in experiments and $K_0 = 1.8 \pm 0.1$ for state-of-the-art high-resolution direct numerical simulation (DNS) (Ishihara *et al.* 2016), it is difficult to say which closure is superior. On the other hand, because of facts such as (i) the so-called DIA families have been mainly limited to the exploration of universal constants of extremely large-Reynolds-number turbulent flow since the closed equation is a complex integro-differential equation having memory effect and (ii) there has recently been a growing tendency to study the effect of finite Reynolds number on the Kolmogorov theory since the clear Kolmogorov's $-5/3$ law is not observed even for state-of-the-art high-resolution DNS (Ishihara *et al.* 2016), practical closure, e.g. eddy-damped quasi-normal Markovianization (EDQNM), has been used in a wide range of turbulence studies (Sagaut & Cambon 2008). One of the advantages of EDQNM is to deepen our understanding of the phenomenology of turbulence including Reynolds number dependency (Bos, Clark & Rubinstein 2007a; Bos, Shao & Bertoglio 2007b), decaying law regardless of Reynolds and Schmidt number (Lesieur & Ossia 2000; Briard *et al.* 2015) and the role of the nonlinear term in comparison with linear theory (Sagaut & Cambon 2008). On the other hand, the disadvantage of EDQNM is that the results quantitatively depend on

adjustable parameters (Bos & Fang 2015). Therefore, it is of great importance for the further development of turbulence theory to develop closure theory derived from the Navier–Stokes equation without the introduction of any adjustable parameters or taking phenomenology into account for closure itself. In this paper, we extend Kaneda’s LRA, which has been recognized as the most sophisticated closure (Sagaut & Cambon 2008), to two-point single-time spectral closure under a few assumptions and legitimate mathematical procedures.

The present paper is organized as follows. In § 2.1, LRA is briefly summarized. The mathematical procedure for the extension to single-time spectral closure is described in § 2.2. In § 3, the numerical methods for closure and DNS are described and the numerical results for single- and two-point statistics are shown in § 4. Discussion and conclusions are presented in § 5.

2. Spectral closure theory

2.1. Lagrangian renormalized approximation

Here, we briefly summarize the LRA (more details can be found in Kaneda (1981) or Kida & Goto (1997)). The equations of incompressible fluid are

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} + \mathbf{f}, \quad (2.1a)$$

$$\nabla \cdot \mathbf{u} = \nabla \cdot \mathbf{f} = 0, \quad (2.1b)$$

where \mathbf{u} is velocity, p is pressure normalized by density, \mathbf{f} is stirring force and ν is kinematic viscosity. Kaneda (1981) introduced the Lagrangian position function defined by

$$\psi(\mathbf{y}, t | \mathbf{x}, s) = \delta(\mathbf{y} - \mathbf{z}(\mathbf{x}, s | t)), \quad (2.2)$$

where δ is the three-dimensional Dirac delta function and $\mathbf{z}(\mathbf{x}, s | t)$ is the position of a fluid element at time t which passes \mathbf{x} at time s . The Lagrangian position function obeys

$$\left(\frac{\partial}{\partial t} + \mathbf{u}(\mathbf{y}, t) \cdot \nabla_{\mathbf{y}} \right) \psi(\mathbf{y}, t | \mathbf{x}, s) = 0, \quad (2.3a)$$

$$\psi(\mathbf{y}, t | \mathbf{x}, t) = \delta(\mathbf{x} - \mathbf{y}). \quad (2.3b)$$

Using the Lagrangian position function, the generalized velocity is defined as

$$\mathbf{v}(\mathbf{x}, s | t) = \int d\mathbf{y} \psi(\mathbf{y}, t | \mathbf{x}, s) \mathbf{u}(\mathbf{y}, t), \quad (2.4)$$

and from (2.1a), (2.1b) and (2.3a), it obeys

$$\begin{aligned} \frac{\partial \mathbf{v}(\mathbf{x}, s | t)}{\partial t} &= \int d\mathbf{y} \psi(\mathbf{y}, t | \mathbf{x}, s) \mathbf{f}(\mathbf{y}, t) + \nu \int d\mathbf{y} \psi(\mathbf{y}, t | \mathbf{x}, s) \Delta_{\mathbf{y}} \mathbf{u}(\mathbf{y}, t) \\ &\quad - \int d\mathbf{y} \psi(\mathbf{y}, t | \mathbf{x}, s) \nabla_{\mathbf{y}} p(\mathbf{y}, t). \end{aligned} \quad (2.5)$$

The closure problem considered here is to determine the subsequent statistical second-order moments such as $\langle \mathbf{u}(\mathbf{x}, t) \mathbf{u}(\mathbf{y}, t) \rangle$ and $\langle \mathbf{v}(\mathbf{x}, s | t) \mathbf{u}(\mathbf{y}, s) \rangle$, where $\langle \cdot \cdot \cdot \rangle$ represents an ensemble average. In LRA, the two-point two-time Lagrangian velocity correlation

\mathbf{Q} and Lagrangian velocity response function \mathbf{G} are chosen as representatives. Here \mathbf{Q} and \mathbf{G} are defined as

$$Q_{ij}(\mathbf{x}, t|\mathbf{x}', s) \equiv \langle P_{i\alpha}(\nabla_x)v_\alpha(\mathbf{x}, s|t)u_j(\mathbf{x}', s) \rangle, \quad t \geq s, \tag{2.6a}$$

$$G_{ij}(\mathbf{x}, t|\mathbf{x}', s) \equiv \left\langle \frac{P_{i\alpha}(\nabla_x)\delta v_\alpha(\mathbf{x}, s|t)}{\delta f_j(\mathbf{x}', s)} \right\rangle, \quad t \geq s, \tag{2.6b}$$

where δ represents the functional derivative. The operator \mathbf{P} plays a role of projecting the arbitrary vector field with respect to \mathbf{x} onto the solenoidal field. This operator plays an important role in LRA to eliminate the irrelevant terms in the expansion. In homogeneous turbulence, the closure theory is formulated in terms of the Fourier representation of (2.1a)–(2.5). After lengthy algebra (the details are shown in appendix A), we have

$$\left(\frac{\partial}{\partial t} + 2\nu k^2 \right) Q(k, t) = S(k, t) + Q_f(k, t), \tag{2.7a}$$

$$\left(\frac{\partial}{\partial t} + \nu k^2 + \eta(k, t, s) \right) Q(k, t, s) = 0, \quad t > s, \tag{2.7b}$$

$$\left(\frac{\partial}{\partial t} + \nu k^2 + \eta(k, t, s) \right) G(k, t, s) = 0, \quad t > s, \tag{2.7c}$$

$$G(k, t, t) = 1, \tag{2.7d}$$

with

$$S(k, t) = 2\pi \int_{t_0}^t ds \iint_{\Delta} dp dq kpqb_{kpq} Q(q, t, s) [G(k, t, s)Q(p, t, s) - Q(k, t, s)G(p, t, s)], \tag{2.8}$$

$$\eta(k, t, s) = \pi \iint_{\Delta} dp dq kpq(1 - y^2)(1 - z^2) \int_s^t ds' Q(q, t, s'), \tag{2.9}$$

where t_0 is the initial time; $Q(k, t) = Q(k, t, t) = E(k, t)/(2\pi k^2)$; $Q_f(k, t)$ is the forcing spectrum; $E(k)$ is the energy spectrum; \iint_{Δ} denotes the integration over all regions of the p – q plane such that the three wavenumber vectors \mathbf{k} , \mathbf{p} and \mathbf{q} form a triangle; x , y and z are the cosines of the angles opposite to k , p and q in this triangle; and b_{kpq} is the geometrical factor given by $b_{kpq} = p/k(xy + z^3)$. The LRA satisfies the fluctuation–dissipation relation $Q(k, t, s) = G(k, t, s)Q(k, s)$ for $t \geq s$. In LRA, an eddy damping term $\eta(k, t, s)$ comes from the Lagrangian acceleration of the pressure. In (2.7b) and (2.7c), contributions from the random force are neglected because they are not important for $t > s$ in the same way as Gotoh, Nagaki & Kaneda (2000) for passive scalar.

2.2. Extension to the single-time Markovianized LRA

When turbulence is quasi-stationary in the sense that the decay with respect to t of $Q(k, t, t)$ is much slower than that of $G(k, t, s)$ for s , then Markovianization $Q(k, t, s) = Q(k, t)G(k, t, s)$ is applicable (Gotoh *et al.* 1988; Gotoh & Kaneda 2000). Applying Markovianization in (2.7a), (2.7c) and (2.7d), we have the Markovianized LRA (MLRA) equations:

$$\left(\frac{\partial}{\partial t} + 2\nu k^2 \right) Q(k, t) = 2\pi \iint_{\Delta} dp dq kpqb_{kpq}\theta_{kpq}(t)Q(q, t)[Q(p, t) - Q(k, t)] + Q_f(k, t), \tag{2.10a}$$

$$\left(\frac{\partial}{\partial \tau} + vk^2 + \eta(k, t, \tau)\right) G(k, t, \tau) = 0, \quad \tau > 0, \tag{2.10b}$$

$$G(k, t, 0) = 1, \tag{2.10c}$$

in which

$$\theta_{kpq}(t) = \int_0^{t-t_0} ds G(k, t, s) G(p, t, s) G(q, t, s), \tag{2.11}$$

$$\eta(k, t, \tau) = \pi \iint_{\Delta} dp dq k p q (1 - y^2)(1 - z^2) Q(q, t) \int_0^{\tau} ds' G(q, t, s'), \tag{2.12}$$

where note that the evolution of G is with respect to time difference $\tau = t - s$. Using (2.10b), (2.10c) and (2.12), when $\tau \rightarrow 0$, the Lagrangian velocity response function is expanded as

$$G(k, t, \tau) = c_0(k, t) + c_1(k, t)\tau + \frac{c_2(k, t)}{2}\tau^2 + \dots, \tag{2.13}$$

where

$$c_0(k, t) = G(k, t, 0) = 1, \tag{2.14a}$$

$$c_1(k, t) = \left. \frac{\partial G(k, t, \tau)}{\partial \tau} \right|_{\tau=0} = -vk^2, \tag{2.14b}$$

$$c_2(k, t) = \left. \frac{\partial^2 G(k, t, \tau)}{\partial \tau^2} \right|_{\tau=0} = (vk^2)^2 - \mu(k, t), \tag{2.14c}$$

with

$$\mu(k, t) = \iint_{\Delta} dp dq \pi k p q (1 - y^2)(1 - z^2) Q(q, t) = 2\pi \int_0^{\infty} dq k q^3 Q(q, t) J\left(\frac{q}{k}\right), \tag{2.15}$$

where the latter expression is derived from the exact integral on p , and $J(x)$ is given by

$$J(x) = \left[\frac{(1-x)^4(1+x)^4}{32x^4} \log \frac{1+x}{|1-x|} - \frac{1+x^2}{48x^3} (3x^4 - 14x^2 + 3) \right]. \tag{2.16}$$

It is noted that $J(x) = J(1/x)$ and $J(x \rightarrow 0) = 8/15x + O(x^3)$. As mentioned by Kaneda (1993) and Gotoh & Kaneda (2000), the short-time behaviours of Q and G have important information about the statistical quantities. For the long-time behaviour of G , we assume

$$G(k, t, \tau) \simeq \exp \left[-C_1(k, t)\tau - \frac{C_2(k, t)}{2}\tau^2 \right] \equiv \mathcal{G}(k, t, \tau), \tag{2.17}$$

where unknown coefficients C_1 and C_2 are chosen to match asymptotic behaviours of (2.17) at $\tau \rightarrow 0$ with the exact short-time behaviours (2.14b) and (2.14c). We then obtain

$$C_1(k, t) = vk^2, \quad C_2(k, t) = \mu(k, t). \tag{2.18a,b}$$

This choice with respect to \mathcal{G} is equivalent to solving the following differential equation:

$$\left(\frac{\partial}{\partial \tau} + \nu k^2 + \mu(k, t)\tau\right) \mathcal{G}(k, t, \tau) = 0, \quad \tau > 0. \tag{2.19}$$

Using (2.17) and (2.18), $\theta_{kpq}(t)$ in the nonlinear term can be calculated in a straightforward way as follows:

$$\begin{aligned} \theta_{kpq}(t) &\simeq \int_0^{t-t_0} ds \mathcal{G}(k, t, s) \mathcal{G}(p, t, s) \mathcal{G}(q, t, s) \\ &= \sqrt{\frac{\pi}{2d_{kpq}(t)}} \exp\left(\frac{c_{kpq}^2}{2d_{kpq}(t)}\right) \\ &\quad \times \left[\operatorname{erf}\left(\frac{c_{kpq}}{\sqrt{2d_{kpq}(t)}} + \sqrt{\frac{d_{kpq}(t)}{2}}(t-t_0)\right) - \operatorname{erf}\left(\frac{c_{kpq}}{\sqrt{2d_{kpq}(t)}}\right) \right], \end{aligned} \tag{2.20}$$

with

$$c_{kpq} = \nu(k^2 + p^2 + q^2), \tag{2.21a}$$

$$d_{kpq} = \mu(k, t) + \mu(p, t) + \mu(q, t). \tag{2.21b}$$

For $t \gg t_0$ or $t_0 = -\infty$, $\theta_{kpq}(t)$ can be simplified as

$$\theta_{kpq}(t) = \sqrt{\frac{\pi}{2d_{kpq}(t)}} \exp\left(\frac{c_{kpq}^2}{2d_{kpq}(t)}\right) \operatorname{erfc}\left(\frac{c_{kpq}}{\sqrt{2d_{kpq}(t)}}\right). \tag{2.22}$$

In the numerical calculation, there are various rational approximations for the error function (Abramowitz & Stegun 1964). We, however, calculated it by means of a built-in function. For $x = c_{kpq}/\sqrt{(2d_{kpq}(t))} \rightarrow \infty$, the integrand is calculated using the asymptotic property of the error function (Gradshteyn & Ryzhik 2014) as follows:

$$\theta_{kpq}(t) = \sqrt{\frac{1}{2d_{kpq}}} \frac{1}{x} \sum_{k=0}^{\infty} (-1)^k \frac{(2k-1)!!}{(2x^2)^k}. \tag{2.23}$$

Thus, equations (2.20), (2.21a) and (2.21b) are used to calculate the nonlinear term in the single-time MLRA. The computational cost of the single-time MLRA by this approximation is of the same order as that of the EDQNM, test field model (Kraichnan 1971) and Lagrangian Markovianized field approximation (Bos & Bertoglio 2013). The present method is also promising for anisotropic turbulence. In EDQNM, an eddy damping term is derived phenomenologically on the dimensional ground and anisotropic effects are not incorporated in an eddy damping term. Thus, the single-time MLRA is more sophisticated compared to the other spectral closures since it can calculate the statistical quantities by theory without introduction of any adjustable parameters.

3. Numerical method

Hereinafter, the single-time MLRA is termed closure for the sake of simplicity. In closure, we solved (2.10a) with (2.22). The time derivative was estimated with a first-order forward time scheme. The viscous, energy transfer and forcing terms are estimated explicitly. We used $Q_F \sim \exp(-k^2/k_f^2)$ in this study and we have confirmed that the form of forcing spectrum is negligible for statistical quantities, where $k_f = 2.5$. The wavenumbers were discretized logarithmically, namely $k_i = k_{min} \times 2^{i-1/F}$, where $k_{min} = 1$, $F = 8$ and $i \in [1, \dots, N]$. Here N was chosen to satisfy $k_{max}\eta \geq 2$ at the statistical steady state. Statistical quantities were evaluated when a criterion $\sqrt{(K(t) - K(t-dt))^2}/K(t) \leq 1 \times 10^{-6}$ is satisfied, where dt is the time step and $K(t)$ is the turbulence kinetic energy (TKE) at time t .

In DNS, the spectral method was used with periodic boundary conditions of periods of 2π in each of the three Cartesian coordinate directions. For time integration, we used the fourth-order Runge–Kutta scheme and the so-called phase shift method was used for de-aliasing, in which the maximum wavenumber of the retained Fourier modes is about $\sqrt{2}/3N$, where N^3 is the number of grid points. In this study, we force the turbulence as

$$f(\mathbf{k}, t) = \frac{\epsilon(t) - (K(t) - K_\infty)/\tau_f}{2K_f(t)} \hat{\mathbf{u}}(\mathbf{k}, t)[H(k) - H(k - k_f)], \quad (3.1)$$

where K_∞ is the ideal TKE, τ_f is the time scale to control forcing, K_f is the TKE composed of wavenumbers lower than forced wavenumber $k_f = 2.5$ and H is the Heaviside step function. Here, the parameters were set as $K_\infty = 1/2$ and $\tau_f = 5dt = O(\tau_\eta)$, where τ_η is the Kolmogorov time scale. Under this condition, TKE is well controlled, e.g. the value is $1/2$ with standard deviation $O(10^{-7})$ for lowest Reynolds number DNS and with $O(10^{-5})$ for highest Reynolds number DNS. Here, $E(k, 0) \sim k^4 \exp(-k^2/k_L^2)$ with $k_L = 2.5$ was used for the initial energy spectrum in all DNSs. We started all runs with the same initial energy spectrum instead of using the final state of the run with the smaller N as the initial state of the new runs as done by Kaneda *et al.* (2003). After 5 eddy-turnover times based on the initial energy spectrum, we carried out DNSs a further 10 eddy-turnover times for calculating the single-point statistics. For two-point statistics, the statistics were calculated every eddy-turnover time, i.e. 10 data gatherings in total. In the present code, hybrid parallelization was implemented using the message-passing interface library and OpenMP.

4. Numerical results

Turbulence characteristics are summarized in table 1. Here, K , ϵ , L , λ and η are the TKE, TKE dissipation rate, integral length scale, Taylor microscale and Kolmogorov scale, respectively. These variables are, in spectral space, defined as $K = \int_0^\infty dkE(k)$, $\epsilon = 2\nu \int_0^\infty dk k^2 E(k)$, $L = 3\pi/(4K) \int_0^\infty dkE(k)/k$, $\lambda^2 = 10\nu K/\epsilon$ and $\eta = (\nu^3/\epsilon)^{1/4}$, respectively (Davidson 2004). Here Re_λ is the turbulence Reynolds number based on the Taylor microscale $Re_\lambda \equiv \sqrt{2K/3}\lambda/\nu$ and C_ϵ is the normalized TKE dissipation rate $C_\epsilon \equiv \epsilon L/(2K/3)^{3/2}$. Figure 1 shows the relation between C_ϵ and Re_λ obtained by closure and DNS together with DNS data of other researchers. The functional form (Doering & Foias 2002) $C_\epsilon = a[1 + \sqrt{1 + (b/Re_\lambda)^2}]$ with $a = 0.217$ and $b = 88.65$ agrees with the present results obtained by closure and DNS. The parameters were evaluated by means of the Levenberg–Marquardt least-squares method for closure results. The fitting curve is within the acceptable range of ± 3 standard deviations for DNS data.




FIGURE 1. Normalized TKE dissipation rate C_ϵ versus turbulence Reynolds number Re_λ with $\pm 3\sigma$ uncertainty error bars for present DNS, where σ is the standard deviation. Also included are DNS data from Jiménez *et al.* (1993), Cao, Chen & Doolen (1999), Gotoh, Fukayama & Nakano (2002) and Kaneda *et al.* (2003).

Method	N	K	ϵ	L	λ	η	Re_λ	C_ϵ
DNS	128	0.500	1.987×10^{-1}	1.973	1.563	2.589×10^{-1}	9.400	2.026
DNS	128	0.500	1.601×10^{-1}	1.827	1.421	2.016×10^{-1}	1.282×10	1.511
DNS	128	0.500	1.338×10^{-1}	1.575	1.099	1.254×10^{-1}	1.983×10	1.088
DNS	128	0.500	1.113×10^{-1}	1.371	8.491×10^{-1}	7.794×10^{-2}	3.064×10	0.791
DNS	128	0.500	9.779×10^{-2}	1.220	6.418×10^{-1}	4.790×10^{-2}	4.632×10	0.616
DNS	128	0.500	8.717×10^{-2}	1.153	4.800×10^{-1}	2.930×10^{-2}	6.929×10	0.521
DNS	128	0.500	8.648×10^{-2}	1.069	3.404×10^{-1}	1.745×10^{-2}	9.827×10	0.480
DNS	256	0.500	7.324×10^{-2}	1.165	2.192×10^{-1}	8.282×10^{-3}	1.808×10^2	0.441
DNS	512	0.500	7.678×10^{-2}	1.068	1.353×10^{-1}	4.116×10^{-3}	2.791×10^2	0.425
DNS	1024	0.500	7.873×10^{-2}	1.026	8.388×10^{-2}	2.031×10^{-3}	4.403×10^2	0.416
Closure	36	1.003	1.414	0.846	5.932×10^{-1}	9.771×10^{-2}	9.771	2.189
Closure	40	0.941	9.458×10^{-1}	0.797	5.215×10^{-1}	6.818×10^{-2}	1.510×10	1.517
Closure	47	0.892	6.109×10^{-1}	0.716	3.921×10^{-1}	3.718×10^{-2}	2.872×10	0.954
Closure	56	0.889	4.684×10^{-1}	0.642	2.543×10^{-1}	1.704×10^{-2}	5.748×10	0.659
Closure	67	0.921	4.188×10^{-1}	0.590	1.423×10^{-1}	6.572×10^{-3}	1.211×10^2	0.514
Closure	78	0.944	4.023×10^{-1}	0.564	7.736×10^{-2}	2.534×10^{-3}	2.407×10^2	0.454
Closure	90	0.964	4.013×10^{-1}	0.550	3.911×10^{-2}	8.959×10^{-4}	4.920×10^2	0.429
Closure	102	0.965	3.994×10^{-1}	0.545	1.960×10^{-2}	3.167×10^{-4}	9.890×10^2	0.421
Closure	114	0.966	4.009×10^{-1}	0.543	9.793×10^{-3}	1.120×10^{-4}	1.975×10^3	0.421
Closure	126	0.960	3.997×10^{-1}	0.543	4.887×10^{-3}	3.960×10^{-5}	3.934×10^3	0.423
Closure	138	0.953	3.973×10^{-1}	0.543	2.439×10^{-3}	1.400×10^{-5}	7.838×10^3	0.426
Closure	150	0.953	3.992×10^{-1}	0.544	1.217×10^{-3}	4.949×10^{-6}	1.562×10^4	0.429

TABLE 1. Turbulence characteristics in the numerical simulations.

Figure 2 shows the normalized energy spectra and compensated energy spectra. As shown in figure 2, the spectra collapse well in the inertial subrange and dissipation range for both DNS and closure. The differences between closure and DNS are




FIGURE 2. Normalized energy spectra for (a) DNS and (b) closure and compensated energy spectra for (c) DNS and (d) closure. Insets in (c) and (d) show the log-linear plots. The dashed grey line in (a) is a closure result for the highest Reynolds number. In (c), DNS data only for the top four highest Reynolds numbers are shown to emphasize the spectral bump.

mainly observed in the energy-containing range due to the different forcing function and the degree of spectral bump. However, the energy spectra obtained in closure qualitatively agree with those of DNS. As shown in figure 2(b), a slight bump is observed in the closure; however, its degree is suppressed compared to EDQNM (Lesieur & Ossia 2000). The estimated Kolmogorov constant is $K_0 = 1.51$ for the closure and is slightly smaller than values in DNSs and theoretical estimation of LRA ($K_0 = 1.72$). Figure 3 shows the energy flux $\Pi(k)$ normalized by ϵ , where energy flux is defined as $\Pi(k) = \int_k^\infty dkT(k) = -\int_0^k dkT(k)$, where $T(k)$ is the energy transfer function. For high-Reynolds-number turbulent flows, $\Pi(k) = \epsilon$ is satisfied in the inertial subrange according to Kolmogorov (1941a). As shown in figure 3, $\Pi(k) = \epsilon$ in the inertial subrange is satisfied for high-Reynolds-number cases in both closure and DNS. The difference between closure and DNS is mainly observed around the energy-containing and bump ranges.

The spatial second- and third-order structure functions are given by

$$S_2(r) = 4 \int_0^\infty dkE(k)f(kr), \quad f(x) = 1 - \frac{\sin x - x \cos x}{x^3}, \quad (4.1a,b)$$

$$S_3(r) = 12r \int_0^\infty dkT(k)g(kr), \quad g(x) = \frac{3 \sin x - 3x \cos x - x^2 \sin x}{x^5}. \quad (4.1c,d)$$




FIGURE 3. Normalized energy fluxes for (a) DNS and (b) closure. For a description, refer to figure 2.




FIGURE 4. Normalized second-order structure functions for (a) DNS and (b) closure and compensated second-order structure functions for (c) DNS and (d) closure. For a description, refer to figure 2. Here, $v_\eta = (\nu\epsilon)^{1/4}$ is the Kolmogorov velocity.

Here $S_2(r) \sim (\epsilon r)^{2/3}$ and $S_3(r) = -4/5\epsilon r$ are believed to be satisfied in the inertial subrange for high-Reynolds-number turbulent flows. These expressions in physical space are equivalent to the $-5/3$ law of energy spectrum and $\Pi(k) = \epsilon$ in wavenumber space. Using Taylor expansions, it is found that $S_2(r \rightarrow 0) = \epsilon r^2/15\nu + O(r^4)$ and $S_3(r \rightarrow 0) = -\frac{2}{35}r^3 \int_0^\infty dk k^2 T(k) + O(r^5)$ for dissipation range. Figure 4 shows the second- and third-order structure functions and skewness function. The Kolmogorov




FIGURE 5. Normalized third-order structure functions for (a) DNS and (b) closure and skewness functions for (c) DNS and (d) closure. For a description, refer to figure 2.

constant in the second-order structure function is given as $C_0 = 27/55\Gamma(1/3)K_0$. The estimated C_0 in the present closure is 2.0, and its value is slightly lower than the DNS value. The Reynolds number dependency of $S_3(r \rightarrow 0)/S_2(r \rightarrow 0)^{3/2} = \langle (\partial u/\partial x)^3 \rangle / \langle (\partial u/\partial x)^2 \rangle^{3/2}$ is observed for both closure and DNS as in Sreenivasan & Antonia (1997). In the present closure, $S_3(r \rightarrow 0)/S_2(r \rightarrow 0)^{3/2}$ approaches -0.6 with an increase of Re_λ and is slightly smaller than for LRA, $S_3(r \rightarrow 0)/S_2(r \rightarrow 0)^{3/2} = -0.66$, for an infinite Reynolds number (Kaneda 1993). Compared to the EDQNM (Bos *et al.* 2007a), the present closure is closer to DNS at moderate Reynolds numbers (Gotoh *et al.* 2002). It is noted that the present closure does not contradict the boundedness of the velocity derivative skewness for high-Reynolds-number turbulent flows (Antonia *et al.* 2015).

As shown in figures 1–5, the present closure yields quantitative agreement with DNS and its extension to anisotropic turbulence will be performed in the future.

Before closing this paper, we discuss the eddy damping and briefly comment on the capability of the intermittency effect. In EDQNM, the response function is regarded as having the form $G(k, t, \tau) = \exp[-(\nu k^2 + \mu^{EDQNM}(k, t))\tau]$, where $\mu^{EDQNM}(k, t) = \alpha \sqrt{\int_0^k dp p^2 E(p, t)}$ is the inverse of the rotation time or that of strain time. In EDQNM, one can adjust K_0 through the parameter α and its relation is given as $K_0 \approx 2.76\alpha^{2/3}$ (Lesieur & Ossia 2000). Figure 6 shows the eddy damping for the present closure and EDQNM. In the inertial subrange using Kolmogorov’s $-5/3$ law, $\mu(k)$ in the present closure takes the form $\mu(k) = 1.06K_0\epsilon^{2/3}k^{4/3}$ (Kaneda 1993), whereas $[\mu^{EDQNM}(k)]^2 = 3/4\alpha^2K_0\epsilon^{2/3}k^{4/3}$ in EDQNM. In the present closure, μ_k is regarded as the inverse of the squared Lagrangian time scale, where the Lagrangian




FIGURE 6. Eddy damping for (a) present closure and (b) EDQNM.

time scale in wavenumber space is given as $\epsilon^{-1/3}k^{-2/3}$ (Sagaut & Cambon 2008). The behaviours of energy-containing range and dissipation range can be understood by analysing the non-local interaction of eddy damping. For $k \ll k_c \leq p \sim q$, we assume that the contribution of integration $\iint_{\Delta} dp dq$ mainly comes from $p \geq k_c$ and $q \geq k_c$, where k_c is the convenient cut-off wavenumber. We then have

$$\mu(k \ll k_c) = 2 \int_{k_c}^{\infty} dp \int_{p-k}^p dq \frac{kp}{q} (x + yz)^2 E(q) \approx \frac{8}{15} k^2 \int_{k_c}^{\infty} dp E(p), \quad (4.2)$$

where the change of variable using $z = (k^2 + p^2 - q^2)/(2kp)$ and Taylor expansions at $k/p \ll 1$ were used in the derivation. It is found from (4.2) that $\mu(k) \sim Kk^2$ at low wavenumber for the present closure, whereas $[\mu^{EDQNM}(k)]^2 \sim k^3 E(k)$ at low wavenumber for EDQNM. In this study, the forcing spectrum is given as k^2 at low wavenumber, so that k^5 behaviours can be seen for the eddy damping in EDQNM as shown in figure 6. In other words, eddy damping of the present closure is independent of the energy spectrum at low wavenumber. On the other hand, for $q \leq k_c \ll k \sim p$ (or $p \leq k_c \ll k \sim q$), we assume that the contribution of integration $\iint_{\Delta} dp dq$ mainly comes from $q \leq k_c$ (or $p \leq k_c$). We then have

$$\mu(k \gg k_c) = 2 \int_0^{k_c} dq \int_{k-q}^{k+q} dp \frac{kp}{q} (x + yz)^2 E(q) \approx \frac{8}{15} \int_0^{k_c} dq q^2 E(q), \quad (4.3)$$

where the change of variable using $y = (k^2 + q^2 - p^2)/(2kq)$ and Taylor expansions at $q/k \ll 1$ were used in the derivation. It is found from (4.3) that $\mu(k)$ approaches $4/15\tau_\eta^2$ for high wavenumber for the present closure, in which $\tau_\eta = \sqrt{\nu/\epsilon}$ is the Kolmogorov time scale. For EDQNM, $[\mu^{EDQNM}(k)]^2 = \alpha^2\tau_\eta^2/2$ for the high-wavenumber limit. Thus, the eddy damping in the present closure qualitatively agrees with squared eddy damping in EDQNM for high wavenumber, whereas there is a qualitative difference for low wavenumber.

It is known that the drawback of spectral closures is neglect of the intermittency effects. However, as discussed by Yoshida, Ishihara & Kaneda (2003), it is non-trivial whether or not the closure theories do not yield anomalous scaling since the present closure is similar in a sense to the exact closure equation for the Kraichnan model (Kraichnan 1994) which has a solution exhibiting anomalous scaling. Thus, the problem related to intermittency still remains unsolved and further studies are required.

5. Discussion and conclusion

In the present study, we derived the two-point single-time spectral closure starting from LRA and the use of assumptions (i) Markovianization and (ii) the Lagrangian velocity response function being expressed by $G(k, \tau) = \exp(-C_1(k)\tau - C_2(k)\tau^2/2)$. The unknown functions $C_1(k)$ and $C_2(k)$ are theoretically derived from being consistent with the exact short-time behaviour of $G(k, \tau)$ and the asymptotic short-time behaviour of assumed exponential form of $G(k, \tau)$. It is found that $C_1(k)$ is the inverse of the viscous time scale and $C_2(k)$ is the inverse of the squared Lagrangian time scale in the present closure. The main difference between the present closure and other single-time spectral closures (EDQNM, test field model and Lagrangian Markovianized field approximation) is in the form of the response function, where $G(k, \tau) = \exp(-C(k)\tau)$ is assumed for the other single-time spectral closures. In the inertial subrange, it is found that $C_2(k)$ and $C(k)^2$ are proportional to $\epsilon^{2/3}k^{4/3}$ regardless of closure. On the other hand, a difference can be seen for eddy damping in the energy-containing range.

More generally speaking, most closures proposed so far take the form of (2.8) for the energy transfer function in homogeneous isotropic turbulence and the differences in closures originate from the Lagrangian velocity response function and Lagrangian velocity correlation. From the perspective of the renormalization theory, these differences come from the different representatives, and a more suitable choice of representatives may improve the level of approximation. However, taking into account attractive points in LRA, e.g. (i) satisfactory nature of the fluctuation–dissipation theorem and (ii) the simple closed equations compared to ALHDIA and SBALHDIA, the present closure based on LRA is not only promising for an extension to closure in anisotropic turbulence but also reduces the computational cost as well as the other two-point single-time closures.

The numerical results show that the single-time MLRA has good agreement with DNS for single- and two-point statistics including the finite Reynolds number dependency. Thus, the single-time MLRA is more sophisticated compared to other two-point single-time spectral closures in the sense that (i) it can calculate the statistical quantities by theory without the introduction of any adjustable parameters, (ii) it involves simple closed equations and (iii) it shows quantitatively good agreements with DNS. An extension of the present closure to anisotropic turbulence will be performed in future work.

Acknowledgements

The author would like to thank colleagues Professor D. Sakaguchi (Nagasaki University), Professor H. Ueki (Nagasaki University) and emeritus Professor M. Ishida (Nagasaki University) for discussions. He would like to acknowledge Professor T. Ishihara (Okayama University) for informing him of the useful paper on closure and to thank Ms H. Kitamura (Nagasaki Nihon University JHS) for checking and correcting the grammatical errors. He is also grateful to the three anonymous referees for interesting discussions and helpful comments on this topic. The DNS was carried out using the computer resource offered by Research Institute for Information Technology, Kyushu University. This work was supported by the Japanese Ministry of Education, Science and Culture through grants-in-aid (no. 19K14891) and Harada Memorial Foundation.

Declaration of interests

The author reports no conflict of interest.

Appendix A

The Fourier coefficient of Lagrangian velocity is defined as

$$\hat{v}_i(\mathbf{k}, s|t) = \frac{1}{(2\pi)^3} \int v_i(\mathbf{x}, s|t) \exp(-i\mathbf{k} \cdot \mathbf{x}) \, d\mathbf{x} \tag{A 1}$$

and that of the Lagrangian position function is defined as

$$\hat{\psi}(\mathbf{k}, t|\mathbf{k}', s) = \frac{1}{(2\pi)^6} \iint \psi(\mathbf{y}, t|\mathbf{x}, s) \exp(-i(\mathbf{k} \cdot \mathbf{y} + \mathbf{k}' \cdot \mathbf{x})) \, d\mathbf{y} \, d\mathbf{x}, \tag{A 2}$$

with an initial condition $\hat{\psi}(\mathbf{k}, s|\mathbf{k}', s) = \delta(\mathbf{k} + \mathbf{k}')/(2\pi)^3$ as understood from (2.3b). Here $i = \sqrt{-1}$. From (2.4), (A 1) and (A 2), we have

$$\hat{v}_i(\mathbf{k}, s|t) = (2\pi)^3 \int d\mathbf{k}' \hat{u}_i(\mathbf{k}', t) \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s), \tag{A 3}$$

with an initial condition $\hat{v}_i(\mathbf{k}, s|s) = \hat{u}_i(\mathbf{k}, s)$. Hereinafter, $\hat{f}(\mathbf{k}, t)$ represents the Fourier representation of $f(\mathbf{x}, t)$. From (2.5) and (A 1), the equation for $\hat{v}(\mathbf{k}, s|t)$ is given as

$$\begin{aligned} \frac{\partial}{\partial t} \hat{v}_i(\mathbf{k}, s|t) &= (2\pi)^3 \int d\mathbf{k}' \hat{f}_i(\mathbf{k}', t) \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s) - \nu(2\pi)^3 \int d\mathbf{k}' k'^2 \hat{u}_i(\mathbf{k}', t) \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s) \\ &+ \alpha i(2\pi)^3 \int d\mathbf{k}' \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s) \frac{k'_i k'_j k'_k}{k'^2} \int_{\mathbf{k}'=\mathbf{p}+\mathbf{q}} \hat{u}_j(\mathbf{p}, t) \hat{u}_k(\mathbf{q}, t) \, d\mathbf{p}, \end{aligned} \tag{A 4}$$

where $\alpha = 1$ is the bookkeeping parameter for the convenience of perturbation expansion and is introduced for the convolution terms. From (A 4), the equations of the Lagrangian velocity correlation $Q_{ij}(\mathbf{k}, t, s) = P_{i\alpha}(\mathbf{k}) \hat{v}_\alpha(\mathbf{k}, s|t) \hat{v}_j(-\mathbf{k}, s|s)$ and the Lagrangian velocity response function $G_{ij} = P_{i\alpha}(\mathbf{k}) \delta \hat{v}_\alpha(\mathbf{k}, s|t) / \delta \hat{f}_j(\mathbf{k}', s)$ are given as

$$\begin{aligned} \frac{\partial}{\partial t} Q_{ij}(\mathbf{k}, t, s) &= (2\pi)^3 P_{i\alpha}(\mathbf{k}) \underbrace{\int d\mathbf{k}' \hat{f}_\alpha(\mathbf{k}', t) \hat{u}_j(-\mathbf{k}, s) \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s)}_A \\ &- \nu(2\pi)^3 P_{i\alpha}(\mathbf{k}) \underbrace{\int d\mathbf{k}' k'^2 \hat{u}_\alpha(\mathbf{k}', t) \hat{u}_j(-\mathbf{k}, s) \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s)}_B \\ &+ \alpha i(2\pi)^3 P_{i\alpha}(\mathbf{k}) \underbrace{\int d\mathbf{k}' \hat{\psi}(-\mathbf{k}', t|\mathbf{k}, s) \frac{k'_\alpha k'_\beta k'_\gamma}{k'^2} \int_{\mathbf{k}'=\mathbf{p}+\mathbf{q}} \hat{u}_j(-\mathbf{k}, s) \hat{u}_\beta(\mathbf{p}, t) \hat{u}_\gamma(\mathbf{q}, t) \, d\mathbf{p}}_C, \end{aligned} \tag{A 5}$$

$$\begin{aligned} \frac{\partial}{\partial t} G_{ij}(\mathbf{k}, t|\mathbf{k}', s) &= (2\pi)^3 P_{i\alpha}(\mathbf{k}) \underbrace{\int d\mathbf{k}'' \delta_{\alpha j} \delta(\mathbf{k}'' + \mathbf{k}') \delta(t-s) \psi(-\mathbf{k}'', t|\mathbf{k}, s)}_D \\ &- \nu(2\pi)^3 P_{i\alpha}(\mathbf{k}) \underbrace{\int d\mathbf{k}'' k''^2 G_{\alpha j}^E(\mathbf{k}'', t|\mathbf{k}', s) \psi(-\mathbf{k}'', t|\mathbf{k}, s)}_E \end{aligned}$$

$$\begin{aligned}
 & + \underbrace{\alpha 2i(2\pi)^3 P_{i\alpha}(\mathbf{k}) \int d\mathbf{k}'' \hat{\psi}(-\mathbf{k}'', t|\mathbf{k}, s) \frac{k''_\alpha k''_\beta k''_\gamma}{k''^2} \int_{\mathbf{k}''=\mathbf{p}+\mathbf{q}} d\mathbf{p} \hat{u}_\beta(\mathbf{p}, t) G_{\gamma j}^E(\mathbf{q}, t|\mathbf{k}', s)}_F \\
 & + \underbrace{(2\pi)^3 P_{i\alpha}(\mathbf{k}) \int d\mathbf{k}'' \hat{f}_\alpha(\mathbf{k}'', t) \Psi_j(-\mathbf{k}'', t|\mathbf{k}, \mathbf{k}', s)}_G \\
 & - \underbrace{\nu(2\pi)^3 P_{i\alpha}(\mathbf{k}) \int d\mathbf{k}'' \hat{u}_\alpha(\mathbf{k}'', t) \Psi_j(-\mathbf{k}'', t|\mathbf{k}, \mathbf{k}', s)}_H \\
 & + \underbrace{\alpha i(2\pi)^3 P_{i\alpha}(\mathbf{k}) \int d\mathbf{k}'' \hat{\psi}_j(-\mathbf{k}'', t|\mathbf{k}, \mathbf{k}', s) \frac{k''_\alpha k''_\beta k''_\gamma}{k''^2} \int_{\mathbf{k}''=\mathbf{p}+\mathbf{q}} d\mathbf{p} \hat{u}_\beta(\mathbf{p}, t) \hat{u}_\gamma(\mathbf{q}, t)}_I
 \end{aligned} \tag{A6}$$

where $G_{\alpha j}^E(\mathbf{k}'', t|\mathbf{k}', s)$ is the Eulerian response function defined as $G_{\alpha j}^E(\mathbf{k}'', t|\mathbf{k}', s) = \delta \hat{u}_\alpha(\mathbf{k}'', t) / \delta \hat{f}_j(\mathbf{k}', s)$ (Kraichnan 1959). Function $\Psi_j(\mathbf{k}'', t|\mathbf{k}, \mathbf{k}', s)$ is the response function of $\hat{\psi}(\mathbf{k}'', t|\mathbf{k}, s)$ for $\hat{f}_j(\mathbf{k}', s)$ and is defined as $\Psi_j(\mathbf{k}'', t|\mathbf{k}, \mathbf{k}', s) = \delta \hat{\psi}(\mathbf{k}'', t|\mathbf{k}, s) / \delta \hat{f}_j(\mathbf{k}', s)$ for $t \geq s$ and zero for $t < s$.

The role of \mathbf{P} is to enforce the dilatational component of Lagrangian velocity to be zero since $\hat{\mathbf{v}}(\mathbf{k}, s|t)$ satisfies the solenoidal condition at $t = s$, but does not necessarily satisfy it at $t \neq s$. As shown later, this operator eliminates the irrelevant terms in the expansion and simplifies the closed equations.

Assuming that variables can be expanded in terms of α , e.g. $\hat{\mathbf{u}}(\mathbf{k}, t) = \sum_{n=0}^\infty \alpha^n \hat{\mathbf{u}}^{(n)}(\mathbf{k}, t)$, $\hat{\psi}(\mathbf{k}, t|\mathbf{k}', s) = \sum_{n=0}^\infty \alpha^n \hat{\psi}^{(n)}(\mathbf{k}, t|\mathbf{k}, s)$, $G_{ij}^E(\mathbf{k}, t|\mathbf{k}', s) = \sum_{n=0}^\infty \alpha^n G_{ij}^{E(n)}(\mathbf{k}, t|\mathbf{k}', s)$ and so on, and substituting the perturbation expansions into each equation and equating powers of α , we find that, for example, $\hat{\psi}^{(0)}(\mathbf{k}, t|\mathbf{k}', s) = \delta(\mathbf{k} + \mathbf{k}') / (2\pi)^3$, $G_{ij}^{E(0)}(\mathbf{k}, t|\mathbf{k}', s) = G_{ij}^{(0)}(\mathbf{k}, t, s) \delta(\mathbf{k} + \mathbf{k}') / (2\pi)^3$ with $G_{ij}^{(0)}(\mathbf{k}, t, s) = P_{ij}(\mathbf{k}) \exp(-\nu k^2(t - s))$ and so on for zeroth-order solutions. The first-order solutions are also obtained using the zeroth-order solutions, e.g. $\hat{u}_i^{(1)}(\mathbf{k}, t) = \int_{t_0}^t ds G_{i\alpha}^{(0)}(\mathbf{k}, t, s) M_{\alpha\beta\gamma}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \hat{u}_\beta^{(0)}(\mathbf{p}, s) \hat{u}_\gamma^{(0)}(\mathbf{q}, s)$ and so on.

In the derivation of closed equations of LRA, the following assumptions are made:

- (a) $\hat{\mathbf{u}}^{(0)}$, $G^{(0)}$, $\hat{\psi}^{(0)}$ and $\hat{\Psi}^{(0)}$ are statistically independent of each other, and
- (b) the probability distribution of $\hat{\mathbf{u}}^{(0)}$ is assumed to be Gaussian, i.e. the odd moments are zero and the even moments can be expressed by the second moments.

It is noted that the basic ideas for the derivations are different between LRA and Lagrangian DIA, although the closed equations of LRA and Lagrangian DIA have the same forms. Here is shown the systematic derivation of LRA equations.

A.1. Two-point single-time Eulerian velocity correlation

After lengthy algebra using the perturbation expansions and assumptions, we have the following equation for the two-point single-time Eulerian velocity correlation $Q_{ij}(\mathbf{k}, t, t)$:

$$\left(\frac{\partial}{\partial t} + 2\nu k^2 \right) Q_{ij}(\mathbf{k}, t, t) = \alpha^2 D_{ij}(\mathbf{k}, t, t) + Q_{ij}^F(\mathbf{k}, t, t) + Q_{ji}^{F*}(\mathbf{k}, t, t), \tag{A7}$$

where

$$D_{ij}(\mathbf{k}, t) = H_{ij}(\mathbf{k}, t) + H_{ji}(-\mathbf{k}, t) + O(\alpha), \tag{A 8}$$

with

$$H_{ij}(\mathbf{k}, t) = M_{i\alpha\beta}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \times \left[2 \int_{t_0}^t ds' G_{j\gamma}^{(0)}(-\mathbf{k}, t, s') M_{\gamma\delta\epsilon}(-\mathbf{k}) Q_{\alpha\delta}^{(0)}(\mathbf{p}, t, s') Q_{\beta\epsilon}^{(0)}(\mathbf{q}, t, s') + 4 \int_{t_0}^t ds' G_{\alpha\gamma}^{(0)}(\mathbf{p}, t, s') M_{\gamma\delta\epsilon}(\mathbf{p}) Q_{j\delta}^{(0)}(-\mathbf{k}, t, s') Q_{\beta\epsilon}^{(0)}(\mathbf{q}, t, s') \right] + O(\alpha), \tag{A 9}$$

where $Q_{ij}^{(0)}(\mathbf{k}, t, s) = \langle \hat{u}_i^{(0)}(\mathbf{k}, t) \hat{u}_j^{(0)}(-\mathbf{k}, s) \rangle$, $Q_{ij}^F(\mathbf{k}, t) = \langle \hat{f}_i(\mathbf{k}, t) \hat{u}_j^*(\mathbf{k}, t) \rangle$, the superscript * denotes the complex conjugate and $M_{i\alpha\beta}(\mathbf{k}) = -i/2[k_\alpha P_{i\beta}(\mathbf{k}) + k_\beta P_{i\alpha}(\mathbf{k})]$. See Leslie (1973) and McComb (1989, 2014) for the derivation of equation of $Q_{ij}(\mathbf{k}, t, t)$.

A.2. Two-point two-time Lagrangian velocity correlation

We consider the ensemble average of (A 5). Substituting the perturbation expansions into (A 5) and taking the ensemble average, $\langle A \rangle$ and $\langle B \rangle$ are given as

$$\langle A \rangle = Q_{ij}^{F(0)}(\mathbf{k}, t, s) + O(\alpha), \tag{A 10}$$

$$\langle B \rangle = -\nu k^2 Q_{ij}^{(0)}(\mathbf{k}, t, s) + O(\alpha). \tag{A 11}$$

After lengthy algebra using similar procedures, $\langle C \rangle$ is given as

$$\langle C \rangle = \alpha^2 4i P_{i\alpha}(\mathbf{k}) \frac{k_\alpha k_\beta k_\gamma}{k^2} \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \int_{t_0}^t ds' G_{\gamma\delta}^{(0)}(\mathbf{q}, t, s') M_{\delta\epsilon\zeta}(\mathbf{q}) Q_{j\epsilon}^{(0)}(\mathbf{k}, s', s) Q_{\beta\zeta}^{(0)}(\mathbf{p}, t, s') + \alpha^2 2i P_{i\alpha}(\mathbf{k}) \frac{k_\alpha k_\beta k_\gamma}{k^2} \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \int_{t_0}^s ds' G_{j\delta}^{(0)}(-\mathbf{k}, s, s') \times M_{\delta\epsilon\zeta}(-\mathbf{k}) Q_{\beta\epsilon}^{(0)}(\mathbf{p}, t, s') Q_{\gamma\zeta}^{(0)}(\mathbf{q}, t, s') - \alpha^2 2P_{i\alpha}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \int_s^t ds' \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} Q_{\gamma\delta}^{(0)}(-\mathbf{q}, t, s') Q_{\beta j}^{(0)}(\mathbf{k}, t, s) + O(\alpha^3), \tag{A 12}$$

where the first and second terms disappear because of $P_{i\alpha}(\mathbf{k})k_\alpha k_\beta k_\gamma/k^2 = 0$. In the derivation of (A 12), symmetries of the integral with respect to \mathbf{p} and \mathbf{q} and $M_{i\alpha\beta}(\mathbf{k})\delta(\mathbf{k}) = 0$ were used. Summarizing the above equations, the equation for two-point two-time Lagrangian velocity correlation is given as

$$\frac{\partial}{\partial t} Q_{ij}^L(\mathbf{k}, t, s) = -\nu X_{ij}(\mathbf{k}, t, s) - \alpha^2 I_{ij}(\mathbf{k}, t, s) + Z_{ij}(\mathbf{k}, t, s), \tag{A 13}$$

where

$$X_{ij}(\mathbf{k}, t, s) = k^2 Q_{ij}^{(0)}(\mathbf{k}, t, s) + O(\alpha), \tag{A 14}$$

$$I_{ij}(\mathbf{k}, t, s) = 2P_{i\alpha}(\mathbf{k}) \int_s^t ds' \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} Q_{\gamma\delta}^{(0)}(-\mathbf{q}, t, s') Q_{\beta j}^{(0)}(\mathbf{k}, t, s) + O(\alpha), \tag{A 15}$$

$$Z_{ij}(\mathbf{k}, t, s) = Q_{ij}^{F(0)}(\mathbf{k}, t, s) + O(\alpha). \tag{A 16}$$

A.3. Lagrangian velocity response function

As in two-point two-time Lagrangian velocity correlation, we consider the ensemble-averaged equation of (A 6) for $\mathbf{k}' = -\mathbf{k}$, i.e. $G_{ij}(\mathbf{k}, t | -\mathbf{k}, s) = G_{ij}(\mathbf{k}, t, s)$. Substituting the perturbation expansions into (A 6), replacing $\mathbf{k}' = -\mathbf{k}$ and taking the ensemble average, $\langle D \rangle$ with $\mathbf{k}' = -\mathbf{k}$ is given as

$$\langle D \rangle = P_{ij}(\mathbf{k})\delta(t - s) + O(\alpha). \tag{A 17}$$

Similarly, $\langle E \rangle$ for $\mathbf{k}' = -\mathbf{k}$ is given as

$$\langle F \rangle = -\nu k^2 G_{ij}^{(0)}(\mathbf{k}, t, s) + O(\alpha). \tag{A 18}$$

After lengthy algebra using similar procedures, $\langle F \rangle$ for $\mathbf{k}' = -\mathbf{k}$ in (A 6) is given as

$$\begin{aligned} \langle F \rangle = & -\alpha^2 2P_{i\alpha}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} \int_s^t ds' Q_{\gamma\delta}^{(0)}(-\mathbf{q}, t, s') G_{\beta j}^{(0)}(\mathbf{k}, t, s) \\ & + \alpha^2 4iP_{i\alpha}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \frac{k_\alpha k_\beta k_\gamma}{k^2} \int_s^t ds' G_{\beta\delta}^{(0)}(\mathbf{p}, t, s') \\ & \times M_{\delta\epsilon\zeta}(\mathbf{p}) Q_{\gamma\epsilon}^{(0)}(\mathbf{q}, t, s') G_{\zeta j}^{(0)}(\mathbf{k}, s', s) + O(\alpha^3), \end{aligned} \tag{A 19}$$

where the second term disappears because of $P_{ij}(\mathbf{k})k_\alpha k_\beta k_\gamma / k^2 = 0$. Substituting perturbation expansions into G and H in (A 6), replacing $\mathbf{k}' = -\mathbf{k}$ and taking the ensemble average, it is found that the zeroth-order terms of $\langle G \rangle$ and $\langle H \rangle$ are zeros, i.e. $\langle G \rangle$ and $\langle H \rangle$ start with $O(\alpha)$. Furthermore, the second-order term of $\langle I \rangle$ is zero, i.e. $\langle I \rangle$ in (A 6) starts with $O(\alpha^3)$.

Summarizing the above equations, the equation for the Lagrangian velocity response function is given as

$$\frac{\partial}{\partial t} G_{ij}(\mathbf{k}, t, s) = -\nu Y_{ij}(\mathbf{k}, t, s) - \alpha^2 J_{ij}(\mathbf{k}, t, s) + W_{ij}(\mathbf{k}, t, s), \tag{A 20}$$

where

$$Y_{ij}(\mathbf{k}, t, s) = k^2 G_{ij}^{(0)}(\mathbf{k}, t, s) + O(\alpha), \tag{A 21}$$

$$J_{ij}(\mathbf{k}, t, s) = 2P_{i\alpha}(\mathbf{k}) \int_{\mathbf{k}=\mathbf{p}+\mathbf{q}} d\mathbf{p} \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} \int_s^t ds' Q_{\gamma\delta}^{(0)}(-\mathbf{q}, t, s') G_{\beta j}^{(0)}(\mathbf{k}, t, s) + O(\alpha), \tag{A 22}$$

$$W_{ij}(\mathbf{k}, t, s) = P_{ij}(\mathbf{k})\delta(t - s) + O(\alpha). \tag{A 23}$$

A.4. Renormalized expansion

In order to close equations by the representatives, the renormalized expansion is used as in Kraichnan (1977) and Kaneda (1981). The procedure is as follows:

- (a) Begin with the primitive perturbation expansion for $\mathbf{Q}(\mathbf{k}, t, t)$, $\mathbf{Q}(\mathbf{k}, t, s)$ and $\mathbf{G}(\mathbf{k}, t, s)$ in terms of $\mathbf{Q}^{(0)}$ and $\mathbf{G}^{(0)}$

$$Q_{ij}(\mathbf{k}, t, t) = Q_{ij}^{(0)}(\mathbf{k}, t, t) + \alpha \mathcal{U}(Q^{(0)}, G^{(0)}) + \dots, \tag{A 24}$$

$$Q_{ij}(\mathbf{k}, t, s) = Q_{ij}^{(0)}(\mathbf{k}, t, s) + \alpha \mathcal{V}(Q^{(0)}, G^{(0)}) + \dots, \tag{A 25}$$

$$G_{ij}(\mathbf{k}, t, s) = G_{ij}^{(0)}(\mathbf{k}, t, s) + \alpha \mathcal{W}(Q^{(0)}, G^{(0)}) + \dots. \tag{A 26}$$

(b) Expand D , I and J in terms of $\mathbf{Q}^{(0)}$ and $\mathbf{G}^{(0)}$

$$D_{ij}(\mathbf{k}, t, s) = \alpha^2 \mathcal{D}(\mathbf{Q}^{(0)}, \mathbf{G}^{(0)}) + O(\alpha^3) + \dots, \tag{A 27}$$

$$I_{ij}(\mathbf{k}, t, s) = \alpha^2 \mathcal{I}(\mathbf{Q}^{(0)}, \mathbf{G}^{(0)}) + O(\alpha^3) + \dots, \tag{A 28}$$

$$J_{ij}(\mathbf{k}, t, s) = \alpha^2 \mathcal{J}(\mathbf{Q}^{(0)}, \mathbf{G}^{(0)}) + O(\alpha^3) + \dots. \tag{A 29}$$

(c) Revert these primitive expansions to obtain $\mathbf{Q}^{(0)}$ and $\mathbf{G}^{(0)}$ as functional power expansions in \mathbf{Q} and \mathbf{G}

$$\mathbf{Q}_{ij}^{(0)}(\mathbf{k}, t, s) = \mathbf{Q}_{ij}(\mathbf{k}, t, s) + \alpha \mathcal{X}(\mathbf{Q}, \mathbf{G}) + \dots, \tag{A 30}$$

$$\mathbf{Q}_{ij}^{(0)}(\mathbf{k}, t, s) = \mathbf{Q}_{ij}(\mathbf{k}, t, s) + \alpha \mathcal{Y}(\mathbf{Q}, \mathbf{G}) + \dots, \tag{A 31}$$

$$\mathbf{G}_{ij}^{(0)}(\mathbf{k}, t, s) = \mathbf{G}_{ij}(\mathbf{k}, t, s) + \alpha \mathcal{Z}(\mathbf{Q}, \mathbf{G}) + \dots. \tag{A 32}$$

(d) Substituting (A 30)–(A 32) into (A 27)–(A 29), D , I and J can be expressed in terms of representatives \mathbf{Q} and \mathbf{G} .

(e) Truncate the renormalized expansion at the lowest order and put α equal to unity.

The following equations are the final equations of LRA:

$$\left(\frac{\partial}{\partial t} + 2\nu k^2 \right) \mathbf{Q}_{ij}(\mathbf{k}, t) = H_{ij}(\mathbf{k}, t) + H_{ji}^*(\mathbf{k}, t) + \mathbf{Q}_{ij}^F(\mathbf{k}, t) + \mathbf{Q}_{ji}^{F*}(\mathbf{k}, t), \tag{A 33}$$

$$\left(\frac{\partial}{\partial t} + \nu k^2 \right) \mathbf{Q}_{ij}(\mathbf{k}, t, s) = I_{ij}(\mathbf{k}, t, s) \quad t > s, \tag{A 34}$$

$$\left(\frac{\partial}{\partial t} + \nu k^2 \right) \mathbf{G}_{ij}(\mathbf{k}, t, s) = J_{ij}(\mathbf{k}, t, s) \quad t > s, \tag{A 35}$$

where

$$\begin{aligned} H_{ij}(\mathbf{k}, t) &= M_{i\alpha\beta}(\mathbf{k}) \int_{k=p+q} d\mathbf{p} \\ &\times \left[2 \int_0^t ds' G_{j\gamma}(-\mathbf{k}, t, s') M_{\gamma\delta\epsilon}(-\mathbf{k}) \mathbf{Q}_{\alpha\delta}(\mathbf{p}, t, s') \mathbf{Q}_{\beta\epsilon}(\mathbf{q}, t, s') \right. \\ &\left. + 4 \int_0^t ds' G_{\alpha\gamma}(\mathbf{p}, t, s') M_{\gamma\delta\epsilon}(\mathbf{p}) \mathbf{Q}_{j\delta}(-\mathbf{k}, s, s') \mathbf{Q}_{\beta\epsilon}(\mathbf{q}, t, s') \right], \tag{A 36} \end{aligned}$$

$$I_{ij}(\mathbf{k}, t, s) = -2P_{i\alpha}(\mathbf{k}) \int_{k=p+q} d\mathbf{p} \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} \int_s^t ds' \mathbf{Q}_{\gamma\delta}(-\mathbf{q}, t, s') \mathbf{Q}_{\beta j}(\mathbf{k}, t, s), \tag{A 37}$$

$$J_{ij}(\mathbf{k}, t, s) = -2P_{i\alpha}(\mathbf{k}) \int_{k=p+q} d\mathbf{p} \frac{P_\alpha P_\beta P_\gamma P_\delta}{p^2} \int_s^t ds' \mathbf{Q}_{\gamma\delta}(-\mathbf{q}, s', s) G_{\beta j}(\mathbf{k}, t, s), \tag{A 38}$$

where the contributions from the random force in (A 35) and (A 35) are neglected because they are not important for $t > s$. In homogeneous isotropic turbulence, the Lagrangian velocity correlation and Lagrangian velocity response function are given as

$$\mathbf{Q}_{ij}(\mathbf{k}, t, s) = \frac{1}{2} \mathbf{Q}(k, t, s) P_{ij}(\mathbf{k}), \tag{A 39}$$

$$\mathbf{G}_{ij}(\mathbf{k}, t, s) = \mathbf{G}(k, t, s) P_{ij}(\mathbf{k}). \tag{A 40}$$

The closed equations for homogeneous isotropic turbulence can be obtained by substituting these expressions into (A 33)–(A 38). For details of the derivation of the closed equation for isotropic turbulence, the reader is referred to Leslie (1973) and McComb (1989, 2014). For anisotropic turbulence, the reader is referred to Sagaut & Cambon (2008).

REFERENCES

- ABRAMOWITZ, M. & STEGUN, I. A. 1964 *Handbook of Mathematical Functions*. Courier Corporation.
- ANTONIA, R. A., TANG, S. L., DJENIDI, L. & DANAILA, L. 2015 Boundedness of the velocity derivative skewness in various turbulent flows. *J. Fluid Mech.* **781**, 727–744.
- ARMSTRONG, J. W., RICKETT, B. J. & SPANGLER, S. R. 1995 Electron density power spectrum in the local interstellar medium. *Astrophys. J.* **443**, 209–221.
- BOS, W. J. T. & BERTOGLIO, J. P. 2013 Lagrangian Markovianized field approximation for turbulence. *J. Turbul.* **14**, 99–120.
- BOS, W. J. T., CLARK, T. T. & RUBINSTEIN, R. 2007a Small scale response and modeling of periodically forced turbulence. *Phys. Fluids* **19**, 055107.
- BOS, W. J. T. & FANG, L. 2015 Dependence of turbulent advection on the Lagrangian correlation time. *Phys. Rev. E* **91**, 043020.
- BOS, W. J. T., SHAO, L. & BERTOGLIO, J. P. 2007b Spectral imbalance and the normalized dissipation rate of turbulence. *Phys. Fluids* **15**, 045101.
- BRIARD, A., GOMEZ, T., SAGAUT, P. & MEMARI, S. 2015 Passive scalar decay laws in isotropic turbulence: Prandtl number effects. *J. Fluid Mech.* **784**, 274–303.
- CAO, N., CHEN, S. & DOOLEN, G. D. 1999 Statistics and structures of pressure in isotropic turbulence. *Phys. Fluids* **11**, 2235–2250.
- DAVIDSON, P. A. 2004 *Turbulence: An Introduction for Scientists and Engineers*. Oxford University Press.
- DOERING, C. R. & FOIAS, C. 2002 Energy dissipation in body-forced turbulence. *J. Fluid Mech.* **467**, 289–306.
- GOTOH, T., FUKAYAMA, D. & NAKANO, T. 2002 Velocity field statistics in homogeneous steady turbulence obtained using a high-resolution direct numerical simulation. *Phys. Fluids* **14**, 1065–1081.
- GOTOH, T. & KANEDA, Y. 2000 A numerical algorithm for efficiently solving spectral closure equations. In *Advances in Turbulence VIII*, pp. 693–696.
- GOTOH, T., KANEDA, Y. & BEKKI, N. 1988 Numerical integration of the Lagrangian renormalized approximation. *J. Phys. Soc. Japan* **57**, 866–880.
- GOTOH, T., NAGAKI, J. & KANEDA, Y. 2000 Passive scalar spectrum in the viscous-convective range in two-dimensional steady turbulence. *Phys. Fluids* **12**, 155–168.
- GRADSHTEYN, I. S. & RYZHIK, I. M. 2014 *Table of Integrals, Series, and Products*. Academic.
- HERRING, J. R. 1965 Self-consistent-field approach to turbulence theory. *Phys. Fluids* **8**, 2219–2225.
- HERRING, J. R. & KRAICHNAN, R. H. 1979 A numerical comparison of velocity-based and strain-based Lagrangian-history turbulence approximations. *J. Fluid Mech.* **91**, 581–597.
- ISHIHARA, T., MORISHITA, K., YOKOKAWA, M., UNO, A. & KANEDA, Y. 2016 Energy spectrum in high-resolution direct numerical simulations of turbulence. *Phys. Rev. Fluids* **1**, 082403.
- JIMÉNEZ, J., WRAY, A. A., SAFFMAN, P. G. & ROGALLO, R. S. 1993 The structure of intense vorticity in isotropic turbulence. *J. Fluid Mech.* **255**, 65–90.
- KANEDA, Y. 1981 Renormalized expansions in the theory of turbulence with the use of Lagrangian position function. *J. Fluid Mech.* **107**, 131–145.
- KANEDA, Y. 1993 Lagrangian and Eulerian time correlations in turbulence. *Phys. Fluids* **5**, 2835–2845.
- KANEDA, Y., ISHIHARA, T., YOKOKAWA, M., ITAKURA, K. & UNO, A. 2003 Energy dissipation rate and energy spectrum in high resolution direct numerical simulations of turbulence in aperiodic box. *Phys. Fluids* **15**, 21–24.

- KIDA, S. & GOTO, S. 1997 A Lagrangian direct-interaction approximation for homogeneous isotropic turbulence. *J. Fluid Mech.* **345**, 307–345.
- KOLMOGOROV, A. N. 1941a Dissipation of energy in the locally isotropic turbulence. *Dokl. Akad. Nauk SSSR* **32**, 15–17.
- KOLMOGOROV, A. N. 1941b The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Dokl. Akad. Nauk SSSR* **31**, 301–305.
- KRAICHNAN, R. H. 1959 The structure of isotropic turbulence at very high Reynolds numbers. *J. Fluid Mech.* **5**, 497–543.
- KRAICHNAN, R. H. 1965 Lagrangian history closure approximation for turbulence. *Phys. Fluids* **8**, 575–595.
- KRAICHNAN, R. H. 1971 An almost-Markovian Galilean-invariant turbulence model. *J. Fluid Mech.* **47**, 513–524.
- KRAICHNAN, R. H. 1977 Eulerian and Lagrangian renormalization in turbulence theory. *J. Fluid Mech.* **83**, 349–374.
- KRAICHNAN, R. H. 1994 Anomalous scaling of a randomly advected passive scalar. *Phys. Rev. Lett.* **72**, 1016–1019.
- KRAICHNAN, R. H. & HERRING, J. R. 1978 A strain based Lagrangian history turbulence theory. *J. Fluid Mech.* **88**, 355–367.
- LESIEUR, M. & OSSIA, S. 2000 3D isotropic turbulence at very high Reynolds numbers: EDQNM study. *J. Turbul.* **1**, 1–25.
- LESLIE, D. C. 1973 *Developments in the Theory of Turbulence*. Oxford University Press.
- MATTHAEUS, W. H. & GOLDSTEIN, L. 1982 Measurement of the rugged invariants of magnetohydrodynamic turbulence in the solar wind. *J. Geophys. Res.* **87**, 6011–6028.
- MCCOMB, W. D. 1989 *The Physics of Fluid Turbulence*. Clarendon Press.
- MCCOMB, W. D. 2014 *Homogeneous, Isotropic Turbulence*. Oxford University Press.
- MCCOMB, W. D. & SHANMUGASUNDARAM, V. 1984 Numerical calculation of decaying isotropic turbulence using the LET theory. *J. Fluid Mech.* **143**, 95–123.
- MONIN, A. S. & YAGLOM, A. M. 1975 *Statistical Fluid Mechanics*, vol. II. MIT Press.
- NASTROM, G. D. & GAGE, K. S. 1985 A climatology of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft. *J. Atmos. Sci.* **42**, 950–960.
- SAGAUT, P. & CAMBON, C. 2008 *Homogeneous Turbulence Dynamics*. Cambridge University Press.
- SREENIVASAN, K. R. 1995 On the universality of the Kolmogorov constant. *Phys. Fluids* **7**, 2778–2784.
- SREENIVASAN, K. R. & ANTONIA, R. A. 1997 The phenomenology of small-scale turbulence. *Annu. Rev. Fluid Mech.* **29**, 435–472.
- TSUJI, Y. 2004 Intermittency effect on energy spectrum in high-Reynolds number turbulence. *Phys. Fluids* **16**, L43–L46.
- YOSHIDA, K., ISHIHARA, T. & KANEDA, Y. 2003 Anisotropic spectrum of homogeneous turbulent shear flow in a Lagrangian renormalized approximation. *Phys. Fluids* **15**, 2385–2397.