

# Impulsivity and aggression in schizophrenia: a neural circuitry perspective with implications for treatment

Matthew J. Hoptman\*

Schizophrenia Research Division, Nathan Kline Institute for Psychiatric Research, Orangeburg, New York, USA; Department of Psychiatry, New York University School of Medicine, New York, New York, USA; Department of Psychology, City University of New York, New York, New York, USA

Elevations of impulsive behavior have been observed in a number of serious mental illnesses. These phenomena can lead to harmful behaviors, including violence, and thus represent a serious public health concern. Such violence is often a reason for psychiatric hospitalization, and it often leads to prolonged hospital stays, suffering by patients and their victims, and increased stigmatization. Despite the attention paid to violence, little is understood about its neural basis in schizophrenia. On a psychological level, aggression in schizophrenia has been primarily attributed to psychotic symptoms, desires for instrumental gain, or impulsive responses to perceived personal slights. Often, multiple attributions can coexist during a single aggressive incident. In this review, I discuss the neural circuitry associated with impulsivity and aggression in schizophrenia, with an emphasis on implications for treatment. Impulsivity appears to account for a great deal of aggression in schizophrenia, especially in inpatient settings. Urgency, defined as impulsivity in the context of strong emotion, is the primary focus of this article. It is elevated in several psychiatric disorders, and in schizophrenia, it has been related to aggression. Many studies have implicated dysfunctional frontotemporal circuitry in impulsivity and aggression in schizophrenia, and pharmacological treatments may act via that circuitry to reduce urgency and aggressive behaviors; however, more mechanistic studies are critically needed. Recent studies point toward manipulable neurobehavioral targets and suggest that cognitive, pharmacological, neuromodulatory, and neurofeedback treatment approaches can be developed to ameliorate urgency and aggression in schizophrenia. It is hoped that these approaches will improve treatment efficacy.

Received 16 January 2015; Accepted 4 March 2015; First published online 22 April 2015

**Key words:** Functional connectivity, impulsivity, MRI, neuroimaging, schizophrenia, treatment, urgency, violence, aggression.

## Introduction

Impulsive behavior is an important feature of a number of serious mental illnesses (SMI), including schizophrenia, bipolar disorder, substance abuse, and borderline personality disorder. In many cases, impulsive behavior can lead to other problematic behaviors, including violence or self-harm. Such violence is a highly significant public health issue because often it is a proximate cause of hospitalization and leads to increased healthcare costs,<sup>1</sup> particularly

because persistently aggressive patients can be difficult to discharge. Moreover, the media often sensationalizes violent acts associated with mental illness, which contributes to stigma.<sup>2</sup> It is becoming increasingly clear that increased violence risk is not fully explained by symptoms characteristic of SMI. Although aggression in SMI is heterogeneous,<sup>3</sup> impulsivity has been posited to play an important role. This article examines the neural circuitry of impulsivity and aggression in schizophrenia, a disorder characterized by positive symptoms (e.g., hallucinations, delusions), negative symptoms (e.g., social withdrawal, blunted affect), and a wide range of cognitive deficits. Because the review focuses specifically on impulsivity and aggression, important work on psychotic and psychopathic bases of aggression is not included herein, largely due to space limitations. To set the context, I first briefly discuss the heterogeneity of violence in schizophrenia and issues related to its measurement.

\* Address for correspondence, Matthew J. Hoptman, PhD, Nathan Kline Institute, 140 Old Orangeburg Rd., Bldg. 35, Orangeburg, NY 10962, USA. (Email: Hoptman@nki.rfmh.org)

I wish to thank Karen A. Nolan, PhD, and Amanda R. McGovern, PhD, for their helpful comments on prior drafts of this article.

This article draws on work supported in part by the National Institute of Mental Health (M.J.H., grant numbers R01MH064783 and R21MH0804031) and NARSAD (M.J.H., Young Investigator Award).

## Body of Review

### Etiology

First and foremost, aggression in schizophrenia is multidetermined.<sup>3</sup> Therefore, attempts to treat aggression in schizophrenia that use a one-size-fits-all approach are doomed to fail. The literature suggests that in schizophrenia, the 3 primary causes of aggression in schizophrenia are psychotic symptoms, psychopathy, and impulsivity.<sup>4</sup> Other factors are also important, including neurological abnormalities, substance use, and poor medication adherence,<sup>5</sup> although these may exert their influence on aggression via the primary causes named above. Thus, in an important study, Nolan *et al*<sup>6</sup> interviewed inpatients with schizophrenia whose violent acts were recorded on videotape. About 20% of assaults were attributed directly to positive symptoms. A factor analysis revealed 2 psychosis-related factors: one related to positive symptoms, and the other related to disorganization/confusion. A third factor differentiated impulsive from psychopathically based assaults. Impulsive versus psychopathically based aggression is often mapped onto a distinction between reactive and instrumental/proactive aggression<sup>7</sup> or impulsive versus predatory aggression.<sup>4</sup> An important conclusion from that study is that patients could not be characterized by the type of violence they tended to commit, and that within an incident, multiple factors are likely to play a role.

The literature consistently shows that at least a plurality of inpatient aggression has an impulsive basis,<sup>8</sup> although it is notable that the comorbidity of psychopathy and schizophrenia is higher in violent than nonviolent patients.<sup>9</sup> However, the most common measures of impulsivity, such as the Barratt Impulsiveness Scale (BIS),<sup>10</sup> have not always shown particularly strong correlations with aggression.<sup>11</sup> This could be, in part, because the theory behind the BIS explicitly separates impulsivity and emotion.<sup>12</sup>

It has long been understood that impulsivity is a multifaceted construct. An important distinction in this regard is between behavioral and trait measures of impulsivity. Regarding the former, a number of studies have examined response inhibition, measured using the go/no-go and stop signal tasks, as a key aspect of impulsivity. Indeed, patients with schizophrenia show significant deficits in the ability to inhibit prepotent responses.<sup>11,13</sup> The circuitry for response inhibition is fairly well understood, and includes the right inferior frontal gyrus, the supplementary motor area, the globus pallidus, the striatum, the thalamus, and the subthalamic nucleus.<sup>14</sup> However, response inhibition is only weakly correlated with aggression,<sup>11,15</sup> suggesting that other aspects of impulsivity may be more relevant to these behaviors in schizophrenia.

A major advance with regard to understanding trait impulsivity regarding aggression was achieved by Whiteside and Lynam.<sup>16</sup> They compared many different measures of impulsivity, and their factor analysis yielded 5 constructs: Urgency, (lack of) Premeditation, (lack of) Perseverance, and Sensation seeking (UPPS). Urgency is defined as impulsive behavior in the context of strong negative emotion. A newer subscale adds the construct of positive urgency, which is defined as impulsivity in the context of strong positive emotion, yielding the UPPS-P.<sup>17</sup> Urgency is elevated in a number of psychiatric populations that have affective dysregulation, including in schizophrenia.<sup>18</sup>

The neural correlates of urgency are thought to include aberrant patterns of activity in the orbitofrontal and ventromedial prefrontal cortex, as well as in the amygdala.<sup>19</sup> Indeed, in a study of social drinkers, Cyders *et al*<sup>20</sup> found that orbitofrontal cortex (OFC) and amygdala activation in response to negative emotional pictures correlated with negative urgency. Further, negative urgency mediated between general risk-taking and both amygdala and OFC activation to negative emotional pictures.

Behavioral and trait impulsivity measures are not strongly related in schizophrenia.<sup>21</sup> These measures are somewhat better correlated in healthy individuals.<sup>22</sup> The lack of correlation in schizophrenia may reflect other factors, including heterogeneity of symptoms or different subtypes. In healthy individuals, variable results may reflect the difference between trait and state aspects of impulsivity. More generally, trait measures (such as urgency) and the most commonly used behavioral measures (stop and go/no-go paradigms) appear to involve somewhat different neural circuitry. Relatively new behavioral tasks that include emotional conditions<sup>23</sup> appear to activate, among other regions, areas thought to be relevant to urgency-based circuitry. However, whether performance on these tasks correlates with trait measures of impulsivity is not yet known.

Individuals with schizophrenia appear to experience emotions at the same level as healthy individuals, albeit with reduced outward expression.<sup>24</sup> Patients with schizophrenia have problems in both emotional control and impulsivity—domains that are highly relevant to urgency. Moreover, urgency appears to have face validity as a correlate of aggression, and as will be discussed below, recent work suggests that it is the specific aspect of impulsivity most clearly linked to aggression in schizophrenia.

### Measurement of aggression

The NIMH Research Domain Criteria (RDoC) initiative<sup>25</sup> points to the importance of examining different units of analysis within a domain. However, the assessment of aggression in schizophrenia so far has focused on only

1 or occasionally 2 levels of measurement. For measurement of aggressive behavior, the most relevant units would include behavior, self-reports, and paradigms. Many studies of violence in schizophrenia use direct observation of behavior.<sup>26</sup> Self-report measures also have been used extensively.<sup>27</sup> Paradigms such as the Taylor Aggression Paradigm (TAP)<sup>28</sup> and the Point Subtraction Aggression Paradigm (PSAP)<sup>29</sup> allow for fine-grained analyses of aggressive phenomena, but these have not been widely applied in schizophrenia. It is thus promising that the PSAP has been included in the RDoC matrix (<http://www.nimh.nih.gov/research-priorities/rdoc/nimh-research-domain-criteria-rdoc.shtml>), which may lead to a broader adoption of this online measure of aggression (classified under Frustrative Nonreward construct within the Negative Valence Systems domain).

### Neural correlates of aggression in schizophrenia

Magnetic resonance imaging (MRI) studies of aggression specifically in schizophrenia have been relatively limited.<sup>30–32</sup> Most studies of the neurobiology of aggression have been based on findings from studies examining mixed diagnostic populations, as well as animal studies that concluded that aggression is associated with dysfunctions in ventral prefrontal and temporolimbic regions, as well as the interaction of these regions.<sup>33</sup>

### Structural MRI

Structural MRI studies have primarily focused on cortical volumetrics. Some of these studies have found that larger caudate,<sup>34</sup> left orbitofrontal volume, and right orbitofrontal white matter volume<sup>35</sup> correlated with aggression in treatment-resistant patients with schizophrenia. These findings may reflect enlargement in cortical volumes due to the iatrogenic effects of long-term treatment with first-generation neuroleptics, given that these agents can increase basal ganglia volumes.<sup>36</sup> Similar findings with regard to orbitofrontal white matter volume were found in patients with schizophrenia who had prior suicide attempts.<sup>37</sup> Narayan *et al.*<sup>38</sup> found reduced cortical thickness in the ventromedial PFC (vmPFC) and lateral sensorimotor areas in violent compared to nonviolent patients. Of the violent patients, only those with antisocial personality disorder (and not schizophrenia) showed a reduction in thickness of the vmPFC. In a diffusion tensor imaging study, Hoptman *et al.*<sup>39</sup> found that increased diffusivity, which may be reflective of atrophy, was correlated with higher levels of aggressive attitudes in men with persistent, long-term forms of schizophrenia. These studies are broadly consistent with the idea that structural abnormalities in ventral prefrontal and some subcortical regions are associated with aggression in schizophrenia.

### Functional studies

There are quite a few functional studies of violence in patients with SMI, although these studies have not necessarily been limited to patients with schizophrenia, and most of them have used radiological methods. For instance, Raine *et al.*<sup>40</sup> conducted an important series of studies using positron emission tomography to examine murderers judged not guilty by reason of insanity who were performing a continuous performance task. These studies found reduced glucose metabolism in prefrontal brain regions. Similarly, Volkow *et al.*<sup>41</sup> found reduced resting regional cerebral metabolism in prefrontal and medial temporal regions in a small sample of psychiatric patients with repetitive violence. Other radiological studies have found generally similar results.<sup>42</sup>

Task-based functional MRI (fMRI) studies of violence in schizophrenia are relatively uncommon and point to the importance of comorbid diagnoses. Joyal *et al.*<sup>43</sup> examined blood oxygen level-dependent (BOLD) responses during a go/no-go task. Patients with comorbid antisocial personality and substance use showed increased activation in motor, premotor, and anterior cingulate cortex, as well as reduced ventral prefrontal activation compared to controls and to violent patients with schizophrenia. Kumari *et al.*<sup>44</sup> used an anticipatory fear task, and found that individuals with schizophrenia and violent history showed hyperactivity in thalamostriatal regions, whereas antisocial patients showed hypoactivity in the same regions. Finally, in a small study of men with schizophrenia, Dolan and Fullam<sup>45</sup> found that patients with schizophrenia and high psychopathy had reduced amygdala activation to fearful faces compared to individuals with schizophrenia with low comorbid psychopathy.

A promising area of study is resting state fMRI. A major advantage of this method is that it avoids performance confounds and can thus be used in most study populations. A key measure using this approach is functional connectivity, which describes the temporal correlation of activation in different brain regions. In an initial study, Hoptman *et al.*<sup>46</sup> examined functional connectivity of the amygdala in patients with persistent forms of schizophrenia. Compared to healthy controls, individuals with schizophrenia showed reduced functional connectivity between the amygdala and ventral prefrontal regions. This reduction correlated with higher levels of aggression in the patients as measured using arrest histories and self-report measures.

In a more recent study, Hoptman *et al.*<sup>48</sup> examined impulsivity and aggression in patients with schizophrenia compared to healthy controls using the UPPS-P, as well as several aggression questionnaires. They found that on the UPPS-P, patients had a selective increase in urgency, and that this increase explained significant

TABLE 1. Potential neuroanatomical substrates and causes of aggression in schizophrenia

Neuroanatomical substrate	Finding(s)
Caudate and orbitofrontal cortex	Increased volume associated with higher aggression in treatment-resistant SZ <sup>33,34</sup>
Orbitofrontal white matter	Increased volume in patients with prior suicide attempts <sup>36</sup>
Ventral prefrontal cortex and lateral sensorimotor areas	Reduced cortical thickness in violent vs nonviolent patients with SZ <sup>27</sup> Reduced cortical thickness and functional connectivity of ventral prefrontal cortex associated with urgency in SZ <sup>18</sup> Reduced activation in SZ + APD + SUD during a go/no-go task compared to SZ only <sup>42</sup>
Motor, premotor, anterior cingulate cortex	Increased activation in SZ + APD + SUD during a go/no-go task compared to SZ only <sup>42</sup>
Ventral prefrontal white matter	Greater diffusivity associated with more aggression <sup>38</sup>
Thalamus and striatum	Hyperactivity during an anticipatory fear task in patients with SZ and violent history <sup>43</sup>
Amygdala	Reduced activation to fear faces in SZ with high psychopathy, <sup>44</sup> reduced resting state functional connectivity to frontal regions associated with aggression in SZ <sup>45</sup>

Note. SZ = schizophrenia, APD = antisocial personality disorder, SUD = substance use disorder.

variance in the also-observed increase in aggressive attitudes. Moreover, the investigators found that impulsive, but not premeditated, aggression was elevated in patients. Higher levels of urgency predicted reduced cortical thickness in ventral prefrontal regions, as well as reduced functional connectivity between ventral prefrontal and both limbic (e.g., rostral anterior cingulate) and executive brain regions. It is clear that a number of frontolimbic and subcortical regions are involved in impulsivity and aggression in schizophrenia. These are shown in Table 1.

### Implications

These findings may have important implications for the theoretical and clinical understanding of aggression in schizophrenia.<sup>18,47</sup> From a theoretical perspective, the findings refine our understanding of the role of impulsivity in aggression. Thus, not all impulsivity is the same, and many aspects of impulsivity are not elevated in chronic schizophrenia. On the basis of self-report measures, only the emotional components (i.e., urgency) of the construct were elevated. Moreover, the neural circuitry related to urgency involves the structural integrity of ventral prefrontal regions and their functional connections to limbic and executive regions. Dysfunction of these circuits, which are implicated in emotional regulation, may predispose an individual toward aggression.<sup>48</sup> The extant literature suggests the importance of emotional dysregulation deficits in schizophrenia.<sup>49</sup>

From a clinical perspective, the results suggest implications for treatment. Targets may include cognitive interventions to enhance regulation of impulsivity and aggression. These could include techniques such as reappraisal of negative emotional experiences, or suppression of negative affect.<sup>50</sup> It is not yet known if these interventions can be used to effectively reduce aggression in patients with schizophrenia. However, their use, including techniques related to acceptance of

emotional experiences,<sup>51</sup> might lessen the impact of distress associated with strong emotional states, and thereby prevent urgency-related aggression.

The literature also points to neural targets. Some antipsychotic medications appear to have specific anti-aggressive effects, particularly clozapine.<sup>52,53</sup> Although second-generation antipsychotic agents have mixed effects on neurotransmitters, they do affect serotonin systems. Abnormalities in these systems have been associated with violent behavior in a number of populations. Of relevance to aggressive behavior, serotonin transporter sites are selectively reduced in ventral prefrontal regions in suicide victims compared to those with depression only.<sup>54</sup> It would be important to examine serotonin transporter distribution patterns in violent patients with schizophrenia. Moreover, it would be interesting to examine the effects of clozapine on urgency in patients with schizophrenia. In the study by Hoptman *et al.*,<sup>18</sup> patients taking clozapine had lower urgency ratings on the UPPS-P than patients taking other medications. Clearly, a better definition of the multi-dimensional construct of aggression may lead to better treatment decisions and outcomes.<sup>55</sup>

Other treatment approaches also may be useful to reduce impulsivity-based aggression. Transcranial magnetic stimulation (TMS) has traditionally been unable to reach cortical depths of greater than 2 cm, thereby rendering it unable to stimulate ventral prefrontal regions. However, we can apply our knowledge from big data approaches such as the Functional Connectomes Project<sup>56</sup> to identify circuits that are functionally connected to ventral prefrontal regions. Thus, it may be efficacious to stimulate regions such as the dorsolateral prefrontal cortex or precuneus in order to modulate activation in ventral prefrontal regions. Alternatively, deep TMS has the potential to reach depths of up to 6 cm from the cortical surface. This technique has shown treatment utility in a number of disorders involving emotional dysregulation, including major depression, hallucinations and delusions



in schizophrenia, bipolar depression, and Asperger's syndrome.<sup>57</sup> Neither of these TMS approaches has been specifically applied to impulsive aggression in schizophrenia, but such methods may prove useful from both mechanistic and treatment perspectives.

Another potentially interesting approach is the use of real-time fMRI neurofeedback<sup>58</sup> to modulate the neural circuitry related to impulsivity and aggression. Real-time fMRI has been applied to the "default mode network" (DMN<sup>59</sup>)—a set of brain regions, including medial prefrontal cortex, posterior cingulate, and lateral parietal regions, that typically shows increased activity during rest, compared to task, conditions. Studies on DMN suggest its involvement in self-directed cognition.<sup>60</sup> Real-time fMRI has had limited application in schizophrenia so far.<sup>61</sup> In principle, one could identify critical circuitry for urgency and train patients to modulate their own circuitry. Recent studies show that orbitofrontal circuitry can be manipulated in this manner in healthy subjects with contamination anxiety,<sup>62</sup> and that anterior insula circuitry can be modulated in criminal psychopaths,<sup>63</sup> which suggests that this approach might be applicable in other populations with abnormalities in such circuitry.

As has been stressed in this article, aggression has multiple causes, and a complete understanding of aggression in schizophrenia will extend the knowledge of impulsive, psychotic, and psychopathically based aggression, as well as interactions among these 3 types of aggression. For example, an important literature is emerging on the role of persecutory delusions in aggression in schizophrenia.<sup>64</sup> Significantly, it appears that delusions that engender anger are those that are most likely to lead to aggression,<sup>65</sup> suggesting a mechanism whereby urgency could interact with psychotic symptoms to yield violent behavior. Finally, the circuitry underlying psychotic, impulsive, and predatory violence likely differs, and an understanding of these distinct circuits and their interactions will inform our ability to understand the genesis of aggression in schizophrenia.<sup>4</sup>

It should be noted that I am not arguing that all aspects of aggression in schizophrenia are determined by urgency. However, given the role that urgency may well play in aggression in schizophrenia, as well as the fact that at least a plurality of inpatient aggression is impulsive in nature, it may be that an assessment of urgency would be useful prior to intervention. Moreover, as noted above, it could be that other causal factors might interact with urgency so that changes in urgency might have as-yet-unknown effects on those factors. It would thus be important to know to what extent treatments typically used to address impulsively based aggression work through the construct of urgency and/or via related circuitry.

More generally, the literature on impulsivity and violence in schizophrenia points to the need for a better

understanding of the regulation of emotionally based impulsivity and aggression in this disorder, as well as the need to better understand the regulation of other problematic behaviors associated with emotionally based impulsivity. This conceptualization places the problem in the context of a phenomenon that we know to be an issue transdiagnostically,<sup>66</sup> in keeping with the NIMH's RDoC mission. It will be important to use information on regulation of impulsivity and aggression more broadly to help understand how it plays out in schizophrenia and other disorders.

### Challenges

There are several important challenges in this line of work. First, there is no known paradigm that ideally probes urgency, which makes it difficult to objectively evaluate this construct in imaging studies. It may be that urgency has such strong trait properties that it cannot be easily modified. Alternatively, behavioral tasks that are typically used to measure impulsivity may have to be adapted to include a strong emotional component in order to better investigate this construct. Even if urgency cannot be easily manipulated, initial approaches might include (a) finding tasks that are closely related to urgency that can be modulated and/or (b) modulating urgency-related circuitry with the goal of examining its downstream effects on impulsively based aggression,

Paradigms such as the TAP or PSAP might be promising. Although the PSAP is part of the RDoC matrix, the task requires substantial motor activity and thus might be difficult to implement in fMRI studies. The TAP has been used in some fMRI studies. In particular, Dambacher *et al.*<sup>67</sup> used the go/no-go task to measure response inhibition and the TAP to measure reactive aggression in healthy men. They found that both failed response inhibition (compared to go trials) and reactive aggression were associated with activation of the anterior insula, suggesting a role for that region in self-control. This same group showed that theta burst TMS in the right anterior insula and superior frontal cortex leads to impairments in both action restraint (go/no-go task) and action cancellation (stop task) in the former, and in action restraint in the latter compared to sham stimulation.<sup>68</sup> By identifying regions indirectly stimulated by TMS in these studies, a better understanding of the circuitry associated with reactive aggression, action restraint, and action cancellation could be derived.

### Conclusion

This article focuses on impulsive aggression. In general, imaging studies point to the role of abnormalities in circuitry involving ventral prefrontal, medial temporal,

and subcortical regions as playing a key role in these behaviors. The recent finding that urgency plays a special role in aggression in schizophrenia suggests several novel approaches to better understand and treat these dysfunctional behaviors, and these approaches may have utility for other problematic behaviors associated with urgency, as well. Nonetheless, aggression in schizophrenia is not homogeneous across individuals, and a complete understanding of these behaviors will consider this heterogeneity. Newer assessments and guidelines for inpatient aggression are explicitly doing just that.<sup>69,70</sup> By improving this understanding, we will be better equipped to manage aggressive behavior in schizophrenia, leading to reductions in harm, trauma, healthcare costs, and stigma.

## Disclosures

Matthew Hoptman has the following disclosures: NIMH, researcher, research support (grants); Kessler Foundation, consultant, consulting fees.

## REFERENCES:

- McEvoy JP. The costs of schizophrenia. *J Clin Psychiatry*. 2007; **68**(Suppl 14): 4-7.
- Torrey EF. Stigma and violence: isn't it time to connect the dots? *Schizophr Bull*. 2011; **37**(5): 892-896.
- Volavka J, Citrome L. Heterogeneity of violence in schizophrenia and implications for long-term treatment. *Int J Clin Pract*. 2008; **62**(8): 1237-1245.
- Stahl SM. Deconstructing violence as a medical syndrome: mapping psychotic, impulsive, and predatory subtypes to malfunctioning brain circuits. *CNS Spectr*. 2014; **19**(5): 357-365.
- Witt K, van Dorn R, Fazel S. Risk factors for violence in psychosis: systematic review and meta-regression analysis of 110 studies. *PLoS One*. 2013; **8**(2): e55942.
- Nolan KA, Czobor P, Roy BB, et al. Characteristics of assaultive behavior among psychiatric inpatients. *Psychiatr Serv*. 2003; **54**(7): 1012-1016.
- Dodge KA, Lochman JE, Harnish JD, Bates JE, Pettit GS. Reactive and proactive aggression in school children and psychiatrically impaired chronically assaultive youth. *J Abnorm Psychol*. 1997; **106**(1): 37-51.
- McDermott BE, Holoyda BJ. Assessment of aggression in inpatient settings. *CNS Spectr*. 2014; **19**(05): 425-431.
- Nolan KA, Volavka J, Mohr P, Czobor P. Psychopathy and violent behavior among patients with schizophrenia or schizoaffective disorder. *Psychiatr Serv*. 1999; **50**(6): 787.
- Patton JH, Stanford MS, Barratt ES. Factor structure of the Barratt impulsiveness scale. *J Clin Psychol*. 1995; **51**(6): 768-774.
- Nolan KA, D'Angelo D, Hoptman MJ. Self-report and laboratory measures of impulsivity in patients with schizophrenia or schizoaffective disorder and healthy controls. *Psychiatry Res*. 2011; **187**(1-2): 301-303.
- Barratt ES. Impulsivity: integrating cognitive, behavioral, biological, and environmental data. In: McCowan W, Johnson J, Shure M, eds. *The Impulsive Client: Theory, Research, and Treatment*. American Psychological Association; 1993: 39-56.
- Lipszyc J, Schachar R. Inhibitory control and psychopathology: a meta-analysis of studies using the stop signal task. *J Int Neuropsychol Soc*. 2010; **16**(6): 1064-1076.
- Chambers CD, Garavan H, Bellgrove MA. Insights into the neural basis of response inhibition from cognitive and clinical neuroscience. *Neurosci Biobehav Rev*. 2009; **33**(5): 631-646.
- Enticott PG, Ogloff JRP, Bradshaw JL, Daffern M. Contrary to popular belief, a lack of behavioural inhibitory control may not be associated with aggression. *Crim Behav Ment Health*. 2007; **17**(3): 179-183.
- Whiteside SP, Lynam DR. The five factor model and impulsivity: using a structural model of personality to understand impulsivity. *Personality and Individual Differences*. 2001; **30**(4): 669-689.
- Cyders MA, Smith GT. Mood-based rash action and its components: positive and negative urgency. *Personality and Individual Differences*. 2007; **43**(4): 839-850.
- Hoptman MJ, Antonius D, Mauro CJ, Parker EM, Javitt DC. Cortical thinning, functional connectivity, and mood-related impulsivity in schizophrenia: relationship to aggressive attitudes and behavior. *Am J Psychiatry*. 2014; **171**(9): 943-948.
- Cyders MA, Smith GT. Emotion-based dispositions to rash action: positive and negative urgency. *Psychol Bull*. 2008; **134**(6): 807-828.
- Cyders MA, Dzemidzic M, Eiler WJ, Coskunpinar A, Karyadi KA, Kareken DA. Negative urgency mediates the relationship between amygdala and orbitofrontal cortex activation to negative emotional stimuli and general risk-taking. *Cereb Cortex*. doi:10.1093/cercor/bhu123.
- Enticott PG, Ogloff JR, Bradshaw JL. Response inhibition and impulsivity in schizophrenia. *Psychiatry Res*. 2008; **157**(1): 251-254.
- Enticott PG, Ogloff JR, Bradshaw JL. Associations between laboratory measures of executive inhibitory control and self-reported impulsivity. *Personality and Individual Differences*. 2006; **41**(2): 285-294.
- Pawliczek CM, Derntl B, Kellermann T, Kohn N, Gur RC, Habel U. Inhibitory control and trait aggression: neural and behavioral insights using the emotional stop signal task. *Neuroimage*. 2013; **79**: 264-274.
- Green MF, Lee J. Neural bases of emotional experience versus perception in schizophrenia. *Biol Psychiatry*. 2012; **71**(2): 96-97.
- Insel TR, Cuthbert BN, Garvey MA, et al. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiatry*. 2010; **167**(7): 748-751.
- Yudofsky SC, Silver JM, Jackson W, Endicott J, Williams D. The Overt Aggression Scale for the objective rating of verbal and physical aggression. *Am J Psychiatry*. 1986; **143**(1): 35-39.
- Buss AH, Perry M. The aggression questionnaire. *J Pers Soc Psychol*. 1992; **63**(3): 452-459.
- Taylor SP. Aggressive behavior and physiological arousal as a function of provocation and the tendency to inhibit aggression. *J Pers*. 1967; **35**(2): 297-310.
- Cherek DR, Moeller FG, Dougherty DM, Rhoades H. Studies of violent and nonviolent male parolees: II. Laboratory and psychometric measurements of impulsivity. *Biol Psychiatry*. 1997; **41**(5): 523-529.
- Soyka M. Neurobiology of aggression and violence in schizophrenia. *Schizophr Bull*. 2011; **37**(5): 913-920.
- Hoptman MJ, Antonius D. Neuroimaging correlates of aggression in schizophrenia: an update. *Curr Opin Psychiatry*. 2011; **24**(2): 100-106.
- Weiss EM. Neuroimaging and neurocognitive correlates of aggression and violence in schizophrenia. *Scientifica*. 2012; **2012**: 158646.
- Volavka J. *Neurobiology of Violence*, 2nd ed. Washington, DC: American Psychiatric Association; 2001.

34. Hoptman MJ, Volavka J, Czobor P, *et al.* Aggression and quantitative MRI measures of caudate in patients with chronic schizophrenia or schizoaffective disorder. *J Neuropsychiatry Clin Neurosci.* 2006; **18**(4): 509–515.
35. Hoptman MJ, Volavka J, Weiss EM, *et al.* Quantitative MRI measures of orbitofrontal cortex in patients with chronic schizophrenia or schizoaffective disorder. *Psychiatry Res.* 2005; **140**(2): 133–145.
36. Chakos MH, Lieberman JA, Bilder RM, *et al.* Increase in caudate nuclei volumes of first-episode schizophrenic patients taking antipsychotic drugs. *Am J Psychiatry.* 1994; **151**(10): 1430–1436.
37. Rüsich N, Spoletini I, Wilke M, *et al.* Inferior frontal white matter volume and suicidality in schizophrenia. *Psychiatry Res.* 2008; **164**(3): 206–214.
38. Narayan VM, Narr KL, Kumari V, *et al.* Regional cortical thinning in subjects with violent antisocial personality disorder or schizophrenia. *Am J Psychiatry.* 2007; **164**(9): 1418–1427.
39. Hoptman MJ, Volavka J, Johnson G, Weiss E, Bilder RM, Lim KO. Frontal white matter microstructure, aggression, and impulsivity in men with schizophrenia: a preliminary study. *Biol Psychiatry.* 2002; **52**(1): 9–14.
40. Raine A, Buchsbaum MS, Stanley J, Lottenberg S, Abel L, Stoddard J. Selective reductions in prefrontal glucose metabolism in murderers. *Biol Psychiatry.* 1994; **36**(6): 365–373.
41. Volkow ND, Tancredi LR, Grant C, *et al.* Brain glucose metabolism in violent psychiatric patients: a preliminary study. *Psychiatry Res.* 1995; **61**(4): 243–253.
42. Wong MTH, Fenwick PBC, Lumsden J, *et al.* Positron emission tomography in male violent offenders with schizophrenia. *Psychiatry Res.* 1997; **68**(2): 111–123.
43. Joyal C, Putkonen A, Mancini-Marie A, *et al.* Violent persons with schizophrenia and comorbid disorders: a functional magnetic resonance imaging study. *Schizophr Res.* 2007; **91**(1): 97–102.
44. Kumari V, Das M, Taylor PJ, *et al.* Neural and behavioural responses to threat in men with a history of serious violence and schizophrenia or antisocial personality disorder. *Schizophr Res.* 2009; **110**(1): 47–58.
45. Dolan MC, Fullam RS. Psychopathy and functional magnetic resonance imaging blood oxygenation level-dependent responses to emotional faces in violent patients with schizophrenia. *Biol Psychiatry.* 2009; **66**(6): 570–577.
46. Hoptman MJ, D'Angelo D, Catalano D, *et al.* Amygdalofrontal functional disconnectivity and aggression in schizophrenia. *Schizophr Bull.* 2010; **36**(5): 1020–1028.
47. Szeszko PR. Aggression in schizophrenia and its relationship to neural circuitry of urgency. *Am J Psychiatry.* 2014; **171**(9): 897–900.
48. Davidson RJ, Putnam KM, Larson CL. Dysfunction in the neural circuitry of emotion regulation—a possible prelude to violence. *Science.* 2000; **289**(5479): 591–594.
49. Kring AM, Werner KH. Emotion regulation and psychopathology. In: Philippot P, Feldman RS, eds. *The Regulation of Emotion.* Mahwah, NJ: Lawrence Erlbaum Associates; 2004: 359–385.
50. Gross JJ, Jazaieri H. Emotion, emotion regulation, and psychopathology: an affective science perspective. *Clinical Psychological Science.* 2014; **2**(4): 387–401.
51. Perry Y, Henry JD, Nangle MR, Grisham JR. Regulation of negative affect in schizophrenia: the effectiveness of acceptance versus reappraisal and suppression. *J Clin Exp Neuropsychol.* 2012; **34**(5): 497–508.
52. Krakowski MI, Czobor P, Citrome L, Bark N, Cooper TB. Atypical antipsychotic agents in the treatment of violent patients with schizophrenia and schizoaffective disorder. *Arch Gen Psychiatry.* 2006; **63**(6): 622–629.
53. Volavka J, Czobor P, Nolan K, *et al.* Overt aggression and psychotic symptoms in patients with schizophrenia treated with clozapine, olanzapine, risperidone, or haloperidol. *J Clin Psychopharmacol.* 2004; **24**(2): 225–228.
54. Arango V, Underwood MD, Mann JJ. Serotonin brain circuits involved in major depression and suicide. *Prog Brain Res.* 2002; **136**: 443–453.
55. Citrome L, Volavka J. The psychopharmacology of violence: making sensible decisions. *CNS Spectr.* 2014; **19**(5): 411–418.
56. Biswal BB, Mennes M, Zuo XN, *et al.* Toward discovery science of human brain function. *Proc Natl Acad Sci U S A.* 2010; **107**(10): 4734–4739.
57. Bersani F, Minichino A, Enticott P, *et al.* Deep transcranial magnetic stimulation as a treatment for psychiatric disorders: a comprehensive review. *Eur Psychiatry.* 2013; **28**(1): 30–39.
58. LaConte SM, Peltier SJ, Hu XP. Real-time fMRI using brain-state classification. *Hum Brain Mapp.* 2007; **28**(10): 1033–1044.
59. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A.* 2001; **98**(2): 676–682.
60. Gusnard DA, Akbudak E, Shulman GL, Raichle ME. Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc Natl Acad Sci U S A.* 2001; **98**(7): 4259–4264.
61. Ruiz S, Lee S, Soekadar SR, *et al.* Acquired self-control of insula cortex modulates emotion recognition and brain network connectivity in schizophrenia. *Hum Brain Mapp.* 2013; **34**(1): 200–212.
62. Hampson M, Stoica T, Saksa J, *et al.* Real-time fMRI biofeedback targeting the orbitofrontal cortex for contamination anxiety. *J Vis Exp.* 2012; (59): 3535.
63. Sitaram R, Caria A, Veit R, Gaber T, Ruiz S, Birbaumer N. Volitional control of the anterior insula in criminal psychopaths using real-time fMRI neurofeedback: a pilot study. *Front Behav Neurosci.* 2014; **8**: 344.
64. Keers R, Ullrich S, DeStavola BL, Coid JW. Association of violence with emergence of persecutory delusions in untreated schizophrenia. *Am J Psychiatry.* 2014; **171**(3): 332–339.
65. Coid JW, Ullrich S, Kallis C, *et al.* The relationship between delusions and violence: findings from the East London first episode psychosis study. *JAMA Psychiatry.* 2013; **70**(5): 465–471.
66. Johnson SL, Carver CS, Joormann J. Impulsive responses to emotion as a transdiagnostic vulnerability to internalizing and externalizing symptoms. *J Affect Disord.* 2013; **150**(3): 872–878.
67. Dambacher F, Sack AT, Lobbstaël J, Arntz A, Brugman S, Schuhmann T. Out of control evidence for anterior insula involvement in motor impulsivity and reactive aggression. *Soc Cogn Affect Neurosci.* doi:10.1093/scan/nsu077.
68. Dambacher F, Sack AT, Lobbstaël J, Arntz A, Brugmann S, Schuhmann T. The role of right prefrontal and medial cortex in response inhibition: Interfering with action restraint and action cancellation using transcranial magnetic brain stimulation. *J Cogn Neurosci.* 2014; **26**(8): 1775–1784.
69. Warburton K. The new mission of forensic mental health services: managing violence as a medical syndrome in an environment that balances treatment and safety. *CNS Spectr.* 2014; **19**(5): 368–373.
70. Stahl SM, Morrisette DA, Cummings M, *et al.* California State Hospital Violence Assessment and Treatment (Cal-VAT). *CNS Spectr.* 2014; **19**(5): 449–465.