

- 2 Baker AB, Knutson J. Psychiatric aspects of uremia. *Am J Psychiatry* 1946; **102**: 683–7.
- 3 Menninger KA. Paranoid psychosis with uremia. *J Nerv Ment Disord* 1924; **60**: 26–34.
- 4 Beal DM, Hudson B, Zaiac M. Amoxicillin-induced psychosis? *Am J Psychiatry* 1986; **143**: 255–6.
- 5 Nordgaard J, Arnfred SM, Handest P, Parnas J. The diagnostic status of first-rank symptoms. *Schizophr Bull* 2008; **34**: 137–54.

**Johannes Huber**, Department of Urology, University of Heidelberg, Germany; **Christian G. Huber**, Department of Psychiatry and Psychotherapy, University Medical Centre Hamburg-Eppendorf, Martinistr. 52, D-20246 Hamburg, Germany. Email: c.huber@uke.uni-hamburg.de; **Martin Lambert**, Department of Psychiatry and Psychotherapy, University Medical Centre Hamburg-Eppendorf, Hamburg; **Markus Hohenfellner**, Department of Urology, University of Heidelberg, Germany

doi: 10.1192/bjp.196.6.496

## Correction

Early intervention in panic: pragmatic randomised controlled trial. *BJP*, 196, 326–331. Table 1 (p. 329): the ranges given for PDSS–SR, HADS–Anxiety, Mobility Inventory and BID–II are the theoretical ranges of these scales.

doi: 10.1192/bjp.196.6.497