

A systematic review of the psychometric properties, usability and clinical impacts of mobile mood-monitoring applications in young people

M. Dubad^{1*}, C. Winsper¹, C. Meyer^{1,2}, M. Livanou¹ and S. Marwaha^{1,3}

¹Mental Health and Wellbeing, Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, UK

²Warwick Manufacturing Group, University of Warwick, Coventry, UK

³Affective Disorders Service, Caludon Centre, Coventry, UK

Background. Mobile mood-monitoring applications are increasingly used by mental health providers, widely advocated within research, and a potentially effective method to engage young people. However, little is known about their efficacy and usability in young populations.

Method. A systematic review addressing three research questions focused on young people: (1) what are the psychometric properties of mobile mood-monitoring applications; (2) what is their usability; and (3) what are their positive and negative clinical impacts? Findings were synthesised narratively, study quality assessed and compared with evidence from adult studies.

Results. We reviewed 25 articles. Studies on the psychometric properties of mobile mood-monitoring applications were sparse, but indicate questionable to excellent internal consistency, moderate concurrent validity and good usability. Participation rates ranged from 30% to 99% across studies, and appeared to be affected by methodological factors (e.g. payments) and individual characteristics (e.g. IQ score). Mobile mood-monitoring applications are positively perceived by youth, may reduce depressive symptoms by increasing emotional awareness, and could aid in the detection of mental health and substance use problems. There was very limited evidence on potential negative impacts.

Conclusions. Evidence for the use of mood-monitoring applications in youth is promising but limited due to a lack of high-quality studies. Future work should explicate the effects of mobile mood-monitoring applications on effective self-regulation, clinical outcomes across disorders and young people's engagement with mental health services. Potential negative impacts in this population should also be investigated, as the adult literature suggests that application use could potentially increase negativity and depression symptoms.

Received 2 September 2016; Revised 19 May 2017; Accepted 23 May 2017; First published online 23 June 2017

Key words: Ecological momentary assessment, mental health, mhealth, mobile application, mood, mood monitoring, young people.

Introduction

Mood is an affective dynamic, which naturally varies across time and contexts (Trull *et al.* 2015). Problems with regulating mood can play a key role in the development and trajectory of a range of psychopathologies (Paris, 2004; Crowell *et al.* 2009; Marwaha *et al.* 2015). Traditionally, mood has been assessed with retrospective measures (Trull *et al.* 2015). This can increase the risk of recall bias subsequently reducing accuracy (Schwartz *et al.* 1999; Reid *et al.* 2009). The relatively

recent use of ecological momentary assessment (EMA) facilitates the real-time assessment of mood by collecting data on multiple occasions throughout the day (Wenze & Miller, 2010). Thus, it may be more suitable for understanding daily mood changes (Cristobal-Narvaez *et al.* 2016; Myin-Germeys *et al.* 2016; van Knippenberg *et al.* 2016).

Various EMA techniques exist, ranging from paper-and-pencil to physiological assessment (Wenze & Miller, 2010) to digital data collection. A number of UK governmental reports (HM Government, 2011; Department of Health, 2013) highlight the benefits of digital tools and Information and Communications Technology (ICT) in aiding the objective, reliable assessment and care of mental health problems. With demand for mental health services outgrowing available resources (Department of Health, 2013), technology might relieve some of this pressure by providing

* Address for correspondence: M. Dubad, BSc, MRes, Mental Health and Wellbeing, Division of Health Sciences, Warwick Medical School, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK.

(Email: m.dubad@warwick.ac.uk)

remote resources that increase access to effective treatment while reducing clinician load.

Applications ('apps') offer great promise to young people who are disproportionately affected by mental illness or may struggle to engage with mental health services (Seko *et al.* 2014). Apps are delivered in a medium young people are familiar with. Figures from Ofcom (2015) indicate that 90% of youth between the ages of 16 and 24 own a smartphone, regardless of sociodemographic domain. Given this widespread ownership and apparent attachment to mobile technology (Ofcom, 2015), youths might feel more comfortable with assessments and treatments utilising mobile apps.

Mental health services increasingly use apps (Olf, 2015), many of which have the capacity for EMA to monitor mood (e.g. Sandstrom *et al.* 2016b). Several reviews with mainly adult studies (e.g. Donker *et al.* 2013; Naslund *et al.* 2015; Nicholas *et al.* 2015; Torous & Powell, 2015; Bakker *et al.* 2016; Faurholt-Jepsen *et al.* 2016; Walsh *et al.* 2016) have appraised evidence for the use of mood-monitoring apps.

Studies included in these reviews provide some evidence for the psychometric properties, e.g. internal consistency (Palmier-Claus *et al.* 2012) and concurrent validity (Faurholt-Jepsen *et al.* 2014) of these apps. There is also evidence for usability (Bardram *et al.* 2013). Participation rates are generally high across studies sampling adults, ranging from 65% (Depp *et al.* 2015) to 88% (Ainsworth *et al.* 2013), though Depp *et al.* (2012) reported much higher completion rates for paper and pencil compared with app measures (82.9% *v.* 42.1%). Evidence also suggests that apps may help people with mental health problems to monitor triggers (Bardram *et al.* 2013), that the capacity to convey experience can be therapeutic, and that apps could be a useful tool for improving patient-clinician communication (Palmier-Claus *et al.* 2013).

Less is known about the use of mental health apps, particularly mood-monitoring apps, in youth (10–24 years). A scoping review by Seko *et al.* (2014) suggested that mood-monitoring apps are positively perceived by youth (Matthews *et al.* 2008a), may improve treatment adherence (Matthews *et al.* 2008b) and possibly improve mental wellbeing (Kauer *et al.* 2012). While intriguing, findings were preliminary due to the low quality of available evidence (NCCMH, 2014), the small number of studies on mood-monitoring apps specifically and the limited number of apps studied ($n=2$) (NCCMH, 2014; Seko *et al.* 2014).

In summary, mood-monitoring apps offer a potentially important step change in the assessment of mood and delivery of youth mental health services. Despite this potential and the widespread advocacy for their use (e.g. Firth *et al.* 2016; Sandstrom *et al.* 2016a), there are no extant reviews examining the

psychometric properties, usability and clinical impacts of mood-monitoring apps in young populations. Therefore, a systematic review was completed to address the following research questions: (1) what are the psychometric properties of mobile mood-monitoring apps; (2) what is their usability; (3) and what are their positive and negative clinical impacts among clinical and non-clinical youth populations? Our secondary aims were to frame our findings within the adult literature, and conduct a quality assessment to examine potential sources of bias.

Method

Following a scoping review, the authors developed the protocol delineating the planned methodology. The review was conducted in adherence to this protocol, and in line with the PRISMA statement (Moher *et al.* 2009).

Information sources and search strategy

The following sources were searched: Medline, EMBASE, PsycINFO, ProQuest Dissertations & Theses, ProQuest SciTech Collection, the Association for Computing Machinery (ACM) Guide to Computing Literature and Web of Science for articles published from 2008 [the year when the first app was launched (Donker *et al.* 2013)]. Search terms were informed by previous reviews (Seko *et al.* 2014), and modified following advice from a medical librarian and field experts. The search was conducted by combining five groups of terms (see online Supplementary Table S1) relating to: type of technology (e.g. 'mhealth'), type of assessment (e.g. 'ambulatory assessment'), mood-related outcome or problem (e.g. 'bipolar disorder'), youth population (e.g. 'youth'), usability/treatment-related outcomes and psychometric properties (e.g. 'reliability', 'validity'). We were interested in all forms of validity potentially examined in the app literature, e.g. concurrent, face or predictive (Faurholt-Jepsen *et al.* 2016), though we anticipated a paucity of studies due to the novelty of the field. We defined the 'usability' of mood-monitoring apps in accordance with the International Organisation for Standardisation (2001) definition of usability, i.e. 'the capability of the software product to be understood, learned, used and attractive to the user, when used under specified conditions'. Consistent with previous systematic reviews (Donker *et al.* 2013), we included young people's participation rates (i.e. compliance, response and completion) and how apps were perceived by youths (including their acceptability – how satisfied they were with the app, whether it could be used with ease) as markers of usability.

MD conducted a hand search of articles published in *Cyberpsychology, Behavior and Social Network*, the *Journal of Medical Internet Research (JMIR)*, the *JMIR Mental Health*, and the *JMIR mHealth and uHealth* over the last 5 years. An additional search of the first 15 pages of Google Scholar was conducted (search terms 'mood', 'phone', 'app' and 'monitoring'). Reference lists and in-text citations of relevant articles were inspected. Finally, subject experts were approached to identify additional articles.

Study selection

Inclusion criteria were:

- (1) Apps must have been developed for, and delivered through, mobile phones or smartphones;
- (2) Participants aged 10–24 years (consistent with the World Health Organisation's definition of young people; World Health Organisation, 1986);
- (3) Studies included published and unpublished research reported in the grey literature;
- (4) Studies must have been published in the English language;
- (5) Studies must have been published in 2008 or later;
- (6) Studies must have included community or clinical populations (to ensure the inclusion of sub-clinical youth, who may subsequently access care).

Screening procedure

Following removal of duplicates, MD and ML independently screened 100% of titles and abstracts for full-text retrieval. MD assessed full-text articles against the inclusion criteria and extracted relevant data.

Quality assessment

MD evaluated the quality of included studies for potential risk of bias using Cochrane's risk of bias tool, in which studies are allocated a rating of high, low or unclear risk of bias (Higgins et al. 2011).

Data synthesis

Quantitative and qualitative data were synthesised narratively.

Results

Study selection

A total of 1747 articles were identified in the initial search, and 19 from the hand search (Fig. 1). Following removal of duplicates, 1176 abstracts were screened, 86 of which were selected for full-text retrieval. There was a high level of agreement between

raters ($\kappa=0.90$). In total, 64 articles were excluded following full-text review. Three additional articles were identified following inspection of included studies. Twenty-five articles were included in the final review.

Study characteristics

Table 1 outlines study methodology, the characteristics and features assessed in the studies, and main findings. Three studies reported on a randomised controlled trial (RCT): one was the primary RCT (Reid et al. 2011), and two reported secondary analyses with the same dataset (Kauer et al. 2012; Reid et al. 2013). The remaining studies were non-experimental or quasi-experimental. The search identified 19 published studies and six unpublished studies (four conference proceedings; two theses). The majority of studies ($n=16$) were quantitative; the remaining nine employed mixed methods.

Sample size ranged from 6 to 1 08 996 participants. Eight studies recruited healthy participants. Eleven studies recruited participants from clinical populations including youth with a range of mental health, emotional or behavioural problems, such as depression ($n=8$), high-functioning autism/Asperger's disorder ($n=2$) and substance or alcohol use ($n=1$). The remaining six studies recruited participants from mixed populations comprising healthy, mentally ill or substance-using individuals. Mean ages across studies ranged from 10.95 to 23.7 years.

Methods across studies varied greatly. For example, some studies lent participants a phone, whereas others let participants use their own device. Please see Table 1 for a description of the different data collection methods used in each study. As observed in the adult literature, terminology also varied greatly across studies (please see Usability section for more details).

Various apps were used, the most frequent of which was the 'Mobiletype' programme (Reid et al. 2009). Mood outcomes were either direct mood assessments, or described mood-related constructs or behaviours (e.g. stress, hostility). Outcomes were monitored over variable time periods. The shortest period was 24 h (Bossmann et al. 2013), the longest 326 days (Matthews & Doherty, 2011). Monitoring schedules also varied, and could comprise hourly, daily or weekly monitoring, or requirements to complete measures a fixed number of times per day (with or without pre-specified time intervals). Reimbursements or incentives were available in 18 studies (e.g. payments, gift vouchers).

Psychometric properties of mood-monitoring apps

Nine studies reported on the reliability or validity of mood-monitoring apps.




Fig. 1. Flowchart of literature search results and selection of studies.

Reliability

The internal consistency (correlation between items within a scale) was assessed in four studies (Dunton *et al.* 2011, 2014; Huh *et al.* 2014; Ansell *et al.* 2015). As demonstrated in Table 2, levels ranged from questionable to excellent (George & Mallery, 2003).

Validity

Concurrent validity. Three studies examined concurrent validity (the correlation between an assessment and a previously validated assessment of the same construct). Concurrent validity was mostly moderate across studies (see Table 1). Khor *et al.* (2014a) compared relationships

between participant and parent-reported data from the retrospective Responses to Stress Questionnaire (Connor-Smith *et al.* 2000) and mobile app data recording participants' responses to stress. In two studies of university students, Ben-Zeev *et al.* (2015) and Wang *et al.* (2014) compared momentary app and retrospective questionnaire data on perceived stress.

Face validity. Two studies described participants' views on the face validity of the 'Mobiletype' app (see Table 1 for numerical details). Reid *et al.* (2012), using a sample with various mental health problems, found that the app was relatively successful in capturing participants' feelings and current situation. Khor *et al.* (2014a), using

Table 1. Study details including the author (year,) study purpose, sample characteristics, intervention details and a summary of the main findings

Author (year)	Study purpose	Sample characteristics	Intervention ^a	Main findings
Ansell <i>et al.</i> (2015)	To explore the effects of marijuana use on impulsivity and hostility in everyday life using smartphone-based EMA	<ul style="list-style-type: none"> • Sample size: $N = 43$ ($M = 23.7$ years) • Population type: young recreational substance users • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: not specified • Operation system: not specified • Accessibility: no web/general/app store access • Device: not specified • Measurements: daily alcohol, tobacco and marijuana use; daily impulsivity and daily interpersonal hostility • Monitoring period: 14 days, monitoring schedule varied. Compliance monitored for irregularities by research staff • Incentive/reimbursement: payments + bonus payment for 95% survey response rate 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • Reliability: acceptable to excellent internal consistency^b <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: impulsivity: 96% completed data; interpersonal interactions: >99% completed data <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for problems with (perceived) interpersonal hostility
Bachmann <i>et al.</i> (2015)	To examine the usability and unobtrusiveness of the Mobile Ambulatory Mood Assessment (MoA ²) app	<ul style="list-style-type: none"> • Sample size: $N = 9$ ($M = 23.4$ years) • Population type: healthy/non-clinical participants • Comparison/control: none • Location: Germany • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: MoA² • Operation system: Android • Accessibility: no web/general/app store access • Device: participants used study phones (Google Nexus 4) or personal Android smartphone • Measurements: mood, tiredness and stress level • Monitoring period: 12 prompts p/day for 4 days • Incentive/reimbursement: no payment 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • Participants' perception: app perceived as user-friendly and convenient <p>Clinical impacts: not studied/reported</p>
Ben-Zeev <i>et al.</i> (2015) ^c	To examine if smartphone sensor data can be used to measure behaviour and mental health	<ul style="list-style-type: none"> • Sample size: $N = 47$ ($M = 22.5$ years) • Population type: students reporting varying levels of depression symptoms • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: StudentLife • Operation system: Android • Accessibility: no web/general/app store access • Device: participants were offered an Android study smartphone – type not specified • Measurements: momentary stress and automated sensor data • Monitoring period: 10 weeks (sensor data gathered automatically; stress ratings completed daily, 5 days per week) • Incentive/reimbursement: (Raffle) prizes 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • Concurrent validity: significant moderate relationship between averaged app-assessed stress ratings and retrospective post-study questionnaire scores on a measure of perceived stress ($r = 0.41$, $p < 0.01$) <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: average weekly response rate was 4.92 days a week (98.4%) <p>Clinical impacts: not studied/reported</p>

Bossmann <i>et al.</i> (2013)	To clarify the relationship between everyday physical activity and affective states over a 1-day period	<ul style="list-style-type: none"> • Sample size: $N = 62$ ($M = 21.4$ years) • Population type: healthy/non-clinical students • Comparison/control: none • Location: Germany • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: MyExperience movisens Edition version 594 • Operation system: Android • Accessibility: web/general access only • Device: participants were provided with an HTC Touch 2 smartphone • Measurements: valence, calmness and energetic arousal • Monitoring period: 1 day – affect measurements every hour after waking up • Incentive/reimbursement: no payment 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> <ul style="list-style-type: none"> ○ Mean completion rate was 10.5 electronic diaries per participant ○ Please note that 15 participants were excluded for missing data <p>Clinical impacts: not studied/reported</p>
Crooke <i>et al.</i> (2013)	To examine the relationship between varying rates of alcohol use and positive and negative mood through EMA	<ul style="list-style-type: none"> • Sample size: $N = 41$ ($M = 15.4$ years) • Population type: young people with varying levels of alcohol intake • Comparison/control: none • Location: Australia • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobiletype • Operation system: not specified • Accessibility: no web/general/app store access • Device: participants were lent a Nokia 6630 • Measurements: activities, company, location, mood, responses to stressful events and coping, and questions on participants' previous evening's alcohol and cannabis use • Monitoring period: 4× p/day on 20 randomised days over the 31-day study period • Incentive/reimbursement: Partial reimbursement/ gift voucher (value: \$25) 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> 58.3% (AM diaries) and 43.8% (PM diaries) completed mood assessments <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for youth alcohol interventions
Dennis <i>et al.</i> (2015)	To assess the feasibility of smartphone-based EMA and recovery support ecological momentary interventions (EMI) via smartphones. The study also assessed the feasibility of using EMA and EMI to predict substance use in the following week	<ul style="list-style-type: none"> • Sample size: $N = 29$ ($M = 16.6$ years) • Population type: adolescents with different clinical problems • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Addiction Comprehensive Health Enhancement Support System (ACHESS) • Operation system: Android • Accessibility: web/general access only • Device: participants provided with a smartphone – type not specified • Measurements: feelings, activities, location and social context, and drug and alcohol related measurements • Monitoring period: 6× p/day for 6 weeks. Compliance monitored for irregularities by research staff • Incentive/reimbursement: payment –up to \$50 per week for adherence to all study requirements 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> 89% of assessments completed • <i>Participants' perception:</i> <ul style="list-style-type: none"> ○ App-based EMA perceived as 'not too long' (95%), 'very easy' or 'easy to learn how to do' (100%), and 'very easy' or 'easy to complete six EMAs per day' (94%) ○ Of note, one participant withdrew early from the study due to frustrations with software problems <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for relapse prevention
Dunton <i>et al.</i> (2014) ^d	Using EMA to bi-directionally explore how affective and physical feeling states are associated with physical activity	<ul style="list-style-type: none"> • Sample size: $N = 119$ ($M = 10.95$ years) • Population type: children with varying body mass index (BMI) levels 	<ul style="list-style-type: none"> • App name: MyExperience • Operation system: Windows • Accessibility: web/general access only • Device: participants were lent an HTC Shadow. 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • <i>Reliability:</i> acceptable to good internal consistency^b

Table 1 (cont.)

Author (year)	Study purpose	Sample characteristics	Intervention ^a	Main findings
Dunton <i>et al.</i> (2011) ^d	To assess if the level and experience of children's leisure-time physical activity vary with social and physical contexts by means of EMA	<ul style="list-style-type: none"> • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • Measurements: main activity type, social context, physical location, mood and enjoyment • Monitoring period: monitoring period: 3–7 random prompts p/day within pre-specified times over two data collection waves (duration: 4 days per wave), separated by 6 months • Incentive/reimbursement: up to \$40 (compensatory) payment • App name: MyExperience • Operation system: Windows • Accessibility: web/general access only • Device: participants were lent an HTC Shadow. • Measurements: main activity type, social context, physical location, mood and enjoyment • Monitoring period: 3–7 random prompts p/day within pre-specified times over 4 days • Incentive/reimbursement: up to \$40 (compensatory) payment 	<p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: 76% of assessments completed on average <p>Clinical impacts: not studied/reported</p> <p>Psychometric properties:</p> <ul style="list-style-type: none"> • Reliability: acceptable to good internal consistency^b <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: 80.3% of assessments completed on average <p>Clinical impacts: not studied/reported</p>
Huh <i>et al.</i> (2014)	To examine the contextual antecedents to smoking in a sample of Korean American young adult smokers through EMA	<ul style="list-style-type: none"> • Sample size: $N = 22$ ($M = 21.23$ years) • Population type: young adult smokers • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: ActiPal (MEI Ltd.) • Operation system: Android • Accessibility: web/general access only (demo app) • Measurements: affect, perceived stress, cigarette craving, and other contextual and environmental measures • Device: Android enabled phones (study phones provided if participants owned iPhones) • Monitoring period: random non-smoking signal contingent (5× p/day for 7 days) + event-contingent prompts over a 7 day period. Compliance closely monitored by research staff • Incentive/reimbursement: not reported 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • Reliability: questionable to acceptable internal consistency^b <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: 92.4% of assessments completed on average • Participants' perception: it should be noted that one participant withdrew from the study due to technical difficulties with the EMA app <p>Clinical impacts: not studied/reported</p>

Kauer <i>et al.</i> (2012) ^e	A secondary analysis that investigated the relationships between self-monitoring, emotional self-awareness, and depression through EMA	<ul style="list-style-type: none"> • Sample size: $N = 69$ ($M = 18.5$ years) • Population type: young people with mild or more severe mental health/emotional problems • Comparison/control: attention comparison ($n = 49$, $M = 17.4$ years). • Location: Australia • Data collection: In-person research 	See Reid <i>et al.</i> (2011)	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> completion rates were 52.9% for the intervention group and 59.6% for the comparison group <p>Clinical impacts: implications for depression symptoms</p>
Kauer <i>et al.</i> (2009) ^f	To assess the feasibility and usefulness of a mobile phone-based EMA app to gather information on alcohol use and related behaviours	<ul style="list-style-type: none"> • Sample size: $N = 18$ [mean ages 15.9 years (females) and 15.8 years (males)] in study 1; $n = 6$ [mean ages 18.3 years (females) and 19.5 years (males)] in study 2 • Population type: healthy/non-clinical students in study 1 and high-risk drinkers in study 2 • Comparison/control: none • Location: Australia • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobiletype • Operation system: not specified • Accessibility: no web/general/app store access • Device: participants were lent a Nokia 6630 • Measurements: activity, mood, stress, alcohol and cannabis use • Monitoring period: 4× p/day for 1 week • Incentive/reimbursement: partial reimbursement/gift voucher (value: \$25) 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> better compliance for school-based adolescents than older adolescent high-risk drinkers <p>Clinical impacts: not studied /reported</p>
Kenny <i>et al.</i> (2015)	To assess the feasibility of the CopeSmart app	<ul style="list-style-type: none"> • Sample size: $N = 43$ ($M = 16.0$ years) • Population type: healthy/non-clinical adolescents • Comparison/control: none • Location: Ireland • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: CopeSmart • Operation system: Android + iOS • Accessibility: no web/general/app store access • Device: app was downloaded on participants' Android or iOS phones • Measurements: happiness, anger, sadness, stress and worries • Monitoring period: 1 week • Incentive/reimbursement: no monetary incentive 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> participants engaged with the app on 4 out of 7 days (57.1%) • <i>Participants' perception:</i> the app's interface layout was liked by 79% of participants. Furthermore, the app was perceived as easy to use (93%); minor technical difficulties with logging on were experienced by 7% of participants; 70% of participants would use the app in the future; 74% believed the app would be used by other young people; and 70% would recommend the app to a friend <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Implications for self-awareness
Khor <i>et al.</i> (2014a) ^g	To assess the utility of the Mobiletype programme to examine adolescents with High-Functioning Autism/Asperger's Disorder's (HFASD) stressors and coping	<ul style="list-style-type: none"> • Sample size: $N = 31$ ($M = 14.46$ years) + parents • Population type: adolescents with HFASD • Comparison/control: none • Location: Australia • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobiletype (adapted) • Operation system: not specified • Accessibility: no web/general/app store access • Device: participants were lent a Sony Ericson 7501i • Measurements: mood, stress, last time and daily stress • Monitoring period: 4× p/day for 2 weeks 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • <i>Concurrent validity:</i> <ul style="list-style-type: none"> ○ Mostly poor to moderate correlations between data from the retrospective Responses to Stress Questionnaire (Connor-Smith <i>et al.</i> 2000) and mobile app data recording participants' responses to stress

Table 1 (cont.)

Author (year)	Study purpose	Sample characteristics	Intervention ^a	Main findings
			<ul style="list-style-type: none"> • Incentive/reimbursement: partial reimbursement (value: \$20) 	<ul style="list-style-type: none"> ○ A significant moderate to strong correlation for the 'involuntary engagement' factor: $r = 0.70$, $p < 0.01$; parent report: $r = 0.48$, $p < 0.01$ ○ A significant strong correlation for the 'primary control engagement coping' factor: $r = 0.53$, $p < 0.05$ • Face validity: <ul style="list-style-type: none"> ○ The face validity was measured by assessing how well the app captured participants' current situation, thoughts and feelings ○ The highest ratings were reported for the app's ability to capture participants' feelings (67%); followed by its ability to capture participants' current situation (63%); and finally its ability to measure participants' thoughts (50%) <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: participants responded to 61.8% of prompts <ul style="list-style-type: none"> ○ Note that a substantial proportion of participants gradually stopped responding throughout the study; while every participant completed at least one entry on the first day, completion rates reduced to 45% on day 14 ○ Also note that there was a significant positive correlation between full scale IQ and compliance rates ($r = 0.46$, $p < 0.01$) <p>Clinical impacts: not studied/reported Psychometric properties: not studied Usability: not studied/reported Clinical impacts:</p> <ul style="list-style-type: none"> • Implications for emotional and behavioural problems <p>Psychometric properties: not studied Usability:</p> <ul style="list-style-type: none"> • Participation rate: different compliance rates across app features –no obvious pattern. Mean normalised compliance for daily registrations of approximately 30%; mean
Khor et al. (2014b) [§]	To investigate how daily hassles, coping, and behaviour and emotional problems are related in adolescents with HFASD	See Khor et al. (2014a)	See Khor et al. (2014a)	
Loventoft et al. (2012)	To find out whether people treated for depression would be interested in using a smartphone app for support in their daily lives	<ul style="list-style-type: none"> • Sample size: $N = 6$ (ages 17–24, no means reported) • Population type: young people with recent depression treatment • Comparison/control: none • Location: Denmark 	<ul style="list-style-type: none"> • App name: Daybuilder • Operation system: Android • Accessibility: no web/general/app store access • Device: participants provided with Android device with app installed 	

		<ul style="list-style-type: none"> • Data collection: in-person research 	<ul style="list-style-type: none"> • Measurements: Weekly Major Depression Inventory; daily mood, appetite and sleep • Monitoring period: 4 weeks • Incentive/reimbursement: payment of 500 DKK (\$95 or 2 h salary) 	<p>normalised compliance for weekly registrations of approximately 50%</p> <ul style="list-style-type: none"> • <i>Participants' perception:</i> user experience negatively affected by technological difficulties; clinicians highlighted the usefulness of self-monitoring when combined with therapy. <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Implications for treatment <p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> 65% response on average <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Implications for treatment
Matthews & Doherty (2011)	To assess the issues around the use of mobile phones for mood charting with the aim to improve adolescent engagement	<ul style="list-style-type: none"> • Sample size: $N = 9$ ($M = 13.78$ years) • Population type: young people with depression, mood disorders, self-harm and anger management • Comparison/control: none • Location: Ireland • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobile Mood Diary (MMD) • Operation system: not specified • Accessibility: no web/general/app store access • Device: app downloaded on clients' phones • Measurements: energy, sleep and mood + free area for thought entries • Monitoring period: min. 1× p/day for two sessions • Incentive/reimbursement: reimbursement where necessary 	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> mobile group significantly more responsive than paper-diary group ($t = -2.324, p < 0.05$) • <i>Participants' perception:</i> participants preferred mobile technology <p>Clinical impacts: not studied/reported</p>
Matthews <i>et al.</i> (2008b)	To explore the effectiveness of mobile phone <i>v.</i> pen-and-paper for mood tracking	<ul style="list-style-type: none"> • Sample size: $N = 73$ ($M = 14.87$ years) • Population type: healthy/non-clinical students • Comparison/control: paper-based diary condition ($n = 52$) • Location: Ireland • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: MMD • Operation system: not specified • Accessibility: no web/general/app store access • Device: app downloaded on students' phones • Measurements: energy, sleep and mood + free area for thought entries • Monitoring period: 1× p/day for 2 weeks • Incentive/reimbursement: none <p>See Kauer <i>et al.</i> (2009)</p>	<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> <ul style="list-style-type: none"> ○ Participants' completed 76% of diaries ○ However, response rates decreased from 91% on day 1 to 67% on day 7 ○ Of note, one-third of the sample stated that they did not always respond honestly to items if a specific response would result in further questioning • <i>Participants' perception:</i> the study's initial response rate suggested mobile technology may not be preferred or adopted by all young people. Nevertheless, the app was
Reid <i>et al.</i> (2009) ^f	A study aimed at developing, piloting and reviewing a youth focused mobile phone programme to track young people's experiences in real time	<ul style="list-style-type: none"> • Sample size: focus group ($n = 11$, mean age not reported) and pilot study [males ($n = 5, M = 15.8$ years) and females ($n = 13, M = 15.9$ years)] • Population type: students • Comparison/control: none • Location: Australia • Data collection: in-person research 		<p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> <ul style="list-style-type: none"> ○ Participants' completed 76% of diaries ○ However, response rates decreased from 91% on day 1 to 67% on day 7 ○ Of note, one-third of the sample stated that they did not always respond honestly to items if a specific response would result in further questioning • <i>Participants' perception:</i> the study's initial response rate suggested mobile technology may not be preferred or adopted by all young people. Nevertheless, the app was

Table 1 (cont.)

Author (year)	Study purpose	Sample characteristics	Intervention ^a	Main findings
Reid <i>et al.</i> (2011) ^e	A randomised controlled trial to investigate some of the mental health benefits of the Mobiletype programme	<ul style="list-style-type: none"> • Sample size: $N = 68$ ($M = 18.5$ years) • Population type: mild/more mental health or emotional problems • Comparison/control: comparison programme ($n = 46$, $M = 17.4$ years) • Location: Australia • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobiletype • Operation system: not specified • Accessibility: no web/general/app store access • Device: participants were lent a Sony Ericsson Z750i mobile phone • Measurements: current activities, company, location, mood, recent stressful events, responses to stressful events, alcohol consumption, cannabis use, and sleep, exercise and diet-related questions • Monitoring period: min. 2×/day for 2–4 weeks • Incentive/reimbursement: partial reimbursement (A\$30) and gift cards (A\$20) for post-questionnaires completion (maximum A\$60) 	<p>overall viewed as youth-friendly and non-invasive</p> <p>Clinical impacts: not studied/reported</p> <p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> response rates for the intervention group: 52.9%; comparison group: 60.9% <p>Clinical impacts:</p> <ul style="list-style-type: none"> • No significant effects on mental health outcomes; potential implications for self-awareness
Reid <i>et al.</i> (2013) ^e	To assess the utility of Mobiletype in a primary care setting (secondary analysis)	See Reid <i>et al.</i> (2011)	See Reid <i>et al.</i> (2011)	<p>Psychometric properties: not studied</p> <p>Usability: not studied/reported</p> <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for treatment and clinicians' understanding of patients
Reid <i>et al.</i> (2012)	To review Mobiletype in clinical settings	<ul style="list-style-type: none"> • Sample size: $n = 47$ ($M = 15.59$ years) • Mental health/clinical status: adolescents with varied (medical) disorders. • Comparison/control: none • Location: Australia • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: Mobiletype • Operation system: not specified • Accessibility: no web/general/app store access • Device: participants were lent a ZTE F851 JAVA MIDP 2.0 phone with \$50 credit • Measurements: location, activity, company, mood, stressful events, responses to stressful events, alcohol and cannabis use, sleep, exercise and diet-related questions • Monitoring period: four random prompts p/day for 2–4 weeks (min. completion: 1× p/day) 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • <i>Face validity:</i> <ul style="list-style-type: none"> ○ The face validity was measured by assessing how well the app captured participants' current situation, thoughts and feelings ○ The highest ratings were reported for the app's ability to capture participants' feelings (86%); followed by its ability to capture participants' current situation (83%); and finally its ability to measure participants' thoughts (57%)

Sacco (2015)	To examine the feasibility and utility of a smartphone app developed to assess five areas of functioning associated with depression	<ul style="list-style-type: none"> • Sample size: $N = 114$ ($M = 19.36$ years) • Population type: students with varying levels of depression symptoms • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • Incentive/reimbursement: none • App name: Android Health and Wellness UDTracker App • Operation system: Android • Accessibility: no web/general/app store access • Device: app installed on participants' own Android enabled phones • Measurements: depression, mood, social functioning, cognitive and lifestyle factors, coping/emotion regulation (daily or weekly) • Monitoring period: 14 days. Assessment times varied across measures: 1× p/evening [e.g. Positive and Negative Affect Scale (Watson <i>et al.</i> 1988)], 1× p/morning [sleep questionnaire, adapted from Pittsburgh Sleep Quality Index (Buysse <i>et al.</i> 1989)], and 1× p/week [e.g. items from the COPE scale (Carver <i>et al.</i> 1989)] • Incentive/reimbursement: extra/research participation course credit 	<p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> Participants completed 91% of entries in week 1 <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for assessment and management <p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participation rate:</i> 85–93% response rate across different measures • <i>Participants' perception:</i> <ul style="list-style-type: none"> ○ App perceived as 'easy to use' (95.6%); 'a little' to 'not at all' irritating (90.3%) ○ The monotony of responding to the same survey questions (15%); the high frequency of the pop-up notifications (9%), and the drain on the phone's battery life (8%) were perceived as irritating. Participants suggested more varied survey questions (23%), fewer crashes, bugs or freezes (9%) and provided suggestions for novel technical features (13%) ○ Some participants also enjoyed the user-friendliness of the app (40%) and the pop-up-reminder feature (17%) <p>Clinical impacts:</p> <ul style="list-style-type: none"> • Potential implications for self-reflection on emotions or behaviours <p>Psychometric properties: not studied</p> <p>Usability:</p> <ul style="list-style-type: none"> • <i>Participants' perception:</i> preference for paper-and-pencil tracking by some participants <p>Clinical impacts: not studied/reported</p>
Scotti (2015)	To assess the efficacy, acceptability and feasibility of the school-based Dialectical Behaviour Therapy skills group for the treatment of adolescent eating disorders and sub-diagnostic problematic eating behaviours	<ul style="list-style-type: none"> • Sample size: High school students ($N = 4$, $M = 16.75$ years) and middle school students ($N = 3$, $M = 13.67$ years) • Population type: students with eating disorder symptoms or body image concerns • Comparison/control: two high school students who had withdrawn ($M = 16.5$ years) • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: not specified • Operation system: not specified • Accessibility: unknown web/general access, no app store access • Device: participants own smartphones – type not specified • Measurements: individual eating disorder-related behaviours and cognitions/feelings • Study/monitoring period: 12 weeks • Incentive/reimbursement: academic credit and/or prize draw • App name: Recovery Record • Operation system: Android + iOS 	<p>Psychometric properties: not studied</p>
Tregarthen <i>et al.</i> (2015)	To describe a smartphone app for the self-monitoring of eating disorder symptoms, evaluate characteristics of app	<ul style="list-style-type: none"> • Sample size: $N = 1\ 08\ 996$ [$M = 22$ years (reported by 48 830 users)] 	<ul style="list-style-type: none"> • App name: Recovery Record • Operation system: Android + iOS 	<p>Psychometric properties: not studied</p>

Table 1 (cont.)

Author (year)	Study purpose	Sample characteristics	Intervention ^a	Main findings
	users and assess the feasibility and utilisation of the app for self-monitoring purposes	<ul style="list-style-type: none"> • Population type: people with varying levels of ED severity • Comparison/control: none • Study Location: USA • Data collection: crowd-sourcing 	<ul style="list-style-type: none"> • Accessibility: general/web and app store access • Device: own (iOS or Android) smartphone – type not specified • Measurements: meals and eating disorder-related behaviours/cognitions/feelings/urges • Monitoring period: overall usage data not available – six monitoring prompts p/day • Incentive: none 	<p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: 89% of participants monitored ≥ 3 meals; 67% continued to monitor at 30 days • Participants' response: app received high user ratings <p>Clinical impacts: not studied/reported</p>
Wang <i>et al.</i> (2014) ^c	To measure university students' mental health, academic performance and behavioural trends using the StudentLife app	<ul style="list-style-type: none"> • Sample size: $N = 48$ ($M = 22.8$ years) • Population type: university students with varying depression scores. • Comparison/control: none • Location: USA • Data collection: in-person research 	<ul style="list-style-type: none"> • App name: StudentLife • Operation system: Android • Accessibility: no web/general/app store access • Device: participants either used their own Android phones (primary users) or were offered an Android Nexus 4a (secondary users) • Measurements: momentary mood, sleep, social, physical exercise, activity, and behaviour; automated sensor data. • Monitoring period: 10 weeks • Incentive: (Raffle) prizes 	<p>Psychometric properties:</p> <ul style="list-style-type: none"> • Concurrent validity: significant moderate relationship between averaged app-assessed stress ratings and retrospective post-study questionnaire scores on a measure of perceived stress ($r = 0.41$, $p < 0.01$) <p>Usability:</p> <ul style="list-style-type: none"> • Participation rate: response rates for participants who used own phones: 65%; response rates for participants who used study phones: 72% <p>Clinical impacts: not studied/reported</p>

^a The accessibility of mood-monitoring apps was assessed through a search of Google and three app stores (iTunes, Google Play and Microsoft store) in June 2016.

^b Please refer to Table 2 for coefficient values.

^c These studies utilised the same data.

^d These studies utilised the same data.

^e These studies utilised the same data.

^f These studies partly utilised the same data.

^g These studies utilised the same data.

Table 2. Internal consistency coefficients across studies and domains

Authors	α Coefficients									Ω Coefficients		
	Positive affect			Negative affect			Perceived stress			Impulsivity		
	O	WS	BS	O	WS	BS	O	WS	BS	O	WS	BS
Ansell <i>et al.</i> (2015)	–	–	–	–	–	–	–	–	–	–	0.78	0.96
Dunton <i>et al.</i> (2011)	0.88	–	–	0.75	–	–	–	–	–	–	–	–
Dunton <i>et al.</i> (2014)	0.87	–	–	0.74	–	–	–	–	–	–	–	–
Huh <i>et al.</i> (2014)	0.65	–	–	0.78	–	–	0.73	–	–	–	–	–

Note: O, Overall, WS, within-subject level, BS, between-subject level. Internal consistency coefficients values interpretation: '>0.9 – excellent, >0.8 – good, >0.7 – acceptable, >0.6 – questionable, >0.5 – poor and <0.5 – unacceptable' (George & Mallery, 2003, pp. 231).

a sample with high-functioning autism and Asperger's found that the app was not quite as successful in these domains. In both studies, the apps were less successful in capturing participants' thoughts.

Usability of mood-monitoring apps

Participation rates

Twenty-one studies examined participation rates, which ranged from 30% to 99%. Average percentages were not computed in four studies. Instead, these studies described the mean number of diary entries per participant (Bossmann *et al.* 2013), between-group differences (Matthews *et al.* 2008b; Kauer *et al.* 2009), or evidence of ongoing compliance (Tregarthen *et al.* 2015). There was some indication that response rates were higher in studies with incentives. For example, Dennis *et al.* (2015) offered an incentive of \$50 per week, and had a participation rate of 89% (see Table 1 for comparative rates and incentive details). Participation rates also appeared to be affected by response fatigue. In Reid *et al.* (2009), for instance, response rates decreased from 91% on day 1 to 67% on day 7. Finally, participation rates were potentially affected by sample-specific characteristics. In a study with high-functioning autistic participants, Khor *et al.* (2014a) found a significant positive correlation between full-scale IQ and compliance rates ($r = 0.46$, $p < 0.01$).

Participants' perceptions

Nine studies considered participants' perceptions of the apps. Three of these studies specifically referred to the 'acceptability' of apps. In Dennis *et al.* (2015), 95% of adolescents felt that the EMA app 'was not too long'. Tregarthen *et al.* (2015) measured app utilisation data as a proxy for acceptability. There were over 100 000 users over a 2-year period (with 89% using the

application at least three times), which the authors interpreted as a demonstration of broad acceptability. While they did not define acceptability specifically, Reid *et al.* (2009) concluded that their app was 'acceptable' based on the data they captured (e.g. completion rates, participants' feedback).

Across studies, 93–100% of respondents found apps easy to learn or use (Dennis *et al.* 2015; Kenny *et al.* 2015; Sacco, 2015). In addition, participants rated apps as useful (Kenny *et al.* 2015), convenient, user-friendly (Bachmann *et al.* 2015), youth-friendly and non-invasive (Reid *et al.* 2009). Despite these positive experiences, technological difficulties (e.g. software crashes, reduced battery life) were reported to negatively affect user experience and participation (Loventoft *et al.* 2012; Huh *et al.* 2014; Dennis *et al.* 2015; Sacco, 2015). Although most young people reported a preference for mobile phone mood charting in comparison to paper diaries (Matthews *et al.* 2008b), not all young people preferred mobile technology (Reid *et al.* 2009; Scotti, 2015). Scotti (2015), e.g. found that several participants from a sub-diagnostic eating disorder sample favoured paper-and-pencil to track their data.

Positive and negative clinical impacts of mood-monitoring apps

Mental health and awareness

Five (two were from the same RCT) studies examined potential clinical impacts of the apps. Reid *et al.* (2011) found a significant improvement in emotional self-awareness, but no significant improvements in depression, anxiety or stress scores in youth with mental health or emotional problems. In a secondary analysis of the same RCT, Kauer *et al.* (2012) reported an indirect association between app use and depression

symptoms via increased emotional self-awareness. The app, however, did not significantly reduce rumination.

Qualitative feedback from two studies also suggested that mood-monitoring apps can help improve self-awareness (Kenny *et al.* 2015), and self-reflection on emotions or behaviours (Sacco, 2015).

Though they did not test this premise directly, Ansell *et al.* (2015) hypothesised that app-based monitoring could have promoted self-awareness in participants subsequently reducing (perceived) interpersonal hostility.

In Khor *et al.* (2014b), parents rated their children with high-functioning autism as showing fewer symptoms of behaviour and emotional problems following use of the self-monitoring app.

Treatment implications

Five studies reported results that could have implications for the prevention and treatment of mental health problems. Mobile app data gathered by Dennis *et al.* (2015) were used to identify high-risk groups for substance use, which could potentially help with relapse prevention. Crooke *et al.* (2013) suggested that mood-monitoring apps could help investigate adolescents' motivations for drinking, thus informing the development of interventions.

Qualitative feedback from therapists suggests that the use of mobile apps could help facilitate engagement with participants suffering from various mental health problems (Matthews & Doherty, 2011). Reid *et al.* (2012) reported that the Mobiletype app facilitated the assessment and management of youth mental health problems and reduced consultation time with paediatricians; the data captured enabled more individually focused consultations, which assisted in rapport building and communication.

In the third of a series of papers detailing their RCT, Reid *et al.* (2013) explored the potential treatment benefits of 'Mobiletype'. In comparison to the control programme, the app significantly increased general practitioners' (GPs) understanding of their patients' health and current functioning, and aided diagnoses, communication, medication and referrals. However, there was no significant effect on doctor's confidence, doctor-patient rapport or pathways to care.

Finally, in a conference paper by Loventoft *et al.* (2012), clinicians highlighted the usefulness of self-monitoring when combined with therapy.

Quality assessment

Please see online Supplementary Fig. S1 for an overall depiction of the risk of bias domains across studies.

Risk of selection bias was difficult to assess in many studies, as they often lacked treatment, control or comparison groups. Three studies (all using the same RCT

data) were deemed at low risk of selection bias due to a clear description of the randomisation and concealment allocation process (Reid *et al.* 2011, 2013; Kauer *et al.* 2012). Two studies were at unclear risk of selection bias because randomised sequence generation and method of allocation concealment were not sufficiently described (Matthews *et al.* 2008b; Reid *et al.* 2009). One study was considered at high risk of selection bias (Scotti, 2015) as there was no random allocation process for the control condition.

Only the RCT study (three publications) addressed the blinding of participants and personnel, and was thus considered at low risk of performance bias (Reid *et al.* 2011, 2013; Kauer *et al.* 2012). The risk of detection bias in these studies was unclear due to a lack of clarity on blinding of outcome assessments.

The risk of attrition bias was difficult to ascertain in three studies. In one study (Kenny *et al.* 2015), a number of participants were not included in the final sample due to restrictions on school access (no other information was available). Bossmann *et al.* (2013) excluded 15 participants from the final sample due to 'missing data', but did not provide further information, including whether any analyses were performed to address missing data. Reid *et al.* (2012) was considered at unclear risk of attrition bias, as there was no information on the participants (21%) lost to follow-up. The remaining studies appeared to be at low risk of attrition bias. There was insufficient information to assess the risk of reporting bias in all studies but those of the RCT, which addressed pre-specified outcomes and appeared to be at low risk (Reid *et al.* 2011, 2013; Kauer *et al.* 2012). All studies appeared to be at unclear or high risk of other types of bias.

Discussion

The aim of this review was to summarise and evaluate evidence for the use of mobile mood-monitoring apps in young people (aged 10–24 years) from clinical and non-clinical populations. We specifically focused on psychometric properties, usability and clinical impacts.

Psychometric properties of mood-monitoring apps

Few studies assessed psychometric properties. There was limited evidence for reliability, with four studies demonstrating questionable to excellent levels of internal consistency. Studies examining concurrent ($n=3$) and face ($n=2$) validity were also sparse, making it difficult to draw firm conclusions. Face validity findings, e.g. could have been moderated by sample characteristics, e.g. reduced insight in participants with autism (Khor *et al.* 2014a).

The limited assessment of psychometric properties observed in the youth literature mirrors the adult literature. Evidence for concurrent validity in adult populations is inconclusive (Depp *et al.* 2012; Palmier-Claus *et al.* 2012; Faurholt-Jepsen *et al.* 2014). Inconsistent methodology across these studies, e.g. momentary (Depp *et al.* 2012) *v.* retrospective assessments (Faurholt-Jepsen *et al.* 2014), varying periods between the event and participants' recollection of the event (Palmier-Claus *et al.* 2012), likely contribute to variable findings. Previous evidence suggests that real-time mood measurement methods (e.g. EMA) only have a modest correlation with retrospective assessments, such as questionnaires (Ebner-Priemer & Trull, 2009). This leads to the conceptual question of whether retrospective measures are the most appropriate comparators when assessing the validity of mood-monitoring apps. Questionnaires measure an individual's retrospective view of their mood state over a number of days. While they are subject to recall bias, this bias incorporates other emotional processing (e.g. contexts) that the more instantaneous assessment of mood (e.g. EMA) may not capture, or at least as richly. Thus, the two assessment methods may be measuring different types of affective experience. As it is difficult to draw robust conclusions about the validity of apps using retrospective assessments, future studies should further examine psychometric properties using other sources of comparative data, e.g. active smartphone app data (i.e. app assessments) with passive sensor smartphone data (Nicholas *et al.* 2015; Sandstrom *et al.* 2016b), associations with clinical rating scales (Faurholt-Jepsen *et al.* 2016).

Usability of mood-monitoring apps

The usability of mood-monitoring apps was more extensively studied, and overall studies suggest that apps are usable for young people. However, there were some within- and between-study differences in participants' perceptions of apps, and participation rates.

Generally, participation rates were lower in studies where participants had mental health difficulties (Reid *et al.* 2011; Kauer *et al.* 2012), problematic drinking patterns (Kauer *et al.* 2009) or autism spectrum disorders – especially those with lower IQ (Khor *et al.* 2014a). In particular, participation levels were low for those living without set routines (Kauer *et al.* 2009). This is an important consideration, as youths with mood-related problems, e.g. borderline personality disorder, often have disorganised daily routines (Fleischer *et al.* 2012). This suggests a need to tailor apps for different clinical populations (Kauer *et al.* 2009).

Some studies indicated that incentives could positively influence participation rates (e.g. Ansell *et al.* 2015; Dennis *et al.* 2015). It may not be financially feasible to offer incentives in non-research settings. However, results tentatively suggest that participation rates may be better for mobile apps than traditional paper-based assessments irrespective of incentives (Matthews *et al.* 2008b). Participation rates for paper-based diaries are as low as 11% (Stone *et al.* 2003) compared with 30–99% for mood-monitoring apps in the current review. This supports that apps could lead to better adherence rates than non-digital assessment tools in young populations. Factors that could improve participation rates include the use of less intensive assessments (e.g. once-daily rather than multiple times), shorter assessments and the incorporation of staff monitoring or automatic reminders (Huh *et al.* 2014).

Studies from the adult literature are somewhat congruent in supporting the usability of mood-monitoring apps (Bardram *et al.* 2013), though evidence suggests that increasing age (e.g. 'middle age') may lower likelihood of mood-monitoring app use (Depp *et al.* 2012). Both adult (Palmier-Claus *et al.* 2013) and adolescent (Bradford & Rickwood, 2014) populations expressed some reservations about using apps due to the perceived risk of reduced personal contact (Palmier-Claus *et al.* 2013).

Overall our review demonstrated that young people positively perceive apps (Reid *et al.* 2009) and would be willing to use this technology in real-life settings (Kenny *et al.* 2015; Tregarthen *et al.* 2015). Very few studies considered clinician perspectives on mood-monitoring apps. Matthews & Doherty (2011) found that therapists' confidence with technology was the biggest barrier to the use of mood apps. More qualitative studies are now needed to further explore young peoples' (and clinicians') perceptions (Hollis *et al.* 2016) to broaden our understanding of factors pertinent to the uptake of mood-monitoring apps in real-life settings.

Positive and negative clinical impacts of mood-monitoring apps

Few of the included studies assessed the clinical impacts of the mood-monitoring apps. Although evidence was generally positive (e.g. facilitating assessment, management and GPs' understanding), most studies relied on subjective participant feedback (Sacco, 2015) rather than RCT methodology with objective outcome measures.

The preliminary evidence (Kauer *et al.* 2012) very tentatively suggests that electronic mood-monitoring apps could function as an intervention tool (Seko

et al. 2014; Olf, 2015; Faurholt-Jepsen et al. 2016). Intriguingly, results from the one RCT indicated that mood-monitoring apps might reduce depression in youths by increasing their levels of emotional awareness (Kauer et al. 2012). Similarly, though in a non-experimental study, Khor et al. (2014b) reported that self-monitoring improved parent-reported behavioural and emotional problems in participants with autism. While these results are promising, they require replication and future studies may further explore the mechanisms via which apps could potentially impact on clinical outcomes. One possibility is that mood apps could have a positive impact on clinical symptoms due to patient/participant expectations regarding their benefits. This phenomenon, coined the digital placebo effect, is an overlooked area, which also merits future investigation (Torous & Firth, 2016).

We were unable to fully examine the potential negative impacts of mood-monitoring apps in youth populations, as they were not directly investigated in studies. However, Reid et al. (2009) found that participants did not always respond to questions truthfully to avoid having to answer further questions. Thus, this type of assessment could potentially lead to the inaccurate assessment (and treatment) of mental health problems.

A small number of adult studies report on the negative effects of mood-monitoring apps. There is some suggestion that apps may increase negative reactivity (Ainsworth et al. 2013), increase focus on negative symptoms and thoughts (Palmier-Claus et al. 2013), and potentially maintain depressive symptoms (Faurholt-Jepsen et al. 2015). Given the evidence from the adult literature, research on the possible harmful effects of app use in youths is needed before these tools are routinely used in clinical practice. Part of this endeavour should seek to identify the optimal balance between a monitoring schedule, which accurately captures affective dynamic processes, while minimising respondent workload (Bolger et al. 2003; Trull et al. 2015). This is particularly important, not only because it affects participation rates, but also because the responsibility of self-monitoring could impose a burden on young people (Shiffman et al. 2008), might result in unnecessary pressure (Lupton, 2013; Seko et al. 2014) and exacerbate mental health problems (Conner & Reid, 2012; Faurholt-Jepsen et al. 2015).

Future work may investigate potential ethical issues surrounding the use of mood-monitoring apps. For example, their use could lead to an over-reliance on technology in young populations, which could exacerbate mental health problems (Thomé et al. 2011). There could also be information security-related risks (e.g. digital theft) that could compromise confidentiality (Prentice & Dobson, 2014). Finally, youths could

use apps as a replacement for treatment and health monitoring (Tregarthen et al. 2015). Considering the importance of the therapeutic alliance for successful treatment outcomes (Karver et al. 2006), the efficacy of smartphone apps could be reduced if they are used without clinicians' involvement (Prentice & Dobson, 2014).

Strengths and limitations

As far as we are aware, this is the first review to systematically examine and quality assess the evidence for the psychometric properties, usability and clinical outcomes of mood-monitoring apps in youth. However, our results should be considered through the lens of a number of limitations.

First, despite undertaking a comprehensive search, there were very few high-quality studies available for inclusion in the review. There was only one primary RCT highlighting the need for more trials on the efficacy of mood-monitoring apps in young people. Indeed, our quality assessment indicated that the majority of studies included some form of bias. For example, many studies were at high or unclear risk of sampling (e.g. self-selected samples) and attrition bias. This could have affected the generalisability of our findings or led to an overestimation of positive effects, e.g. our findings may only apply to individuals with less severe psychopathology who are more likely to engage with services.

Second, studies demonstrated a great variability in terminology (especially for implementation outcomes, e.g. acceptability) making interpretations and cross-study comparisons difficult (inconsistent terminology is also a common feature of the adult app literature). For example, we found that 'acceptability' was defined very differently across studies, ranging from proxy markers, i.e. utilisation data (Tregarthen et al. 2015) to participants' experience of burden (Dennis et al. 2015). This highlights the need for more careful delineation and measurement of implementation outcomes in future work (Proctor et al. 2011).

Third, there were large variations in samples and methodologies, again making cross-study comparisons difficult and quantitative synthesis (i.e. meta-analysis) impossible. Thus, some of our conclusions remain tentative pending further rigorous, higher quality research (e.g. RCTs).

Fourth, it should be noted that studies in this review often used apps that were specifically developed for the study, and therefore not publically available through app platforms (e.g. iTunes). Thus, there is a need for more research to assess the evidence for apps that are freely downloaded and used by youth,

and whether their use can be incorporated into clinical care (Nicholas *et al.* 2015).

Clinical and research implications

Mood-monitoring apps could potentially have positive effects in both clinical and sub-clinical youth populations. Indeed, mood-monitoring apps may help youth identify and address burgeoning mental health and substance use problems (Dennis *et al.* 2015), and possibly utilise more adaptive coping strategies (Kauer *et al.* 2012). Further research is needed to examine the effects of these apps in samples with serious mental disorders, such as bipolar disorder (Grunerbl *et al.* 2015), borderline personality disorder (Lederer *et al.* 2014) and psychosis (Ben-Zeev *et al.* 2014; Palmier-Claus *et al.* 2014).

Evidence, though limited, suggests that mood-monitoring apps could potentially aid diagnosis and treatment decision-making (Reid *et al.* 2013). Future studies should explore whether this technology could aid in the assessment of disorders that can be difficult to differentiate [e.g. borderline personality disorder, bipolar disorder (Yen *et al.* 2015)] by providing rich data about the timing and extent of mood fluctuations.

As technological innovations have been endorsed at a government level, integrating mood-monitoring apps within mental health services may improve access and relieve some of the strain these services are currently experiencing [e.g. by improving access to mental health treatment (Department of Health, 2013)]. However, to date, the potential positive and negative impacts of apps have not been sufficiently investigated in youth.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717001659>.

Acknowledgements

MD is funded by an Economic and Social Research Council (ESRC) Collaborative Award Studentship – ES/J500203/1. Funding for open access publication was kindly provided by Research Councils UK (RCUK).

References

Ainsworth J, Palmier-Claus JE, Machin M, Barrowclough C, Dunn G, Rogers A, Buchan I, Barkus E, Kapur S, Wykes T, Hopkins RS, Lewis S (2013). A comparison of two delivery modalities of a mobile phone-based assessment for serious mental illness: native smartphone application vs

text-messaging only implementations. *Journal of Medical Internet Research* **15**, e60.

- Ansell EB, Laws HB, Roche MJ, Sinha R (2015). Effects of marijuana use on impulsivity and hostility in daily life. *Drug and Alcohol Dependence* **148**, 136–142.
- Bachmann A, Klebsattel C, Budde M, Riedel T, Beigl M, Reichert M, Santangelo P, Ebner-Priemer U (2015). How to use smartphones for less obtrusive ambulatory mood assessment and mood recognition. In *Adjunct Proceedings of the 2015 ACM International Joint Conference on Pervasive and Ubiquitous Computing and Proceedings of the 2015 ACM International Symposium on Wearable Computers*, pp. 693–702. ACM: Osaka.
- Bakker D, Kazantzis N, Rickwood D, Rickard N (2016). Mental health smartphone apps: review and evidence-based recommendations for future developments. *Journal of Medical Internet Research Mental Health* **3**, e7.
- Bardram JE, Frost M, Szántó K, Faurholt-Jepsen M, Vinberg M, Kessing LV (2013). Designing mobile health technology for bipolar disorder: a field trial of the monarca system. In *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*, pp. 2627–2636. ACM: Paris.
- Ben-Zeev D, Brenner CJ, Begale M, Duffecy J, Mohr DC, Mueser KT (2014). Feasibility, acceptability, and preliminary efficacy of a smartphone intervention for schizophrenia. *Schizophrenia Bulletin* **40**, 1244–1253.
- Ben-Zeev D, Scherer EA, Wang R, Xie H, Campbell AT (2015). Next-generation psychiatric assessment: using smartphone sensors to monitor behavior and mental health. *Psychiatric Rehabilitation Journal* **38**, 218–226.
- Bolger N, Davis A, Rafaeli E (2003). Diary methods: capturing life as it is lived. *Annual Review of Psychology* **54**, 579–616.
- Bossmann T, Kanning M, Koudela-Hamila S, Hey S, Ebner-Priemer U (2013). The association between short periods of everyday life activities and affective states: a replication study using ambulatory assessment. *Frontiers in Psychology* **4**, 102.
- Bradford S, Rickwood D (2014). Adolescent's preferred modes of delivery for mental health services. *Child and Adolescent Mental Health* **19**, 39–45.
- Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ (1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research* **28**, 193–213.
- Carver CS, Scheier MF, Weintraub JK (1989). Assessing coping strategies: a theoretically based approach. *Journal of Personality and Social Psychology* **56**, 267–283.
- Conner TS, Reid KA (2012). Effects of intensive mobile happiness reporting in daily life. *Social Psychological and Personality Science* **3**, 315–323.
- Connor-Smith JK, Compas BE, Wadsworth ME, Thomsen AH, Saltzman H (2000). Responses to stress in adolescence: measurement of coping and involuntary stress responses. *Journal of Consulting and Clinical Psychology* **68**, 976–992.
- Cristobal-Narvaez P, Sheinbaum T, Ballestri S, Mitjavila M, Myin-Germeys I, Kwapil TR, Barrantes-Vidal N (2016). Impact of adverse childhood experiences on psychotic-like symptoms and stress reactivity in daily life in nonclinical young adults. *PLoS ONE* **11**, e0153557.

- Crooke AH, Reid SC, Kauer SD, McKenzie DP, Hearps SJ, Khor AS, Forbes AB (2013). Temporal mood changes associated with different levels of adolescent drinking: using mobile phones and experience sampling methods to explore motivations for adolescent alcohol use. *Drug and Alcohol Review* 32, 262–268.
- Crowell SE, Beauchaine TP, Linehan MM (2009). A biosocial developmental model of borderline personality: elaborating and extending Linehan's theory. *Psychological Bulletin* 135, 495–510.
- Dennis ML, Scott CK, Funk RR, Nicholson L (2015). A pilot study to examine the feasibility and potential effectiveness of using smartphones to provide recovery support for adolescents. *Substance Abuse* 36, 486–492.
- Department of Health (2013). Annual Report of the Chief Medical Officer 2013, Public Mental Health Priorities: Investing in the Evidence. (https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/413196/CMO_web_doc.pdf).
- Depp CA, Ceglowski J, Wang VC, Yaghouti F, Mausbach BT, Thompson WK, Granholm EL (2015). Augmenting psychoeducation with a mobile intervention for bipolar disorder: a randomized controlled trial. *Journal of Affective Disorders* 174, 23–30.
- Depp CA, Kim DH, de Dios LV, Wang V, Ceglowski J (2012). A pilot study of mood ratings captured by mobile phone versus paper-and-pencil mood charts in bipolar disorder. *Journal of Dual Diagnosis* 8, 326–332.
- Donker T, Petrie K, Proudfoot J, Clarke J, Birch MR, Christensen H (2013). Smartphones for smarter delivery of mental health programs: a systematic review. *Journal of Medical Internet Research* 15, e247.
- Dunton GF, Huh J, Leventhal AM, Riggs N, Hedeker D, Spruijt-Metz D, Pentz MA (2014). Momentary assessment of affect, physical feeling states, and physical activity in children. *Health Psychology* 33, 255–263.
- Dunton GF, Liao Y, Intille S, Wolch J, Pentz MA (2011). Physical and social contextual influences on children's leisure-time physical activity: an ecological momentary assessment study. *Journal of Physical Activity and Health* 8 (Suppl. 1), S103–S108.
- Ebner-Priemer UW, Trull TJ (2009). Ecological momentary assessment of mood disorders and mood dysregulation. *Psychological Assessment* 21, 463–475.
- Faurholt-Jepsen M, Frost M, Ritz C, Christensen EM, Jacoby AS, Mikkelsen RL, Knorr U, Bardram JE, Vinberg M, Kessing LV (2015). Daily electronic self-monitoring in bipolar disorder using smartphones – the MONARCA I trial: a randomized, placebo-controlled, single-blind, parallel group trial. *Psychological Medicine* 45, 2691–2704.
- Faurholt-Jepsen M, Frost M, Vinberg M, Christensen EM, Bardram JE, Kessing LV (2014). Smartphone data as objective measures of bipolar disorder symptoms. *Psychiatry Research* 217, 124–127.
- Faurholt-Jepsen M, Munkholm K, Frost M, Bardram JE, Kessing LV (2016). Electronic self-monitoring of mood using IT platforms in adult patients with bipolar disorder: a systematic review of the validity and evidence. *BioMed Central Psychiatry* 16, 1.
- Firth J, Torous J, Yung AR (2016). Ecological momentary assessment and beyond: the rising interest in e-mental health research. *Journal of Psychiatric Research* 80, 3–4.
- Fleischer M, Schäfer M, Coogan A, Häßler F, Thome J (2012). Sleep disturbances and circadian CLOCK genes in borderline personality disorder. *Journal of Neural Transmission* 119, 1105–1110.
- George D, Mallery P (2003). *SPSS for Windows Step by Step: A Simple Guide and Reference, 11.0 Update*. Allyn & Bacon: Boston.
- Grunerbl A, Muaremi A, Osmani V, Bahle G, Ohler S, Troster G, Mayora O, Haring C, Lukowicz P (2015). Smartphone-based recognition of states and state changes in bipolar disorder patients. *Institute of Electrical and Electronics Engineers Journal of Biomedical and Health Informatics* 19, 140–148.
- Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savović J, Schulz KF, Weeks L, Sterne JAC, Cochrane Bias Methods Group, Cochrane Statistical Methods Group (2011). The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 343, d5928.
- HM Government (2011). No Health without Mental Health: a Cross-Government Mental Health Outcomes Strategy for People of All Ages. (<https://www.rcpsych.ac.uk/PDF/No%20Health%20without%20Mental%20Health.pdf>).
- Hollis C, Falconer CJ, Martin JL, Whittington C, Stockton S, Glazebrook C, Davies EB (2016). Annual research review: digital health interventions for children and young people with mental health problems – a systematic and meta-review. *Journal of Child Psychology and Psychiatry* 58, 474–503.
- Huh J, Shin H, Leventhal AM, Spruijt-Metz D, Abramova Z, Cerrada C, Hedeker D, Dunton G (2014). Momentary negative moods and being with friends precede cigarette use among Korean American emerging adults. *Nicotine and Tobacco Research* 16, 1248–1254.
- ISO/IEC 9126-1 (2001). *Software Engineering – Product Quality – Part 1: Quality Model*. International Organization for Standardization: Geneva.
- Karver MS, Handelsman JB, Fields S, Bickman L (2006). Meta-analysis of therapeutic relationship variables in youth and family therapy: the evidence for different relationship variables in the child and adolescent treatment outcome literature. *Clinical Psychology Review* 26, 50–65.
- Kauer SD, Reid SC, Crooke AHD, Khor A, Hearps SJ, Jorm AF, Sanci L, Patton G (2012). Self-monitoring using mobile phones in the early stages of adolescent depression: randomized controlled trial. *Journal of Medical Internet Research* 14, e67.
- Kauer SD, Reid SC, Sanci L, Patton GC (2009). Investigating the utility of mobile phones for collecting data about adolescent alcohol use and related mood, stress and coping behaviours: lessons and recommendations. *Drug and Alcohol Review* 28, 25–30.
- Kenny R, Dooley B, Fitzgerald A (2015). Feasibility of 'CopeSmart': a telemental health app for adolescents. *Journal of Medical Internet Research Mental Health* 2, e22.
- Khor AS, Gray KM, Reid SC, Melvin GA (2014a). Feasibility and validity of ecological momentary assessment in

- adolescents with high-functioning autism and Asperger's disorder. *Journal of Adolescence* **37**, 37–46.
- Khor AS, Melvin GA, Reid SC, Gray KM** (2014b). Coping, daily hassles and behavior and emotional problems in adolescents with high-functioning autism/Asperger's disorder. *Journal of Autism and Developmental Disorders* **44**, 593–608.
- Lederer N, Grechenig T, Baranyi R** (2014). unCUT: bridging the gap from paper diary cards towards mobile electronic monitoring solutions in borderline and self-injury. In *2014 IEEE 3rd International Conference on Serious Games and Applications for Health (SeGAH)*, pp. 1–8. IEEE: Rio de Janeiro.
- Loventoft PK, Norregaard LB, Frokjaer E** (2012). Designing daybuilder: an experimental app to support people with depression. In *Proceedings of the 12th Participatory Design Conference: Exploratory Papers, Workshop Descriptions, Industry Cases – vol. 2*, pp. 1–4. ACM: Roskilde.
- Lupton D** (2013). Quantifying the body: monitoring and measuring health in the age of mHealth technologies. *Critical Public Health* **23**, 393–403.
- Marwaha S, Balbuena L, Winsper C, Bowen R** (2015). Mood instability as a precursor to depressive illness: a prospective and mediational analysis. *Australian and New Zealand Journal of Psychiatry* **49**, 557–565.
- Matthews M, Doherty G** (2011). In the mood: engaging teenagers in psychotherapy using mobile phones. In *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*, pp. 2947–2956. ACM: Vancouver.
- Matthews M, Doherty G, Coyle D, Sharry J** (2008a). Designing mobile applications to support mental health interventions. In *Handbook of Research on User Interface Design and Evaluation for Mobile Technology* (ed. J. Lumsden), pp. 635–656. IGI Global: Hershey.
- Matthews M, Doherty G, Sharry J, Fitzpatrick C** (2008b). Mobile phone mood charting for adolescents. *British Journal of Guidance and Counselling* **36**, 113–129.
- Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group** (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine* **151**, 264–269, w264.
- Myin-Germeys I, Klippel A, Steinhart H, Reininghaus U** (2016). Ecological momentary interventions in psychiatry. *Current Opinion in Psychiatry* **29**, 258–263.
- Naslund JA, Marsch LA, McHugo GJ, Bartels SJ** (2015). Emerging mHealth and eHealth interventions for serious mental illness: a review of the literature. *Journal of Mental Health* **24**, 321–332.
- NCCMH** (2014). *E-therapies Systematic Review for Children and Young People with Mental Health Problems*. NCCMH: London.
- Nicholas J, Larsen ME, Proudfoot J, Christensen H** (2015). Mobile apps for bipolar disorder: a systematic review of features and content quality. *Journal of Medical Internet Research* **17**, e198.
- Ofcom** (2015). The Communications Market Report. (https://www.ofcom.org.uk/__data/assets/pdf_file/0022/20668/cmr_uk_2015.pdf).
- Olff M** (2015). Mobile mental health: a challenging research agenda. *European Journal of Psychotraumatology* **6**, 27882.
- Palmier-Claus JE, Ainsworth J, Machin M, Barrowclough C, Dunn G, Barkus E, Rogers A, Wykes T, Kapur S, Buchan I, Salter E, Lewis SW** (2012). The feasibility and validity of ambulatory self-report of psychotic symptoms using a smartphone software application. *BioMed Central Psychiatry* **12**, 172.
- Palmier-Claus JE, Rogers A, Ainsworth J, Machin M, Barrowclough C, Lavery L, Barkus E, Kapur S, Wykes T, Lewis SW** (2013). Integrating mobile-phone based assessment for psychosis into people's everyday lives and clinical care: a qualitative study. *BioMed Central Psychiatry* **13**, 34.
- Palmier-Claus JE, Taylor PJ, Ainsworth J, Machin M, Dunn G, Lewis SW** (2014). The temporal association between self-injurious thoughts and psychotic symptoms: a mobile phone assessment study. *Suicide and Life-Threatening Behavior* **44**, 101–110.
- Paris J** (2004). Borderline or bipolar? Distinguishing borderline personality disorder from bipolar spectrum disorders. *Harvard Review of Psychiatry* **12**, 140–145.
- Prentice JL, Dobson KS** (2014). A review of the risks and benefits associated with mobile phone applications for psychological interventions. *Canadian Psychology/Psychologie Canadienne* **55**, 282–290.
- Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R, Hensley M** (2011). Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Administration and Policy in Mental Health* **38**, 65–76.
- Reid SC, Kauer SD, Dudgeon P, Sancu LA, Shrier LA, Patton GC** (2009). A mobile phone program to track young people's experiences of mood, stress and coping: development and testing of the mobiletype program. *Social Psychiatry and Psychiatric Epidemiology* **44**, 501–507.
- Reid SC, Kauer SD, Hearps SJC, Croke AHD, Khor AS, Sancu LA, Patton GC** (2011). A mobile phone application for the assessment and management of youth mental health problems in primary care: a randomised controlled trial. *BioMed Central Family Practice* **12**, 131.
- Reid SC, Kauer SD, Hearps SJC, Croke AHD, Khor AS, Sancu LA, Patton GC** (2013). A mobile phone application for the assessment and management of youth mental health problems in primary care: health service outcomes from a randomised controlled trial of mobiletype. *BioMed Central Family Practice* **14**, 84.
- Reid SC, Kauer SD, Khor AS, Hearps SJC, Sancu LA, Kennedy AD, Patton GC** (2012). Using a mobile phone application in youth mental health: an evaluation study. *Australian Family Physician* **41**, 711–714.
- Sacco GR** (2015). *Developing a Smartphone Application for Depression: Tracking Risk and Wellness Factors*. M.A. University of Delaware.
- Sandstrom GM, Lathia N, Mascolo C, Rentfrow PJ** (2016a). Opportunities for smartphones in clinical care: the future of mobile mood monitoring. *Journal of Clinical Psychiatry* **77**, e135–e137.
- Sandstrom GM, Lathia N, Mascolo C, Rentfrow PJ** (2016b). Putting mood in context: Using smartphones to examine

- how people feel in different locations. *Journal of Research in Personality* (<http://dx.doi.org/10.1016/j.jrp.2016.06.004>).
- Schwartz JE, Neale J, Marco C, Shiffman SS, Stone AA** (1999). Does trait coping exist? A momentary assessment approach to the evaluation of traits. *Journal of Personality and Social Psychology* **77**, 360–369.
- Scotti JF** (2015). *School-based DBT skills groups for adolescent eating disorders and body image concerns: a pilot study*. Ph.D. University of Montana.
- Seko Y, Kidd S, Wiljer D, McKenzie K** (2014). Youth mental health interventions via mobile phones: a scoping review. *Cyberpsychology, Behavior and Social Networking* **17**, 591–602.
- Shiffman S, Stone AA, Hufford MR** (2008). Ecological momentary assessment. *Annual Review of Clinical Psychology* **4**, 1–32.
- Stone AA, Shiffman S, Schwartz JE, Broderick JE, Hufford MR** (2003). Patient compliance with paper and electronic diaries. *Controlled Clinical Trials* **24**, 182–199.
- Thomé S, Härenstam A, Hagberg M** (2011). Mobile phone use and stress, sleep disturbances, and symptoms of depression among young adults – a prospective cohort study. *BioMed Central Public Health* **11**, 66.
- Torous J, Firth J** (2016). The digital placebo effect: mobile mental health meets clinical psychiatry. *Lancet Psychiatry* **3**, 100–102.
- Torous J, Powell AC** (2015). Current research and trends in the use of smartphone applications for mood disorders. *Internet Interventions* **2**, 169–173.
- Tregarthen JP, Lock J, Darcy AM** (2015). Development of a smartphone application for eating disorder self-monitoring. *International Journal of Eating Disorders* **48**, 972–982.
- Trull TJ, Lane SP, Koval P, Ebner-Priemer UW** (2015). Affective dynamics in psychopathology. *Emotion Review* **7**, 355–361.
- van Knippenberg RJM, de Vugt ME, Ponds RW, Myin-Germeys I, van Twillert B, Verhey FRJ** (2016). Dealing with daily challenges in dementia (deal-id study): an experience sampling study to assess caregiver functioning in the flow of daily life. *International Journal of Geriatric Psychiatry* (<https://dx.doi.org/10.1002/gps.4552>).
- Walsh S, Golden E, Priebe S** (2016). Systematic review of patients' participation in and experiences of technology-based monitoring of mental health symptoms in the community. *BMJ Open* **6**, e008362.
- Wang R, Chen F, Chen Z, Li T, Harari G, Tignor S, Zhou X, Ben-Zeev D, Campbell AT** (2014). StudentLife: assessing mental health, academic performance and behavioral trends of college students using smartphones. In *Proceedings of the 2014 ACM International Joint Conference on Pervasive and Ubiquitous Computing*, pp. 3–14. ACM: Seattle.
- Watson D, Clark LA, Tellegen A** (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology* **54**, 1063–1070.
- Wenze SJ, Miller IW** (2010). Use of ecological momentary assessment in mood disorders research. *Clinical Psychology Review* **30**, 794–804.
- World Health Organisation** (1986). *Young people's health: a challenge for society – Report of a WHO Study Group on Young People and Health for All by the Year 2000*. World Health Organisation Technical Report Series 731. World Health Organisation: Geneva.
- Yen S, Frazier E, Hower H, Weinstock LM, Topor DR, Hunt J, Goldstein TR, Goldstein BI, Gill MK, Ryan ND, Strober M, Birmaher B, Keller MB** (2015). Borderline personality disorder in transition age youth with bipolar disorder. *Acta Psychiatrica Scandinavica* **132**, 270–280.