

ARTICLE

Relationship quality and functional limitations among older adults with cardiovascular disease in the United States of America

Yan-Liang Yu^{1*} and Zhenmei Zhang²

¹Department of Sociology and Criminology, Howard University, Washington, DC, USA and ²Department of Sociology, Michigan State University, East Lansing, Michigan, USA

*Corresponding author. Email: yanliang.yu@howard.edu

(Accepted 22 January 2019; first published online 14 March 2019)

Abstract

Substantial research shows that cardiovascular disease is a major cause of disability in the United States of America (USA) and worldwide. Despite the well-documented significance of intimate partnerships for cardiovascular health and disease management, how relationship quality contributes to the functional health of older adults diagnosed with cardiovascular disease is much less understood than mental health and mortality risk. Informed by the disablement process model and the lifecourse perspective, this study examines the association between relationship quality and functional limitations among partnered older adults aged 50 years and older diagnosed with cardiovascular disease in the USA. Data are from the Health and Retirement Study, 2006–2012 (N = 1,355). Multi-level linear regression analyses show that baseline negative relationship quality is significantly associated with increased functional limitations over the two- and four-year follow-ups. Additionally, the link between negative relationship quality and functional limitations is stronger among older adults with lower household income over a two-year span, compared to their higher-income counterparts, suggesting that these older adults are doubly disadvantaged by higher relationship strains and limited economic resources. Our findings demonstrate the significance of relationship quality for the functional health of older adults with cardiovascular disease and shed light on the importance of marriage/partnerships as an important social context for a critical stage in the disablement process (*i.e.* functional limitations).

Keywords: marriage; relationship quality; functional health; disease progression; Health and Retirement Study

Introduction

Substantial research has shown that marriage contributes to cardiovascular functioning (Zhang and Hayward, 2006) and that such health benefits depend largely on the quality of marital relationships (Liu and Waite, 2014). With growing prevalence of cardiovascular disease (CVD) in the United States of America (USA) and

worldwide (Yusuf *et al.*, 2001), increasing attention has been paid to how marital context contributes to the wellbeing of people living with CVD. While the associations of relationship quality with cardiovascular risks, mental health and survival of individuals diagnosed with CVD are well documented (Coyne *et al.*, 2001; Rohrbaugh *et al.*, 2002; Liu and Waite, 2014), much less is known about how relationship quality affects their functional health.

The disablement process model posits that functional limitations are a critical pathway through which chronic disease pathology develops into disability and emphasises the salience of social contexts for modifying this process (Verbrugge and Jette, 1994). Disability is a major health complication of CVD (Masoudi *et al.*, 2004) that undermines patients' quality of life and independence, and causes substantial financial burdens on individuals and the US health-care system (Newschaffer *et al.*, 2010). As marriage is an important social context for health, particularly among older adults (Umberson and Williams, 2005), it is critical to examine how relationship quality contributes to the progression of functional health after disease onset so that interventions for disease management can be better implemented. To this end, we used data from the 2006–2012 Health and Retirement Study (HRS) to investigate the link between relationship quality and functional limitations among older partnered adults diagnosed with CVD in the USA. Our study consists of three research objectives. First, we examine the associations of baseline positive and negative relationship quality with subsequent functional limitations in two-year and four-year follow-ups. Second, guided by previous work on marital quality and health, we evaluate health behaviours and psychological distress as possible explanations for the observed associations (Kiecolt-Glaser and Newton, 2001). Lastly, informed by the status–resource interaction model of the cumulative advantage/disadvantage process (Choi and Marks, 2013), we assess whether the association between relationship quality and functional limitations differs by individuals' household income.

Relationship quality, CVD and functional health

The disablement process: functional limitations as a critical stage

The disablement process model provides a useful framework for conceptualising the development of physical limitations and disability after the onset of disease pathology, and how contextual factors, such as relationship quality, can shape the course of this disease process. In essence, the model postulates that disease pathology leads to physiological impairments in specific body systems, which in turn leads to functional limitations that ultimately progress into disability (Verbrugge and Jette, 1994). It is also suggested that functional limitations are a critical stage for the progression from disease to disability (Lawrence and Jette, 1996).

The disablement process model not only delineates the pathways from illness to disability, but also accentuates the role of social contexts in modifying the course of the postulated pathways (Verbrugge and Jette, 1994; Jette *et al.*, 1997). Given the significance of marriage and intimate partnerships for population health, assessing how relationship quality influences the development of functional limitations

among older adults with CVD can inform us of how the progression into disability can be delayed or even prevented.

Why relationship quality matters

Relationship quality can have direct physiological impacts on the functional health of CVD patients. Both clinical and population-based studies have shown that relationship distress can cause physiological insults on the cardiovascular, neuroendocrine and immune systems (Robles *et al.*, 2014). For example, marital strains were associated with greater cardiovascular reactivity such as heightened blood pressure and faster heart rate (Robles and Kiecolt-Glaser, 2003; Liu and Waite, 2014), both of which are precursors of CVD and risk factors for functional decline (Karlman et al., 2002). Additionally, recent studies reported significant associations between marital quality and metabolic disorders such as central obesity, high blood pressure and dyslipidemia (Whisman and Uebelacker, 2012), all of which are associated with a higher risk of functional limitations (Penninx *et al.*, 2009). Lower marital quality has also been linked to higher levels of inflammation (Donoho *et al.*, 2013), another risk factor for functional disability (Kuo *et al.*, 2012).

Additionally, relationship quality can affect the functional health of older adults with CVD via its impacts on psychological wellbeing (Kiecolt-Glaser and Newton, 2001). Studies have shown that better relationship quality can help CVD patients develop psychological adjustments and buffer depression resulting from coping with the disease, thereby maintaining better functional health. In parallel, relationship distress may exacerbate adverse outcomes of CVD via psychological distress (Brecht *et al.*, 1994; Rohrbaugh *et al.*, 2002; Roijers *et al.*, 2016). Recent developments in behavioural cardiology have established psychological distress as a major risk factor for adverse cardiovascular outcomes, including decline in functional capacity (Das and O'Keefe, 2006).

Lastly, relationship quality can influence the functional health of CVD patients via change in health behaviours. Studies have shown that relationship strains can induce or exacerbate unhealthy habits such as smoking and excessive drinking (Kiecolt-Glaser and Newton, 2001), both of which can negatively affect the functional health of CVD patients (Fried and Guralnik, 1997; Lin *et al.*, 2011). Furthermore, it is well-documented that marriage benefits health through spousal health monitoring and the effectiveness of such health controls depends much on relationship quality (Waite and Gallagher, 2000). Recent evidence also showed that greater health-related spousal support, such as listening to one's health concerns and assistance in health care, can encourage the adoption of healthy behaviours such as engagement in physical activity and healthy diet among heart disease patients (Franks *et al.*, 2006), whereas relationship distress can hinder medication adherence (Cornwell and Waite, 2012). Given the importance of self-care for managing CVD (Gallant, 2003), we expect that relationship quality could also affect the functional health of CVD patients via behavioural changes.

In general, the empirical work shows that better relationship quality is significantly associated with better mental health (Brecht *et al.*, 1994; Roijers *et al.*, 2016) and greater survival prospects among CVD patients (Orth-Gomer *et al.*, 2000; Coyne *et al.*, 2001).

In contrast, little is known about how relationship quality contributes to the development of functional limitations among CVD patients despite its significant role in the disablement process. However, a small but growing number of studies have shown that relationship quality is significantly associated with functional health in the general population. Overall, the studies found empirical support for both positive and negative relationship quality as risk/protective factors for disability. For example, Bookwala (2005) reported that greater negative marital quality was significantly associated with more physical disabilities among older adults. Choi and Marks (2006) showed that marital conflicts significantly increased functional impairments at a later time. As for positive marital exchanges, Warner and Kelley-Moore (2012) demonstrated that positive marital quality buffered against the stressful effect of loneliness on functional limitations while Choi *et al.* (2016) found that higher positive marital quality of older adults and their spouses was significantly linked to fewer physical disabilities later. Both theoretical and empirical work on relationship quality, CVD and functional health suggest that relationship quality should play a significant role in the functional health progression of CVD patients after the disease onset. Building on the existing work, we propose the following hypothesis:

- Hypothesis 1: Higher baseline positive relationship quality is significantly associated with fewer functional limitations while higher baseline negative relationship quality is significantly linked to more functional limitations among older adults diagnosed with CVD over two-year and four-year follow-ups.

Additionally, past research showed that psychological distress and health behaviours were significant risk factors for adverse outcomes of CVD (Das and O'Keefe, 2006) and linked to the development of functional disability (Fried and Guralnik, 1997). These two factors are also theoretical pathways through which relationship quality affects physical health (Kiecolt-Glaser and Newton, 2001). Thus, we also examine them as possible explanations for the observed associations. We hypothesise that:

- Hypothesis 2: Psychological distress and health behaviours partially explain the link between relationship quality and functional limitations among older adults diagnosed with CVD.

The moderating role of household income

The principle of linked lives in the lifecourse framework maintains that individuals' health is influenced by their socially significant others like spouses or co-habiting partners (Moen and Hernandez, 2009). Viewed from this perspective, relationship quality can be seen as a resource/constraint that married or co-habiting couples develop over time and can influence the course of their health status (Umberson and Montez, 2010; Warner and Kelley-Moore, 2012). Furthermore, the significance of important social relationships for health may depend on other macrostructural contexts such as income or race/ethnicity (Choi and Marks, 2013).

The status–resource interaction model of cumulative advantage/disadvantage proposed by DiPrete and Eirich (2006) suggests that when examining the effects of individual resources or constraints on lifecourse outcomes such as health, one needs to also consider how these resources or constraints interact with structural status to have differential impacts on the outcomes. Guided by this model, we consider how the link between relationship quality and functional limitations among older adults with CVD may depend on household income as income, an important form of socio-economic resource, is consistently shown to impact health and disease outcomes, and is particularly crucial for shaping disease progression compared to education (Zimmer and House, 2003; Herd *et al.*, 2007). Following the status–resource interaction model and cumulative advantage theory, we expect that the association between relationship quality and functional health may be different across income levels where those with higher income benefit more from relationship support and those with lower income are more disadvantaged by relationship strains. Examinations of relationship quality, income and health are scarce. A recent study by Choi and Marks (2013) provides a good example. They found that increases in marital happiness were associated with improvements in physical health primarily for higher-educated individuals whereas increases in marital conflicts were linked with more physical limitations for individuals with lower income. Thus, we propose that

- Hypothesis 3: The association between higher positive relationship quality and fewer functional limitations among older adults with CVD is stronger for those with higher household income while the link between higher negative relationship quality and more functional limitations is stronger for those with lower income.

Methods

Data

We used data from the HRS 2006–2012, a multi-stage longitudinal household survey with a probability sample representative of non-institutionalised civilian adults aged 50 years and older in the USA with over-samples of Blacks, Hispanics and residents of the state of Florida. The HRS routinely collects rich information on respondents' socio-demographic and socio-economic characteristics and their health conditions biennially. Variables used in this study are from individual HRS waves, the 2014 tracker file and the RAND HRS Data Version N (RAND Center for the Study of Aging, 2016).

To understand better the psycho-social experience of US older adults, the HRS launched a pilot study in 2004 and officially included a module in 2006 to collect psycho-social information of the respondents with self-administered leave-behind questionnaires. The HRS conducts the leave-behind questionnaires on a rotating basis with a random half of the total 2006 sample surveyed in 2006, and the other half in 2008, and collects longitudinal information on the relationship quality every four years (Smith *et al.*, 2013). Measures of relationship quality used in this study are from the baseline psycho-social modules in 2006 and 2008 (Time 1 (T1)) and their respective four-year follow-up in 2010 and 2012 (T3).

Our study focuses on partnered older adults in the US diagnosed with CVD. Respondents with CVD were identified between 1992 and 2008 through their self-reported heart problems (e.g. heart attack, coronary heart disease or heart failure) or stroke (including transient ischaemic attack) diagnosed by a doctor. As our primary objective was to examine the health implications of relationship quality, we did not distinguish between union types and the term 'spouse' was used throughout the text to refer to either marital spouses or co-habiting partners. Among the 2006 and 2008 HRS respondents who were diagnosed with CVD, randomly selected and returned the psycho-social questionnaires, 1,475 older adults reported being partnered, of whom 1,414 were married and 61 were co-habitators. Our analytic samples came from these 1,475 individuals. Respondents' union types at T1 were controlled for in all analytic models. Figure 1 shows our detailed sample selection procedures.

Our analyses consisted of two parts, the two-year (2006/2008–2008/2010) and four-year follow-ups (2006/2008–2010/2012). The two-year follow-up analysis examined how relationship quality at T1 (2006/2008) was prospectively associated with functional limitations at T2 (2008/2010). The sample used in this analysis included 1,355 older adults who were at least 50 years old at T1 and alive at T2. The four-year follow-up analysis examined how relationship quality at T1 was prospectively associated with functional limitations at T3 (2010/2012), controlling for relationship quality at T3. A total of 1,080 older adults who were alive and remained partnered at T3 were used in the four-year analysis.

Measures

Functional limitations was created by totalling respondents' answers across 12 questions on difficulty in performing the following tasks (no difficulty = 0 versus at least some difficulty = 1): 'walking several blocks', 'jogging one mile', 'walking one block', 'sitting for about two hours', 'getting up from a chair after sitting for long periods', 'climbing several flights of stairs without resting', 'climbing one flight of stairs without resting', 'lifting or carrying weights over 10 lbs [approximately 4.5 kilograms]', 'stooping, kneeling or crouching', 'reaching arms above shoulder level', 'pushing or pulling large objects', and 'picking up a dime [coin] from the table'. The scores ranged from 0 to 12.

Relationship quality was measured by seven questions that tapped into respondents' perceived support from their spouses, an important dimension of relationship quality. Three questions assessed respondents' perceptions of spouses' positive support: (a) 'How much do they really understand the way you feel about things?', (b) 'How much can you rely on them if you have a serious problem?' and (c) 'How much can you open up to them if you need to talk about your worries?' Four questions evaluate respondents' perceptions of spouses' negative behaviours: (a) 'How often do they make too many demands on you?', (b) 'How much do they criticise you?', (c) 'How much do they let you down when you are counting on them?' and (d) 'How much do they get on your nerves?' The response categories for these seven questions were (1) a lot, (2) some, (3) a little and (4) at not all. Respondents' raw values were reverse-coded so that higher values indicated more positive/negative behaviours from spouses. The Cronbach's alpha for the




Figure 1. Flow chart of sample selection procedure for the current study.
 Note: HRS: Health and Retirement Study.

positive dimension items was 0.78 and 0.76 for negative items in our sample, indicating satisfactory internal consistency across the items.

Past research suggested that marital quality consisted of both positive and negative dimensions, two distinct constructs not exclusive of each other (Fincham and Linfield, 1997). Following the suggestion of the HRS manual (Smith *et al.*, 2013) and past research (*e.g.* Choi *et al.*, 2016), we created two summary scores for positive and negative relationship quality by summing up the values across the items

and taking the average respectively for the positive and negative dimensions. The two summary measures were centred at their respective mean values when entered in the models.

Health behaviours were indexed by drinking, smoking, physical activity and body weight. *Drinking* was recoded into three categories: abstainers, light to moderate drinkers (one to two drinks per day; reference category) and heavy drinkers (three drinks or more) (Zhang and Hayward, 2006). *Smoking* included the following categories: non-smokers (reference category), past smokers and current smokers. *Physical activity* was a binary indicator of whether respondents engaged in vigorous physical activity at least more than once a week (no = 0). *Body weight*, measured by the Body Mass Index, was included as an indicator for health behaviour because it was not only a significant risk factor for CVD but also reflected one's health behaviour and lifestyle. All health behaviours were measured at baseline.

Psychological distress was indexed by depressive symptoms and diagnosed emotional problems. *Depressive symptoms* were measured by a short eight-item version of the Center for Epidemiological Studies Depression Scale asking whether respondents experienced the following feelings in the past week: 'depressed', 'everything is an effort', 'restless sleep', 'lonely', 'sad', 'could not get going', 'happy' and 'enjoying life'. We used the summary score created by the RAND HRS data-set, ranging from 0 to 8, with higher values indicating more depressive symptoms. *Emotional problems* were measured by respondents' self-reported diagnosis of any emotional, nervous or psychiatric problems (no = 0). The two psychological indicators were measured at baseline.

We controlled for a series of covariates, including socio-demographic and socio-economic variables and respondents' social integration across all the models. *Gender* was a binary indicator (male = 0). *Age at baseline* (centred at age 50) and *age of diagnosis* were controlled for their close links to functional ability and CVD severity. *Race/ethnicity* included non-Hispanic white (reference category), non-Hispanic black, non-Hispanic other races and Hispanic. *Birthplace* was measured by a dummy indicator of South versus non-South regions (reference category). *Immigration status* was a binary indicator (immigrants = 0). *Current union status* at baseline included the first-time married (reference category), remarried and co-habiting. Two chronic conditions at baseline were also controlled: respondents' self-reported diagnosis of *hypertension* (no = 0) and *diabetes* (no = 0). The analysis of the two-year follow-up also controlled for whether respondents remained partnered after two years (no = 0). Approximately 93.75 per cent (N = 1,331) remained partnered after two years among the 1,355 respondents at baseline.

Baseline socio-economic conditions were controlled with the following four indicators. *Education* was measured by years of formal schooling. *Annual household income* was measured in nominal dollars by the sum of all income from the respondent and the spouse, not including other household members. *Net household assets* were measured by the sum of all wealth components (excluding the second residency) minus all debts. Missing cases on household income and net assets were imputed by the RAND HRS. The imputation procedures were implemented in a wave-specific and progressive fashion, depending on the type of missing cases. Readers interested in detailed imputation methodology can refer to the latest report contributed by the RAND Corporation (Pantoja *et al.*, 2018).

Both income and net assets were adjusted for household size and naturally logged. The adjusted household income was further centred at its median value. *Insurance status* was coded as '0' if respondent reported not having any kind of public or private health insurance and '1' if at least one type of health insurance. To control for respondents' other social integration, all the models were adjusted for whether respondents had *living children* (no = 0), *siblings* (no = 0) and *religious preference* (no = 0). Lastly, *baseline survey wave* (year 2006 = 0 versus 2008 = 1) and *baseline functional limitations* were controlled in all the models.

The proportion of missing values in our analytic sample is less than 4 per cent. We handled missing values with multiple imputation. Ten multiply imputed datasets were produced using the PROC MI procedure in SAS 9.2, and the model results were consolidated with PROC MIANALYZE.

Analytic approach

We used a lagged dependent variable approach to examine the links between baseline relationship quality (*i.e.* T1) and subsequent functional limitations in two (*i.e.* T2) and four years (*i.e.* T3). Baseline functional limitations, socio-demographic, socio-economic and social integration covariates, and survey years were controlled in all the models. Model 1 examined the association between baseline positive/negative relationship quality and functional limitations at T2 and T3, respectively. Models 2 and 3, respectively, adjusted for health behaviours and psychological distress to assess these two factors as possible explanations. Model 4 controlled for all the covariates. The final model added interaction terms to test whether the link between relationship quality and functional limitations varied by household income. Additionally, in our analysis of the four-year follow-up, we also controlled for change in positive and negative relationship quality over the four-year span in all the models.

To control for potential bias introduced by sample attrition due to premature mortality (as couples with poor relationship quality may have higher risk of death than their counterparts with good relationship quality), we adopted a two-stage Heckman approach to control for mortality selection (Liu, 2012; Umberson *et al.*, 2009). First, we used the entire HRS sample at T1 to predict T2/T3 probability of death by estimating discrete hazard models with baseline covariates including age, gender, race/ethnicity, education, household income, insurance status, smoking and depression. Then, we included the predicted probability of death at T2/T3 as a control variable in all the models.

Multi-level linear regression models were employed to assess the association between relationship quality and functional limitations to control for clustering effects at the household level because more than 15 per cent of our respondents were partnered couples in the same household. Results from the unconditional model (not shown here) indicated that approximately 24 per cent of the total variance in T2 functional limitations occurred at the household level and 15 per cent in T3 functional limitations, lending empirical support for our using multi-level models.

Results

Table 1 shows that the average age at baseline for our sample was about 70 years old, 63 per cent of the respondents were men and, on average, they had at least

Table 1. Weighted sample characteristics

	Percentage or mean	Standard deviation
Positive relationship quality at T1	3.46	0.018
Negative relationship quality at T1	2.01	0.022
Change in positive relationship quality, T1–T3 ¹	–0.19	0.085
Change in negative relationship quality, T1–T3 ¹	–0.06	0.119
Number of functional limitations at T2	3.82	0.102
Number of functional limitations at T3 ¹	4.77	0.311
Drinking (%):		
Abstainer	65.43	
Light/moderate drinker	32.09	
Heavy drinker	2.48	
Smoking (%):		
Non-smoker	38.13	
Past smoker	51.05	
Current smoker	10.82	
Physically active (%) (no = 0)	22.17	
Body Mass Index score	28.57	0.212
Depression score	1.46	0.073
Diagnosed emotional problems (%) (no = 0)	17.16	
Age at T1	69.24	0.304
Age of diagnosis	64.86	0.270
Female (%) (male = 0)	36.97	
Race/ethnicity (%):		
Non-Hispanic white	87.24	
Non-Hispanic black	5.47	
Non-Hispanic other races	2.23	
Hispanics	5.06	
Born in the South (%) (other regions = 0)	29.18	
Born in the USA (%) (immigrant = 0)	93.67	
Marital status at T1 (%):		
First-time married	68.44	
Remarried	28.69	
Co-habiting	2.87	
Diagnosed with hypertension (%) (no = 0)	67.54	
Diagnosed with diabetes (%) (no = 0)	26.14	

(Continued)

Table 1. (Continued.)

	Percentage or mean	Standard deviation
With living children (%)	96.16	
With living siblings (%)	84.38	
With religious preference (%)	93.44	
Partnered at T2 (%) (no = 0)	94.28	
Number of functional limitations at T1	3.57	0.097
Probability of death at T2	0.07	0.003
Survey wave (%) (2006 = 0)	48.37	
Years of formal education	12.78	0.122
Annual household income	85,539.65	13,764.76
Net asset	564,671.70	35,868.51
Having health insurance (%) (no = 0)	96.41	

Notes: N = 1,355. T1: Time 1. T2: Time 2. T3: Time 3. USA: United States of America. 1. The sample size for these measures is 1,080, *i.e.* the four-year follow-up sample.

a high school degree. Most respondents were former smokers and light/moderate drinkers at baseline, and physically inactive. The average number of depressive symptoms was approximately 1.5 out of a total of 8, and over 17 per cent of the respondents reported having emotional problems at baseline. More than two-thirds of the respondents were diagnosed with hypertension and over a quarter with diabetes, both of which are highly correlated with CVD. Most remained partnered two years later at T2.

Relationship quality and functional limitations of older adults with CVD

Two-year follow-up

Model 1 in Table 2 shows that positive relationship quality at T1 was not significantly associated with T2 functional limitations among older adults diagnosed with CVD ($\beta = 0.121$, $p = 0.271$). In contrast, negative relationship quality was significantly linked to more functional limitations two years later ($\beta = 0.273$, $p = 0.006$). Model 2 tests health behaviours as a possible explanation for the association observed in Model 1. The results show that the estimated coefficient for negative relationship quality only changed slightly with additional adjustments for health behaviour indicators. Model 3 assesses psychological distress as another explanation. Adjusting for psychological distress reduced the estimated coefficient of negative relationship quality by more than 15 per cent ($[(0.273 - 0.230)/0.273] \times 100$), suggesting that psychological distress was likely an explanation for the link between negative relationship quality and functional limitations among older adults with CVD. Negative relationship quality remained significantly associated with functional limitations after the model adjusted for psychological distress ($\beta = 0.230$, $p = 0.02$). Model 4 controlled for all the covariates, and the association between negative relationship quality and functional limitations remained statistically significant ($\beta = 0.224$, $p = 0.023$).

Table 2. Multi-level models of relationship quality and functional limitations among US older adults with cardiovascular disease, two-year follow-up

	Model 1	Model 2	Model 3	Model 4	Model 5
	<i>β (standard deviation)</i>				
Baseline PRQ (mean-centred)	0.121 (0.110)	0.117 (0.109)	0.181 (0.111)	0.177 (0.111)	0.145 (0.112)
Baseline NRQ (mean-centred)	0.273 (0.098)**	0.266 (0.098)**	0.230 (0.099)*	0.224 (0.099)*	0.204 (0.099)*
Baseline PRQ × Household income					−0.180 (0.139)
Baseline NRQ × Household income					−0.265 (0.108)*
Drinking (light/moderate = 0):					
Abstainer		0.051 (0.135)		0.044 (0.134)	0.057 (0.134)
Heavy drinker		0.753 (0.372)*		0.727 (0.370)*	0.698 (0.370)
Smoking (non-smoker = 0):					
Past smoker		−0.027 (0.126)		−0.035 (0.126)	−0.033 (0.126)
Current smoker		0.227 (0.220)		0.203 (0.219)	0.163 (0.219)
Physically active (no = 0)		−0.186 (0.146)		−0.196 (0.145)	−0.178 (0.145)
Body Mass Index score		0.013 (0.011)		0.013 (0.011)	0.011 (0.011)
Depression score			0.062 (0.036)	0.061 (0.036)	0.063 (0.036)
Emotional problems (no = 0)			0.482 (0.169)**	0.476 (0.169)**	0.499 (0.169)**
Age at T1 (centred at 50 years old)	0.030 (0.016)	0.036 (0.016)*	0.034 (0.016)*	0.039 (0.016)*	0.040 (0.016)*
Age of diagnosis	−0.016 (0.014)	−0.014 (0.014)	−0.014 (0.014)	−0.012 (0.014)	−0.012 (0.014)
Female (male = 0)	0.364 (0.122)**	0.370 (0.126)**	0.327 (0.122)**	0.332 (0.126)**	0.323 (0.126)*
Race/ethnicity (non-Hispanic White = 0):					
Non-Hispanic Black	−0.187 (0.226)	−0.197 (0.227)	−0.135 (0.226)	−0.146 (0.228)	−0.139 (0.227)

(Continued)

Table 2. (Continued.)

	Model 1	Model 2	Model 3	Model 4	Model 5
Non-Hispanic other races	0.130 (0.435)	0.099 (0.435)	0.161 (0.433)	0.133 (0.433)	0.145 (0.432)
Hispanics	-0.274 (0.312)	-0.261 (0.312)	-0.263 (0.310)	-0.251 (0.310)	-0.280 (0.309)
Born in the South (other regions = 0)	-0.025 (0.130)	-0.0002 (0.130)	-0.020 (0.129)	0.003 (0.129)	-0.0001 (0.129)
Born in the USA (immigrant = 0)	0.466 (0.282)	0.415 (0.283)	0.502 (0.281)	0.453 (0.282)	0.453 (0.282)
Marital status at T1 (first-time married = 0):					
Remarried	-0.013 (0.131)	-0.028 (0.131)	-0.031 (0.130)	-0.043 (0.130)	-0.052 (0.130)
Co-habiting	0.198 (0.299)	0.193 (0.300)	0.114 (0.299)	0.110 (0.299)	0.115 (0.298)
Diagnosed with hypertension (no = 0)	0.086 (0.126)	0.069 (0.126)	0.083 (0.124)	0.067 (0.125)	0.081 (0.125)
Diagnosed with diabetes (no = 0)	0.067 (0.134)	0.051 (0.139)	0.066 (0.134)	0.050 (0.139)	0.047 (0.138)
Whether partnered at T2 (no = 0)	-0.131 (0.239)	-0.059 (0.241)	-0.098 (0.238)	-0.029 (0.239)	-0.025 (0.239)
Functional limitations at T1	0.645 (0.023)***	0.638 (0.023)***	0.622 (0.024)***	0.615 (0.024)***	0.618 (0.024)***
Probability of death at T2	3.636 (0.897)***	2.843 (0.987)**	3.037 (0.908)**	2.532 (0.998)*	2.605 (0.998)**
Survey wave (2006 = 0)	0.213 (0.116)	0.183 (0.116)	0.219 (0.115)	0.190 (0.116)	0.196 (0.115)
Years of formal education	-0.052 (0.022)*	-0.047 (0.022)*	-0.047 (0.022)*	-0.042 (0.022)	-0.045 (0.022)*
Annual household income (median-centred)	-0.112 (0.084)	-0.111 (0.084)	-0.110 (0.084)	-0.110 (0.084)	1.104 (0.652)
Net asset	-0.133 (0.109)	-0.116 (0.110)	-0.146 (0.109)	-0.129 (0.109)	-0.141 (0.109)
Having health insurance (no = 0)	-0.357 (0.383)	-0.321 (0.383)	-0.309 (0.382)	-0.273 (0.381)	-0.234 (0.382)
Whether have child at T1 (no = 0)	-0.380 (0.367)	-0.371 (0.366)	-0.292 (0.366)	-0.285 (0.365)	-0.318 (0.364)
Whether have living siblings at T1 (no = 0)	-0.124 (0.155)	-0.098 (0.155)	-0.129 (0.154)	-0.104 (0.154)	-0.108 (0.154)

Whether have religious preference (no = 0)	0.106 (0.257)	0.088 (0.257)	0.153 (0.256)	0.133 (0.256)	0.142 (0.256)
Intercept	3.404 (1.841)	2.519 (1.915)	2.911 (1.837)	2.077 (1.910)	2.434 (1.916)
Variance components:					
Level 1 residual	3.312 (0.394)***	3.259 (0.392)***	3.301 (0.382)***	3.252 (0.380)***	3.280 (0.383)***
Level 2 variance, intercept	0.796 (0.387)*	0.824 (0.387)*	0.762 (0.373)*	0.786 (0.373)*	0.738 (0.375)*
−2 Log-likelihood	5,755.90	5,747.14	5,741.04	5,732.50	5,726.28
AIC	5,811.90	5,815.14	5,801.04	5,804.50	5,802.28
BIC	5,955.52	5,989.54	5,954.92	5,989.15	5,997.20

Notes: N = 1,355. T1: Time 1. T2: Time 2. PRQ: positive relationship quality. NRQ: negative relationship quality. USA: United States of America. AIC: Akaike information criterion. BIC: Bayesian information criterion. Annual household income and net asset have been naturally logged and adjusted for household size.

Significance levels: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Four-year follow-up

Model 1 in [Table 3](#) demonstrates the link between baseline relationship quality and functional limitations over a four-year span. The results are consistent with our analysis of the two-year follow-up. While baseline negative relationship quality was significantly associated with increased functional limitations among older adults with CVD ($\beta = 0.363$, $p = 0.02$), there was no significant link between positive relationship quality and functional limitations ($\beta = 0.204$, $p = 0.212$). Model 2 shows that adjusting for health behaviours barely changed the estimated coefficient of negative relationship quality whereas Model 3 suggests that controlling for psychological distress further reduced the estimated coefficient of negative relationship quality by approximately 7 per cent ($[(0.363 - 0.337) / 0.363] \times 100$). Model 4 shows that negative relationship quality was still significantly associated with increased functional limitations in four years ($\beta = 0.332$, $p = 0.03$) after adjustments for all the covariates.

Comparing the results in Models 2 and 3 in both [Tables 2](#) and [3](#) suggests that psychological distress played a more important role than health behaviours in explaining the link between negative relationship quality and functional limitations among older adults with CVD.

Does the association between relationship quality and functional limitations vary by levels of household income?

Model 5 in both [Tables 2](#) and [3](#) tests whether the association between relationship quality and functional limitations among older adults with CVD varied by levels of annual household income. The results in [Table 2](#) show that while there was no significant interaction effect between positive relationship quality and annual household income on functional limitations in a two-year follow-up, the link between negative relationship quality and functional limitations significantly varied by household income. While the main effect of negative relationship quality was significant and positive ($\beta = 0.204$, $p = 0.039$), the interaction effect was significant and negative ($\beta = -0.265$, $p = 0.014$), suggesting that baseline negative relationship quality was associated with worse functional limitations more strongly for older adults with lower household income over the two-year follow-up. In other words, higher household income buffered the effects of negative relationship on functional limitations. In contrast to the analysis of the two-year follow-up, Model 5 in [Table 3](#) shows that there was no significant interaction between either positive or negative relationship quality at baseline and household income in relation to subsequent functional limitations over the four-year span.

To illustrate this interactive relationship, we plotted estimated numbers of functional limitations at different levels of negative relationship quality (*i.e.* mean, mean ± 1 standard deviation (SD) and mean ± 2 SD) by low- versus high-income groups (25th versus 75th percentile) in [Figure 2](#) based on the results of Model 5 in [Table 2](#). The figure clearly shows that the lower-income group (*i.e.* 25th percentile) had a steeper increase in functional limitations with increasing negative relationship quality compared to the higher-income group (75th percentile).

Table 3. Multi-level models of relationship quality and functional limitations among US older adults with cardiovascular disease, four-year follow-up

	Model 1	Model 2	Model 3	Model 4	Model 5
	<i>β (standard deviation)</i>				
Baseline PRQ (mean-centred)	0.204 (0.164)	0.198 (0.164)	0.243 (0.165)	0.236 (0.165)	0.238 (0.165)
Baseline NRQ (mean-centred)	0.363 (0.151)*	0.357 (0.151)*	0.337 (0.152)*	0.332 (0.152)*	0.334 (0.153)*
Baseline PRQ × Household income					0.074 (0.162)
Baseline NRQ × Household income					0.018 (0.129)
Drinking (light/moderate = 0):					
Abstainer		0.011 (0.156)		0.004 (0.156)	0.005 (0.156)
Heavy drinker		0.321 (0.462)		0.302 (0.461)	0.312 (0.461)
Smoking (non-smoker = 0):					
Past smoker		-0.152 (0.149)		-0.163 (0.150)	-0.159 (0.150)
Current smoker		0.302 (0.264)		0.285 (0.264)	0.291 (0.265)
Physically active (no = 0)		-0.110 (0.168)		-0.123 (0.168)	-0.124 (0.168)
Body Mass Index score		0.015 (0.013)		0.015 (0.013)	0.015 (0.013)
Depression score			0.026 (0.045)	0.022 (0.045)	0.022 (0.045)
Emotional problems (no = 0)			0.316 (0.209)	0.332 (0.209)	0.332 (0.210)
Age at T1	0.005 (0.019)	0.012 (0.020)	0.007 (0.019)	0.014 (0.020)	0.014 (0.020)
Age of diagnosis	0.022 (0.017)	0.024 (0.017)	0.023 (0.017)	0.025 (0.017)	0.025 (0.017)
Female (male = 0)	0.537 (0.152)***	0.515 (0.155)***	0.517 (0.152)***	0.493 (0.155)**	0.493 (0.156)**
Race/ethnicity (non-Hispanic White = 0):					
Non-Hispanic Black	-0.008 (0.264)	-0.056 (0.268)	0.030 (0.265)	-0.017 (0.269)	-0.023 (0.269)

(Continued)

Table 3. (Continued.)

	Model 1	Model 2	Model 3	Model 4	Model 5
Non-Hispanic other races	-0.912 (0.501)	-0.968 (0.502)	-0.877 (0.500)	-0.927 (0.501)	-0.928 (0.501)
Hispanics	-0.270 (0.367)	-0.273 (0.367)	-0.243 (0.367)	-0.246 (0.367)	-0.236 (0.368)
Born in the South (other regions = 0)	0.023 (0.154)	0.053 (0.154)	0.026 (0.153)	0.056 (0.154)	0.053 (0.154)
Born in the USA (immigrant = 0)	0.004 (0.319)	-0.055 (0.321)	0.028 (0.319)	-0.029 (0.321)	-0.023 (0.321)
Marital status at T1 (first-time married = 0):					
Remarried	-0.095 (0.154)	-0.106 (0.154)	-0.107 (0.153)	-0.116 (0.153)	-0.111 (0.153)
Co-habiting	0.676 (0.367)	0.635 (0.368)	0.631 (0.368)	0.591 (0.368)	0.591 (0.368)
Diagnosed with hypertension (no = 0)	-0.078 (0.144)	-0.094 (0.146)	-0.086 (0.144)	-0.102 (0.146)	-0.098 (0.146)
Diagnosed with diabetes (no = 0)	0.043 (0.159)	0.039 (0.165)	0.033 (0.160)	0.028 (0.166)	0.028 (0.166)
Functional limitations at T1	0.636 (0.028)***	0.631 (0.028)***	0.624 (0.029)***	0.619 (0.029)***	0.620 (0.029)***
Probability of death at T2	2.520 (0.787)**	2.222 (0.909)*	2.380 (0.801)**	2.106 (0.924)*	2.076 (0.926)*
Survey wave (2006 = 0)	-0.126 (0.134)	-0.160 (0.135)	-0.132 (0.134)	-0.166 (0.135)	-0.167 (0.135)
Years of formal education	-0.051 (0.026)	-0.050 (0.026)	-0.048 (0.026)	-0.047 (0.026)	-0.046 (0.026)
Annual household income (median-centred)	-0.020 (0.098)	-0.015 (0.098)	-0.023 (0.098)	-0.018 (0.098)	-0.316 (0.764)
Net asset	-0.157 (0.116)	-0.144 (0.117)	-0.169 (0.116)	-0.155 (0.117)	-0.157 (0.117)
Having health insurance (no = 0)	-0.503 (0.437)	-0.518 (0.436)	-0.479 (0.437)	-0.490 (0.436)	-0.477 (0.438)
Whether have child at T1 (no = 0)	0.173 (0.430)	0.162 (0.431)	0.242 (0.432)	0.230 (0.432)	0.227 (0.432)
Whether have living siblings at T1 (no = 0)	-0.024 (0.187)	-0.011 (0.187)	-0.018 (0.186)	-0.007 (0.187)	-0.008 (0.187)
Whether have religious preference (no = 0)	0.133 (0.299)	0.091 (0.298)	0.155 (0.299)	0.113 (0.298)	0.123 (0.299)
Change in PRQ, T1-T3	-0.194 (0.171)	-0.192 (0.173)	-0.194 (0.172)	-0.192 (0.174)	-0.195 (0.174)
Change in NRQ, T1-T3	-0.048 (0.169)	-0.044 (0.170)	-0.051 (0.169)	-0.046 (0.169)	-0.046 (0.169)

Intercept	1.583 (2.067)	1.027 (2.153)	1.357 (2.070)	0.823 (2.155)	0.777 (2.155)
Variance components:					
Level 1 residual	4.386 (0.193)***	4.361 (0.192)***	4.370 (0.192)***	4.345 (0.191)***	4.344 (0.191)***
Level 2 variance, intercept	-	-	-	-	-
-2 Log-likelihood	4,658.87	4,652.75	4,655.13	4,648.89	4,648.52
AIC	4,714.87	4,720.75	4,715.13	4,720.89	4,724.52
BIC	4,852.26	4,887.57	4,862.33	4,897.53	4,910.97

Notes: N = 1,080. T1: Time 1. T2: Time 2. T3: Time 3. PRQ: positive relationship quality. NRQ: negative relationship quality. USA: United States of America. AIC: Akaike information criterion. BIC: Bayesian information criterion. Annual household income and net asset have been naturally logged and adjusted for household size.
Significance levels: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Figure 2. Estimated number of functional limitations and negative relationship quality: 25th versus 75th percentile household (HH) income, two-year follow-up.

Notes: 1SD: one standard deviation. 2SD: two standard deviations.

Discussion and conclusion

Guided by the disablement process model and the lifecourse paradigm, the current study examined how relationship quality, both positive and negative, was associated with the development of functional limitations in two and four years among US partnered older adults diagnosed with CVD, and assessed health behaviours and psychological distress as possible explanations. Additionally, we considered whether and how the link between relationship quality and change in functional limitations varied by levels of annual household income. Findings from our study make several important contributions to the literature.

Relationship quality matters

First, given that little is known of how relationship quality impacts the functional health of individuals with CVD, we contribute to the literature by demonstrating that baseline negative relationship quality, but not positive, is significantly associated with increased functional limitations subsequently over a two- and four-year span among older adults with CVD, providing robust empirical evidence from a national sample. Additional analyses on gender differences (not shown here) show that these significant associations are comparable for older men *versus* women. Our findings here suggest that frequent unsupportive behaviours such as criticising or making excessive demands (*i.e.* measures of negative relationship quality) are detrimental to the functional health of older adults living with a major chronic illness like CVD over time.

Our results are consistent with findings from previous research that also showed greater importance of negative relationship quality for physical health than positive relationship quality (Kiecolt-Glaser and Newton, 2001; Bookwala, 2005). Additionally, it aligns with the larger literature that demonstrates a stronger link between negative exchanges and wellbeing than positive ones in interpersonal

relationships (Newsom *et al.*, 2005). This greater salience of negative relationship quality also fits with research that shows individuals are more attuned to negative experiences in daily life (Taylor, 1991; Umberson *et al.*, 2006). In a similar vein, experimental studies also show immediate physiological changes in response to relationship conflicts or discords such as heightened blood pressure or faster heart rates (Robles and Kiecolt-Glaser, 2003).

Household income moderates the association

Additionally, informed by the status–resource interaction model in the lifecourse paradigm (DiPrete and Eirich, 2006), we examined whether the link between relationship quality and functional limitations was moderated by annual household income. The results provide partial support for our hypothesis. Given that higher negative relationship quality was significantly associated with increased functional limitations, we found that this association was stronger among older adults with lower household income than those with higher over the two-year follow-up, suggesting that older adults with higher income were partially protected by their greater economic resources against the physical burden of being in a bad relationship, whereas those with lower income were doubly disadvantaged by negative relationship quality and limited economic resources while managing their disease. Our findings also suggest that the buffering effect of household income is only temporary. In this respect, we contribute further empirical evidence to the literature on how the relationship between social contexts and the disablement process can be further modified by important structural statuses like income, although only temporarily. Since our sample consists of older adults living with CVD, our finding here has implications for public health interventions in CVD management.

Psychological distress partially explains the link

We also examined health behaviours and psychological distress as possible explanations. Our results showed that psychological distress played a greater role in explaining the association between relationship strains and deteriorating functional health among US older adults with CVD, contributing additional evidence on mental health as an important risk factor for managing CVD (Das and O’Keefe, 2006). However, even after adjusting for all the covariates, negative relationship quality was still significantly associated with functional limitations. Future research should assess other aspects of psycho-social attributes as explanations such as anxiety, pessimism or anger (Rozanski, 2014). A dyadic approach to the psycho-social distress of both patients and spouses as potential care-givers would also inform the current scholarship on the link between relationship strains and chronic disease progression.

Although we did not find evidence for health behaviours as a possible explanation, we encourage researchers to investigate other behavioural explanations given the significance of self-care in managing a chronic condition like CVD (Sayers *et al.*, 2008). In contrast to our general behavioural measures, future research should examine behaviours specific to disease management such as medication adherence or dietary adjustments to understand better how relationship strains impact chronic disease management via behavioural pathways.

Limitations

Despite our robust findings that are consistent with the literature, a few limitations need to be addressed. First, this study only examined a specific dimension of relationship quality – *i.e.* supportive and unsupportive behaviours – due to data limitation. In light of the significance of relationship quality for chronic disease management, future research should employ other validated measures such as the Dyadic Adjustment Scale and the Marital Adjustment Test to investigate how other dimensions of relationship quality contribute to the functional health of individuals with CVD and other chronic illnesses. Additionally, our analyses may have underestimated the impact of relationship quality due to the selectivity of our sample. Given that relationship quality and the mortality risk of CVD are closely linked (Coyne *et al.*, 2001), our sample, by definition, is a select group of older adults who survived CVD during our observation window. Our adjustment for mortality selection reduces this bias but does not eradicate it. Furthermore, as relationship quality is indicative of partnership stability (Amato and Rogers, 1997) and relationships of very low quality may have already dissolved before we began our observation, our sample may be in relatively more robust partnerships that would have otherwise broken up had they been worse and more fragile. In light of this sample selectivity, our findings could be considered conservative estimates. It is of theoretical interest to also compare the association of relationship quality and functional limitations between the divorced/separated *versus* the partnered sample, given that divorced/separated couples tend to have lower relationship quality than the married. The fairly small sample size of divorced/separated older adults who were diagnosed with CVD and reported relationship quality (less than 1%) prevents us from pursuing this analysis.

Here, we only considered how relationship quality affected subsequent functional health among older adults with CVD. However, the health burden of disease management and care-giving for both patients and their partners can exert a toll on relationship quality (Choi and Marks, 2006). Future research should consider the reciprocal relationships between relationship quality and chronic disease progression to gain more insight into the causal dynamics. Lastly, our study did not distinguish between co-habitation and marriage due to the small sample size of co-habitors. In light of co-habitation as a growing alternative partnership among older adults in addition to marriage (Brown *et al.*, 2006), future research should examine whether the link between relationship quality and chronic disease progression differs by union types.

The limitations notwithstanding, our study sheds light on the importance of relationship quality for a critical phase of the disablement process – functional limitations – after the onset of CVD and showcases how income inequality may exacerbate the impact of negative relationship quality for the less fortunate. Therefore, our findings call for greater attention to older people who manage CVD in a partnership characterised by negative experiences, and more importantly, interventional efforts need to be directed particularly to those with limited economic resources.

References

- Amato PR and Rogers SJ** (1997) A longitudinal study of marital problems and subsequent divorce. *Journal of Marriage and the Family* **59**, 612–624.
- Bookwala J** (2005) The role of marital quality in physical health during the mature years. *Journal of Aging and Health* **17**, 85–104.
- Brecht M-L, Dracup K, Moser DK and Riegel B** (1994) The relationship of marital quality and psychosocial adjustment to heart disease. *Journal of Cardiovascular Nursing* **9**, 74–85.
- Brown SL, Lee GR and Bulanda JR** (2006) Cohabitation among older adults: a national portrait. *Journals of Gerontology: Psychological Sciences and Social Sciences* **61B**, S71–S79.
- Choi H and Marks NF** (2006) Transition to caregiving, marital disagreement, and psychological well-being: a prospective US national study. *Journal of Family Issues* **27**, 1701–1722.
- Choi H and Marks NF** (2013) Marital quality, socioeconomic status and physical health. *Journal of Marriage and Family* **75**, 903–919.
- Choi H, Yorgason JB and Johnson DR** (2016) Marital quality and health in middle and later adulthood: dyadic associations. *Journals of Gerontology: Psychological Sciences and Social Sciences* **71B**, 154–164.
- Cornwell EY and Waite LJ** (2012) Social network resources and management of hypertension. *Journal of Health and Social Behavior* **53**, 215–231.
- Coyne JC, Rohrbaugh MJ, Shoham V, Sonnega JS, Nicklas JM and Cranford JA** (2001) Prognostic importance of marital quality for survival of congestive heart failure. *American Journal of Cardiology* **88**, 526–529.
- Das S and O’Keefe JH** (2006) Behavioral cardiology: recognizing and addressing the profound impact of psychosocial stress on cardiovascular health. *Current Atherosclerosis Reports* **8**, 111–118.
- DiPrete TA and Eirich GM** (2006) Cumulative advantage as a mechanism for inequality: a review of theoretical and empirical developments. *Annual Review of Sociology* **32**, 271–297.
- Donoho CJ, Crimmins EM and Seeman TE** (2013) Marital quality, gender, and markers of inflammation in the MIDUS cohort. *Journal of Marriage and Family* **75**, 127–141.
- Fincham FD and Linfield KJ** (1997) A new look at marital quality: can spouses feel positive and negative about their marriage? *Journal of Family Psychology* **11**, 489–502.
- Franks MM, Stephens MAP, Rook KS, Franklin BA, Keteyian SJ and Artinian NT** (2006) Spouses’ provision of health-related support and control to patients participating in cardiac rehabilitation. *Journal of Family Psychology* **20**, 311–318.
- Fried LP and Guralnik JM** (1997) Disability in older adults: evidence regarding significance, etiology, and risk. *Journal of the American Geriatrics Society* **45**, 92–100.
- Gallant MP** (2003) The influence of social support on chronic illness self-management: a review and directions for research. *Health Education & Behavior* **30**, 170–195.
- Herd P, Goesling B and House JS** (2007) Socioeconomic position and health: the differential effects of education versus income on the onset versus progression of health problems. *Journal of Health and Social Behavior* **48**, 223–238.
- Jette DU, Manago D, Medved E, Nickerson A, Warzycha T and Bourgeois MC** (1997) The disablement process in patients with pulmonary disease. *Physical Therapy* **77**, 385–394.
- Karlamangla AS, Singer BH, McEwen BS, Rowe JW and Seeman TE** (2002) Allostatic load as a predictor of functional decline: MacArthur studies of successful aging. *Journal of Clinical Epidemiology* **55**, 696–710.
- Kiecolt-Glaser JK and Newton TL** (2001) Marriage and health: his and hers. *Psychological Bulletin* **127**, 472–503.
- Kuo H-K, Al Snih S, Kuo Y-F and Raji MA** (2012) Chronic inflammation, albuminuria, and functional disability in older adults with cardiovascular disease: the National Health and Nutrition Examination Survey, 1999–2008. *Atherosclerosis* **222**, 502–508.
- Lawrence RH and Jette AM** (1996) Disentangling the disablement process. *Journals of Gerontology: Psychological Sciences and Social Sciences* **51B**, S173–S182.
- Lin JC, Guerrieri JG and Moore AA** (2011) Drinking patterns and the development of functional limitations in older adults: longitudinal analyses of the Health and Retirement Survey. *Journal of Aging and Health* **23**, 806–821.

- Liu H** (2012) Marital dissolution and self-rated health: age trajectories and birth cohort variations. *Social Science & Medicine* **74**, 1107–1116.
- Liu H and Waite L** (2014) Bad marriage, broken heart? Age and gender differences in the link between marital quality and cardiovascular risks among older adults. *Journal of Health and Social Behavior* **55**, 403–423.
- Masoudi FA, Rumsfeld JS, Havranek EP, House JA, Peterson ED, Krumholz HM, Spertus JA, for the Cardiovascular Outcomes Research Consortium (CORC)** (2004) Age, functional capacity, and health-related quality of life in patients with heart failure. *Journal of Cardiac Failure* **10**, 368–373.
- Moen P and Hernandez E** (2009) Social convoys: studying linked lives in time, context, and motion. In Elder GH and Giele JZ (eds), *The Craft of Life Course Research*. New York, NY: Guilford Press. pp. 258–279.
- Newschaffer CJ, Liu L and Sim A** (2010) Cardiovascular disease. In Remington PL, Brownson RC and Wegner MV (eds), *Chronic Disease Epidemiology and Control*, 3rd Edn. Washington, DC: American Public Health Association, chap. 13, pp. 383–428.
- Newsom JT, Rook KS, Nishishiba M, Sorkin DH and Mahan TL** (2005) Understanding the relative importance of positive and negative social exchanges: examining specific domains and appraisals. *Journals of Gerontology: Psychological Sciences and Social Sciences* **60B**, P304–P312.
- Orth-Gomer K, Wamala SP, Horsten M, Schenck-Gustafsson K, Schneiderman N and Mittleman MA** (2000) Marital stress worsens prognosis in women with coronary heart disease: the Stockholm Female Coronary Risk Study. *JAMA: The Journal of the American Medical Association* **284**, 3008–3014.
- Pantoja P, Bugliari D, Campbell N, Chan C, Hayden O, Hayes J, Hurd M, Main R, Mallet J, McCullough C, Meijer E, Moldoff M, Rohwedder S and St. Clair P** (2018) *RAND HRS Detailed Imputations File 2014 (V2) Documentation*. Santa Monica, CA: RAND Corporation.
- Penninx BW, Nicklas BJ, Newman AB, Harris TB, Goodpaster BH, Satterfield S, de Rekeneire N, Yaffe K, Pahor M, Kritchevsky SB, for the Health ABC Study** (2009) Metabolic syndrome and physical decline in older persons: results from the Health, Aging and Body Composition Study. *Journals of Gerontology: Biological Sciences and Medical Sciences* **64A**, 96–102.
- RAND Center for the Study of Aging** (2016) *RAND HRS Data, Version P*. Santa Monica, CA: RAND Center for the Study of Aging.
- Robles TF and Kiecolt-Glaser JK** (2003) The physiology of marriage: pathways to health. *Physiology & Behavior* **79**, 409–416.
- Robles TF, Slatcher RB, Trombello JM and McGinn MM** (2014) Marital quality and health: a meta-analytic review. *Psychological Bulletin* **140**, 140–187.
- Rohrbaugh MJ, Cranford JA, Shoham V, Nicklas JM, Sonnega JS and Coyne JC** (2002) Couples coping with congestive heart failure: role and gender differences in psychological distress. *Journal of Family Psychology* **16**, 3–13.
- Roijers J, Sunamura M, Utens EM, Dulfer K, ter Hoeve N, van Geffen M, Draaijer J, Steenaard R and van Domburg RT** (2016) Marital quality and loneliness as predictors for subjective health status in cardiac rehabilitation patients following percutaneous coronary intervention. *European Journal of Preventive Cardiology* **23**, 1245–1251.
- Rozanski A** (2014) Behavioral cardiology: current advances and future directions. *Journal of the American College of Cardiology* **64**, 100–110.
- Sayers SL, Riegel B, Pawlowski S, Coyne JC and Samaha FF** (2008) Social support and self-care of patients with heart failure. *Annals of Behavioral Medicine* **35**, 70–79.
- Smith J, Fisher G, Ryan L, Clarke P, House J and Weir D** (2013) *Psychosocial and Lifestyle Questionnaire, 2006–2010: Documentation Report, Core Section LB*. Ann Arbor, MI: Survey Research Center, University of Michigan, pp. 1–62
- Taylor SE** (1991) Asymmetrical effects of positive and negative events: the mobilization-minimization hypothesis. *Psychological Bulletin* **110**, 67–85.
- Umberson D, Liu H and Powers D** (2009) Marital status, marital transitions, and body weight. *Journal of Health and Social Behavior* **50**, 327–343.
- Umberson D and Montez JK** (2010) Social relationships and health a flashpoint for health policy. *Journal of Health and Social Behavior* **51**, S54–S66.
- Umberson D and Williams K** (2005) Marital quality, health, and aging: gender equity? *Journals of Gerontology: Psychological Sciences and Social Sciences* **60B**, S109–S113.

- Umberson D, Williams K, Powers DA, Liu H and Needham B** (2006) You make me sick: marital quality and health over the life course. *Journal of Health and Social Behavior* **47**, 1–16.
- Verbrugge LM and Jette AM** (1994) The disablement process. *Social Science & Medicine* **38**, 1–14.
- Waite LJ and Gallagher M** (2000) *The Case for Marriage: Why Married People Are Happier, Healthier and Better Off Financially?* New York, NY: Doubleday Books.
- Warner DF and Kelley-Moore J** (2012) The social context of disablement among older adults: does marital quality matter for loneliness? *Journal of Health and Social Behavior* **53**, 50–66.
- Whisman MA and Uebelacker LA** (2012) A longitudinal investigation of marital adjustment as a risk factor for metabolic syndrome. *Health Psychology* **31**, 80–86.
- Yusuf S, Reddy S, Ōunpuu S and Anand S** (2001) Global burden of cardiovascular diseases. Part II: variations in cardiovascular disease by specific ethnic groups and geographic regions and prevention strategies. *Circulation* **104**, 2855–2864.
- Zhang Z and Hayward MD** (2006) Gender, the marital life course, and cardiovascular disease in late mid-life. *Journal of Marriage and Family* **68**, 639–657.
- Zimmer Z and House JS** (2003) Education, income, and functional limitation transitions among American adults: contrasting onset and progression. *International Journal of Epidemiology* **32**, 1089–1097.

Cite this article: Yu Y-L, Zhang Z (2020). Relationship quality and functional limitations among older adults with cardiovascular disease in the United States of America. *Ageing & Society* **40**, 1694–1717. <https://doi.org/10.1017/S0144686X19000163>