


cambridge.org/ber

Lvquan Zhao¹ , Wei Wang¹, Ying Qiu¹ and Alex S. Torson²

Research Paper

Cite this article: Zhao L, Wang W, Qiu Y, Torson AS (2021). Plasticity of nutrient accumulation patterns in diapausing fall webworm pupae. *Bulletin of Entomological Research* **111**, 637–644. <https://doi.org/10.1017/S0007485321000201>

Received: 19 May 2020

Revised: 8 January 2021

Accepted: 26 February 2021

First published online: 31 March 2021

Keywords:

Diapause; energy reserves; enzyme activity; feeding efficiency; Lepidoptera; photoperiod; plasticity

Author for correspondence:Lvquan Zhao, Email: zhaolvquan@njfu.edu.cn

¹Co-innovation Center for Sustainable Forestry in Southern China, College of Forestry, Nanjing Forestry University, Nanjing 210037, China and ²Department of Biology, The University of Western Ontario, London, ON N6A 5B7, Canada

Abstract

The accumulation of nutrients during diapause preparation is crucial because any lack of nutrition will reduce the likelihood of insects completing diapause, thereby decreasing their chances of survival and reproduction. The fall webworm, *Hyphantria cunea*, diapause as overwintering pupae and their diapause incidence and diapause intensity are regulated by the photoperiod. In this study, we test the hypothesis that photoperiod influences energy reserve accumulation during diapause preparation in fall webworm. We found that the body size and mass, lipid and carbohydrate content of pupae with a short photoperiod during the diapause induction phase were significantly greater than those of pupae with a relatively short photoperiod, and the efficiency of converting digested food and ingested food into body matter was greater in the short-photoperiod diapause-destined larvae than the relatively short-photoperiod diapause-destined larvae. We also observed higher lipase and amylase activities in short-photoperiod diapause-destined larvae relative to the counterparts. However, no obvious difference was found in protein and protease in the pupae with a short photoperiod during the diapause induction phase and short-photoperiod diapause-destined larvae compared with the counterparts. Therefore, we conclude that the energy reserve patterns of diapausing fall webworm pupae are plastic and that short-photoperiod diapause-destined larvae increase their energy reserves by improving their feeding efficiency and increase their lipid and carbohydrate stores by increasing the lipase and amylase activities in the midgut.

Introduction

Diapause is a seasonal, programmed state of developmental and/or metabolic arrest, which can occur at various stages of an insect's life, including egg, larval, pupa, and adult (Tauber *et al.*, 1986; Denlinger, 1991, 2002). Insect diapause consists of several phases: induction, preparation, initiation, maintenance, termination, and post-diapause quiescence (Košťál, 2006). Each diapause phase may be influenced by environmental factors as well as insect genetics (Han and Denlinger, 2009). Changes in the photoperiod provide reliable seasonal signals for insects and other organisms, thereby allowing them to ensure that their life history is synchronized with seasonal changes; thus, the photoperiod is a key factor related to the regulation of seasonal development and life history in temperate species (Tauber *et al.*, 1986; Tanaka *et al.*, 1999). For example, *Velarifictorus parvus* Chopard (Orthoptera: Gryllidae) larvae develop faster under long photoperiod conditions, whereas larval development is inhibited under short photoperiod conditions, and the diapause status of larvae ends rapidly after moving from short photoperiod conditions to long-day sunshine conditions (Tanaka *et al.*, 1999).

In addition to controlling diapause induction, photoperiod also influences diapause incidence and diapause intensity in insects. For example, in *Ostrinia furnacalis* Guenée (Lepidoptera: Crambidae), the diapause incidence increases significantly as the photoperiod becomes shorter, and a short photoperiod below the critical day length can induce entry into deeper diapause than did the relatively short photoperiod below the critical day length (Yang *et al.*, 2014). In order to adapt to local seasonal changes, the diapause characteristics of widespread insects have also changed significantly during long-term evolution; thus, the critical photoperiod for inducing diapause will be shorter for low-latitude populations compared with high-latitude populations, and the diapause intensity is deeper (Danks, 1987; Chen *et al.*, 2013). The deeper diapause intensity plays an important role in avoiding an untimely exit from diapause due to high temperatures before winter for low-latitude populations (Tanaka and Zhu, 2008).

In the diapause state, most insects are unable to move and feed, so the energy reserves stored during the diapause preparation phase must meet the needs for both basic metabolism during the diapause period and for their development after diapause (Hahn and Denlinger, 2011; Sinclair, 2015). During diapause induction phase, token stimuli initiate physiological changes that lead to increased energy storage and preparation for diapause (Košťál, 2006). For example, in *Colaphellus bowringi* Baly (Coleoptera: Chrysomelidae), the diapause-destined

© The Author(s), 2021. Published by Cambridge University Press

CAMBRIDGE
UNIVERSITY PRESS

adults will store more lipids in response to short photoperiod; by contrast, *C. bowringi* will store more carbohydrates and protein when the adult senses the signal of long photoperiod (Tan *et al.*, 2016).

The fall webworm, *Hyphantria cunea* (Drury) (Lepidoptera: Arctiinae), is native to North America and was introduced into Japan around 1970 and was subsequently introduced into China (Ji *et al.*, 2003). Due to its wide range of food plants and high adaptability, fall webworm has caused severe forestry production losses in China. Fall webworm completes one generation each year in the North American region, but two generations each year in the Tokyo area of Japan, and three generations or more can occur each year toward the south of Japan (Gomi, 1997). Fall webworm was introduced into China in 1979 in Dandong city (about 40°N), Liaoning province and it spread to areas such as Nanjing city (about 32°N), Jiangsu Province (Liu *et al.*, 2016). When fall webworm was first introduced into China, it completed two generations each year, but three generations now occur each year in most parts of China (Ji *et al.*, 2003; Wei *et al.*, 2006).

Fall webworm overwinters as diapausing pupae from October to May of next year in Shanghe County of Shandong Province. Photoperiodic cues during early larval stage determine whether diapause is induced (Masaki *et al.*, 1968). The critical photoperiod for inducing diapause is 14 h 30 min to 14 h 45 min at 25°C (Gomi, 1997). The critical photoperiod is shorter as temperature increases during the induction period, and the diapause-inducing short day lengths of 8, 10, and 12 h evoke greater intensities of diapause than did 13 and 14 h (Chen *et al.*, 2014). We know that energy reserve patterns differ in the diapause pupae and non-diapause pupae of the fall webworm and that diapause pupae reserve more lipid and carbohydrate content relative to the non-diapause pupae (Zhao *et al.*, unpublished). However, it is still unclear whether there is variation in energy accumulation in fall webworm when it experiences a shortened photoperiod during diapause induction phase.

The present study aimed to determine whether the energy reserve patterns change during the diapause preparation phase when fall webworm is exposed to different photoperiods below the critical photoperiod during the diapause induction phase. We tested the hypothesis that photoperiod influences energy reserve accumulation during diapause preparation in fall webworm. We predicted that fall webworm larvae that experience a short photoperiod will increase their feeding efficiency or food intake, or both, to increase their energy reserve during the diapause preparation period. We also predicted that body size, dry mass, and energy reserves would be significantly greater in diapausing pupae exposed to a shorter photoperiod during the diapause induction phase as larvae than relatively short photoperiod diapausing pupae. Furthermore, we expected that, if short-photoperiod diapausing pupae store more energy, the short photoperiod would promote increased energy storage in larvae during diapause preparation via increased feeding efficiency, food intake, or both. As a result, we expected that a short photoperiod would induce the higher lipase, amylase, and protease activities in the midgut of diapause-destined larvae in order to increase the storage of lipid, carbohydrates, and protein.

Materials and methods

Insect rearing

Fall webworm larvae were obtained from poplar trees (*Populus deltoids*) in Shanghe County (37.35°N, 117.34°E) of Shandong

Province in May 2018 and reared in the laboratory for three generations under non-diapause-inducing conditions (25°C, L:D 16:8) by the methods described by Zhao *et al.* (unpublished). The newly laid eggs were moved to a Petri dish (7 cm diameter) containing absorbent cotton and moist qualitative filter paper at the bottom to maintain humidity. We checked the eggs every day after the initiation of larvae hatching. The newly hatched larvae were divided into two groups after hatching. The first group was transferred into a chamber at 25°C with a photoperiod of L:D 13:11, and the second group was kept at 25°C with a photoperiod of L:D 10:14. All the larvae were raised in aggregate during the 1st to 4th instars, subsequently they were raised separately after entering the 5th instar with artificial diet consisting of wheat germ (12%), casein (4.5%), sugar (2%), agar (1.5%), and other dietary ingredients (Cao *et al.*, 2014). Given that fall webworms were exposed to two groups of photoperiods below the critical photoperiod during the diapause induction phase, these larvae were destined to enter the diapause state after pupation.

Body size and dry mass

On the day of pupation, we transferred pupae to Petri dishes containing absorbent cotton and moist filter paper at the bottom. The Petri dishes with pupae were transferred to chambers at a constant temperature of 25 ± 1°C with a photoperiod of L:D 13:11 and L:D 10:14. The sex of each pupa was determined by the presence (female) or absence (male) of a line intersecting the first abdominal sternite (Loewy *et al.*, 2013; Williams *et al.*, 2015). To investigate the effect of photoperiod variation on body size and dry mass of diapausing pupae, on the 2nd day after pupation, we measured the lengths and widths of the pupae with a micrometer (0.1 cm) and dry mass with a balance (AL104 Mettler-Toledo; Mettler-Toledo, $d = 0.0001$ g) after drying on a slide glass at 110°C for 24 h according to the method of Tanaka (1991).

Body composition

To examine the effect of photoperiod variation below the critical photoperiod on the energy reserve during the diapause preparation phase, we extracted the lipid, protein, and glycogen contents from pupae on the 1st, 3rd, and 5th days after pupation according to the method described by Lorenz (2003). The lipid contents were determined using the sulfophosphovanillin method (adapted from Lorenz, 2003). Samples (20 µl) were added to 980 µl sulfuric acid, and then the mixtures were incubated at 100°C for 10 min. After cooling to room temperature, 5 ml of 0.2% phosphovanillin in 57% ortho-phosphoric acid was added to the mixtures. Samples were measured at 530 nm and concentrations were determined against cholesterol standards. The protein was quantified by Coomassie blue staining. Samples (25 µl) were added to 125 µl phosphate-buffered saline, then mixed with 2.85 µl Coomassie blue fast staining solution. Absorbance was measured against bovine serum albumin standards at 595 nm. Carbohydrate contents were quantified using the anthrone method (Carroll *et al.*, 1956). We mixed the sample (100 µl) with 900 µl distilled water, then added 4 ml of the 0.1% (1 g anthrone in 1000 ml 80% sulfuric acid) anthrone to the mixture, and heated the mixture to 100°C for 10 min. Absorbance was measured at 620 nm using the known glucose concentration as standards.

Feeding experiment

Since early-instar larvae were aggregated (Rehnberg, 2006), we only measured the food utilization in the 5th to 6th instar in

the short-photoperiod diapause-destined and relatively short-photoperiod diapause-destined larvae. After molting into the 5th instar, we transferred larvae to a new container to rear independently. On day 2 after entering the 5th instar, we fed larvae *ad libitum* with a pre-weighed diet that was replaced every 2 days. The remaining diet and frass were weighed after each diet change. The dry mass of the food and larvae was regressed against the wet mass of food and larvae to calculate the wet-to-dry conversion equation of linear regression according to the method described by Zhao *et al.* (2017) (the diet: dry mass = $0.313 \times$ wet mass -0.003 , $r^2 = 0.99$; short-photoperiod diapause-destined larvae: dry mass = $0.148 \times$ wet mass $+ 0.00004$, $r^2 = 0.99$; relatively short-photoperiod diapause-destined larvae: dry mass = $0.157 \times$ wet mass $+ 0.00003$, $r^2 = 0.99$). We used growth, relative growth, consumption, approximate digestibility (AD), the efficiency of converting ingested food into body matter (ECI), and the efficiency of conversion of digested food into body matter (ECD) to analyze the consumption and feeding efficiency. The formulae used to calculate these nutritional indices are as follows (adapted from Mole and Zera, 1993): growth = [dry body weight at the end day – initial dry body weight]; relative growth = [growth/initial dry body weight]; consumption = dry weight of food ingested during the feeding trial; AD = [(dry weight of food ingested – dry weight of feces)/dry weight of food ingested] $\times 100\%$; ECI = (dry weight gain/dry feed ingested); and ECD = growth/(dry weight of food ingested – dry weight of feces).

Enzyme activities

Two hours after the beginning of the photophase on the 3rd day after molting, the 5th and 6th instar larvae were transferred to an empty plastic container without food. At the beginning of the photophase on the 4th day after molting, the larvae, which had been starved for 22 h, were fed *ad libitum* for 6 h. Groups of larvae from both instars were then transferred to -40°C . To dissect the midgut, frozen larvae were removed from the freezer, thawed on ice, and pinned ventral side up on a silicon surface and cut longitudinally. The digestive tract was then transferred to a glass slide. The midgut was isolated and placed in a 1.5 ml microcentrifuge tube containing 200 μl of 0.15 mol l^{-1} NaCl. The midgut tissue was disrupted by sonification and centrifuged at 10,500 g for 15 min at 4°C . The supernatant was used in the enzyme assays. Amylase activity was measured by the dinitrosalicylic acid (DNS) method of Bernfeld (1955). Enzyme stock solution (20 μl) was added to 2 ml of phosphate buffer (pH 7.2) containing 250 μg of soluble starch (Sigma-Aldrich, Louis), and 2 ml of the DNS reagent was then added to the solution. The solution was heated to 100°C for 10 min. After cooling in water, absorbance was measured at 530 nm. To assay the lipase activity, the enzyme stock solution (20 μl) was mixed with 1.5 ml of phosphate buffer (pH 7.2) containing 80 nmol triglyceride. The concentration of free fatty acids released was measured using a test kit (Nanjing Jiancheng Bioengineering Institute, Jiangsu, China) and measuring the absorbance at 550 nm. The protease activities were assayed using a test kit (Nanjing Jiancheng Bioengineering Institute) according to the manufacturer's instructions and measuring the absorbance at 660 nm.

Statistical analysis

The experimental data were analyzed in SPSS 22.0 (IBM Inc., New York, USA). General linear model (GLM) was used to

analyze the body size and dry mass between short-photoperiod diapause and relatively short-photoperiod diapause pupae. Student's *t*-test was used to compare lipase, amylase, and protease activity of short-photoperiod diapause-destined larvae and relatively short-photoperiod diapause-destined larvae within the same instars. The lipid and glycogen between short-photoperiod diapause and relatively short-photoperiod diapause pupae were analyzed using an analysis of covariance (ANCOVA) with total protein (mg protein per individual) as a covariate (Williams *et al.*, 2012) and followed by Tukey's multiple range test. The content of protein between short-photoperiod diapause and relatively short-photoperiod diapause pupae was analyzed using analysis of variance with post-hoc Tukey's multiple range test. Growth, relative growth, consumption, ECI, ECD, and AD were analyzed with ANCOVA with the numerator of the ratio as the independent variable and the denominator as the covariate (Raubenheimer and Simpson, 1992; Mole and Zera, 1993).

Results

Body size and dry mass

The result of GLM analysis revealed that photoperiod and sex had a significant effect on body size and dry mass (width: $F_{1,171} = 47.2$ and 124.5 for photoperiod and sex; length: $F_{1,171} = 25.2$ and 90.34 for photoperiod and sex; body mass: $F_{1,171} = 53.7$ and 180.3 for photoperiod and sex; $P < 0.001$ for all treatments, fig. 1), and body size and dry mass of short-photoperiod diapause pupae were significantly greater than those of the relatively short-photoperiod diapause pupae ($P < 0.001$ for all treatments, fig. 1).

Body composition

We observed significant differences in lipid content between short-photoperiod diapause pupae and relatively short-photoperiod diapause pupae (female: $F_{1,64} = 74.12$, $P < 0.001$; male: $F_{1,74} = 92.34$, $P < 0.001$, fig. 2a), time (days after pupation) had no effect on the content of lipid (female: $F_{2,70} = 0.016$, $P = 0.985$; male: $F_{2,80} = 0.11$, $P = 0.89$, fig. 2a), and there was no interaction between pupae development status and time (female: $F_{2,69} = 1.11$, $P = 0.34$; male: $F_{2,79} = 2.71$, $P = 0.073$, fig. 2a). The lipid content within treatments for female diapause pupae did not change significantly within 5 days after pupation, but lipid content of short-photoperiod diapause pupae was, on average, 26% higher than the relatively short-photoperiod group ($P < 0.001$ for all days, fig. 2a). Similar to the female diapause pupae, the lipid contents of males in the two groups of diapause pupae remained stable within 5 days after pupation, but the lipid content of short-photoperiod diapause pupae was, on average, 39% higher than the relatively short-photoperiod group ($P < 0.001$ in all days, fig. 2a).

We observed significantly higher glycogen content in short-photoperiod diapause pupae relative to relatively short-photoperiod diapause pupae (female: $F_{1,69} = 100.87$, $P < 0.001$; male: $F_{1,94} = 22.86$, $P < 0.001$, fig. 2b), time (days after pupation) had no effect on the content of glycogen (female: $F_{2,75} = 0.075$, $P = 0.93$; male: $F_{2,100} = 0.084$, $P = 0.92$, fig. 2b), and there was no interaction between pupae development status and time (female: $F_{2,74} = 0.04$, $P = 0.96$; male: $F_{2,99} = 0.049$, $P = 0.95$, fig. 2b). The glycogen content of two groups of female diapause pupae did not change significantly and was maintained at a relatively high and low level, respectively, but a significant difference was detected between them at 1st, 3rd, and 5th days after pupation




Figure 1. Comparison of the body size and dry mass of relatively short photoperiod and short photoperiod diapause pupae. (a) Dry mass (in mg). (b) Body length (cm). (c) Body width (cm). We measured the width of the pupae at the basal margin of the abdomen. White: short photoperiod diapause pupae; gray: relatively short photoperiod diapause pupae. Different letters indicate significant differences between relatively short photoperiod diapause and short photoperiod diapause pupae at $P < 0.05$ according to *t*-tests ($n = 40$ for short photoperiod female diapause pupae; $n = 45$ for relatively short photoperiod female diapause pupae, relatively short and short photoperiod male diapause, respectively).




Figure 2. Lipid (a), carbohydrate (b), and protein (c) contents on the 1st, 3rd, and 5th days after pupation in relatively short photoperiod and short photoperiod diapause pupae. White: short photoperiod diapause pupae; gray: relatively short photoperiod diapause pupae. Asterisks indicate significant differences between relatively short photoperiod diapause and short photoperiod diapause pupae of the same age ($P < 0.05$), with NS indicating no significant difference ($P > 0.05$) ($n = 10-30$).




Figure 3. Ingested food during 8th day treatment in relatively short-photoperiod diapauses and short-photoperiod diapauses larvae. White: short-photoperiod diapauses; gray: relatively short-photoperiod diapauses. Asterisks indicate significant differences between relatively short-photoperiod diapauses and short-photoperiod diapauses larvae ($P < 0.05$), with NS indicating no significant difference ($P > 0.05$) (relatively short-photoperiod diapauses larval: $n = 26$; short-photoperiod diapauses larval: $n = 25$).

($P < 0.001$ in all days, [fig. 2b](#)). Similar to the female diapauses pupae, the glycogen content of two groups of male diapauses pupae did not change significantly and was maintained at relatively high and low levels, respectively, with significantly higher glycogen levels observed in the short-photoperiod male diapauses pupae compared to the relatively short-photoperiod male diapauses pupae on the 1st, 3rd, and 5th days after pupation (day 1: $P = 0.010$; day 3: $P = 0.015$; day 5: $P = 0.027$, [fig. 2b](#)).

The protein contents of two groups of female diapauses pupae, as well as the two groups of male diapauses pupae, remained stable within 5 days after pupation. There is no significant difference between short-photoperiod diapauses pupae and relatively short-photoperiod diapauses pupae (female: $F_{1,73} = 2.19$, $P = 0.143$; male: $F_{1,93} = 1.29$, $P = 0.28$, [fig. 3c](#)), time (days after pupation) had no effect on the content of protein (female: $F_{2,79} = 1.98$, $P = 0.144$; male: $F_{2,99} = 0.089$, $P = 0.92$, [fig. 3c](#)), and there was no interaction between pupae development status and time (female: $F_{2,78} = 0.23$, $P = 0.79$; male: $F_{2,98} = 0.56$, $P = 0.57$, [fig. 2c](#)).

Food utilization

During the 8th day experimental treatment, the growth of relatively short-photoperiod diapauses larvae was only 81.7% of that of short-photoperiod diapauses larvae ($F_{1,48} = 128.24$, $P < 0.001$, [table 1](#)). Relative growth of short-photoperiod diapauses larvae was also higher than that

of the relatively short-photoperiod diapauses larvae ($F_{1,48} = 246.52$, $P < 0.001$, [table 1](#)). Similar to the growth and relative growth, the ECI and ECD were significantly higher in the short-photoperiod diapauses larvae than the relatively short-photoperiod diapauses larvae (ECI: $F_{1,48} = 6.46$, $P = 0.0087$; ECD: $F_{1,48} = 9.82$, $P = 0.0054$, [table 1](#)). However, AD did not differ between relatively short-photoperiod diapauses larvae and short-photoperiod diapauses larvae ($F_{1,48} = 1.76$, $P = 0.55$, [table 1](#)).

The total food consumption of short-photoperiod diapauses larvae was similar to that of the relatively short-photoperiod diapauses larvae ($F_{1,48} = 0.27$, $P = 0.31$, [fig. 3](#)), but there was an obvious difference in the pattern of food intake between the relatively short-photoperiod diapauses larvae and short-photoperiod diapauses larvae. The consumption by relatively short-photoperiod diapauses larvae and short-photoperiod diapauses larvae declined from 0 to 4 days, and then increased during days 4–8. During 0–2 and 2–4 days, the consumption by short-photoperiod diapauses larvae was significantly higher than that of relatively short-photoperiod diapauses larvae (0–2 days: $F_{1,48} = 12.46$, $P = 0.0041$; 2–4 days: $F_{1,48} = 6.52$, $P = 0.022$, [fig. 3](#)); by contrast, consumption was significantly greater in relatively short-photoperiod diapauses larvae than the short-photoperiod diapauses larvae during 4–6 and 6–8 days, respectively (4–6 days: $F_{1,48} = 5.54$, $P = 0.036$; 6–8 days: $F_{1,48} = 11.44$, $P = 0.0052$, [fig. 3](#)).

Digestive enzyme activities

Both the lipase and amylase activities in the midguts of the short-photoperiod diapauses larvae and the relatively short-photoperiod diapauses larvae increased after the 5th instar larvae molted to the 6th instar larvae. However, the lipase and amylase activities in the midguts of the 5th and 6th instar of short-photoperiod diapauses larvae were significantly higher than those of the relatively short-photoperiod diapauses larvae (5th instar: $t_{45} = 2.43$, $P = 0.019$; 6th instar: $t_{44} = 3.46$, $P = 0.0017$ for lipase activity; 5th instar: $t_{45} = 2.16$, $P = 0.036$; 6th instar: $t_{44} = 2.11$, $P = 0.023$ for amylase activity, [figs 4a, b](#)). The protease activities in the midguts of the short-photoperiod diapauses larvae and relatively short-photoperiod diapauses larvae 5th instar larvae underwent a slight decline after the 5th instar larvae molted to the 6th instar larvae but the protease activities did not differ significantly between the two groups of larvae at either instar (5th instar: $t_{45} = 0.77$, $P = 0.44$; 6th instar: $t_{44} = 1.24$, $P = 0.22$, [fig. 4c](#)).

Discussion

Consistent with our predictions, the diapausing pupae exposed to a short photoperiod during larval development had significantly

Table 1. Mean nutritional indices in relatively short-photoperiod diapauses larvae and short-photoperiod diapauses larvae of fall webworm

Larval development	Growth (mg)	Relative growth (mg mg^{-1})	ECI (mg mg^{-1})	ECD (mg mg^{-1})	AD (%)
SPDD	32.3 ± 2.4a	6.5 ± 0.4a	11.8 ± 0.6a	14.6 ± 0.9a	81.9 ± 0.8a
RPDD	26.4 ± 1.8b	4.3 ± 0.31b	9.3 ± 0.7b	11.6 ± 0.2b	81.4 ± 0.9a

SPDD, short-photoperiod diapauses; RPDD, relatively short-photoperiod diapauses. Different letters after means within a column indicate significant differences between the two treatments (relatively short-photoperiod diapauses larval: $n = 26$; short-photoperiod diapauses larval: $n = 25$).




Figure 4. Comparison of digestive enzyme activities in the midguts of relatively short-photoperiod diapause-destined and short-photoperiod diapause-destined 5th and 6th instar larvae. (a) Amylase activity (in U mg^{-1}). (b) Lipase activity (in U mg^{-1}). (c) Protease activity (in U mg^{-1}). White: short-photoperiod diapause-destined larvae; gray: relatively short-photoperiod diapause-destined larvae. Asterisks indicate significant differences between relatively short-photoperiod diapause-destined and short-photoperiod diapause-destined larvae of the same age ($P < 0.05$), with NS indicating no significant difference ($P > 0.05$) ($n = 24$ for relatively short-photoperiod diapause-destined 5th instar larvae; $n = 23$ for short-photoperiod diapause-destined 5th instar larvae; $n = 23$ for relatively short-photoperiod and short-photoperiod diapause-destined 6th instar larvae, respectively).

higher body size and dry mass, as well as higher lipid and carbohydrate content, relative to those exposed to a longer photoperiod, suggesting that the short-photoperiod diapausing pupae reserved more energy. These results indicate that the energy reserve pattern of diapausing fall webworm pupae is plastic in response to photoperiod during diapause induction phase.

Diapause is promoted by environmental factors during the diapause induction phase. Insects that are destined for diapause undergo a suite of physiological changes, such as changes in body size and energy reserves, during the diapause preparation phase (Tauber *et al.*, 1986; Danks, 1987; Denlinger *et al.*, 2005). Our results showed that body size and dry mass were significantly higher for both female and male fall webworm diapause pupae subjected to a short photoperiod compared to those that had a relatively short photoperiod during the diapause induction phase. The larger size of the diapause individuals may provide more energy stores to meet the requirements for surviving under harsh environmental conditions and completing metamorphosis in the spring (Hahn and Denlinger, 2007; Shin *et al.*, 2012). We found that the lipid and carbohydrate contents were significantly greater in female diapause pupae exposed to a short photoperiod during the diapause induction phase than female diapause pupae exposed to a relatively short photoperiod. We observed a similar trend in male diapause pupae. These results indicate that the energy storage mode in fall webworm diapause pupae is influenced by photoperiod during the diapause induction phase, where a short photoperiod below the critical day length can promote increased storage of energy for diapause.

Insects use two strategies in order to meet energy consumption requirements during diapause: (1) storing energy reserves during the diapause preparation phase, and (2) inhibiting metabolism during diapause (Hahn and Denlinger, 2011). Diapause can reduce the energy consumed by insects, but an increase in the temperature during diapause accelerates the metabolic rate and increases the energy consumed while wintering, thereby decreasing the survival rate of insects during wintering or lowering the fitness of the population post-diapause (Irwin and Lee, 2003; Schulte *et al.*, 2011; Williams *et al.*, 2012). Therefore, the accumulation of nutrients in the diapause preparation phase is crucial because any lack of nutrition will reduce the likelihood of insects completing diapause, thereby decreasing their chances of survival and reproduction (Denlinger, 2002). The energy reserves stored before diapause mainly comprise lipid and carbohydrates (Hahn and Denlinger, 2007; Sinclair, 2015; Sinclair and Marshall, 2018).

Fall webworm was introduced into Dandong city (about 40°N), China during 1979 and it spread to Nanjing city (about 32°N) in Jiangsu Province. The winter temperature exhibits a clear geographical latitudinal gradient from Dandong to Nanjing (Liu *et al.*, 2016). Therefore, after fall webworm spread from the north to south following its introduction into China, the southern population likely experienced a strong selective pressure due to an increased metabolic rate and higher energy consumption caused by warmer winters. We consider that the short photoperiod induces fall webworm pupae to store more lipid and carbohydrates in order to increase their energy stores before overwintering and this change will increase the fitness of the population after overwintering following the spread from north to south.

In order to store more energy, diapause individuals must ingest more food, increase their food utilization efficiency, divert nutrients from supporting somatic cell growth to storage, or a combination of these processes (Hahn and Denlinger, 2007). Our results

showed that the short-photoperiod diapause-destined larvae stored more lipid and carbohydrates compared with the relatively short-photoperiod diapause-destined larvae. The growth rate, relative growth rate, ECI, and ECD were significantly higher for the short-photoperiod diapause-destined larvae, but there was no significant difference in the food intake between the two larval treatments. These results indicate that the short photoperiod under the critical photoperiod stimulated the diapause-destined larvae to enhance their energy reserves by increasing ECI and ECD rather than increasing their food intake during diapause preparation. This is possibly an adaptive strategy in response to changes in the quantity and quality of food that is likely to occur in late autumn as plants begin to senesce. Thus, if the final generation of fall webworm (i.e. third generation of trivoltine populations) could not enhance its energy reserves by increasing the intake of food, then improving the efficiency of food utilization could also provide sufficient resources to successfully overwinter. In addition, variation in the pattern of food intake between the relatively short-photoperiod diapause-destined larvae and short-photoperiod diapause-destined larvae reflected the different feeding strategies employed by the two groups of larvae. The use of artificial diet in our experiment did not reflect the natural changes in the food quality over time or the gradual fading of autumn leaves under natural environmental conditions, but the increased early feed intake by the diapause-destined larvae with a short photoperiod was a better strategy that allowed them to cope with the declining leaf quality in the autumn.

The short-photoperiod diapause pupae stored more lipid and carbohydrates, and the ECI and ECD values were significantly higher for the short-photoperiod diapause-destined larvae than the relatively short-photoperiod larvae. Therefore, we suggest that the different energy reserve patterns of the short-photoperiod and relatively short-photoperiod diapause pupae may have been related to differences in the activities of digestive enzyme in the midgut. In insects, digestive enzyme activities are affected by the type of food consumed, larval density, larval stage, and photoperiod (Woodring *et al.*, 2007; Kong *et al.*, 2012; Weidlich *et al.*, 2012, 2013; Zhao *et al.*, unpublished). The elevated midgut lipase and amylase activities that we observed in short-photoperiod diapause-destined larvae are consistent with the higher lipid and carbohydrate contents in diapausing pupae. Thus, diapause-destined larvae exposed to a short photoperiod increased the lipase and amylase activities in their midgut in order to store more lipid and carbohydrates.

Our results support the hypothesis that photoperiod influences energy reserve accumulation during diapause preparation in fall webworm. Consistent with our expectations, the body size and dry mass were significantly larger in the short photoperiod diapause pupae than the relatively short photoperiod pupae. Moreover, the lipid and carbohydrate contents of the short photoperiod diapause pupae were significantly higher than those of the relatively short photoperiod diapause pupae. The increased energy reserves were due to changes in the lipase and protease activities in the midgut of short photoperiod diapause-destined larval. The results obtained in this study indicate that the photoperiod can regulate energy accumulation in fall webworm. A shorter photoperiod, below the critical photoperiod, promoted larvae to store more lipid and carbohydrates during diapause preparation. These increased energy reserves will likely help this species to overcome increased energetic demands in locations with warmer winters.

Acknowledgements. We thank Jia-heYan for his assistance on the collection of the fall webworm. This study was supported by the Natural Science

Foundation of Jiangsu Province (BK20181399) and the Highly Educated Talents Foundation in Nanjing Forestry University (GXL2017001).

Author contributions. L.Z. designed the experimental protocol. W.W. and Y.Q. carried out the experimental work. L.Z. analyzed the data. L.Z. and A.S.T. wrote the manuscript.

References

- Bernfeld MM (1955) Amylases, a and b. In Colowick SP and Kaplan N (eds), *Methods in Enzymology*, vol. I. New York, NY, USA: Academic Press, 149–158.
- Cao LJ, Yang F, Tang SY and Chen M (2014) Development of an artificial diet for three lepidopteran insects. *Chinese Journal of Applied Entomology* 51, 1376–1386.
- Carroll NV, Longley RW and Roe JH (1956) Determination of glycogen in liver and muscle by use of anthrone reagent. *Journal of Biological Chemistry* 220, 583–593.
- Chen YS, Chen C, He HM, Xia QW and Xue FS (2013) Geographic variation in diapause induction and termination of the cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae). *Journal of Insect Physiology* 59, 855–862.
- Chen C, Wei XT, Xiao HJ, He HM, Xia QW and Xue FS (2014) Diapause induction and termination in *Hyphantria cunea* (Drury) (Lepidoptera: Arctiinae). *PLoS ONE* 9, e98145. doi: 10.1371/journal.pone.0098145
- Danks HV (1987) *Insect Dormancy: An Ecological Perspective*. Biological Survey of Canada (Terrestrial Arthropods).
- Denlinger DL (1991) Relationship between cold hardiness and diapause. In Lee Jr. RE and Denlinger DL (eds), *Insects at Low Temperature*. New York: Chapman & Hall, 174–198.
- Denlinger DL (2002) Regulation of diapause. *Annual Review of Entomology* 47, 93–122.
- Denlinger DL, Yocum GD and Rinehart JL (2005) Hormonal control of diapause. In Gilbert LI, Iatrou K and Gill SS (eds), *Comprehensive Molecular Insect Science*, vol. 3. Amsterdam: Elsevier Press, 615–650.
- Gomi T (1997) Geographic variation in critical photoperiod for diapause induction and its temperature dependence in *Hyphantria cunea* Drury (Lepidoptera: Arctiidae). *Oecologia* 111, 160–165.
- Hahn DA and Denlinger DL (2007) Meeting the energetic demands of insect diapause: nutrient storage and utilization. *Journal of Insect Physiology* 53, 760–773.
- Hahn DA and Denlinger DL (2011) Energetics of insect diapause. *Annual Review of Entomology* 56, 103–121.
- Han B and Denlinger DL (2009) Mendelian inheritance of pupal diapause in the flesh fly, *Sarcophaga bullata*. *Journal of Heredity* 100, 251–255.
- Irwin JT and Lee Jr RE (2003) Cold winter microenvironments conserve energy and improve overwintering survival and potential fecundity of the goldenrod gall fly, *Eurosta solidaginis*. *Oikos* 100, 71–78.
- Ji R, Xi BY, Li XH, Gao ZX and Li DM (2003) Research progress on the invasive species *Hyphantria cunea*. *Entomological Knowledge* 40, 13–18.
- Kong HL, Luo LZ, Jiang XF, Zhang L, Yang ZL and Hu Y (2012) Effects of larval density on food utilization efficiency and digestive enzyme activity of the beet webworm, *Loxostege sticticalis* (Lepidoptera: Pyralidae). *Acta Entomologica Sinica* 55, 361–366.
- Košťál V (2006) Eco-physiological phases of insect diapause. *Journal of Insect Physiology* 52, 113–127.
- Liu J, Ye LQ, Cheng G, Xiong DB and Chen ZY (2016) The occurrence of forestry pests in 2015 and its occurrence trend prediction in 2016 in Jiangsu Province. *Journal of Jinling Institute of Technology* 32, 64–67.
- Loewy K, Flansburg AL, Grenis K and Murphy SM (2013) Life history traits and rearing techniques for fall webworms (*Hyphantria cunea* Drury) in Colorado. *Journal of the Lepidopterists' Society* 67, 196–205.
- Lorenz MW (2003) Adipokinetic hormone inhibits the formation of energy stores and egg production in the cricket *Gryllus bimaculatus*. *Comparative Biochemistry and Physiology B* 136, 197–206.
- Masaki S, Umeya K, Sekiguchi Y and Kawasaki R (1968) Biology of *Hyphantria cunea* Drury (Lepidoptera : Arctiidae) in Japan III. Photoperiodic induction of diapause in relation to the seasonal life cycle. *Applied Entomology and Zoology* 3, 55–66.

- Mole S and Zera AJ** (1993) Differential allocation of resources underlies the dispersal-reproduction trade-off in the wingdimorphic cricket, *Gryllus rubens*. *Oecologia* **93**, 121–127.
- Raubenheimer D and Simpson SJ** (1992) Analysis of covariance: an alternative to nutritional indices. *Entomologia Experimentalis et Applicata* **62**, 221–231.
- Rehnberg BG** (2006) Temperature profiles inside webs of the fall webworm, *Hyphantria cunea* (Lepidoptera: Arctiidae): influence of weather, compass orientation, and time of day. *Journal of Thermal Biology* **31**, 274–279.
- Schulte PM, Healy TM and Fangue NA** (2011) Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology* **51**, 691–702.
- Shin SM, Akram W and Lee JJ** (2012) Effect of body size on energy reserves in *Culex pipiens pallens* females (Diptera: Culicidae). *Entomological Research* **42**, 163–167.
- Sinclair BJ** (2015) Linking energetics and overwintering in temperate insects. *Journal of Thermal Biology* **54**, 5–11.
- Sinclair BJ and Marshall KE** (2018) The many roles of fats in overwintering insects. *Journal of Experimental Biology* **221**, jeb161836.
- Tan QQ, Feng L, Liu W, Zhu L, Lei CL and Wang XP** (2016) Differences in the pre-diapause and pre-oviposition accumulation of critical nutrients in adult females of the beetle *Colaphellus bowringi*. *Entomologia Experimentalis et Applicata* **160**, 117–125.
- Tanaka S** (1991) De-alation and its influences on egg production and flight muscle histolysis in a cricket (*Velarifictorus parvus*) that undergoes inter-reproductive migration. *Journal of Insect Physiology* **37**, 517–523.
- Tanaka S and Zhu DH** (2008) Geographic variation of embryonic diapause, cold-hardiness and life cycles in the migratory locust, *Locusta migratoria* (Orthoptera: Acrididae), in China. *Entomological Science* **11**, 327–339.
- Tanaka S, Arai T and Tanaka K** (1999) Nymphal development, diapause and cold-hardiness in a nymph-overwintering cricket. *Entomological Science* **2**, 173–182.
- Tauber MJ, Tauber CA and Masaki S** (1986) *Seasonal Adaptations of Insects*. New York: Oxford University Press, 411.
- Wei XT, Xiao HJ, Yang D, Wu SH and Xue FS** (2006) Several important biological characters in the fall webworm, *Hyphantria cunea*. *Plant Quarantine* **1**, 14–17.
- Weidlich S, Huster J, Hoffmann KH and Woodring J** (2012) Environmental control of trypsin secretion in the midgut of the two-spotted field cricket, *Gryllus bimaculatus*. *Journal of Insect Physiology* **58**, 1477–1484.
- Weidlich S, Müller S, Hoffmann KH and Woodring J** (2013) Regulation of amylase, cellulase and chitinase secretion in the digestive tract of the two-spotted field cricket, *Gryllus bimaculatus*. *Archives of Insect Biochemistry and Physiology* **83**, 69–85.
- Williams CM, Chick WD and Sinclair BJ** (2015) A cross-seasonal perspective on local adaptation: metabolic plasticity mediates responses to winter in a thermal-generalist moth. *Functional Ecology* **29**, 549–561.
- Williams CM, Marshall KE, MacMillan HA, Dzurisin JDK, Hellmann JJ, et al.** (2012) Thermal variability increases the impact of autumnal warming and drives metabolic depression in an overwintering butterfly. *PLoS ONE* **7**, e34470. doi: doi: 10.1371/journal.pone.0034470
- Woodring J, Hoffmann K and Lorenz M** (2007) Activity, release and flow of digestive enzymes in the cricket, *Gryllus bimaculatus*. *Physiological Entomology* **32**, 56–63.
- Yang HZ, Tu XY, Xia QW, He HM, Chen C and Xue FS** (2014) Photoperiodism of diapause induction and diapause termination in *Ostrinia furnacalis* (Lepidoptera: Crambidae). *Entomologia Experimentalis et Applicata* **153**, 34–46.
- Zhao LQ, Liao HY, Zeng Y, Wu HJ and Zhu DH** (2017) Food digestion capability and digestive enzyme activity in female adults of the wing-dimorphic cricket *Velarifictorus ornatus*. *Entomologia Experimentalis et Applicata* **163**, 35–42.
- Zhao LQ, Wang W, Qiu Y and Torson AS**. Physiological mechanisms of variable nutrient accumulation patterns between diapausing and non-diapausing fall webworm (Lepidoptera: Arctiidae) pupae (unpublished).