

ON PSEUDO-NULLITY OF THE FINE MORDELL–WEIL GROUP

MENG FAI LIM[✉], CHAO QIN[✉] and JUN WANG

(Received 6 September 2024; accepted 26 December 2024)

Abstract

Let E be an elliptic curve defined over \mathbb{Q} with good ordinary reduction at a prime $p \geq 5$ and let F be an imaginary quadratic field. Under appropriate assumptions, we show that the Pontryagin dual of the fine Mordell–Weil group of E over the \mathbb{Z}_p^2 -extension of F is pseudo-null as a module over the Iwasawa algebra of the group \mathbb{Z}_p^2 .

2020 Mathematics subject classification: primary 11G05; secondary 11R23, 11S25.

Keywords and phrases: fine Mordell–Weil group, pseudo-nullity.

1. Introduction

At the turn of the millennium, Coates and Sujatha [4] and, a little later, Wuthrich [22] initiated a systematic study on the fine Selmer group of an elliptic curve E . The fine Selmer group is a subgroup of the classical p -primary Selmer group defined by stricter local conditions at primes above p . Analogous to the usual Selmer group, this fine Selmer group $R(E/F)$ sits in the middle of the short exact sequence

$$0 \longrightarrow \mathcal{M}(E/F) \longrightarrow R(E/F) \longrightarrow \mathcal{XK}(E/F) \longrightarrow 0,$$

where $\mathcal{M}(E/F)$ and $\mathcal{XK}(E/F)$ are the fine Mordell–Weil group and fine Tate–Shafarevich group, respectively (defined in the sense of Wuthrich [23]), which can be thought as the ‘fine’ counterparts of the usual Mordell–Weil group and Tate–Shafarevich group.

In [4, Conjecture B], Coates and Sujatha proposed the following conjecture.

CONJECTURE 1.1 (Conjecture B). Let E be an elliptic curve defined over a number field F . Suppose that F_∞ is a p -adic Lie extension of F for which $\text{Gal}(F_\infty/F)$ has dimension ≥ 2 containing the cyclotomic \mathbb{Z}_p -extension F^{cyc} . Then, $R(A/F_\infty)^\vee$ is pseudo-null over $\mathbb{Z}_p[[\text{Gal}(F_\infty/F)]]$.

Chao Qin’s research is supported by the National Natural Science Foundation of China under Grant No. 12001546, Heilongjiang Province under Grant No. 3236330122 and Harbin Engineering University under Grant No. GK0000020127. Jun Wang’s research is supported by the National Natural Science Foundation of China under Grant No. 12331004.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.



Here, a finitely generated $\mathbb{Z}_p[[\text{Gal}(F_\infty/F)]]$ -module M is said to be pseudo-null if

$$\text{Ext}_{\mathbb{Z}_p[[\text{Gal}(F_\infty/F)]]}^i(M, \mathbb{Z}_p[[\text{Gal}(F_\infty/F)]]) = 0 \quad \text{for } i = 0, 1.$$

This conjecture remains wide open but we refer readers to [2, 7, 11, 12] for some discussion and numerical examples in support of the conjecture. The goal of this note is to provide further theoretical support for Conjecture B. We will prove the following result.

THEOREM 1.2. *Let E be an elliptic curve defined over \mathbb{Q} with good ordinary reduction at a prime $p \geq 5$, and let F be an imaginary quadratic field. Suppose that E has no complex multiplication and that the discriminant of F is coprime to the conductor of E . In the event that the root number $\epsilon(E/F, 1)$ equals -1 , assume further that p does not divide the class number of F . Suppose that $\text{Sel}(E/F^{\text{cyc}})$ is cofinitely generated over \mathbb{Z}_p . Then, the Pontryagin dual $\mathcal{M}(E/F_\infty)^\vee$ of the fine Mordell–Weil group is a pseudo-null $\mathbb{Z}_p[[\text{Gal}(F_\infty/F)]]$ -module, where F_∞ is the \mathbb{Z}_p^2 -extension of F .*

As an application, we can prove the following result, where we note that every \mathbb{Z}_p -extension of F is necessarily contained in F_∞ .

COROLLARY 1.3. *Retain the settings of Theorem 1.2. Then, for every \mathbb{Z}_p -extension L_∞ of F , the Pontryagin dual $\mathcal{M}(E/L_\infty)^\vee$ is torsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/L)]]$.*

It is conjectured that $R(E/L_\infty)^\vee$ should always be torsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/L)]]$ (see [13, 18, 21]). The above corollary thus constitutes partial evidence towards this conjecture. Furthermore, if one is willing to assume the finiteness of the fine Tate–Shafarevich group, we have the following observation.

COROLLARY 1.4. *Retain the settings of Theorem 1.2. Suppose that L_∞ is a \mathbb{Z}_p -extension of F with the property that $\text{XK}(E/L)$ is finite for every finite extension L of F contained in L_∞ . Then, $R(E/L_\infty)^\vee$ is torsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/L)]]$.*

The proofs of the preceding theorem and corollaries will be given in Section 4.

2. Selmer group

We now introduce the Selmer group in a slightly more general context. As a start, we let F be an arbitrarily fixed number field and E an elliptic curve defined over F . Let S be a finite set of primes of F which contains all the primes above p , the infinite primes and the primes of bad reduction of E . We shall also write S_p for the set of primes of F above p . Denote by F_S the maximal algebraic extension of F which is unramified outside S . If \mathcal{L} is a (possibly infinite) extension of F contained in F_S and $S' \subseteq S$, we write $S'(\mathcal{L})$ for the set of primes of \mathcal{L} above S' .

For each $v \in S$ and a finite extension L of F , set

$$J_v(E/L) = \bigoplus_{w|v} H^1(L_w, E)_{p^\infty}.$$

If \mathcal{L} is an infinite extension of F contained in F_S , we define

$$J_v(E/\mathcal{L}) = \varinjlim_L J_v(E/L),$$

where L runs through all finite extensions of F contained in \mathcal{L} .

The classical (p -primary) Selmer group of E over \mathcal{L} is defined by

$$\text{Sel}(E/\mathcal{L}) = \ker \left(H^1(G_S(\mathcal{L}), E_{p^\infty}) \longrightarrow \bigoplus_{v \in S} J_v(E/\mathcal{L}) \right),$$

where we write $G_S(\mathcal{L}) = \text{Gal}(F_S/\mathcal{L})$. The Pontryagin dual of $\text{Sel}(E/\mathcal{L})$ is then denoted by $X(E/\mathcal{L})$.

From now on, we will always assume that the elliptic curve E has good ordinary reduction at all primes of F above p . For each prime v of F above p , denote by \hat{E}_v and \tilde{E}_v the formal group of E at v and the reduced curve of E at v , respectively. Furthermore, we have a short exact sequence

$$0 \longrightarrow \hat{E}_{v,p^\infty} \longrightarrow E_{p^\infty} \longrightarrow \tilde{E}_{v,p^\infty} \longrightarrow 0$$

of discrete $\text{Gal}(\bar{F}_v/F_v)$ -modules. Since E is assumed to have good ordinary reduction, both \hat{E}_{v,p^∞} and \tilde{E}_{v,p^∞} are cofree \mathbb{Z}_p -modules of corank 1.

It is convenient to work with an equivalent description of the local terms $J_v(E/\mathcal{L})$, following an insight of Coates and Greenberg [3]. Let \mathcal{L} be an algebraic extension of F . For every non-Archimedean prime w of \mathcal{L} , write \mathcal{L}_w for the union of the completions at w of the finite extensions of F contained in \mathcal{L} . If w is a prime above p , we write $E_w = E_v$, where v is a prime of F below w . Finally, we shall always denote by F^{cyc} the cyclotomic \mathbb{Z}_p -extension of F . With these in hand, we have the following lemma.

LEMMA 2.1. *Let \mathcal{L} be an algebraic extension of F^{cyc} which is unramified outside a set of finite primes of F . Then, we have an isomorphism*

$$J_v(E/\mathcal{L}) \cong \begin{cases} \varinjlim_{\mathcal{L}'} \bigoplus_{w|v} H^1(\mathcal{L}'_w, \tilde{E}_{w,p^\infty}) & \text{if } v \text{ divides } p, \\ \varinjlim_{\mathcal{L}'} \bigoplus_{w|v} H^1(\mathcal{L}'_w, E_{p^\infty}) & \text{if } v \text{ does not divide } p, \end{cases}$$

where the direct limit is taken over all finite extensions \mathcal{L}' of F^{cyc} contained in \mathcal{L} .

PROOF. See [3, Propositions 4.1, 4.7 and 4.8] or [10, Lemma 4.1]. □

We now establish a control theorem for the Selmer group over a \mathbb{Z}_p^2 -extension.

PROPOSITION 2.2. *Let E be an elliptic curve defined over F which has good ordinary reduction at each prime of F above p . Suppose that F_∞ is a \mathbb{Z}_p^2 -extension of F which contains F^{cyc} . Write $H = \text{Gal}(F_\infty/F^{\text{cyc}})$. Then, the restriction map*

$$\text{Sel}(E/F^{\text{cyc}}) \longrightarrow \text{Sel}(E/F_\infty)^H$$

has finite kernel and cokernel.

PROOF. The proposition should definitely be well known. For the convenience of the readers, we shall supply an argument here. Consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Sel}(E/F^{\text{cyc}}) & \longrightarrow & H^1(G_S(F^{\text{cyc}}), E_{p^\infty}) & \longrightarrow & \bigoplus_{v \in S} J_v(E/F^{\text{cyc}}) \longrightarrow 0 \\
 & & \downarrow & & \downarrow h & & \downarrow \oplus g_v \\
 0 & \longrightarrow & \text{Sel}(E/F_\infty)^H & \longrightarrow & H^1(G_S(F_\infty), E_{p^\infty})^H & \longrightarrow & \left(\bigoplus_{v \in S} J_v(E/F_\infty) \right)^H
 \end{array}$$

with exact rows. Via the snake lemma, it suffices to show that $\ker h$, $\ker g_v$ and $\text{coker } h$ are finite. To begin with, we show that h is surjective with a finite kernel. Indeed, since $H \cong \mathbb{Z}_p$, the restriction-inflation sequence tells us that the map h is surjective with kernel $H^1(H, E(F_\infty)_{p^\infty})$. However,

$$0 = \text{rank}_{\mathbb{Z}_p[[H]]} (E(F_\infty)_{p^\infty})^\vee = \text{rank}_{\mathbb{Z}_p} ((E(F_\infty)_{p^\infty})^\vee)_H - \text{rank}_{\mathbb{Z}_p} ((E(F_\infty)_{p^\infty})^\vee)^H,$$

where the second equality follows from [17, Proposition 5.3.20]. Nevertheless, observe that

$$((E(F_\infty)_{p^\infty})^\vee)_H = ((E(F_\infty)_{p^\infty})^H)^\vee = (E(F^{\text{cyc}})_{p^\infty})^\vee,$$

and the latter is finite by a theorem of Imai [6]. Hence, $((E(F_\infty)_{p^\infty})^\vee)^H$ is also finite. However, this group is precisely $H^1(H, E(F_\infty)_{p^\infty})^\vee$, and so we have our claim.

It remains to show that g_v has finite kernel for every v . Again, by the restriction-inflation sequence, we have $\ker g_v = \bigoplus_{w|v} H^1(H_w, D)$, where here the sum runs over all the primes of F^{cyc} above v , H_w is the decomposition group of w in H and D denotes either \tilde{E}_{v,p^∞} or E_{p^∞} according as to whether v divides p or not. In particular, if $H_w = 1$, then $H^1(H_w, D) = 0$. This is indeed the case when v does not divide p for a \mathbb{Z}_p^2 -extension that is unramified outside p . It remains to consider primes w which divide p and for which H_w is nontrivial. Since $H \cong \mathbb{Z}_p$, it then follows that $H_w \cong \mathbb{Z}_p$. We may then apply the same argument as in the preceding paragraph to conclude that $H^1(H_w, \tilde{E}_{v,p^\infty})$ is finite. The proof of the proposition is now complete. \square

The following is a corollary of the preceding proposition.

COROLLARY 2.3. *Retain the setting as in Proposition 2.2. Then, $\text{Sel}(E/F^{\text{cyc}})$ is cofinitely generated over \mathbb{Z}_p if and only if $\text{Sel}(E/F_\infty)$ is cofinitely generated over $\mathbb{Z}_p[[H]]$.*

3. Mordell–Weil group over a \mathbb{Z}_p^2 -extension

We continue to suppose that F_∞ is a \mathbb{Z}_p^2 -extension of F which contains F^{cyc} . Write $H = \text{Gal}(F_\infty/F^{\text{cyc}})$ and denote by H_n the unique subgroup of H with index p^n . The fixed field of H_n is in turn denoted by K_n . The following hypothesis will be in full force for the remainder of the section.

(Fg) $\text{Sel}(E/F^{\text{cyc}})$ is cofinitely generated over \mathbb{Z}_p .

Write $\Lambda = \mathbb{Z}_p[[H]]$. We shall identify this latter ring with the power series ring $\mathbb{Z}_p[[T]]$ in one variable. By abuse of notation, we shall also write Λ for the ring $\mathbb{Z}_p[[T]]$. Denote by Φ_n the p^n th-cyclotomic polynomial which is viewed as an element in Λ .

PROPOSITION 3.1. *Suppose that hypothesis (Fg) is valid. Write $\Lambda = \mathbb{Z}_p[[H]]$.*

(a) *There is an injective Λ -homomorphism*

$$(E(F_\infty) \otimes \mathbb{Q}_p/\mathbb{Z}_p)^\vee \longrightarrow \Lambda^{\oplus r} \oplus \left(\bigoplus_{n \geq 0} (\Lambda/\Phi_n)^{\oplus t_n} \right)$$

with a finite cokernel, where $\{t_n\}$ is a sequence of nonnegative integers with $t_n = 0$ for $n \gg 0$.

(b) *Define*

$$e_n = \begin{cases} \frac{\text{rank}_{\mathbb{Z}} E(K_n) - \text{rank}_{\mathbb{Z}} E(K_{n-1})}{p^n - p^{n-1}} & \text{if } n \geq 1, \\ \text{rank}_{\mathbb{Z}} E(F^{\text{cyc}}) & \text{if } n = 0. \end{cases}$$

Then,

$$T_p(E(K_n) \otimes \mathbb{Q}_p/\mathbb{Z}_p) \cong \bigoplus_{j=0}^n (\Lambda/\Phi_j)^{\oplus e_j}.$$

PROOF. By hypothesis (Fg), $\text{Sel}(E/K_n)$ is cofinitely generated over \mathbb{Z}_p for every n (for instance, see [5, Corollary 3.4]). Thus, it follows that $E(K_n) \otimes \mathbb{Q}_p/\mathbb{Z}_p$ is also cofinitely generated over \mathbb{Z}_p for every n . In view of this observation, we may apply a similar argument to that of Lee in [8, Theorem 2.1.2] to obtain the conclusion of statement (a). Once, we have statement (a), statement (b) will follow by carrying out the argument in [15, Proposition 3.8]. □

4. Fine Selmer groups

Let L be a finite extension of F contained in F_S . Recall that the fine Selmer group of E over L is defined by

$$R(E/L) = \ker \left(H^1(G_S(L), E_{p^\infty}) \longrightarrow \bigoplus_{v \in S(L)} H^1(L_v, E_{p^\infty}) \right).$$

The fine Selmer group and the classical Selmer group are related by the following exact sequence.

LEMMA 4.1. *We have an exact sequence*

$$0 \longrightarrow R(E/L) \longrightarrow \text{Sel}(E/L) \longrightarrow \bigoplus_{v \in S_p(L)} E(L_v) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p.$$

In particular, the definition of the fine Selmer group does not depend on the choice of the set S .

PROOF. See [16, Lemma 4.1] or [23, Section 2]. □

Following Wuthrich [23], the fine Mordell–Weil group $\mathcal{M}(E/L)$ is defined by

$$\mathcal{M}(E/L) = \ker \left(E(L) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p / \mathbb{Z}_p \longrightarrow \bigoplus_{v \in S_p(L)} E(L_v) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p / \mathbb{Z}_p \right),$$

where $S_p(L)$ denotes the set of primes of L above p . This fits into the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M}(E/L) & \longrightarrow & E(L) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p / \mathbb{Z}_p & \longrightarrow & \bigoplus_{v \in S_p(L)} E(L_v) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p / \mathbb{Z}_p \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & R(E/L) & \longrightarrow & \text{Sel}(E/L) & \longrightarrow & \bigoplus_{v \in S_p(L)} E(L_v) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p / \mathbb{Z}_p \end{array}$$

with exact rows, where the leftmost vertical map is induced by the middle vertical map. Following Wuthrich [23], the fine Tate–Shafarevich group $\check{\mathcal{K}}(A/L)$ is given by

$$\check{\mathcal{K}}(E/L) = \text{coker}(\mathcal{M}(E/L) \longrightarrow R(E/L)).$$

Since the middle vertical map in the above diagram is injective, so is the leftmost vertical map. Consequently, a snake lemma argument yields a short exact sequence

$$0 \longrightarrow \mathcal{M}(E/L) \longrightarrow R(E/L) \longrightarrow \check{\mathcal{K}}(E/L) \longrightarrow 0$$

with $\check{\mathcal{K}}(E/L)$ injecting into $\text{III}(E/L)_{p^\infty}$, the p -primary part of the usual Tate–Shafarevich group.

We can now give the proof of our main theorem.

PROOF OF THEOREM 1.2. Set C_n and C_∞ to be the respective cokernels of $\mathcal{M}(E/K_n) \longrightarrow E(K_n) \otimes_{\mathbb{Q}_p} \mathbb{Z}_p$ and $\mathcal{M}(E/F_\infty) \longrightarrow E(F_\infty) \otimes_{\mathbb{Q}_p} \mathbb{Z}_p$. From the argument in [19, Proposition 3.14], we see that

$$\text{rank}_{\mathbb{Z}_p[[H]]} (E(F_\infty) \otimes_{\mathbb{Q}_p} \mathbb{Z}_p)^\vee = \begin{cases} 0 & \text{if } \epsilon(E/F, 1) = +1; \\ 1 & \text{if } \epsilon(E/F, 1) = -1. \end{cases} \tag{4.1}$$

Plainly, if $\epsilon(E/F, 1) = +1$, then $\mathcal{M}(E/F_\infty)$ has trivial $\mathbb{Z}_p[[H]]$ -corank, and so by a result of Venjakob [20] (or see [12, Lemma 5.1]), $\mathcal{M}(E/F_\infty)^\vee$ is pseudo-null over $\mathbb{Z}_p[[G]]$.

Now suppose that we have $\epsilon(E/F, 1) = -1$. In view of (4.1) and the tautological short exact sequence

$$0 \longrightarrow \mathcal{M}(E/F_\infty) \longrightarrow E(F_\infty) \otimes_{\mathbb{Q}_p} \mathbb{Z}_p \longrightarrow C_\infty \longrightarrow 0,$$

it therefore remains to show that $\text{rank}_{\mathbb{Z}_p[[H]]} C_\infty > 0$.

Now, consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathcal{M}(E/K_n) & \longrightarrow & E(K_n) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p & \longrightarrow & \bigoplus_{w \in S_p(K_n)} E(K_{n,w}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p \\
 & & \downarrow \alpha_n & & \downarrow \beta_n & & \downarrow \gamma_n \\
 0 & \longrightarrow & \mathcal{M}(E/F_\infty)^{H_n} & \longrightarrow & (E(F_\infty) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p)^{H_n} & \longrightarrow & \left(\bigoplus_{u \in S_p(F_\infty)} E(F_{\infty,u}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p \right)^{H_n}
 \end{array}$$

with exact rows. Since $\ker \beta_n$ is contained in $H^1(H_n, E(F_\infty)_{p^\infty})$, it follows from the argument in Proposition 2.2 that the latter, and hence $\ker \beta_n$, is finite. Similarly, we see that $\ker \gamma_n$ is finite. Consequently, the map $C_n \rightarrow (C_\infty)^{H_n}$ also has a finite kernel.

We claim that $\text{corank}_{\mathbb{Z}_p} C_n \geq p^n - p^{n-1}$ for sufficiently large n . Suppose that this claim holds. Since we have seen above that the map $C_n \rightarrow (C_\infty)^H$ has finite kernel, it follows that $\text{corank}_{\mathbb{Z}_p} (C_\infty)^H \geq p^n - p^{n-1}$. However, since

$$\text{corank}_{\mathbb{Z}_p} (C_\infty)^H = (\text{corank}_{\mathbb{Z}_p[[H]]} C_\infty) p^n + O(1),$$

we therefore have

$$\text{corank}_{\mathbb{Z}_p[[H]]} C_\infty > 0,$$

which is what we want to show.

It therefore remains to establish our claim. For this, we follow an idea of Lei [9]. Consider the natural map

$$f_n : E(K_n) \otimes \mathbb{Q}_p/\mathbb{Z}_p \longrightarrow \bigoplus_{w \in S_p(K_n)} E(K_{n,w}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p/\mathbb{Z}_p,$$

where $\text{im } f_n = C_n$. Write F^{ac} for the anticyclotomic \mathbb{Z}_p -extension of F , and L_n for the intermediate subextension of F^{ac}/F with $|L_n : F| = p^n$. Since $\epsilon(E/F, 1) = -1$,

$$\text{rank}_{\mathbb{Z}} E(L_n) = p^n + O(1)$$

by [1, Proposition 7.6]. As $L_n \subseteq K_n$, this in turn implies that $e_n \geq 1$ for sufficiently large n , where e_n is defined as in Proposition 3.1. For each such n , choose an element $x_n \in E(K_n) \setminus E(K_{n-1})$ which is of infinite order. Then, for a prime w of K_n above p , the image of x_n in $E(K_{n,w})$ is still of infinite order and so cannot be divisible by an arbitrary power of p . Therefore, the image of x_n in $E(K_{n,w}) \otimes \mathbb{Q}_p/\mathbb{Z}_p$ is nontrivial.

As before, write Φ_n for the p^n -th-cyclotomic polynomial viewed as an element in $\mathbb{Z}_p[[H]]$. For brevity, we write

$$V_n = T_p(E(K_n) \otimes \mathbb{Q}_p/\mathbb{Z}_p) \otimes \mathbb{Q}_p.$$

Then, the map f_n induces the $\mathbb{Q}_p[[H/H_n]]$ -homomorphism

$$V_n[\Phi_n] \hookrightarrow V_n \longrightarrow T_p(E(K_{n,w}) \otimes \mathbb{Q}_p/\mathbb{Z}_p) \otimes \mathbb{Q}_p,$$

which is nonzero. Since Φ_n is irreducible over $\mathbb{Q}_p[[T]]$, it follows that the image of this homomorphism contains at least a copy of $\mathbb{Q}_p[[T]]/\Phi_n$, and since $\dim_{\mathbb{Q}_p}(\mathbb{Q}_p[[T]]/\Phi_n) = p^n - p^{n-1}$, this image has \mathbb{Q}_p -dimension at least $p^n - p^{n-1}$. Consequently, one has

$\text{corank}_{\mathbb{Z}_p}(C_n) \geq p^n - p^{n-1}$ for sufficiently large n . This establishes our claim and completes the proof of the theorem. \square

We end the paper with the proofs of the two corollaries stated in the introductory section.

PROOF OF COROLLARY 1.3. For each \mathbb{Z}_p -extension L_∞ of F , we write $H_{L_\infty} = \text{Gal}(F_\infty/L_\infty)$. Consider the commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathcal{M}(E/L_\infty) & \longrightarrow & R(E/L_\infty) \\ & & \downarrow \alpha & & \downarrow \delta \\ 0 & \longrightarrow & \mathcal{M}(E/F_\infty)^{H_{L_\infty}} & \longrightarrow & R(E/F_\infty)^{H_{L_\infty}} \end{array}$$

with exact rows. From the proof of [14, Proposition 3.8], we see that the kernel of δ is cofinitely generated over \mathbb{Z}_p and hence so is the kernel of α . However, combining Theorem 1.2 with [14, Lemma 2.1(ii)], we see that $\mathcal{M}(E/F_\infty)^{H_{L_\infty}}$ is cotorsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/F)]]$. Therefore, it follows that $\mathcal{M}(E/L_\infty)$ is cotorsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/F)]]$ as required. \square

PROOF OF COROLLARY 1.4. Under the finiteness of $\mathcal{X}(E/L)$, it is shown in [13, Proposition 4.1] that $\mathcal{X}(E/L_\infty)$ is cotorsion over $\mathbb{Z}_p[[\text{Gal}(L_\infty/F)]]$. The conclusion therefore follows from this, Corollary 1.3 and the tautological short exact sequence

$$0 \longrightarrow \mathcal{M}(E/L_\infty) \longrightarrow R(E/L_\infty) \longrightarrow \mathcal{X}(E/L_\infty) \longrightarrow 0. \quad \square$$

Acknowledgements

We would like to thank Jishnu Ray for his interest and comments on our paper. Some part of the research of this article took place when the first named author was visiting Harbin Engineering University and Harbin Institute of Technology, and he would like to acknowledge the hospitality and conducive working conditions provided by these institutes. The second named author would especially like to thank Chenyan Wu for her invaluable support and guidance during his visit to the University of Melbourne. The second named author is also grateful to the University of Melbourne for providing a stimulating research environment and the necessary resources, which greatly contributed to the completion of this work.

References

- [1] M. Bertolini, ‘Iwasawa theory for elliptic curves over imaginary quadratic fields’, *J. Théor. Nombres Bordeaux* **13**(1) (2001), 1–25.
- [2] A. Bhawe, ‘Analogue of Kida’s formula for certain strongly admissible extension’, *J. Number Theory* **122** (2007), 100–120.
- [3] J. Coates and R. Greenberg, ‘Kummer theory for abelian varieties over local fields’, *Invent. Math.* **124**(1–3) (1996), 129–174.
- [4] J. Coates and R. Sujatha, ‘Fine Selmer groups of elliptic curves over p -adic Lie extensions’, *Math. Ann.* **331**(4) (2005), 809–839.

- [5] Y. Hachimori and K. Matsuno, ‘An analogue of Kida’s formula for the Selmer groups of elliptic curves’, *J. Algebraic Geom.* **8** (1999), 581–601.
- [6] H. Imai, ‘A remark on the rational points of abelian varieties with values in cyclotomic \mathbb{Z}_p -extensions’, *Proc. Japan Acad.* **51** (1975), 12–16.
- [7] S. Jha, ‘Fine Selmer group of Hida deformations over non-commutative p -adic Lie extensions’, *Asian J. Math.* **16**(2) (2012), 353–366.
- [8] J. Lee, ‘Structure of the Mordell–Weil group over the \mathbb{Z}_p -extensions’, *Trans. Amer. Math. Soc.* **373**(4) (2020), 2399–2425.
- [9] A. Lei, ‘Algebraic structure and characteristic ideals of fine Mordell–Weil groups and plus/minus Mordell–Weil groups’, *Math. Z.* **303**(1) (2023), Article no. 14, 17 pages.
- [10] M. F. Lim, ‘A remark on the $\mathfrak{W}_H(G)$ -conjecture and Akashi series’, *Int. J. Number Theory* **11**(1) (2015), 269–297.
- [11] M. F. Lim, ‘On the pseudo-nullity of the dual fine Selmer groups’, *Int. J. Number Theory* **11**(7) (2015), 2055–2063.
- [12] M. F. Lim, ‘Notes on the fine Selmer groups’, *Asian J. Math.* **21**(2) (2017), 337–362.
- [13] M. F. Lim, ‘On the control theorem for fine Selmer groups and the growth of fine Tate–Shafarevich groups in \mathbb{Z}_p -extensions’, *Doc. Math.* **25** (2020), 2445–2471.
- [14] M. F. Lim, ‘Structure of fine Selmer groups over \mathbb{Z}_p -extensions’, *Math. Proc. Cambridge Philos. Soc.* **176**(2) (2024), 287–308.
- [15] M. F. Lim, ‘On fine Mordell–Weil groups over \mathbb{Z}_p -extensions of an imaginary quadratic field’, *Ann. Math. Qué.*, to appear. Published online (24 December 2024).
- [16] M. F. Lim and V. K. Murty, ‘The growth of fine Selmer groups’, *J. Ramanujan Math. Soc.* **31**(1) (2016), 79–94.
- [17] J. Neukirch, A. Schmidt and K. Wingberg, *Cohomology of Number Fields*, 2nd edn, Grundlehren der mathematischen Wissenschaften, 323 (Springer-Verlag, Berlin, 2008).
- [18] B. Perrin-Riou, *p -adic L -functions and p -adic Representations*, SMF/AMS Texts and Monographs, 3 (American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2000), translated from the 1995 French original by Leila Schneps and revised by the author.
- [19] J. Van Order, ‘Some remarks on the two-variable main conjecture of Iwasawa theory for elliptic curves without complex multiplication’, *J. Algebra* **350** (2012), 273–299.
- [20] O. Venjakob, ‘A noncommutative Weierstrass preparation theorem and applications to Iwasawa theory’, *J. reine angew. Math.* **559** (2003), 153–191.
- [21] C. Wuthrich, *The Fine Selmer Group and Height Pairings*. PhD Thesis (University of Cambridge, 2004).
- [22] C. Wuthrich, ‘Iwasawa theory of the fine Selmer group’, *J. Algebraic Geom.* **16**(1) (2007), 83–108.
- [23] C. Wuthrich, ‘The fine Tate–Shafarevich group’, *Math. Proc. Cambridge Philos. Soc.* **142**(1) (2007), 1–12.

MENG FAI LIM, School of Mathematics and Statistics,
Central China Normal University, Wuhan 430079, PR China
e-mail: limmf@cnu.edu.cn

CHAO QIN, College of Mathematical Sciences,
Harbin Engineering University, Harbin 150001, PR China
e-mail: qinchao@hrbeu.edu.cn

JUN WANG, Institute for Advanced Study in Mathematics of HIT,
Harbin Institute of Technology, Harbin 150001, PR China
e-mail: junwangmath@hit.edu.cn