

# The influence of host ecology and biogeography on the helminth species richness of freshwater fishes in Mexico

L. GARRIDO-OLVERA<sup>1</sup>, H. T. ARITA<sup>2</sup> and G. PÉREZ-PONCE DE LEÓN<sup>3\*</sup>

<sup>1</sup> *Posgrado en Ciencias Biológicas, Instituto de Ecología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Apartado Postal 70-275, C. P. 04510, Distrito Federal, México*

<sup>2</sup> *Centro de Investigaciones en Ecosistemas, Universidad Nacional Autónoma de México, Antigua Carretera a Pátzcuaro 8701, Ex-Hacienda de San José de La Huerta, C. P. 58190, Morelia, Michoacán, México*

<sup>3</sup> *Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Apartado Postal 70-153, C. P. 04510, Distrito Federal, México*

(Received 19 December 2011; revised 17 April and 9 May 2012; accepted 12 May 2012; first published online 19 July 2012)

## SUMMARY

Freshwater fish helminths, the most well known Mexican vertebrate parasites, include approximately 260 species (platyhelminthes, acanthocephalans, nematodes, and hirudineans). The distribution patterns of adult helminth diversity (throughout parasite and host groups and hydrological regions) are described and the effects of host traits and environmental and geographical factors on diversity are evaluated. Adult helminths include 160 species, parasitizing 149 fish species of 23 families distributed in 21 regions. Nematoda was the most species-rich (> 50 species). Cichlidae harboured rich helminth assemblages, with widespread parasites. By contrast, Atherinopsidae and Goodeidae showed relatively poor helminth assemblages, including specific parasites with narrow distribution. Helminth richness in southeastern Mexico was higher than northern or central regions. Non-parametric richness estimators were used to avoid confusion in comparisons with unequal sampling efforts. Bootstrap values, the method with the best performance, indicated that estimated richness shows the same distribution pattern that observed richness. Non-phylogenetic and phylogenetic analyses were used to determine the role of different factors in the parasite diversification. The distribution range was the most important richness predictor (widespread fishes harbour richer parasite assemblages), although interactions between this variable and others such as trophic level, latitude, habitat temperature and precipitation are also important. Likewise, biogeographical factors can also affect parasite diversity.

Key words: geographical range, parasites, diversity, independent contrasts, Nearctic region, Neotropical region, hydrological regions, Osteichthyes.

## INTRODUCTION

The evolutionary events determining the structure of a parasite assemblage are relatively well understood. A parasite species may have been inherited by the host species from its ancestor, may be the result of an intra-host speciation event, and/or may have colonized the host species from another sympatric host species (host switching) (Paterson and Gray, 1997). Therefore, it is possible to search for the key factors that have caused certain parasite assemblages to diversify more than others over evolutionary time (Poulin, 1998a; Page, 2003). The studies attempting to determine which host features may promote the diversification of parasite assemblages are based on 2 theoretical frameworks. First, following island biogeography theory (MacArthur and Wilson, 1967;

Kuris *et al.* 1980), host features that promote high rates of parasite speciation or colonization, and low probabilities of extinction, by parasite species should be associated with high parasite diversity (e.g. large body size, broad geographical range, breadth of habitats). Epidemiological modelling represents the second theoretical source of predictions regarding parasite diversity (Dobson and Roberts, 1994; Roberts *et al.* 2002). These models suggest that host population density, which regulates the contact rate between parasite infective stages and hosts, is the key factor determining whether a parasite species can invade and persist in a host population. Comparing different host species, those occurring at higher population density (e.g. schooling fish species) should harbour more species of parasites, because they exceed the persistence threshold of more parasite species than hosts with low population density.

Several studies have investigated the factors potentially controlling the number of parasite species in freshwater and marine fishes (Bell and Burt, 1991; Guégan and Kennedy, 1993; Poulin, 1995; Poulin and Rohde, 1997; Sasal *et al.* 1997; Morand *et al.* 2000; Luque and Poulin, 2004, 2008). However, the

\* Corresponding author: Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Apartado Postal 70-153, C. P. 04510, Distrito Federal, México. Tel: +56229131. Fax: +55500164. E-mail: ppdleon@ibunam2.ibiologia.unam.mx

results available to date show very little consistency, since some studies claim that a certain variable is a predictor of parasite species richness, either host body size, host geographical range, host diet, etc., but no consensus on a particular variable as the main determinant has been reached.

Freshwater fish helminths are undoubtedly the most well known group of vertebrate parasites in Mexico, because fishes have been studied more than any other group (Pérez-Ponce de León and Choudhury, 2010; Pérez-Ponce de León *et al.* 2011). The helminth fauna of these hosts includes approximately 260 species (Oceguera-Figueroa *et al.* 2010; Pérez-Ponce de León and Choudhury, 2010). The first descriptions of distributional patterns of fish-parasite and drainage associations indicated that the parasite fauna is largely circumscribed by higher levels of monophyletic host taxa, especially at the level of fish family. Likewise, areas within a certain biogeographical region and consequently with similar fish composition possess more similar parasite faunas compared to areas with less similar fish faunal composition (Pérez-Ponce de León and Choudhury, 2005). Recently, Pérez-Ponce de León and Choudhury (2010) evaluated the progress made on the inventory of freshwater fish helminths of Mexico and suggested that following a traditional approach, the inventory is nearing completion for most helminth groups (except monogeneans). These authors suggested that host species and geographical areas that would be targeted in the future could be expected, with very few exceptions, to have helminth faunas that are consistent with the fish composition in those areas. However, they argued that in the future DNA-based taxonomic methods have the potential to alter drastically the estimates of helminth diversity in freshwater fishes, because of the existence of cryptic species (morphologically indistinguishable but genetically distinct, see Poulin, 2011; Nadler and Pérez-Ponce de León, 2011) in several lineages of helminths in Mexico. Despite this fact, they described species richness patterns and pointed out that the diversity is distributed heterogeneously. These patterns were described by considering all freshwater helminth parasites irrespective of their developmental stage i.e. larval and adult forms, under a strict definition of what constitutes a freshwater species and by including both native and introduced fish species. However, Pérez-Ponce de León and Choudhury (2010) did not use the proper methods to analyse data with unequal sampling effort that might act as a confounding factor. In addition, potential factors that cause the heterogeneous distribution of helminth species richness were not explored. For these reasons, our aims in this paper are to describe the distribution patterns of adult helminth diversity throughout parasite and host groups as well as across hydrological regions, and to investigate the host traits and/or environmental and geographical factors that determine the uneven

diversification of assemblages of freshwater fish helminths.

#### MATERIALS AND METHODS

A database updated to November, 2011, with distribution records for Mexican freshwater fish helminths (trematodes, monogeneans, cestodes, acanthocephalans, nematodes, and hirudineans) was constructed. Our study was restricted to adult helminths of the native freshwater fishes, which are unable to disperse across brackish and marine waters (i.e., presumably non-diadromous). Larval helminths were excluded because of the confounding effects of dispersal caused by their definitive hosts in any distribution pattern. Also, if we consider larval stages, the parasite species count could be underestimated because they often cannot be identified to the species level and one putative taxon might actually consist of more than one species.

The data set was mostly obtained from monographs (e.g. Pérez-Ponce de León *et al.* 1996, 2007; Garrido-Olvera *et al.* 2006; Kohn *et al.* 2006; Salgado-Maldonado, 2006; García-Prieto *et al.* 2010; Oceguera-Figueroa *et al.* 2010) and specimens deposited in the Colección Nacional de Helminths (CNHE), Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City. Nomenclature was updated using specialized literature (e.g. Scholz *et al.* 1997; Moravec, 1998; Scholz *et al.* 2001; Vidal-Martínez *et al.* 2001; Aguirre-Macedo and Scholz, 2005; Caspeta-Mandujano, 2005; Pérez-Ponce de León *et al.* 2007). To avoid cases of synonymies among fish species, valid species names were adopted according to FishBase (Froese and Pauly, 2011).

The geographical distribution patterns were recognized through distribution maps for each taxon, which were prepared using ArcView GIS 3.2 (ESRI, 1999). Sample-based rarefaction curves considering the number of localities studied as a measure of sampling effort were used to compare the helminth species richness (the number of parasite species in an assemblage) between host families and hydrological regions at comparable sampling effort (Colwell *et al.* 2004). Non-parametric species richness estimators are also used to analyse data with unequal sampling effort. Therefore, total helminth species richness for each data set was calculated using 5 estimation methods: ICE, Chao2, Jackknife1, Jackknife2, and Bootstrap (Poulin, 1998b; Walther and Morand, 1998). In addition, an evaluation of the performance of these estimators with the unscaled measures of bias, precision, and accuracy was conducted (Walther and Moore, 2005). EstimateS version 8.2 was used to obtain rarefaction curves and total species richness (Colwell, 2006).

In order to examine the possible factors determining the helminth species richness in an assemblage,

we gathered detailed information on the fish species and geographical areas. The host traits were obtained from Miller *et al.* (2005) and Froese and Pauly (2011) (Table 1). The environmental and geographical data for each hydrological region were obtained from the Comisión Nacional del Agua (2008) and Fernández-Eguiarte *et al.* (2010) (Table 2).

The helminths include various taxonomic groups, with different biological characteristics related with life-cycle dynamics and transmission modes. Therefore, these parasite taxa may be subjected to different factors potentially influencing their species richness. For this reason, helminth species richness for each assemblage was estimated in 3 ways: for all parasites, for endoparasites only (trematodes, cestodes, acanthocephalans, and nematodes), and for ectoparasites only (monogeneans and hirudineans).

To analyse factors influencing differences in species richness among assemblages, the information on the fish species and their distribution regions was combined. In the initial analyses, fish species were treated as independent observations. The relationships between helminth species richness and continuous variables were tested through correlations, while the influence of categorical variables was assessed by inspecting mean values. Next, we determined which factors played a role in the diversification of parasite assemblages, i.e., we tested whether any of those associations were significant and whether there were interactions between the explanatory variables by performing generalized linear models, where species richness (total parasites, ectoparasites, and endoparasites, independently) was the response variable and host traits were explanatory variables. The models were fitted with a log link (to ensure that the fitted values are bounded below) and Poisson errors (to account for the non-normality) (Crawley, 2007).

Parasite assemblages of closely related host species are not truly independent statistical observations. Therefore, we used Felsenstein's (1985) comparative method to control for the effects of phylogenetic association between host species. The PDAP: PDTREE software (Midford *et al.* 2011), implemented in Mesquite Modular System for Evolutionary Analysis, Mesquite version 1.12 (Maddison and Maddison, 2011) was used to compute independent contrasts. Host phylogeny was constructed from mainly molecular studies and the phylogenetic relationships of the orders were based on Nelson (2006), as follows: ((((((*Ophisternon aenigmaticum*, (*Eugerres mexicanus*, *Etheostoma* sp., (*Lepomis megalotis*, *L. macrochirus*), *Micropterus salmoides*), *Aplodinotus grunniens*, (*Awaous banana*, *Sicydium multipunctatum*), (*Cichlasoma geddesi*, *C. mayorum*, (((*C. beani*, *C. istlanum*), (*Parachromis friedrichsthalii*, *C. trimaculatum*), (*Petenia splendida*, *C. urophthalmus*))), (*Rocio octofasciata*, *Amphilophus robertsoni*), ((*C. salvini*, (*Thorichthys helleri*,

*T. pasionis*), (*T. meeki*, (*T. ellioti*, *T. callolepis*))), ((*Herichthys labridens*, (*H. minckleyi*, *H. cyanoguttatus*)), ((*C. pearsei*, (*Theraps lentiginosus*, *Vieja intermedia*)), ((*V. fenestrata*, (*V. bifasciata*, *V. hartwegi*)), (*V. synspila*, *V. argentea*))))))))) ((((*Cyprinodon meeki*, *C. nazas*), ((((*Poecilia butleri*, *P. mexicana*), (*P. sphenops*, *P. catemacensis*)), (*P. velifera*, (*P. petenensis*, *P. latipunctata*))), ((*Poeciliopsis catemaco*, *P. gracilis*), (*P. infans*, *P. baenschii*), *P. balsas*), ((*Xiphophorus hellerii*, *X. variatus*), (*Heterandria bimaculata*, (*Belonesox belizanus*, ((*Gambusia vittata*, *G. marshi*), *G. senilis*, *G. yucatanana*))))))))) ((*Profundulus hildebrandi*, *P. punctatus*, *P. labialis*), ((*Characodon audax*, *C. lateralis*), (((*Ilyodon cortesae*, *I. whitei*, *I. furcoidens*), (*Xenotaenia resolanae*, ((*Allodontichthys tamazulae*, *A. hubbsi*), *A. zonistius*))), (((((((*Chapalichthys encaustus*, *C. pardalis*), *Ameca splendens*), *Xenotoca variata*), *Allophorus robustus*), *Zoogoneticus quitzeoensis*), (*X. eiseni*, *X. melanosoma*)), ((*Ataeniobius toweri*, (*Goodea atripinnis*, *G. gracilis*)), (((*Girardinichthys multiradiatus*, *G. viviparus*), (*Hubbsina turneri*, (*Skiffia bilineata*, *S. multipunctata*, *S. lermiae*))), ((((*Allotoca catarinae*, (*A. diazi*, *A. meeki*)), *A. zacapuensis*), *A. dugesii*), *A. maculata*), *A. regalis*))))))))) (*Strongylura* sp., *Hyporhamphus mexicanus*)), ((*Atherinella crystallina*, *A. balsana*), ((((((*Poblana letholepis*, *P. squamata*), *P. alchichica*), *Chirostoma riojai*), *Ch. arge*), (*Ch. attenuatum*, ((*Ch. jordani*, *Ch. labarcae*), (*Ch. lucius*, (*Ch. humboldtianum*, (*Ch. grandocule*, *Ch. estor*))))))))) *Typhliasina pearsei*), ((*Rhamdia guatemalensis*, (*Potamarius nelsoni*, (*Ameiurus melas*, (*Pyloodictis olivaris*, ((*Ictalurus furcatus*, *I. punctatus*), *I. balsanus*, *I. mexicanus*, *I. pricei*, *I. dugesii*))), ((*Ictiobus meridionalis*, *Catostomus nebuliferus*), ((((*Algansea lacustris*, *A. tincella*), *A. monticola*), *Gila conspersa*), (*Campostoma ornatum*, (*Dionda ipni*, (*Pimephales promelas*, ((((*Notropis calientis*, *Aztecucula sallaei*), *Yuriria alta*), *Hybopsis boucardi*), *N. nazas*, *N. chihuahua*), (*Codoma ornata*, ((*Cyprinella garmani*, *C. lutrensis*), *C. xanthicara*))))))))) ((((*Astyanax aeneus*, *A. mexicanus*), *A. fasciatus*), *Bramocharax caballeroi*), *Brycon guatemalensis*)), ((*Dorosoma cepedianum*, *D. petenense*), *D. anale*)), *Atractosteus tropicus*); (Ptacek and Breden, 1998; Breden *et al.* 1999; Harris and Mayden, 2001; Reznick *et al.* 2002; Miya *et al.* 2003; Schönhuth and Doadrio 2003; Simons *et al.* 2003; Cunha *et al.* 2002; Doadrio and Dominguez, 2004; Wilcox *et al.* 2004; Near *et al.* 2005; Hulsey *et al.* 2006; Chakrabarty, 2007; Concheiro Pérez *et al.* 2007; Hrbek *et al.* 2007; Hardman and Hardman, 2008; Hertwig, 2008; Lavoué *et al.* 2008; Rícan *et al.* 2008; Bloom *et al.* 2009, 2012; Mirande 2009; Pérez-Rodríguez *et al.* 2009; McMahan *et al.* 2010; Schönhuth and Mayden, 2010).

True branch lengths are not available in this tree, so all branch lengths were set to unity, which adequately standardized contrasts of all variables.

Table 1. Summary information to family level of freshwater fish species recorded as hosts of adult helminths in Mexico

(The number of fish species was obtained from Miller *et al.* (2005) and Froese and Pauly (2011).)

| Family          | Fishes | Helminths     | Size $\pm$ s.d.   | Regions | Environment | Geographical range | Trophic level $\pm$ s.d. | Climate   |
|-----------------|--------|---------------|-------------------|---------|-------------|--------------------|--------------------------|-----------|
| Ariidae*        | 3/1    | 4/-, 1, 3     | 39                | 1       | D           | M                  | 3.58                     | Tr        |
| Atherinopsidae  | 39/14  | 10/12, 2, 8   | 13.91 $\pm$ 9.53  | 4       | B, P        | M                  | 3.06 $\pm$ 0.15          | Tr        |
| Belontiidae*    | 4/1    | 4/-, -, 4     | 49.8              | 1       | P           | MCA                | 4.13                     | Tr        |
| Bythitidae*     | 1/1    | 1/-, -, 1     | 9.7               | 1       | D           | M                  | 3.3                      | Tr        |
| Catostomidae    | 18/2   | 8/10, -, 8    | 39.95 $\pm$ 34.72 | 4       | P, D        | M, MCA             | 2.94 $\pm$ 0.24          | Tr        |
| Centrarchidae   | 10/3   | 16/21, 8, 8   | 37.67             | 5       | B           | MNA                | 3.55                     | S, Tm     |
| Characidae      | 10/5   | 28/34, 9, 19  | 17.26 $\pm$ 13.46 | 11      | B           | M, MSA, MCA, MNA   | 2.81 $\pm$ 0.34          | S, Tr     |
| Cichlidae       | 55/27  | 58/69, 12, 44 | 18.56 $\pm$ 7.07  | 12      | B, D        | M, MCA, MNA        | 3.12 $\pm$ 0.73          | S, Tr     |
| Clupeidae       | 11/3   | 9/11, 2, 7    | 26.13 $\pm$ 8.52  | 3       | P, PN       | MCA, MNA, NAMCA    | 2.96 $\pm$ 0.51          | Tr, S     |
| Cyprinidae      | 69/23  | 14/17, 4, 10  | 10.02 $\pm$ 6.26  | 7       | B, D        | M, MNA             | 2.815 $\pm$ 0.27         | S, Tm, Tr |
| Cyprinodontidae | 28/2   | 5/6, 1, 4     | 4.75 $\pm$ 0.21   | 2       | B           | M                  | 2.94 $\pm$ 0             | Tr        |
| Gerreidae*      | 11/1   | 3/-, 2, 1     | 20.8              | 2       | D           | MCA                | 3.35                     | Tr        |
| Gobiidae*       | 4/2    | 3/-, -, 3     | 21.5              | 3       | B, D        | NAMCASA            | 2.025                    | Tr        |
| Goodeidae       | 41/33  | 24/28, 5, 19  | 7.72 $\pm$ 3.01   | 7       | B, D, P     | M                  | 2.09 $\pm$ 0.17          | S, Tm, Tr |
| Hemiramphidae*  | 4/1    | 1/-, -, 1     | 16.1              | 1       | P           | MCA                | 3.02                     | Tr        |
| Heptapteridae   | 4/1    | 24/28, 6, 18  | 30 $\pm$ NA       | 4       | B           | MSA                | 3.16 $\pm$ NA            | Tr        |
| Ictaluridae     | 14/8   | 35/40, 7, 28  | 67.23 $\pm$ 26.99 | 13      | B, D        | M, MNA, NAMCA      | 3.63 $\pm$ 0.15          | S, Tm, Tr |
| Lepisosteidae   | 4/1    | 6/7, 1, 5     | 85.9 $\pm$ NA     | 2       | D           | MCA                | 4.23 $\pm$ NA            | Tr        |
| Percidae*       | 6/1    | 1/-, 1, -     | 5                 | 1       | B           | M                  | 3.17                     | Tr        |
| Poeciliidae     | 82/20  | 22/26, 5, 17  | 7.55 $\pm$ 3.28   | 12      | B, D        | M, MCA, MSA, MNA   | 2.72 $\pm$ 0.48          | S, Tr     |
| Profundulidae*  | 5/3    | 4/-, -, 4     | 10.4              | 2       | B           | MCA                | 3.073                    | Tr        |
| Sciaenidae*     | 1/1    | 5/-, 3, 2     | 50                | 3       | D           | NAMCA              | 3.36                     | S         |
| Synbranchidae*  | 3/1    | 7/-, -, 7     | 20.8              | 1       | D           | MCA                | 3.25                     | Tr        |

\* Data were insufficient for estimating total parasite species richness. Fishes, total/known host fish species; Helminths, observed/estimated total species richness, ectoparasites, endoparasites; Size  $\pm$  s.d., mean body size (standard length in cm)  $\pm$  standard deviation; Regions, number of regions where fish family is distributed; Environment: B, Benthopelagic, D, Demersal, P, Pelagic, PN, Pelagic-neritic; Trophic level  $\pm$  s.d. = mean trophic level  $\pm$  standard deviation; Climate: S, subtropical, Tr, tropical, Tm, temperate; Geographical range: M, Mexico, MCA, Mexico and Central America, MNA, Mexico and North America, MSA, Mexico and South America, NAMCA, North America, Mexico and Central America, NAMCASA, North America, Mexico, Central America, and South America.

Table 2. Summary information on Mexican hydrological regions, where adult helminth species have been recorded as parasites of freshwater fishes (The number of fish species is result of a review of databases of fish collections and fish lists of Miller *et al.* (2005) and Froese and Pauly (2011).)

| Region | Fishes | Helminths        | MxL   | MnL   | ML    | Length | Area   | Flow   | Basins | MAP    | PDM   | PWM    | MTWM  | MTCM  | Slope |
|--------|--------|------------------|-------|-------|-------|--------|--------|--------|--------|--------|-------|--------|-------|-------|-------|
| 9*     | 34/4   | 3/-, -, 3        | 31·33 | 26·6  | 28·97 | 39635  | 139370 | 4935   | 48     | 507·2  | 0·11  | 22·8   | 34·55 | 7·21  | P     |
| 11     | 33/13  | 16/19·73, 2, 14  | 25·18 | 21·69 | 23·44 | 19704  | 51717  | 7956   | 9      | 815·2  | 7·36  | 215·91 | 32·88 | 5·88  | P     |
| 12     | 74/41  | 36/44·03, 12, 24 | 23·42 | 19·06 | 21·24 | 51096  | 132916 | 13637  | 8      | 723·2  | 0·37  | 81·02  | 34·33 | 9·51  | P     |
| 14*    | 25/6   | 4/-, -, 4        | 21·28 | 20·07 | 20·68 | 6132   | 12255  | 2236   | 1      | 122·7  | 4·34  | 92·2   | 32·49 | 3·21  | P     |
| 15*    | 17/1   | 3/-, -, 3        | 20·45 | 18·88 | 19·67 | 6307   | 12967  | 3684   | 45     | 1185·5 | 16·3  | 153·35 | 34·47 | 8·83  | P     |
| 16     | 29/11  | 14/18·05, -, 14  | 20·45 | 18·59 | 19·52 | 7393   | 17628  | 3882   | 8      | 911·3  | 7·28  | 71·58  | 30·69 | 4·16  | P     |
| 18     | 54/27  | 33/40·01, 3, 30  | 19·99 | 17·02 | 18·51 | 45372  | 118268 | 1757   | 3      | 949·7  | 15·22 | 184·34 | 30·82 | 7·72  | P     |
| 19*    | 16/1   | 1/-, -, 1        | 18·14 | 16·7  | 17·42 | 4992   | 12132  | 691    | 41     | 1232   | 5·27  | 182·06 | 29·94 | 4·83  | P     |
| 20*    | 24/3   | 4/-, 1, 3        | 17·62 | 15·98 | 16·8  | 12907  | 39936  | 18714  | 21     | 1393·1 | 42·26 | 288·22 | 30·45 | 11·11 | P     |
| 23*    | 17/1   | 1/-, -, 1        | 16·54 | 14·53 | 15·54 | 3908   | 12293  | 12554  | 25     | 2352·7 | 5·46  | 261·16 | 31·43 | 8·25  | P     |
| 24     | 101/14 | 27/33·81, 7, 20  | 31·77 | 24·85 | 28·31 | 77898  | 229740 | 5156   | 2      | 448·5  | 2·82  | 318·3  | 33·27 | 13·73 | A     |
| 25*    | 41/4   | 9/-, 4, 5        | 25·76 | 22·28 | 24·02 | 16729  | 54961  | 4328   | 27     | 758·5  | 4·27  | 224·06 | 31·37 | 10·11 | A     |
| 26     | 77/21  | 24/28·74, 4, 20  | 23·95 | 19·04 | 21·5  | 34263  | 96989  | 20329  | 2      | 889·2  | 34·07 | 200·23 | 33·66 | 16·87 | A     |
| 27*    | 38/3   | 10/-, 1, 9       | 22·2  | 19·48 | 20·84 | 10139  | 26592  | 14306  | 38     | 1423·2 | 4·08  | 216·68 | 32·42 | 10·13 | A     |
| 28     | 67/23  | 49/58·6, 11, 38  | 19·74 | 16·93 | 18·34 | 19382  | 57355  | 49951  | 31     | 1447·1 | 22·85 | 220·73 | 35·28 | 15·79 | A     |
| 29*    | 52/19  | 28/-, 11, 17     | 18·44 | 16·63 | 17·54 | 8866   | 30217  | 39482  | 10     | 1953·8 | 28·39 | 344·94 | 30·99 | 12·36 | A     |
| 30     | 86/37  | 61/71·03, 19, 42 | 18·9  | 15·3  | 17·1  | 23249  | 102465 | 117546 | 4      | 1708·9 | 38·47 | 326·82 | 32·86 | 15·17 | A     |
| 31*    | 32/7   | 12/-, 3, 9       | 19·97 | 17·81 | 18·89 | 1704   | 25443  | 591    | 6      | 1227·4 | 2·78  | 272·12 | 31·39 | 12·94 | A     |
| 32     | 42/15  | 40/45·98, 17, 23 | 21·61 | 19·63 | 20·62 | 523    | 58135  | 0      | 1      | 192·4  | 45·93 | 424·76 | 33·66 | 16·71 | A     |
| 33     | 37/10  | 19/22·31, 6, 13  | 20·1  | 17·81 | 18·96 | 2926   | 38308  | 1989   | 2      | 1239·8 | 2·8   | 323·08 | 33·58 | 17·24 | A     |
| 36     | 33/14  | 9/11·24, -, 9    | 26·6  | 22·67 | 24·64 | 31046  | 9332   | 1912   | 1      | 422·1  | 7·3   | 457·02 | 34·1  | 16·7  | I     |

\* Data were insufficient for estimating total species richness. Region, 9, Sonora Sur; 11, Presidio-San Pedro; 12, Lerma-Santiago; 14, Río Ameca; 15, Costa de Jalisco; 16, Armería-Coahuayana; 18, Balsas; 19, Costa Grande de Guerrero; 20, Costa Chica de Guerrero; 23, Costa de Chiapas; 24, Bravo-Conchos; 25, San Fernando-Soto La Marina; 26, Pánuco; 27, Norte de Veracruz; 28, Papaloapan; 29, Coatzacoalcos; 30, Grijalva-Usumacinta; 31, Yucatán Oeste; 32, Yucatán Norte; 33, Yucatán Este; 36, Nazas-Aguanaval; Fishes, total/known host fish species; Helminths, observed/estimated total species richness, ectoparasites, endoparasites; MxL, maximum latitude; MnL, minimum latitude; ML, mean latitude; Length, total length of rivers (in km); Area, area (in km<sup>2</sup>); Flow, mean current flow (in hm<sup>3</sup>/year); Basins, basin number in the region; MAP, mean annual precipitation (mm); PDM, precipitation of driest month (mm); PWM, precipitation of wettest month (mm); MTWM, maximum temperature of warmest month (°C); MTCM, minimum temperature of coldest month (°C); Slope, A, Atlantic, I, Interior, P, Pacific.

The program can only compute independent contrasts in continuous variables. In order to obtain independent contrasts of categorical variables, we treated these variables as continuous with states of 1, 2, 3 or more as recommended by Midford *et al.* (2011) in the PDAP:PDTREE manual.

The relationships among independent contrasts were assessed using least-squares regressions forced through the origin (Garland *et al.* 1992). The possibility that multiple host traits influence parasite diversity was also addressed and then contrasts in independent variables (host traits) were entered into multiple regression methods (computed through the origin) to predict contrasts in species richness (total parasites, ectoparasites, and endoparasites, independently) (Crawley, 2007).

In both generalized linear models and multiple regression analyses, the minimal adequate models were obtained with categorical and continuous explanatory variables, separately (i.e., models without redundant parameters or factor levels). We achieved this by fitting a maximal model and then simplifying it by stepwise deletion: non-significant terms were left out, and significant terms were added back (Crawley, 2007). All statistical analyses were conducted in the software R version 2.11.0 for Windows (<http://www.r-project.org/>).

## RESULTS

The adult helminth fauna parasitizing freshwater fishes in Mexico is composed by 160 species belonging to 6 groups. The most numerous group was Nematoda with more than 50 species, followed by Trematoda and Monogenea, with 39 species each. The remaining groups were composed by no more than 10 species each (4 Hirudinea, 6 Acanthocephala, and 9 Cestoda). No parasite species occurred in all hydrological regions, but the cestode *Bothriocephalus acheilognathi* and the acanthocephalan *Neoechinorhynchus golvani* were distributed in 14 and 11 regions, respectively. In contrast, more than 100 species (approximately 60% of total parasite species) were recorded from only 1 region. The most generalist helminth species were the cestode *B. acheilognathi* and the nematode *Rhabdochona kidderi* parasitizing 11 and 10 host families, respectively. However, approximately 120 species (75% of total parasite species) infected only 1 fish family each.

In total, 149 fish species belonging to 23 families have been recorded as hosts of adult helminths. This means that approximately 40% of Mexican freshwater fishes have been recorded as hosts of these parasites. Helminths were found infecting 8% to 100% of the species included in each host family, though the highest percentages correspond to monotypical families. With the exception of the Cyprinodontidae and Catostomidae, the more species-rich host groups (i.e., Poeciliidae, Cyprinidae, Cichlidae, Goodeidae,

Atherinopsidae, and Ictaluridae, which account for more than 80% of the Mexican ichthyofauna) have been sampled with relatively high intensity, since at least 24%, 33%, 49%, 81%, 36%, and 57%, of their species, respectively, have been recorded as hosts of helminths (Table 1).

In terms of geographical distribution, freshwater fish adult helminths were found in 198 localities pertaining to 21 of the 37 Mexican hydrological regions. The number of records was reduced in most of the regions located in the Nearctic biogeographical region and in coastal areas. In contrast, Lerma-Santiago, Balsas, Grijalva-Usumacinta, Nazas-Aguanaval, and Presidio-San Pedro were the best known hydrological regions, considering the number of fish species recorded as hosts and the actual number of fish species occurring in each region as well as the number of localities studied (Fig. 1, Table 2).

Although the sample-based rarefaction curves did not reach the asymptote, the majority showed a certain stability indicating that the sample size in each data set was large enough to consider that their adult helminth fauna was properly known. The comparisons of rarefaction curves at comparable sampling effort indicated that the difference in helminth species richness was significant among both host families and hydrological regions. The cichlids and ictalurids harboured the highest helminth species richness, whereas an intermediate number of species parasitized characids, goodeids, heptapterids, and poeciliids. The remaining host families did not harbour helminth faunas with more than 10 species (rarefaction curves not shown). At the species level, most fish species (60%) harboured poor helminth faunas (i.e., fewer than 5 species per host species). In contrast, only 6% of the fish species were parasitized by helminth faunas comprising more than 10 parasite species. The parasite species richness among hydrological regions also varied significantly. In southeastern Mexico, particularly in the Grijalva-Usumacinta and Papaloapan river basins, the highest helminth species richness was found. In contrast, with the exception of the regions situated in the Yucatán Peninsula, the hydrological regions with intermediate or low helminth species richness were always located in northern and central areas of the country (e.g. the Balsas, Lerma-Santiago, Nazas-Aguanaval and Pánuco systems) (rarefaction curves not shown).

We only considered the estimations of species richness produced by the bootstrap method, which achieved the best performance measures in all data sets. Based on these values, the minimum number of missing species remaining to be found varies from 1 to 11 and from 2 to 10 in the 13 host families and 11 hydrological regions analysed, respectively. Interestingly, the estimated species richness was distributed in the same way as the observed species




Fig. 1. Distribution of adult helminth parasites of freshwater fishes in Mexico. The country is divided in 37 hydrological regions and numbers indicate the regions where data are available: 9, Sonora Sur; 11, Presidio-San Pedro; 12, Lerma-Santiago; 14, Río Ameca; 15, Costa de Jalisco; 16, Armería-Coahuayana; 18, Balsas; 19, Costa Grande de Guerrero; 20, Costa Chica de Guerrero; 23, Costa de Chiapas; 24, Bravo-Conchos; 25, San Fernando-Soto La Marina; 26, Pánuco; 27, Norte de Veracruz; 28, Papaloapan; 29, Coatzacoalcos; 30, Grijalva-Usumacinta; 31, Yucatán Oeste; 32, Yucatán Norte; 33, Yucatán Este; 36, Nazas-Aguanaval. Shaded regions are those for which the data were sufficient for comparative analysis.

richness in both host families and hydrological regions (Tables 1 and 2).

The analyses across fish species values, not corrected for possible phylogenetic effects, revealed that the extent of their geographical distribution correlated positively with species richness of ectoparasites, endoparasites, and total parasites. The  $r$  values obtained by considering the number of sites as well as the regions in which a host species occurs (within its regional distribution range) were 0.73 and 0.77, 0.78 and 0.70, and 0.83 and 0.78,  $N=149$ ,  $P=<2.2e-16$ , respectively. In addition, the total area of hydrological regions where fishes are distributed correlated positively with ectoparasite species richness ( $r=0.70$ ,  $N=149$ ,  $P=<2.2e-16$ ). Based on main effect means of categorical variables, we also found that the distribution range of the host is an important variable and it seems that neotropical fishes have higher parasite species richness than those occurring in nearctic ones (Table 3).

In the generalized linear models performed on fish species values, only the following variables, geographical distribution, trophic level, temperature,

precipitation and latitude were retained, with the distribution range being the most important explanatory variable. The aforementioned trend is significant and there is compelling evidence that fish species with a broad distribution range tend to harbour more parasite species. In addition, these models showed significant interactions between distribution range and temperature and between latitude and precipitation in determining the number of endoparasites (Table 4).

After controlling for phylogenetic influences by using the independent contrasts method (Felsenstein, 1985), the same factors (the number of sites and regions in which a host species occurs) were correlated with the helminth species richness (ectoparasites  $r=0.66$  and 0.74, endoparasites  $r=0.74$  and 0.65, and total parasites 0.79 and 0.74, respectively,  $N=148$ ,  $P=<2.2e-16$ ).

The results of the regression analyses corroborated that the geographical range was a key determinant of the variability in parasite species richness among host species (Figs 2–4). These findings suggest that, on average, widespread fish species harbour more species

Table 3. Main effect means of categorical variables on helminth richness

(See Tables 1 and 2 for details of variables.)

| Variable           | Level | Ectoparasites | Endoparasites | Total parasites |
|--------------------|-------|---------------|---------------|-----------------|
| Environment        | B     | 2.10          | 4.67          | 6.77            |
|                    | D     | 1.65          | 4.46          | 6.10            |
|                    | P     | 1.38          | 3.25          | 4.63            |
|                    | PN    | 2             | 3.5           | 5.5             |
| Geographical range | M     | 1.51          | 3.72          | 5.23            |
|                    | MCA   | 2.5           | 5.62          | 8.12            |
|                    | MNA   | 2.07          | 4.47          | 6.53            |
|                    | MSA   | 6             | 12.67         | 18.67           |
|                    | NAMCA | 3.33          | 9.33          | 12.67           |
| Climate            | S     | 2.65          | 5.96          | 8.61            |
|                    | Tm    | 1.43          | 2.71          | 4.14            |
|                    | Tr    | 1.80          | 4.34          | 6.13            |
| Slope              | A     | 2.14          | 4.93          | 7.07            |
|                    | AI    | 3             | 8             | 11              |
|                    | AP    | 3.56          | 6.94          | 10.50           |
|                    | I     | 1             | 2.4           | 3.4             |
|                    | IP    | 1             | 4.2           | 5.2             |
|                    | P     | 1.33          | 3.52          | 4.85            |
|                    | PIA   | 2             | 4             | 6               |

of helminth parasites and that this pattern is explained mainly by recent processes. Although, some results were similar when phylogenetic relationships were considered, there were some changes in the relationships between helminth species richness and their explanatory variables. For example, trophic level and latitude were not important in determining the number of endoparasites. In addition, a difference in parasite species richness between fish species with larger *vs* smaller body size was detected when data were corrected for host phylogeny (Table 4).

#### DISCUSSION

In this paper, we only considered adult helminth species found in native Mexican freshwater fishes unable to disperse across the sea (i.e., presumably non-diadromous), because of the fact that the parasite fauna of exotic fishes is not necessarily one they have acquired over evolutionary time in their area of origin. The species count provided in this study for adult helminths parasitizing freshwater fishes in Mexico (160) is not in agreement with species counts presented in previous studies. Part of the problem is the challenge in defining what is a freshwater fish (Miller *et al.* 2005). Salgado-Maldonado (2006) quantified approximately 180 adult parasite species of freshwater fishes, but this author listed some helminth species found in marine or brackish water fishes that regularly enter freshwater at some life-history stage. In a recent publication, after a detailed consideration of the status of the freshwater fish fauna where objective criteria were followed, Pérez-Ponce de León and Choudhury (2010) considered that the

adult helminth fauna of freshwater fish in Mexico consists of 177 species (37 trematodes, 62 monogeneans, 15 cestodes, 6 acanthocephalans, and 54 nematodes); however, they considered both native and exotic species of fishes. In this context, the inclusion of either marine and brackish water fish species or exotic species could obscure not only any biodiversity pattern, but also biogeographical patterns, since the processes that determine the distribution of parasites are different in the sea than in freshwater.

An obvious pattern uncovered in this study after the analysis of the information is the asymmetrical distribution of the species richness per parasite group. The group with the highest species richness is the nematodes, and even though we cannot establish at this point the reason for this, we argue that it is due to some of the nematode genera occurring in Mexican freshwater fishes (e.g. *Rhabdochona*, with 12 species, see Aguilar-Aguilar *et al.* 2010) not conforming to monophyletic assemblages (Mejía-Madrid *et al.* 2007). This means that their diversification is not associated strictly with vicariance/dispersal events from the same ancestor. Instead, they seem to derive from different lineages from fishes whose affinities are found in both the Nearctic and Neotropical biogeographical regions.

Recent papers have discussed the appropriateness of taxonomic distinctness of parasite assemblages as an alternative measure of parasite diversity in comparative analyses (Luque *et al.* 2004; Luque and Poulin, 2008; Ponlet *et al.* 2011). However, we decided to use species richness since we were only looking for factors that influence species number in an assemblage and not the causes of a narrow or broad



Table 4. Coefficients obtained from generalized linear models performed on fish species values and results of multiple regressions of standardized independent contrasts (estimated through the origin) predicting species richness

(See Tables 1 and 2 for details of variables.)

| Variable             | Total parasites |          |           |         | Ectoparasites |         |           |          | Endoparasites |         |           |         |
|----------------------|-----------------|----------|-----------|---------|---------------|---------|-----------|----------|---------------|---------|-----------|---------|
|                      | Species         |          | Contrasts |         | Species       |         | Contrasts |          | Species       |         | Contrasts |         |
|                      | Estimate        | z value  | Estimate  | t value | Estimate      | z value | Estimate  | t value  | Estimate      | z value | Estimate  | t value |
| (Intercept)          | 0.83            | 7.25***  |           |         | 2.002e-02     | 0.20    |           |          | 2.84          | 1.93    |           |         |
| Size                 |                 |          |           |         |               |         | 2.47e-02  | 4.55***  |               |         |           |         |
| Regions              | 0.23            | 6.26***  | 0.92      | 4.12*** | 2.14e-01      | 4.56*** | -1.14e-01 | -0.77    | -5.28e-02     | -0.77   | 0.29      | 1.42    |
| Localities           | 0.05            | 6.43***  | 0.33      | 9.48*** |               |         | 5.41e-02  | 4.68***  | 7.50e-02      | 7.24*** | 0.28      | 9.02*** |
| Trophic level        |                 |          |           |         |               |         |           |          | 1.74e-01      | 2.64**  |           |         |
| MxL                  |                 |          |           |         |               |         | -1.48e-01 | -3.25**  |               |         |           |         |
| MnL                  |                 |          |           |         |               |         | 1.79e-01  | 2.80**   | -8.03e-02     | -2.26*  |           |         |
| ML                   |                 |          |           |         |               |         | na        | NA       |               |         |           |         |
| Length               |                 |          |           |         |               |         | 1.59e-06  | 0.28     |               |         |           |         |
| Area                 |                 |          |           |         |               |         | 6.74e-06  | 3.38***  |               |         |           |         |
| Flow                 |                 |          |           |         | 2.74e-06      | 2.45*   | 9.58e-06  | 4.34***  | 5.47e-06      | 3.58*** |           |         |
| Basins               |                 |          |           |         |               |         | -2.68e-04 | -0.04    | 5.96e-03      | 2.22*   |           |         |
| MAP                  | 0.001           | 4.22***  | 0.002     | 2.76**  |               |         | -5.10e-05 | -0.14    |               |         | 0.002     | 2.65**  |
| PDM                  | -0.01           | -2.59**  | -0.05     | -2.36*  |               |         | 4.78e-04  | 0.06     | -1.24e-01     | -2.70** | -0.05     | -2.47*  |
| MTWM                 |                 |          |           |         |               |         | 9.23e-02  | 1.31     | -2.08e-02     | -0.51   |           |         |
| MTCM                 |                 |          | 0.01      | 0.07    |               |         | -1.36e-02 | -0.44    |               |         | 0.01      | 0.10    |
| Flow:Basins          |                 |          |           |         |               |         | 1.36e-06  | 9.61***  |               |         |           |         |
| Basins:MAP           |                 |          |           |         |               |         | -1.33e-04 | -7.20*** |               |         |           |         |
| Size:MTCM            |                 |          |           |         |               |         | -2.10e-02 | -6.91*** |               |         |           |         |
| Localities:MTCM      |                 |          | 0.05      | 5.09*** |               |         | 3.14e-02  | 4.32***  |               |         | 0.04      | 4.73*** |
| MnL:PDM              |                 |          |           |         |               |         |           |          | 6.09e-03      | 2.40*   |           |         |
| Localities:PDM       |                 |          |           |         |               |         | -7.18e-03 | -4.29*** |               |         |           |         |
| Regions:Localities   | -0.01           | -3.82*** |           |         |               |         |           |          | 3.32e-01      | 4.23*** |           |         |
| Regions:Basins       |                 |          |           |         |               |         | -2.37e-02 | -7.09*** |               |         |           |         |
| Size:MTWM            |                 |          |           |         |               |         | -1.40e-02 | -2.20*   |               |         |           |         |
| Localities:MTWM      |                 |          |           |         |               |         | -3.55e-02 | -2.55*   |               |         |           |         |
| Area:PDM             |                 |          |           |         |               |         | -4.38e-07 | -2.40*   |               |         |           |         |
| Length:PDM           |                 |          |           |         |               |         | 9.36e-07  | 1.99*    |               |         |           |         |
| Localities:Flow      |                 |          |           |         |               |         | -6.25e-07 | -2.70**  |               |         |           |         |
| Localities:MnL       |                 |          |           |         |               |         | -2.49e-02 | -2.60*   |               |         |           |         |
| Regions:PDM          |                 |          | -0.05     | -2.26*  |               |         |           |          |               |         | -0.03     | -1.60   |
| Regions:MAP          |                 |          | -0.002    | -2.88** |               |         |           |          |               |         | -0.002    | -2.85** |
| Size:Localities:Area |                 |          |           |         |               |         | -3.18e-08 | -3.21**  |               |         |           |         |
| Size:Regions:Basins  |                 |          |           |         |               |         | 2.11e-03  | 5.43***  |               |         |           |         |
| Basins:MAP:MTCM      |                 |          |           |         |               |         | -1.77e-05 | -3.15**  |               |         |           |         |
| Area:Basins:MTWM     |                 |          |           |         |               |         | 1.92e-07  | 4.57***  |               |         |           |         |






Fig. 2. Relationship between the total parasite species richness and the geographical range of the Mexican freshwater fish species ( $r=0.6285$ ,  $p<2.2e-16$ ). Points are phylogenetically independent contrasts positimized as suggested by Garland *et al.* (1992).




Fig. 4. Relationship between the endoparasite species richness and the geographical range of the Mexican freshwater fish species ( $r=0.5497$ ,  $p<2.2e-16$ ). Points are phylogenetically independent contrasts positimized as suggested by Garland *et al.* (1992).




Fig. 3. Relationship between the ectoparasite species richness and the geographical range of the Mexican freshwater fish species ( $r=0.5507$ ,  $p<2.2e-16$ ). Points are phylogenetically independent contrasts positimized as suggested by Garland *et al.* (1992).

host size, precipitation, temperature and latitude, appear to influence the number of parasite species exploiting a freshwater fish species. The host geographical range (the number of sites in which a host species occurs) was the main predictor of the species richness in our study. This characteristic correlated positively with parasite richness, i.e., fishes with broad geographical distributions have richer assemblages than those with limited distribution. This relationship suggests that new parasite species can be acquired in various geographical regions

because fishes living across many localities feed on a wider array of prey species and therefore are exposed to colonization by more parasite species than fishes restricted to a narrow distribution. Under this scenario, fishes are also exposed to colonization by more ectoparasite species with the consequent increase in species richness.

A positive correlation between host body size and ectoparasite species richness was also found. This finding can be explained in the light of island biogeography theory (Kuris *et al.* 1980), where larger hosts offer a larger number of habitats for parasite colonization than smaller ones.

Another pattern detected in the present study was the correlation between the parasite species richness and latitude as well as precipitation and temperature of the distribution area of the fish species. This relation has been addressed in several earlier papers dealing with species richness and latitudinal gradients. Rohde (1992) and Rohde *et al.* (1995) observed that marine fish from tropical latitudes typically harbour richer ectoparasite communities than fish from temperate latitudes. Later, Poulin and Rohde (1997) suggested that the relationship between temperature and the marine-fish ectoparasite community richness was an indicator of the importance of temperature in the diversification of fish parasites in the tropics. Rohde and Heap (1998) confirmed that pattern, but they did not find a correlation between temperature and endoparasite species richness in marine fishes, suggesting that biological differences between ecto- and endoparasites may explain this difference. In accordance with Rohde (1992) the increase in parasite species diversity towards tropical areas is the result of higher diversification rates or

effective evolutionary time, which are correlated with temperature. In the case of freshwater fishes a negative relationship between parasite species richness and temperature has been observed (Choudhury and Dick, 2000; Poulin, 2001). However, in a more recent study Luque and Poulin (2008) found a positive correlation between the parasite taxonomic distinctness of freshwater fishes from the Neotropical Region and temperature for all parasites and for ectoparasites and endoparasites treated separately. Apparently, in marine and freshwater environments, temperature is a major force driving not only the increase in the number of parasite species but also the increase of their taxonomic complexity over evolutionary time, mediated by greater rates of speciation and host colonization (Poulin and Morand, 2004).

Trophic level (host feeding habits, host diet or trophic category) influenced endoparasite species richness, but only when the analyses were conducted with fish species values, i.e., fish species at higher trophic levels may not only acquire more parasites through their diet, but also they may be exposed to a broader range of different parasite taxa. This pattern was only evident when endoparasite assemblages were analysed separately, because helminth endoparasites are usually acquired by ingestion due to their indirect life cycle. For instance, predatory fish (fishes with higher trophic levels) should be exposed to more infective helminth larvae in their diet than planktivores. Over evolutionary time, this should translate in higher parasite colonization rates in predatory fish than in planktivorous fish, pushing up the equilibrium of parasite species richness.

Another host trait that was evaluated in this paper was the position of the host in the water column and its relation with parasite species richness. In particular, no correlation was found between these two variables even though previous studies have shown that benthic fish may harbour more directly transmitted parasite species than pelagic fish. The reason for that pattern has not been identified for freshwater fish parasites, although it has been argued that the benthic boundary layer is a focal point for parasite exchange among faunas inhabiting different vertical zones in marine habitats (Marcogliese, 2002).

Finally, on a larger scale, several other important factors can influence the parasite species distribution and the diversification of parasite assemblages in Mexican fish (e.g. high levels of endemism in certain river basins (Domínguez-Domínguez *et al.* 2006; Huidobro *et al.* 2006) and a significant number of exotic species introduced in the region). Therefore, the patterns uncovered in this study support the contention that parasite diversity, in this case parasite species richness, is largely affected by both host ecology and biogeography.

Although our approach uses the best available data for the helminth parasites of freshwater fishes, our attempts to identify the factors that determine

parasite species richness are preliminary and provide but a glimpse of the patterns and processes that determine the species distribution of the helminth fauna of freshwater fish. Further sampling needs to be conducted in certain host families and particular river basins along the country, even though, as stated by Pérez-Ponce de León and Choudhury (2010), the inventory of the helminth parasite fauna in freshwater fishes in Mexico may be nearing completion (with the exception of the monogeneans).

#### ACKNOWLEDGEMENTS

We thank L. Garcia-Prieto and E. V. Vega-Peña for efficient technical support and statistical advice, respectively. Special thanks to Robert Poulin for helpful observations and comments on an earlier draft of this manuscript and to José Luis Luque and Saúl López Alcaide for suggesting the use of PDAP to perform the phylogenetic contrast method.

#### FINANCIAL SUPPORT

L.G.O. was supported by a scholarship from the Consejo Nacional de Ciencia y Tecnología (CONACyT). This paper fulfills the requirement of L.G.O. to obtain her Ph.D. degree within the Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México. The study was partly funded by the Consejo Nacional de Ciencia y Tecnología (G. P. P. L., grant number CONACyT 83043 and 178830); and the Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica (G. P. P. L., grant number PAPIIT-UNAM IN202111).

#### REFERENCES

- Aguilar-Aguilar, R., Rosas-Valdez, R. and Pérez-Ponce de León, G.** (2010). *Rhabdochona ictaluri* sp. nov. (Nematoda, Rhabdochonidae) from ictalurid catfishes in Mexico. *Acta Parasitologica* **55**, 276–280.
- Aguirre-Macedo, M.L. and Scholz, T.** (2005). *Culuwiya cichlidorum* n. sp. (Digenea: Haploporidae) from the black-belt cichlid *Vieja maculicauda* (Pisces: Cichlidae) from Nicaragua. *Journal of Parasitology* **91**, 1379–1384.
- Bell, G. and Burt, A.** (1991). The comparative biology of parasite species diversity: intestinal helminths of freshwater fishes. *Journal of Animal Ecology* **60**, 1046–1063.
- Bloom, D.D., Piller, K.R., Lyons, J., Mercado-Silva, N. and Medina-Nava, M.** (2009). Systematics and Biogeography of the Silverside Tribe Menidiini (Teleostomi: Atherinopsidae) Based on the Mitochondrial ND2 Gene. *Copeia* **2**, 408–417. doi:10.1643/CI-07-151.
- Bloom, D.D., Unmack, P.J., Gosztonyi, A.E., Piller, K.R. and Lovejoy, N.R.** (2012). It's a family matter: Molecular phylogenetics of Atheriniformes and the polyphyly of the surf silversides (Family: Notocheiridae). *Molecular Phylogenetics and Evolution* **62**, 1025–1030. doi:10.1016/j.ympev.2011.12.006.
- Breden, F., Ptacek, M.B., Rashed, M., Taphorn, D. and Figueiredo, C.A.** (1999). Molecular Phylogeny of the Live-Bearing Fish Genus *Poecilia* (Cyprinodontiformes: Poeciliidae). *Molecular Phylogenetics and Evolution* **12**, 95–104.
- Caspeta-Mandujano, J.M.** (2005) *Nematode Parasites of Freshwater Fish in Mexico: Key to Species, Descriptions and Distribution*. Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, México.
- Chakraborty, P.** (2007). A Morphological Phylogenetic Analysis of Middle American Cichlids with Special Emphasis on the Section 'Nandopsis' sensu Regan. *Miscellaneous Publications Museum of Zoology, University of Michigan* **198**, 1–31.
- Choudhury, A. and Dick, T.A.** (2000). Richness and diversity of helminth communities in tropical freshwater fishes: empirical evidence. *Journal of Biogeography* **27**, 935–956. doi:10.1046/j.1365-2699.2000.00450.x.

- Colwell, R. K. (2006). *EstimateS: Statistical Estimation of Species Richness and Shared Species from Samples. Version 8*. Persisten URL <purl.oclc.org/estimates>
- Colwell, R. K., Mao, C. X. and Chang, J. (2004). Interpolating, extrapolating, and comparing incidence-based species accumulation curves. *Ecology* **85**, 2717–2727.
- Comisión Nacional del Agua (2008). *Estadísticas del Agua en México*. Secretaría del Medio Ambiente y Recursos Naturales, México, D. F.
- Concheiro Pérez, G. A., Rícan, O., Ortí, G., Bermingham, E., Doadrio, I. and Zardoya, R. (2007). Phylogeny and biogeography of 91 species of heroine cichlids (Teleostei: Cichlidae) based on sequences of the cytochrome b gene. *Molecular Phylogenetics and Evolution* **43**, 91–110. doi:10.1016/j.ympev.2006.08.012.
- Crawley, M. J. (2007). *The R Book*. John Wiley & Sons, Ltd, Imperial College London at Silwood Park, UK.
- Cunha, C., Mesquita, N., Dowling, T. E., Gilles, A. and Coelho, M. M. (2002). Phylogenetic relationships of Eurasian and American cyprinids using cytochrome b sequences. *Journal of Fish Biology* **61**, 929–944. doi:10.1006/jfbi.2002.2105.
- Doadrio, I. and Domínguez, O. (2004). Phylogenetic relationships within the fish family Goodeidae based on cytochrome b sequence data. *Molecular Phylogenetics and Evolution* **31**, 416–430. doi:10.1016/j.ympev.2003.08.022.
- Dobson, A. P. and Roberts, M. (1994). The population dynamics of parasitic helminth communities. *Parasitology* **109** (Suppl.) S97–S108.
- Domínguez-Domínguez, O., Doadrio, I. and Pérez-Ponce de León, G. (2006). Historical biogeography of some river basins in central Mexico evidenced by their goodeine freshwater fishes: a preliminary hypothesis using secondary Brooks parsimony analysis. *Journal of Biogeography* **33**, 1437–1447. doi:10.1111/j.1365-2699.2006.01526.
- Espinosa-Huerta, E., García-Prieto, L. and Pérez-Ponce de León, G. (1996). Helminth community structure of *Chirostoma attenuata* (Osteichthyes: Atherinidae) in two Mexican lakes. *Southwestern Naturalist* **41**, 288–292.
- ESRI (1999). *Arc View GIS, Version 3.2*. Environmental Systems Research Institute, Redlands, CA, USA.
- Felsenstein, J. (1985). Phylogenies and the comparative method. *American Naturalist* **125**, 1–15.
- Fernandez-Eguarte, A., Zavala-Hidalgo, J. and Romero-Centeno, R. (2010). Atlas Climático Digital de México. Centro de Ciencias de la Atmósfera, UNAM. <http://uniatmos.atmosfera.unam.mx/>
- Froese, R. and Pauly, D. (2011). *FishBase*. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), version (11/2011).
- García-Prieto, L., García-Varela, M., Mendoza-Garfias, B. and Pérez-Ponce de León, G. (2010). Checklist of the Acanthocephala in wildlife vertebrates of Mexico. *Zootaxa* **2419**, 1–50.
- Garland, T. Jr., Harvey, P. H. and Ives, A. R. (1992). Procedures for the analysis of comparative data using phylogenetically independent contrasts. *Systematic Biology* **41**, 18–32.
- Garrido-Olvera, L., García-Prieto, L. and Pérez-Ponce de León, G. (2006). Checklist of the adult nematode parasites of fishes in freshwater localities from Mexico. *Zootaxa* **1201**, 1–45.
- Guégan, J.-F. and Kennedy, C. R. (1993). Maximum local helminth parasite community richness in British freshwater fish: a test of the colonization time hypothesis. *Parasitology* **106**, 61–100.
- Hardman, M. and Hardman, L. M. (2008). The relative importance of body size and paleoclimatic change as explanatory variables influencing lineage diversification rate: an evolutionary analysis of bullhead catfishes (Siluriformes: Ictaluridae). *Systematic Biology* **57**, 16–130. doi:10.1080/10635150801902193.
- Harris, P. M. and Mayden, R. L. (2001). Phylogenetic relationships of major clades of Catostomidae (Teleostei: Cypriniformes) as inferred from mitochondrial SSU and LSU rDNA sequences. *Molecular Phylogenetics and Evolution* **20**, 225–237. doi:10.1006/mpev.2001.0980.
- Hertwig, S. T. (2008). Phylogeny of the Cyprinodontiformes (Teleostei, Atherinomorpha): the contribution of cranial soft tissue characters. *Zoologica Scripta* **37**, 141–174. doi:10.1111/j.1463-6409.2007.00314.x.
- Hrbek, T., Seckinger, J. and Meyer, A. (2007). A phylogenetic and biogeographic perspective on the evolution of poeciliid fishes. *Molecular Phylogenetics and Evolution* **43**, 986–998. doi:10.1016/j.ympev.2006.06.009.
- Huidobro, L., Morrone, J. J. and Alvarez, J. L. (2006). Distributional patterns of freshwater taxa (fishes, crustaceans and plants) from the Mexican transition zone. *Journal of Biogeography* **33**, 731–741. doi:10.1111/j.1365-2699.2005.01400.x.
- Hulsey, C. D., García de León, F. J. and Rodiles-Hernández, R. (2006). Micro- and macroevolutionary decoupling of cichlid jaws: a test of Liem's key innovation hypothesis. *Evolution* **60**, 2096–2109.
- Kohn, A., Cohen, S. C. and Salgado-Maldonado, G. (2006). Checklist of Monogenea parasites of freshwater and marine fishes, amphibians and reptiles from Mexico, Central America and Caribbean. *Zootaxa* **1289**, 1–114.
- Kuris, A. M., Blaustein, A. R. and Alió, J. J. (1980). Hosts as islands. *American Naturalist* **116**, 570–586.
- Lavoué, S., Miya, M., Kawaguchi, A., Yoshino, T. and Nishida, M. (2008). The phylogenetic position of an undescribed paedomorphic clupeiform taxon: mitogenomic evidence. *Ichthyological Research* **55**, 328–334. doi:10.1007/s10228-008-0044-3.
- Luque, J. L., Mouillot, D. and Poulin, R. (2004). Parasite biodiversity and its determinants in coastal marine teleost fishes of Brazil. *Parasitology* **128**, 671–682.
- Luque, J. L. and Poulin, R. (2004). Use of fish as intermediate hosts by helminth parasites: a comparative analysis. *Acta Parasitologica* **49**, 353–361.
- Luque, J. L. and Poulin, R. (2008). Linking ecology with parasite diversity in Neotropical fishes. *Journal of Fish Biology* **72**, 189–204. doi:10.1111/j.1095-8649.2007.01695.x.
- MacArthur, R. H. and Wilson, E. O. (1967). *The Theory of Island Biogeography*. Princeton University Press, Princeton, NJ, USA.
- Maddison, W. P. and Maddison, D. R. (2011). *Mesquite: A Modular System for Evolutionary Analysis. Version 2.75*. <http://mesquiteproject.org>
- Marcogliese, D. J. (2002). Food webs and the transmission of parasites to marine fish. *Parasitology* **124**, S83–S99.
- McMahan, C. D., Geheber, A. D. and Piller, K. R. (2010). Molecular systematics of the enigmatic Middle American genus Vieja (Teleostei: Cichlidae). *Molecular Phylogenetics and Evolution* **57**, 1293–1300. doi:10.1016/j.ympev.2010.09.005.
- Mejía-Madrid, H., Vázquez-Domínguez, E. and Pérez-Ponce de León, G. (2007). Phylogeography and freshwater basins in Central Mexico: recent history as revealed by the fish parasite *Rhabdochona lichtenfelsi* (Nematoda). *Journal of Biogeography* **34**, 787–801.
- Midford, P. E., Garland, T. Jr. and Maddison, W. P. (2011). *PDAP Package of Mesquite. Version 1.16*. <http://mesquiteproject.org/pdap-mesquite/index.html>
- Miller, R. R., Minckley, W. L. and Norris, S. M. (2005). *Freshwater Fishes of México*, 1st Edn. The University of Chicago Press, Chicago, IL, USA.
- Mirande, J. M. (2009). Weighted parsimony phylogeny of the family Characidae (Teleostei: Characiformes). *Cladistics* **25**, 574–613. doi:10.1111/j.1096-0031.2009.00262.x.
- Miya, M., Takeshima, H., Endo, H., Ishiguro, N. B., Inoue, J. G., Mukai, T., Satoh, T. P., Yamaguchi, M., Kawaguchi, A., Mabuchi, K., Shirai, S. M. and Nishida, M. (2003). Major patterns of higher teleostean phylogenies: a new perspective based on 100 complete mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* **26**, 121–138.
- Morand, S., Cribb, T. H., Kulbicki, M., Rigby, M. C., Chauvet, C., Dufour, V., Faliex, E., Galzin, R., Lo, C. M., Lo-Zat, A., Pichelin, S. and Sasal, P. (2000). Endoparasite species richness of New Caledonian butterfly fishes: host density and diet matter. *Parasitology* **121**, 65–72.
- Moravec, F. (1998). *Nematodes of Freshwater Fishes of the Neotropical Region*. Academia, Praha, Czech Republic.
- Nadler, S. and Pérez-Ponce de León, G. (2011). Integrating molecular and morphological approaches for characterizing parasite cryptic species: implications for parasitology. *Parasitology* **138**, 1688–1709. doi:10.1017/S003118201000168X.
- Near, T. J., Bolnick, D. I. and Wainwright, P. C. (2005). Fossil calibrations and molecular divergence time estimates in centrarchid fishes (Teleostei: Centrarchidae). *Evolution* **59**, 1768–1782.
- Nelson, J. S. (2006). *Fishes of the World*, 4th Edn. John Wiley & Sons, Inc. Hoboken, New Jersey, USA.
- Oceguera-Figueroa, A., Sidall, M. E. and García-Prieto, L. (2010). Sanguijuelas. *Biodiversitas* **90**, 1–5.
- Page, R. D. M. (2003). *Tangled Trees: Phylogeny, Cospeciation, and Coevolution*. University of Chicago Press, Chicago, IL, USA.
- Paterson, A. M. and Gray, R. D. (1997). Host–parasite cospeciation, host switching, and missing the boat. In *Host–Parasite Evolution: General Principles and Avian Models* (ed. Clayton, D. H. and Moore, J.), pp. 236–250. Oxford University Press, Oxford, UK.
- Pérez-Ponce de León, G. and Choudhury, A. (2005). Biogeography of helminth parasites of freshwater fish in Mexico: The search for patterns and processes. *Journal of Biogeography* **32**, 645–659.
- Pérez-Ponce de León, G. and Choudhury, A. (2010). Parasite inventories and DNA-based taxonomy: Lessons from helminths of freshwater fishes in a megadiverse country. *Journal of Parasitology* **96**, 236–244.
- Pérez-Ponce de León, G., García-Prieto, L., León-Régagnon, V. and Choudhury, A. (2000). Helminth communities of native and introduced fishes in Lake Pátzcuaro, Michoacán, México. *Journal of Fish Biology* **57**, 303–325.

- Pérez-Ponce de León, G., García-Prieto, L. and Mendoza-Garfias, B.** (2007). Trematode parasites (Platyhelminthes) of wildlife vertebrates in Mexico. *Zootaxa* **1534**, 1–247.
- Pérez-Ponce de León, G., García-Prieto, L. and Mendoza-Garfias, B.** (2011). Describing Parasite Biodiversity: The Case of the Helminth Fauna of Wildlife Vertebrates in Mexico. In *Changing Diversity in Changing Environment* (ed. Grill, O. and Gianfranco, V.), pp. 33–54. INTECH open. Croatia. Available from: <http://www.intechopen.com/articles/show/title/describing-parasite-biodiversity-the-case-of-the-helminth-fauna-of-wild-life-vertebrates-in-mexico>.
- Pérez-Ponce de León, G., García-Prieto, L., Osorio-Sarabia, D. and León-Régagnon, V.** (1996). *Listados Faunísticos de México VI. Helminthos Parasitos de Peces de Aguas Continentales de México*, 1st Edn. Instituto de Biología, Universidad Nacional Autónoma de México, México.
- Pérez-Rodríguez, R., Domínguez-Domínguez, O., Pérez-Ponce de León, G. and Doadrio, I.** (2009). Phylogenetic relationships and biogeography of the genus *Algansea* Girard (Cypriniformes: Cyprinidae) of central Mexico inferred from molecular data. *BMC Evolutionary Biology* **9**, 223. doi:10.1186/1471-2148-9-223.
- Ponlet, N., Chaisiri, K., Claude, J. and Morand, S.** (2011). Incorporating parasite systematics in comparative analyses of variation in spleen mass and testes sizes of rodents. *Parasitology* **138**, 1804–1814. doi:10.1017/S003118201100028X.
- Poulin, R.** (1995). Phylogeny, ecology, and the richness of parasite communities in vertebrates. *Ecological Monographs* **65**, 283–302.
- Poulin, R.** (1998a). *Evolutionary ecology of parasites*. Chapman and Hall, New York.
- Poulin, R.** (1998b). Comparison of three estimators of species richness in parasite component communities. *Journal of Parasitology* **84**, 485–490.
- Poulin, R.** (2001). Another look at the richness of helminth communities in tropical freshwater fish. *Journal of Biogeography* **28**, 737–743. doi:10.1046/j.1365-2699.2001.00570.x.
- Poulin, R.** (2011). Uneven distribution of cryptic diversity among higher taxa of parasitic worms. *Biology Letters* **7**, 241–244. doi:10.1098/rsbl.2010.0640.
- Poulin, R. and Morand, S.** (2004). *Parasite Biodiversity*. Washington, DC: Smithsonian Books.
- Poulin, R. and Rohde, K.** (1997). Comparing the richness of metazoan communities of marine fishes: controlling for host phylogeny. *Oecologia* **110**, 278–283.
- Ptacek, M. B. and Breden, F.** (1998). Phylogenetic relationships among the mollies (Poeciliidae: *Poecilia: Mollinnesia* group) based on mitochondrial DNA sequences. *Journal of Fish Biology* **53** (Suppl. A), 64–81.
- R Development Core Team.** (2010). *R: Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <http://www.r-project.org/>
- Reznick, D. N., Mateos, M. and Springer, M. S.** (2002). Independent Origins and Rapid Evolution of the Placenta in the Fish Genus *Poeciliopsis*. *Science* **298**, 1018–1020. doi:10.1126/science.1076018.
- Rícan, O., Zardoya, R. and Doadrio, I.** (2008). Phylogenetic relationships of Middle American cichlids (Cichlidae, Heroini) based on combined evidence from nuclear genes, mtDNA, and morphology. *Molecular Phylogenetics and Evolution* **49**, 941–957. doi:10.1016/j.ympev.2008.07.022.
- Roberts, M. G., Dobson, A. P., Arneberg, P., De Leo, G. A., Krecek, R. C., Manfredi, M. T., Lanfranchi, P. and Zaffaroni, E.** (2002). Parasite community ecology and biodiversity. In *The Ecology of Wildlife Diseases* (ed. Hudson, P. J., Rizzoli, A., Grenfell, B. T., Heesterbeek, H. and Dobson, A. P.), pp. 63–82. Oxford University Press, Oxford, UK.
- Rohde, K.** (1992). Latitudinal gradients in species diversity: the search for the primary cause. *Oikos* **65**, 514–527.
- Rohde, K., Hayward, C. and Heap, M.** (1995). Aspects of the ecology of metazoan ectoparasites of marine fishes. *International Journal for Parasitology* **25**, 945–970.
- Rohde, K. and Heap, M.** (1998). Latitudinal differences in species and community richness and in community structure of metazoan endo- and ectoparasites of marine teleost fish. *International Journal for Parasitology* **28**, 461–474.
- Rojas, E., Pérez-Ponce de León, G. and García-Prieto, L.** (1997). Helminth community structure of some freshwater fishes from Pátzcuaro, Michoacán, Mexico. *Tropical Ecology* **38**, 121–131.
- Salgado-Maldonado, G.** (2006). Checklist of helminth parasites of freshwater fishes from Mexico. *Zootaxa* **1324**, 1–357.
- Salgado-Maldonado, G., Aguilar-Aguilar, R., Cabañas-Carranza, G., Soto-Galera, E. Y. and Mendoza-Palmero, C.** (2005). Helminth parasites in freshwater fish from the Papaloapan river basin, Mexico. *Parasitology Research* **96**, 69–89.
- Salgado-Maldonado, G. and Kennedy, C. R.** (1997). Richness and similarity of helminth communities in the tropical cichlid fish *Cichlasoma urophthalmus* from the Yucatán Peninsula, Mexico. *Parasitology* **114**, 581–590.
- Sánchez-Nava, P., Salgado-Maldonado, G., Soto-Galera, E. and Jaimes-Cruz, B.** (2004). Helminth parasites of *Girardinichthys multiradiatus* (Pisces: Goodeidae) in the upper Lerma River sub-basin, Mexico. *Parasitology Research* **93**, 396–402.
- Sasal, P., Morand, S. and Guégan, J.-F.** (1997). Parasite species richness for fish of the Mediterranean Sea. *Marine Ecology Progress Series* **149**, 61–71.
- Scholz, T., Aguirre-Macedo, M. L. and Salgado-Maldonado, G.** (2001). Trematodes of the family Heterophyidae (Digenea) in Mexico: a review of species and new host and geographical records. *Journal of Natural History* **35**, 1733–1772.
- Scholz, T., Vargas-Vázquez, J., Aguirre-Macedo, L. and Vidal-Martínez, V. M.** (1997). Species of Ascocotyle Looss, 1899 (Digenea: Heterophyidae) of the Yucatan Peninsula, Mexico, and notes on their life-cycles. *Systematic Parasitology* **36**, 161–181.
- Schönhuth, S. and Doadrio, I.** (2003). Phylogenetic relationships of Mexican minnows of the genus *Notropis* (Actinopterygii, Cyprinidae). *Biological Journal of the Linnean Society* **80**, 323–337.
- Schönhuth, S. and Mayden, R. L.** (2010). Phylogenetic relationships in the genus *Cyprinella* (Actinopterygii: Cyprinidae) based on mitochondrial and nuclear gene sequences. *Molecular Phylogenetics and Evolution* **55**, 77–98. doi:10.1016/j.ympev.2009.10.030.
- Simková, A., Morand, S., Matejusová, I., Jurajda, P. and Gelnar, M.** (2001). Local and regional influences on patterns of parasite species richness of central European fishes. *Biodiversity and Conservation* **10**, 511–525.
- Simons, A. M., Berendzen, P. B. and Mayden, R. L.** (2003). Molecular systematics of North American phoxinin genera (Actinopterygii: Cyprinidae) inferred from mitochondrial 12S and 16S ribosomal RNA sequences. *Zoological Journal of the Linnean Society* **139**, 63–80.
- Takemoto, R. M., Pavanelli, G. C., Lizama, M. A. P., Luque, J. L. and Poulin, R.** (2005). Host population density as the major determinant of endoparasite species richness in floodplain fishes of the upper Paraná River, Brazil. *Journal of Helminthology* **79**, 75–84.
- Vidal-Martínez, V. M.** (1995). Processes structuring the helminth communities of native cichlid fishes from Southern Mexico. Ph.D. thesis, University of Exeter, Exeter, UK.
- Vidal-Martínez, V. M., Aguirre-Macedo, M. L., Scholz, T., González-Solís, D. and Mendoza-Franco, E. F.** (2001). *Atlas of the Helminth Parasites of Cichlid Fish of Mexico*. Academia, Praha, Czech Republic.
- Walther, B. A. and Moore, J. L.** (2005). The concepts of bias, precision and accuracy, and their use in the performance of species richness estimators, with a literature review of estimator performance. *Ecography* **28**, 815–829.
- Walther, B. A. and Morand, S.** (1998). Comparative performance of species richness estimation methods. *Parasitology* **116**, 395–405.
- Wilcox, T. P., García de León, F. J., Hendrickson, D. A. and Hillis, D. M.** (2004). Convergence among cave catfishes: long-branch attraction and a Bayesian relative rates test. *Molecular Phylogenetics and Evolution* **31**, 1101–1113. doi:10.1016/j.ympev.2003.11.006.