

# IRISH JOURNAL OF PSYCHOLOGICAL MEDICINE

VOL 21 NO 3 SEP 2004

ISSN 0790-9667



**'Tuscan landscape'** by Jane Willoughby, 2004. Detail from large mural (c.70ft x 11ft).  
Private commission, Co. Kilkenny.



In a broad  
spectrum of  
patients...

...See the difference  
**GEODON** can make

Abbreviated Prescribing Information for Geodon (ziprasidone). Republic of Ireland Geodon™. Presentation: Capsules containing ziprasidone hydrochloride monohydrate equivalent to 20, 40, 60 and 80mg ziprasidone. Indications: Treatment of schizophrenia Dosage: Acute treatment - 40mg twice daily with food. Maximum dosage of 80mg twice daily may be reached by day 3 of treatment. Maintenance treatment - use the lowest effective dose. In elderly: A lower starting dose should be considered for patients over 65 where clinical factors warrant. In children: Caution as no evaluation under 18 years of age. In renal impairment: No dosage adjustment required. In hepatic impairment: Consider lower doses in hepatic insufficiency. Caution in severe hepatic insufficiency. Contra-indications: Known hypersensitivity to any ingredient of the product. Known QT-interval prolongation. Congenital long QT syndrome. Recent acute myocardial infarction. Uncompensated heart failure. Arrhythmias treated with class IA and III antiarrhythmic drugs. Concomitant treatment with medicines known to prolong the QT interval. Special warnings: A medical history, family history and physical examination should be undertaken to identify patients for whom ziprasidone is not recommended. Mild to moderate dose-related QT-interval prolongation, therefore, do not give together with medicinal products known to prolong the QT interval. Caution in patients with significant bradycardia. Before treatment is started - correct electrolyte disturbances; and as with other drugs which prolong QT interval, consider ECG review in patients with stable cardiac disease. If cardiac symptoms occur, consider the possibility of a malignant cardiac arrhythmia and perform a cardiac evaluation, including an ECG. It is recommended to stop treatment if the QT interval is >500msec. No cases of Neuroleptic Malignant Syndrome (NMS) seen in clinical trials, but potential risk cannot be excluded. Management of NMS should include immediate withdrawal of all antipsychotic drugs. Potential to cause tardive dyskinesia, if signs appear consider dose reduction or discontinuation. Caution in patients with a history of seizures. Interactions: ziprasidone should not be given with medicinal products known to prolong the QT interval (see SPC for details). Caution in combination with other centrally acting drugs and alcohol. Ziprasidone is unlikely to cause clinically important drug interactions mediated by CYP3A4 or CYP2D6 (see SPC for details). Pregnancy and lactation: Not recommended unless the expected benefit outweighs the risk. Women of childbearing potential should use an appropriate method of contraception. Avoid breastfeeding. Driving: Ziprasidone may cause somnolence, therefore caution patients likely to drive or operate machines. Undesirable effects: In short term placebo controlled trials: >1/10 somnolence; >1/100, <1/100 asthenia, headache, constipation, dry mouth, dyspepsia, increased salivation, nausea, vomiting, agitation, akathisia, dizziness, dystonia, extrapyramidal syndrome, hypertonia, tremor, abnormal vision; >1/1000, <1/100 pain, postural hypotension, tachycardia, flatulence, thirst, joint disorder, leg cramps, cogwheel rigidity, paresthesia, speech disorder, tardive dyskinesia, rhinitis, rash, urticaria. In long term maintenance trials: elevated prolactin levels, returning to normal without cessation of treatment and rare reports of clinical manifestation (gynaecomastia and breast enlargement). Legal Category: POM. Package quantities: blister packs containing 56 capsules. Further information on request: Pfizer (Ireland) Limited, Parkway House, Ballymount Road Lower, Dublin 12, Republic of Ireland. Marketing Authorisation numbers: PA 19/52/5. Date of first authorisation/renewal of the authorisation:

MAKE THE SWITCH

**GEODON**™

(ziprasidone HCl)

See the difference



**Editor-in-Chief:** Brian Lawlor

**Production Editor:**

Anne Henrichsen

**Advertising Manager:**

Helen Martin

**Administrator:**

Andrea McAdam

**Founding Editor:** Mark Hartman

**Associate Editors:**

Ted Dinan (Cork), David King (Belfast)

**Editorial Board:** Ken Brown (Belfast),

Patricia Casey (Dublin), Anthony Clare

(Dublin), Stephen Cooper (Belfast),

Thomas Fahy (Galway), Michael

Fitzgerald (Dublin), Brian Leonard

(Galway), Roy McClelland (Belfast),

Aidan McGennis (Dublin), Ciaran

O'Boyle (Dublin), Eadbhard

O'Callaghan (Dublin), Art O'Connor

(Dublin), Ethna O'Gorman (Belfast),

Brian O'Shea (Wicklow), Ian Pullen

(Edinburgh), David Sheehan (Tampa),

Philip Snaith (Leeds), Hugh Staunton

(Dublin), John Waddington (Dublin),

Richard Williams (Victoria)

**Statistical Editor:**

Ronan Conroy (Dublin)

**Submissions & correspondence to:**

The Editor,

Irish Journal of Psychological Medicine,

25 Adelaide Street,

Dun Laoghaire,

Co Dublin, Ireland.

**Telephone:** 00-353-1-2803967

**Fax:** 00-353-1-2807076

**Email:** psych@medmedia.ie

**Website:** www.ijpm.org

**Publisher**

MedMedia Ltd.

25 Adelaide Street, Dun Laoghaire,

Co Dublin, Ireland.

**Printing:** W&G Bairds Ltd

**Subscriptions**

Rates per volume of four issues

(Mar, Jun, Sept, Dec) Price Regions:

EU countries: €107, Stg65

Rest of World: €126, \$111

Incl. airmail postage internationally.

**Subscription enquiries, orders and cheques made payable to:**

Extenza Turpin

Blackhorse Road, Letchworth

SG6 1HN, England.

Tel : +44 01462 672555

Fax: + 44 01462 480947

Email: custserv@extenza-turpin.com

www.extenza-turpin.com

**Circulation**

2,200 to 54 countries.

The Journal participates in the World

Health Organisation project to improve

distribution of scientific materials on

mental health. Publication does not

imply endorsement. Limited

photocopying authorisation granted for

a fee to Copyright Clearance Center,

27 Congress Street, Salem, MA 01970,

USA, or to appropriate Reproduction

Rights Organisation; isolated non-profit,

academic photocopying excepted.

**Editorial**

**77 Cannabis abuse and psychiatric disorder**

John Tobin

**Original Papers**

**78 Adolescents with suicidal behaviour: attendance at A&E and six month follow-up**

Sharon Bolger, Peter O'Connor, Kevin Malone, Carol Fitzpatrick

**85 Withdrawal of antipsychotic drugs from adults with intellectual disabilities**

Claire Stevenson, Lovely Rajan, Gavin Reid, Craig Melville, Robin McGilp, Sally-Ann Cooper

**Brief reports**

**91 A brief adherence therapy: training issues and patients' satisfaction**

Marie Boilson, Jane M Murdoch, Alastair M Hull, Ross J Hamilton, Jan Scott

**Reviews**

**95 Evidence for use of depot neuroleptic medication**

MacDara McCauley, Gerard Connolly

**100 Sertraline as a treatment for post-traumatic stress disorder: a systematic review and meta-analysis**

Paul Mooney, Janette Oakley, Michael Ferriter, Raymond Travers

**Perspective**

**104 Interpersonal disturbance in OCD and its place in the professional consultation**

John O'Connor, Ray Fuller, Mary Fell

**88a John Dunne Medal**

**94 Guidelines for Authors**

**99 Subscriptions**

Indexed and abstracted by BIOLOGICAL ABSTRACTS (BIOSIS Previews); CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE/INIST: PASCAL; EXCERPTA MEDICA/EMBASE; INSTITUTE FOR SCIENTIFIC INFORMATION: CURRENT CONTENTS/ Social & Behavioural Sciences (Social Science CITATION INDEX, Research Alert); PSYCHOLOGICAL ABSTRACTS (PsycINFO/PsycLIT); Cumulative Index to Nursing & Allied Health Literature, Current AIDS Literature (CAB Abstracts), International Pharmaceutical Abstracts, Linguistics & Language Behaviour Abstracts, Nutrition Abstracts and Reviews, (CAB Abstracts), Referativnyi Zhurnal, Social Planning/Policy & Development Abstracts, Social Work Research & Abstracts, Sociological Abstracts.


Microfilm, microfiche & article copies from **University Microfilms International**, 300 North Zeeb Rd., Ann Arbor, MI 48106, USA. Journal included in the **Adonis** service, whereby article copies can be printed out from compact disks (CD-ROM) on demand; explanatory leaflet available from ADONIS BV, PO Box 639, 1000 AV Amsterdam, The Netherlands. Journal listed in **Ulrich's** International Periodicals Directory (**Bowker** International Serials Database), **EBSCO's** Selected Periodicals for the Medical and Health Sciences, & **EBSCO's** Librarians' Handbook.



# In Depression & Anxiety...



... Patients Get Better and Stay Better with



**LUSTRAL**<sup>TM</sup>  
sertraline



Abbreviated Prescribing Information: LUSTRAL<sup>TM</sup> (sertraline) Presentation: Tablets containing 50mg or 100mg sertraline. Indications: Treatment of symptoms of depressive illness, including accompanying symptoms of anxiety. Prevention of relapse or recurrence of depressive episodes, including accompanying symptoms of anxiety. Obsessive compulsive disorder (OCD) in adults and children. Panic disorder, with or without agoraphobia. Post-traumatic stress disorder (PTSD). Dosage: Lustral should be given as a single daily dose. The initial dose in depression and OCD is 50mg and the usual antidepressant dose is 50mg. The initial dose in panic disorder and PTSD is 25mg, increasing to 50mg after one week. Dosage can be further increased, if appropriate, to a maximum of 200mg daily. Changes in dose should not be made more frequently than once per week given the 24 hour elimination half life of sertraline. Patients should be maintained on the lowest effective dose. Use in children (aged 6-17): Age 7-16: 25-75mg daily. Age 17: Usual adult dose increasing to 50 mg/day after 1 week. Ages 13-17: Usual adult

dose. Consider generally lower body weights of children to avoid overdosing. Do not increase doses at intervals of less than 1 week. Use in the elderly: Usual adult dose. Contra-Indications: Hypersensitivity to this group of drugs. Hepatic insufficiency, unstable epilepsy and convulsant disorders, pregnancy and lactation. Do not use with, or within two weeks of ending treatment with, MAOIs. At least 14 days should elapse before starting any MAOI following discontinuation of Lustral. Do not use with pimozide. Precautions, warnings: Renal insufficiency, ECT, epilepsy, driving. Lustral should be discontinued in a patient who develops seizures. Lustral should not be administered with benzodiazepines or other tranquilizers in patients who drive or operate machinery. Serotonergic drugs such as tryptophan or fenfluramine should be used with caution. Patients should be closely supervised for the possibility of suicide attempt or activation of mania/hypomania. The use of Lustral in children and adolescents under 18 with major depressive disorder is not recommended, as benefit/risk has not been established. Drug Interactions: Caution with other centrally active medication. Lithium levels should be monitored. Increased plasma levels have been demonstrated when co-administered with Lustral, therefore Lustral should not

be taken together with pimozide. Although Lustral has been shown to have no adverse interaction with alcohol, concomitant use with alcohol is not recommended. The potential for Lustral to interact with other highly protein bound drugs should be borne in mind. Interactions with e.g. warfarin, diazepam, tolbutamide and cimetidine have not been fully assessed. With warfarin prothrombin time should be monitored when Lustral is initiated or stopped. Side-Effects: Dry mouth, nausea, diarrhoea/loose stools, ejaculatory delay, tremor, increased sweating, dizziness, insomnia, somnolence, headache, anorexia and dyspepsia. Rarely, abnormal LFTs, hyponatraemia. Additionally, the following adverse events were observed in clinical trials in paediatric OCD patients: anorexia, weight decrease, fatigue, chest pain, fever, malaise, hyperkinesia, tremor, urinary incontinence, nausea, insomnia, nervousness, agitation, impaired concentration, manic reaction, anxiety, emotional lability, abnormal thinking, breast pain, dysmenorrhoea, menstrual disorder, epistaxis, rash, skin disorder, purpura and headache. The following have been reported with Lustral but may have no causal relationship: vomiting, abdominal pain, movement disorders, convulsions, menstrual irregularities, hyperprolactinaemia, galactorrhoea, rash and alopecia. Rarely, pancre-

atitis, serious liver events, altered platelet function, abnormal bleeding and purpura. As with other serotonin re-uptake inhibitors rare reports of agitation, confusion, depersonalisation, hallucinations, nervousness, postural hypotension, hypo/hypertension, tachycardia and arrhythmias. Withdrawal reactions have been reported with Lustral. Common symptoms include dizziness, paraesthesia, headache, anxiety and nausea. Abrupt discontinuation of treatment with Lustral should be avoided. The majority of symptoms experienced on withdrawal of Lustral are non-serious and self-limiting. Legal Category: S1A. Package Quantities: 50mg tablet (PA 822/1/4) Calendar pack of 28; 100mg tablet (PA 822/1/5) Calendar pack of 28. Product Authorisation Holder: Pfizer Healthcare Ireland, Parkway House, Ballymount Road Lower, Dublin 12, Republic of Ireland. Further information on request: Pfizer Healthcare Ireland. Date last revised: January 2004

