

Homomorphisms between diffeomorphism groups

KATHRYN MANN

Department of Mathematics, University of Chicago, 5734 University Avenue,
Chicago, IL 60637, USA
(e-mail: mann@math.uchicago.edu)

(Received 28 November 2012 and accepted in revised form 28 March 2013)

Abstract. For $r \geq 3$, $p \geq 2$, we classify all actions of the groups $\text{Diff}_c^r(\mathbb{R})$ and $\text{Diff}_+^r(S^1)$ by C^p -diffeomorphisms on the line and on the circle. This is the same as describing all non-trivial group homomorphisms between groups of compactly supported diffeomorphisms on 1-manifolds. We show that all such actions have an elementary form, which we call *topologically diagonal*. As an application, we answer a question of Ghys in the 1-manifold case: if M is any closed manifold, and $\text{Diff}^\infty(M)_0$ injects into the diffeomorphism group of a 1-manifold, must M be one-dimensional? We show that the answer is yes, even under more general conditions. Several lemmas on subgroups of diffeomorphism groups are of independent interest, including results on commuting subgroups and flows.

1. Introduction

For a manifold M (assumed connected and without boundary but not necessarily closed), let $\text{Diff}_c^r(M)$ denote the group of compactly supported, orientation-preserving, C^r -diffeomorphisms of M isotopic to the identity. We study all non-trivial homomorphisms $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ between two such diffeomorphism groups. Here are some easy examples to keep in mind.

Example 1.1.

- If M_1 is an open submanifold of M_2 , there is a natural inclusion $\text{Diff}_c^r(M_1) \hookrightarrow \text{Diff}_c^r(M_2)$ by extending any compactly supported diffeomorphism of M_1 to be the identity on $M_2 \setminus M_1$.
- A C^r -diffeomorphism $f : M_1 \rightarrow M_2$ induces an isomorphism $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ defined by $\Phi(g) = fgf^{-1}$.
- Combining the two examples above, a diffeomorphism f from M_1 to an open subset of M_2 can be used to define a natural injection $\Phi : \text{Diff}_c^r(M_1) \hookrightarrow \text{Diff}_c^r(M_2)$ by

$$\Phi(g)(x) = \begin{cases} fgf^{-1}(x) & \text{if } x \in f(M_1), \\ x & \text{otherwise.} \end{cases}$$

If f is instead C^p for some $p < r$, then the image of Φ will lie in $\text{Diff}_c^p(M_2)$.

- (d) Finally, there are some special cases where $M_1 \subset M_2$ is a non-open submanifold and diffeomorphisms of M_1 still extend to M_2 in a natural way. This happens, for example, if $M_2 = M_1 \times N$ for a compact manifold N , or if M_2 is the unit tangent bundle of M_1 .

Our goal is to find conditions on M_1 and M_2 that guarantee the existence of non-trivial homomorphisms $\text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$, and to describe as much as possible what these homomorphisms look like. Note that even though all the examples of homomorphisms above are continuous, we make no assumptions on continuity.

The analogous question for *isomorphisms* rather than *homomorphisms* between diffeomorphism groups is completely answered by a theorem of Filipkiewicz. It says that case (b) of Example 1.1 is all that can occur.

THEOREM 1.2. (Filipkiewicz [5]) *Let M_1 and M_2 be smooth manifolds and suppose that there is an isomorphism $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$. Then M_1 and M_2 are diffeomorphic. In fact, more is true: $r = p$ and Φ is induced by a C^r -diffeomorphism $f : M_1 \rightarrow M_2$.*

Filipkiewicz's original statement was for isomorphisms between the identity components of the full diffeomorphism groups, but his theorem holds for the group of compactly supported diffeomorphisms as well. More recently, Filipkiewicz's theorem has been generalized to isomorphisms between other groups of diffeomorphisms (e.g. symplectomorphisms, contact diffeomorphisms) by Banyaga, Rybicki, and others (see, for example, [1, 15]). The spirit of all these results is the same: given a 'sufficiently large' class of diffeomorphisms \mathcal{C} , the existence of an isomorphism between $\mathcal{C}(M_1)$ and $\mathcal{C}(M_2)$ implies that M_1 and M_2 are diffeomorphic and the isomorphism is induced by a diffeomorphism.

By contrast, we know almost nothing about homomorphisms between diffeomorphism groups. As an example of our ignorance, consider the following question of Ghys. Here $\text{Diff}^\infty(M)_0$ denotes the identity component of the group of C^∞ -diffeomorphisms of M .

Question 1.3. (Ghys [6]) *Let M_1 and M_2 be closed manifolds, and suppose that there is an (injective) homomorphism $\text{Diff}^\infty(M_1)_0 \rightarrow \text{Diff}^\infty(M_2)_0$. Does it follow that $\dim(M_1) \leq \dim(M_2)$?*

In fact, $\text{Diff}^\infty(M)_0$ is a simple group, so any non-trivial homomorphism is necessarily injective. That $\text{Diff}^\infty(M)_0$ is simple follows from a deep result due to Mather and Thurston, non-trivial even in the case of 1-manifolds (!). Simplicity holds for a larger class of groups as well; for any connected manifold M and any $r \neq \dim(M) + 1$, the group $\text{Diff}_c^r(M)_0$ is simple [9, 10]. Thus, it makes sense to ask Question 1.3 for non-compact manifolds, replacing $\text{Diff}^r(M_i)_0$ with $\text{Diff}_c^r(M_i)$.

Ghys's question appeared in print in 1991, and to our knowledge, this paper is the first to give even a partial answer in a special case.

1.1. Our results. We study the special case of homomorphisms $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ when M_2 is a 1-manifold. The reader will see the difficulty of the problem and some of the richness in examples already apparent at the one-dimensional level. However, we are able to give a complete answer to the question of which manifolds M_1 admit non-trivial homomorphisms $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ or $\Phi : \text{Diff}^r(M_1)_0 \rightarrow \text{Diff}_c^r(M_2)$ for

some 1-manifold M_2 , and describe precisely what all such homomorphisms look like. Essentially, M_1 must be one-dimensional, and Φ described by a slight generalization of the obvious embedding of diffeomorphism groups given in Example 1.1(c). This generalization is constructed by taking multiple embeddings $M_1 \hookrightarrow M_2$ and having diffeomorphisms of M_1 act ‘diagonally’ on M_2 . We call this a *topologically diagonal embedding*.

Definition 1.4. (Topologically diagonal embedding) Let M_1 and M_2 be manifolds, and suppose that $\{f_1, f_2, f_3, \dots\}$ is a collection (finite or infinite) of C^r embeddings $M_1 \rightarrow M_2$ whose images are pairwise disjoint and contained in some compact subset of M_2 . Define $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ by

$$\Phi(g)(x) = \begin{cases} f_i g f_i^{-1}(x) & \text{if } x \in f_i(M_1), \\ x & \text{otherwise.} \end{cases}$$

A C^r -topologically diagonal embedding is a homomorphism $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ obtained by this construction.

A *topologically diagonal embedding* is an embedding $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ obtained by the same construction when the embeddings f_i are only required to be continuous.

Our main theorem says that these define basically all homomorphisms between 1-manifold diffeomorphism groups.

THEOREM 1.5. *Let $r \geq 3$, $p \geq 2$ and let M_1 and M_2 be 1-manifolds. Every homomorphism $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ is topologically diagonal. If $r = p$, then Φ is C^r -topologically diagonal.*

We also have a stronger version.

THEOREM 1.6. *Let $r \geq 3$, $p \geq 2$, let M_1 be any manifold and M_2 a 1-manifold. Suppose that $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ is an injective homomorphism. Then $\dim(M_1) = 1$ and Φ is topologically diagonal. If $r = p$, then Φ is C^r -topologically diagonal.*

In particular, this answers the 1-manifold case of Ghys’s Question 1.3 in the affirmative.

1.2. *The algebra–topology link.* There are two points to note regarding Theorems 1.5 and 1.6. First, our results parallel that of Filipkiewicz by establishing a relationship between the topology of a manifold and the algebraic structure of its group of diffeomorphisms. In fact, our proof strategy will be to gradually pin down this relationship as closely as possible.

Second, our results say that the richness of the algebraic structure of diffeomorphism groups *forces* continuity: group homomorphisms $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$, where M_2 is a 1-manifold, are *continuous* (assuming appropriate constraints on p and r). This need not be true if $\text{Diff}_c^r(M_1)$ is replaced with another topological group! Consider the following example.

Example 1.7. (A non-continuous, injective group homomorphism $\mathbb{R} \rightarrow \text{Diff}_c^r(M)$) Let M be a manifold, and let $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ be a non-continuous, injective additive group

homomorphism. Such a homomorphism may be constructed by permuting the elements of a basis for \mathbb{R} over \mathbb{Q} and extending linearly to \mathbb{R} . Then for any compactly supported C^r flow ψ^t on M , the map $t \mapsto \psi^{\alpha(t)}$ is a non-continuous, injective homomorphism from \mathbb{R} to $\text{Diff}_c^r(M)$.

For an even more pathological example, consider a second flow ϕ^t with support disjoint from ψ^t . Then

$$t \mapsto \begin{cases} \psi^t & \text{for } t \in \mathbb{Q}, \\ \phi^{\alpha(t)} & \text{for } t \notin \mathbb{Q}, \end{cases}$$

also gives a non-continuous, injective homomorphism from \mathbb{R} to $\text{Diff}_c^r(M)$.

A key step in our proof will be showing that this example cannot occur within the context of diffeomorphism groups: if $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ is a homomorphism between 1-manifold diffeomorphism groups, then any topological \mathbb{R} -subgroup of $\text{Diff}_c^r(M_1)$ maps continuously into $\text{Diff}_c^p(M_2)$.

1.3. Outline. In §2 we establish a preliminarily algebraic–topological relationship using commuting subgroups. This is done by developing the elementary theory of 1-manifold diffeomorphism groups stemming from Kopell’s lemma and Hölder’s theorem.

In §3 we attack continuity. Using the tools of §2, we show that \mathbb{R} -subgroups of diffeomorphisms of a 1-manifold (flows) behave well under homomorphisms of diffeomorphism groups. This rules out the kind of behavior in Example 1.7.

We prove Theorem 1.5 in §4, strengthening the algebraic–topological relationship from §2 and using continuity of \mathbb{R} -subgroups to construct continuous maps f_i from M_1 to M_2 that define a topologically diagonal embedding.

In §5 we then compare 1-manifold diffeomorphism groups to n -manifold diffeomorphism groups and show that $\dim(M) = 1$ can be detected as an algebraic invariant of a diffeomorphism group. This is used to prove Theorem 1.6, answering the 1-manifold case of Ghys’s question.

Finally, §6 concludes with some remarks on the necessity of our hypotheses (e.g. compact supports) and further questions.

2. Commuting groups of diffeomorphisms

Algebraic–topological associations. Our first goal is to associate the algebraic data of a group of diffeomorphisms of a manifold to the topological data of the manifold. One example of the kind of association we have in mind appears in Filipkiewicz’s proof of Theorem 1.2. Given an isomorphism between the diffeomorphism groups of two manifolds, Filipkiewicz builds a map between the manifolds by associating points of each manifold with their stabilizers—the subgroups $\{g \in \text{Diff}_c^r(M) : g(x) = x\}$. Point stabilizers are maximal subgroups of $\text{Diff}_c^r(M)$, so this gives a loose association between an algebraic property (maximality) and a topological object (point). This association is eventually used to build a point-to-point map out of the algebraic data of a group isomorphism.

However, Filipkiewicz’s point/maximal subgroup association will not work for our purposes. The main problem is that the maximality is not preserved under group homomorphisms. Whereas an *isomorphism* $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ maps maximal

subgroups of $\text{Diff}_c^r(M_1)$ to maximal subgroups of $\text{Diff}_c^r(M_2)$, a *homomorphism* may not. To prove that group homomorphisms $\text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^r(M_2)$ have nice topological properties, we need to make use of an algebraic–topological correspondence robust under homomorphisms. We will look at commuting, rather than maximal, subgroups. Our analysis starts with a trivial observation.

Observation 2.1. Let M be a manifold, $N \subset M$ a submanifold, and H the subgroup of $\text{Diff}^r(M)$ that fixes N pointwise. Then H commutes with the subgroup of diffeomorphisms supported on N . In other words, the centralizer of H in $\text{Diff}^r(M)$ is ‘large’.

It turns out that when M is one-dimensional, this property is *characteristic* of subgroups of $\text{Diff}_c^r(M)$ that fix a submanifold. Specifically, we have the following proposition.

PROPOSITION 2.2. *If G is a non-abelian subgroup of $\text{Diff}_c^r(\mathbb{R})$ or $\text{Diff}^r(I)$ and G has non-abelian centralizer, then there is an open interval contained in $\text{Fix}(G)$.*

Here, and in the following, $\text{Fix}(G)$ denotes the set $\{x \in M : g(x) = x \text{ for all } g \in G\}$. Similarly, for a single diffeomorphism g , we set $\text{Fix}(g) := \{x \in M : g(x) = x\}$.

A similar statement to Proposition 2.2 holds for S^1 ; it is stated as Proposition 2.12 below. But before we consider the S^1 case, let us first develop some of the necessary background for the proof of Proposition 2.2.

2.1. Hölder–Kopell theory. To prove Proposition 2.2 and its counterpart for diffeomorphisms of S^1 , we extend the theory of commuting diffeomorphisms of 1-manifolds that stems from two classical theorems (Theorems 2.3 and 2.4 below) due to Hölder and Kopell. This theory for 1-manifolds is already surprisingly rich—one can use Theorems 2.3 and 2.4 to prove, for instance, that all nilpotent groups of $\text{Diff}_+^2(S^1)$ are abelian, and that certain dynamically characterized subgroups of $\text{Diff}_+^2(\mathbb{R})$ are all conjugate into the affine group. Details and some background can be found in [4] and in [14, Ch. 4]. Here we restrict our attention to results needed to prove Proposition 2.2.

THEOREM 2.3. (Kopell’s lemma [8]) *Let g and h be commuting C^2 -diffeomorphisms of the half-open interval $I := [0, 1)$. If g is fixed point-free on the interior of I and h is not the identity, then h is also fixed-point-free on $\text{int}(I)$.*

THEOREM 2.4. (Hölder [7]) *Let G be a group of orientation-preserving homeomorphisms of S^1 or of the open interval. If G acts freely (i.e. no element other than the identity has a fixed point), then G is abelian.*

Both of these theorems admit elementary proofs. A nice proof of Theorem 2.3 is given in [14, §4.1.1], and also in [4, Theorem 2.4]. The result most useful to us is the following combination of Hölder’s theorem and Kopell’s lemma.

COROLLARY 2.5. (Hölder–Kopell lemma) *Let $g \in \text{Diff}_+^2(I)$. If g has no interior fixed points, then the centralizer of g in $\text{Diff}_+^2(I)$ is abelian.*

Proof. Each element h in the centralizer $C(g)$ commutes with g , so by Kopell’s lemma h is either the identity or acts without fixed points on $\text{int}(I)$. Thus, $C(g)$ acts freely and so by Hölder’s theorem $C(g)$ is abelian. \square

Remark 2.6. Corollary 2.5 also follows from a more general theorem of Szekeres (Theorem 3.2 in §3). We do not need the generality of Szekeres's theorem here, and as it is a deeper result than Hölder's theorem and Kopell's lemma we postpone its introduction to §3.

We now make use of Corollary 2.5 to prove Proposition 2.2. We note that this proposition can be easily deduced from [3, Lemma 3.2 and Theorem 3.5] (which also proves some similar results for S^1), but we give a self-contained proof here for completeness.

Proof of Proposition 2.2. We will repeatedly use the following elementary fact.

FACT 2.7. $\text{Fix}(hgh^{-1}) = h \text{Fix}(g)$. In particular, if g and h commute, then $\text{Fix}(g)$ is h -invariant.

Let G be a non-abelian subgroup of $\text{Diff}_+^r(I)$ with $r \geq 2$. Let H be the centralizer of G in $\text{Diff}_+^r(I)$, and assume that H is non-abelian.

First, note that each element $g \in G$ must have a fixed point in $(0, 1)$, for if g is fixed-point-free, then Corollary 2.5 says that $C(g)$ is abelian, but H (which we assumed to be non-abelian) is a subset of $C(g)$.

We claim that in fact every endpoint of each component of $I \setminus \text{Fix}(g)$ is fixed by each $h \in H$. To see this, let $h \in H$. If some point $x \in \text{Fix}(g)$ is not fixed by h , then it lies in some connected component J of $I \setminus \text{Fix}(h)$. Since g and h commute, $\text{Fix}(h)$ is g -invariant. In particular, g permutes the connected components of $I \setminus \text{Fix}(h)$. Since $x \in J$ is fixed by g , the interval J must be g -invariant. Apply the Hölder–Kopell lemma to $g|_J$. Since h has no fixed points in J , but g does, we conclude that $g|_J = \text{id}$. Thus, the only fixed points of g not necessarily fixed by h are interior points of $\text{Fix}(g)$.

Since the endpoints of each component of $I \setminus \text{Fix}(g)$ are fixed by each $h \in H$, each component of $I \setminus \text{Fix}(g)$ is H -invariant. Consider again a particular connected component J of $I \setminus \text{Fix}(g)$. Kopell's lemma implies that $H|_J$ is abelian. Thus, if we let $\text{supp}(g)$ denote the closure of $I \setminus \text{Fix}(g)$, we know that H is abelian when restricted to $\bigcup_{g \in G} \text{supp}(g)$, and hence on the closure of $\bigcup_{g \in G} \text{supp}(g)$. It follows that either there is some open set fixed by each $g \in G$ or H is abelian everywhere.

The proof for $G \subset \text{Diff}_c^r(\mathbb{R})$ follows quickly from the following observation.

Observation 2.8. There is an embedding $\iota : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}^r(I)$ whose image is the set of diffeomorphisms of I supported away from the endpoints of I . In particular, for $x \in \text{int}(I)$ we may define $\iota(g)(x)$ by $fg(f^{-1}(x))$, where $f : \mathbb{R} \rightarrow \text{int}(I)$ is a C^∞ -diffeomorphism.

Hence, if $G \subset \text{Diff}_c^r(\mathbb{R})$ has non-abelian centralizer, then so does $\iota(G)$ as in Observation 2.8. Our proof above shows that $\text{Fix}(\iota(G))$ contains an open interval, hence $\text{Fix}(G)$ does as well. \square

A schematic picture of what fixed sets of non-abelian commuting G and H may look like is given in Figure 1. In particular, we note the following corollary of the proof.




FIGURE 1. Fix sets of G and H indicated above and below the line. If H has a fixed point in J and is not abelian on J , then $J \subset \text{Fix}(H)$.

COROLLARY 2.9. *Let $G \subset \text{Diff}_+^r(I)$ be as in Proposition 2.2, and let H be the centralizer of G . Then G and H have a common fixed point in $\text{int}(I)$. Similarly, if $G \subset \text{Diff}_c^r(\mathbb{R})$ and H is the centralizer of G , then G and H have a common fixed point.*

Proof. We know that $\text{Fix}(G)$ contains a non-trivial interval, and the proof of Proposition 2.2 implies that each endpoint of this interval is fixed by all $h \in H$. \square

As another immediate consequence of Proposition 2.2, we have the following corollary.

COROLLARY 2.10. *Let $r \geq 2$ and let G and H be commuting, non-abelian subgroups of $\text{Diff}_+^r(I)$ or $\text{Diff}_c^r(\mathbb{R})$. If H is perfect, then H acts trivially on the complement of $\text{Fix}(G)$.*

Proof. We showed above that for any $g \in G$, the restriction of its centralizer to $\mathbb{R} \setminus \text{Fix}(g)$ is abelian. Since H is contained in the centralizer, its restriction to $\mathbb{R} \setminus \text{Fix}(g)$ is abelian as well. If H is perfect, this restriction must be trivial. \square

Since the groups $\text{Diff}_c^r(M)$ and $\text{Diff}^r(M)_0$ are perfect as long as $r \neq \dim(M) + 1$, we will be able to apply Corollary 2.10 in the following.

We would like to have an analog of Proposition 2.2 for diffeomorphisms of the circle. To do this, we develop a version of the Hölder–Kopell lemma for S^1 .

2.2. Circle diffeomorphisms. A major tool for studying homeomorphisms (or diffeomorphisms) of S^1 is the *rotation number*. This is an \mathbb{R}/\mathbb{Z} -valued number, well defined for each $g \in \text{Homeo}_+(S^1)$. If we consider S^1 as \mathbb{R}/\mathbb{Z} , the rotation number of an element g is given by $\lim_{n \rightarrow \infty} g^n(0)/n$, taken mod \mathbb{Z} . A good exposition of the basic theory can be found in [14]. We need only three facts here.

- (1) A homeomorphism has rotation number 0 if and only if it has a fixed point.
- (2) Homeomorphisms with rational rotation number of the form m/k are precisely those with periodic points of period k .
- (3) (Denjoy’s theorem) If g is a C^2 -diffeomorphism of the circle with irrational rotation number θ , then g is topologically conjugate to rotation by θ .

Using fact (2), we see that if g is a diffeomorphism with rational rotation number, then some iterate g^k has a fixed point, hence can be viewed as a diffeomorphism of the interval. This puts us in a situation where Kopell’s lemma applies. If instead g has irrational rotation number, fact (3) tells us that g is conjugate to an irrational rotation. Since any homeomorphism that commutes with an irrational rotation must also be a rotation (see [14]), we conclude the following corollary.

COROLLARY 2.11. *Let $g \in \text{Diff}_+^2(S^1)$. Then there is either some $k \in \mathbb{N}$ such that g^k has a fixed point, or the centralizer of g in $\text{Diff}_+^2(S^1)$ is abelian.*

This is the analog of the Hölder–Kopell lemma for S^1 . Now we state and prove our analog of Proposition 2.2.

PROPOSITION 2.12. *Let $r \geq 2$ and let $G \subset \text{Diff}_+^r(S^1)$ be non-abelian. Let H be contained in the centralizer of G . Then either:*

- (a) *(H is abelian after powers) for each pair of elements $h_1, h_2 \in H$, there are integers k_1 and k_2 such that $h_1^{k_1}$ and $h_2^{k_2}$ commute; or*
- (b) *(G fixes an interval after powers) there is a subinterval $J \subset S^1$ and for each $g \in G$ an integer k_g such that J is fixed pointwise by g^{k_g} .*

Proof. Let G and H be as in the statement of Proposition 2.12. Let $g \in G$. Since H is non-abelian and centralizes g , Corollary 2.11 implies that g has rational rotation number. Thus, after replacing g with g^{k_g} if necessary we can assume that g has a fixed point. We may also assume that g was chosen so that $g^{k_g} \neq 1$, for if every element of G has finite order, then case (b) holds. (In fact, in this case one can use Hölder’s theorem to show that G is abelian.)

Now let h_1 and h_2 be any two elements of H . Since H and G commute, Corollary 2.11 implies again that h_1 and h_2 have rational rotation numbers, so there are integers k_1 and k_2 such that $h_1^{k_1}$ and $h_2^{k_2}$ have fixed points. The same argument as in Proposition 2.2 shows that endpoints of components of $S^1 \setminus \text{Fix}(g^{k_g})$ are fixed by each $h_i^{k_i}$. Cutting S^1 at any one of the points of $\text{Fix}(g^{k_g})$ reduces the action of g^{k_g} , $h_1^{k_1}$ and $h_2^{k_2}$ to an action on the interval, and the proof of Proposition 2.2 again shows that $h_1^{k_1}$ and $h_2^{k_2}$ must commute when restricted to $S^1 \setminus \text{Fix}(g^{k_g})$.

Thus, if $h_1^{k_1}$ and $h_2^{k_2}$ do not commute everywhere, then there is some open set $J \subset S^1$ that is fixed pointwise by some iterate g^{k_g} for every $g \in G$. This is exactly what we wanted to prove. \square

Remark 2.13. Note that the collection of elements g^{k_g} that fix J in case (b) of Proposition 2.12 are precisely the elements of G with rotation number zero. In general, a collection of elements of $\text{Diff}_+^2(S^1)$ that all have rotation number zero need not be a group, since the composition of two diffeomorphisms with fixed points need not have a fixed point. But in this case the set of elements with rotation number zero is equal to the subgroup of diffeomorphisms that fix J pointwise.

We remark also that Proposition 2.12 fails without the ‘after powers’ hypothesis—there exist examples of two non-abelian, commuting subgroups of S^1 , neither of which fixes any interval pointwise. However, it would be interesting to know whether the ‘after powers’ condition could be replaced by ‘virtually’ (i.e. after taking a finite index subgroup).

3. Flows and continuity

Example 1.7 in the introduction shows that not all \mathbb{R} -actions on a manifold are continuous. Our goal in this section is to see (roughly) that whenever M_1 and M_2 are 1-manifolds, and $\text{Diff}_c^r(M_1)$ includes in $\text{Diff}_c^r(M_2)$, then \mathbb{R} -subgroups of $\text{Diff}_c^r(M_1)$ do act continuously—at least on part of M_2 .

Definition 3.1. By a *flow* on a manifold M , we mean a continuous family of C^r -diffeomorphisms $\phi(t, x) : \mathbb{R} \times I \rightarrow I$; in other words, an \mathbb{R} -subgroup of $\text{Diff}^r(M)$

or $\text{Diff}_c^r(M)$. We denote $\phi(t, x)$ by $\phi^t(x)$, and by $\{\phi^t\}$ we mean the full \mathbb{R} -subgroup $\{\phi^t \in G : t \in \mathbb{R}\}$.

We will use the tools we developed in §2, as well as the following further result in the spirit of Hölder–Kopell theory. Keeping our notation from §2, let $I := [0, 1)$. We will also use $C_G(H)$ to denote the centralizer of a subgroup H in a group G and $C_G(g)$ for the centralizer of $g \in G$.

THEOREM 3.2. (Szekeres, see [14, Theorem 4.1.11]) *Let $r \geq 2$ and let $\psi \in \text{Diff}_+^r(I)$ be a diffeomorphism with no interior fixed points. Then there is a unique flow ψ^t of class C^{r-1} on $(0, 1)$ and C^1 on I such that $\psi = \psi^1$ and the centralizer of ψ in $\text{Diff}_+^1(I)$ is $\{\psi^t\}$.*

Compare this theorem with the Hölder–Kopell lemma (Corollary 2.5). Under the same hypotheses, the Hölder–Kopell lemma concludes that the centralizer of ψ in $\text{Diff}_+^r(I)$ is abelian. Using Szekeres, we can conclude that $C_{\text{Diff}_+^r(I)}(\psi)$ is a subgroup of a C^1 flow on I .

3.1. Flows on the interval. Though our ultimate goal is a theorem about $\text{Diff}_+^r(S^1)$ and $\text{Diff}_c^r(\mathbb{R})$, we first prove a result about flows in $\text{Diff}_+^r(I)$. This allows us to use Szekeres’s theorem in a straightforward way, and to make the simplifying assumption that our flows have full support.

PROPOSITION 3.3. *Let $p, r \geq 2$ and let $\Phi : \text{Diff}_+^p(I) \rightarrow \text{Diff}_+^p(I)$ be an injective group homomorphism. Let $\{\phi^t\} \subset \text{Diff}_+^r(I)$ be a flow that is fixed-point-free on $(0, 1)$, and suppose that there is some s such that $\Phi(\phi^s)$ is fixed-point-free on $(0, 1)$. Then $\Phi(\{\phi^t\})$ is continuous in t . Moreover, there is a flow $\{\psi^t\}$ such that $\Phi(\phi^t) = \psi^{\lambda t}$ for some $\lambda \in \mathbb{R}$.*

The proof of Proposition 3.3 has several steps. We use Szekeres to find a candidate flow $\{\psi^t\}$ and reduce the question to continuity of a group homomorphism $\mathbb{R} \rightarrow \mathbb{R}$. We next establish a simple criterion for continuity, and then relate this criterion to fixed sets and commutators and apply our work from §2. Our main results of this section (Propositions 3.8 and Corollary 3.12) are essentially built from this proof with the assumption of full supports removed.

Proof of Proposition 3.3. To simplify notation let $G = \text{Diff}_+^p(I)$. Suppose that $\Phi : G \rightarrow G$ is injective. Let $\{\phi^t\} \subset G$ be a flow with no fixed points in $(0, 1)$ and assume that $\psi := \Phi(\phi^s)$ has no fixed points in $(0, 1)$. Since $\{\phi^t\}$ is contained in $C_G(\phi^1)$, its image $\{\Phi(\phi^t) : t \in \mathbb{R}\}$ is contained in $C_{\Phi(G)}(\psi)$. By Szekeres’s theorem, there is a C^1 flow $\{\psi^t\}$ on I such that $\{\Phi(\phi^t)\} \subset \{\psi^t\}$. Moreover, that Φ is a group homomorphism implies that $\Phi(\phi^t) = \psi^{\alpha(t)}$ for some additive group homomorphism $\alpha : \mathbb{R} \rightarrow \mathbb{R}$. Thus, we now only need to show that α is continuous.

We can detect continuity in a very simple way using intersections of open sets. Essentially, α being continuous just means that if t is small enough then $\phi^{\alpha(t)}$ will not push an open set off itself. Formally, we have the following lemma.

LEMMA 3.4. (Continuity criterion) *Let $\{\psi^t\}$ be a flow on $(0, 1)$ such that ψ^t is fixed-point-free for some (hence all) $t \neq 0$. Let $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ be a group homomorphism, and*

let $U \subset (0, 1)$ be an open set that is bounded away from 0 and 1. Then α is continuous if and only if there is some $\delta > 0$ such that $(\psi^{\alpha(t)}(U)) \cap U \neq \emptyset$ for all $t < \delta$.

Proof. One direction is almost immediate: if α is a continuous $\mathbb{R} \rightarrow \mathbb{R}$ homomorphism, then $\alpha(t) = \lambda t$ for some $\lambda \in \mathbb{R}$ (easy exercise). Given δ sufficiently small and U open, for any $t < \delta$ we have $\psi^{\lambda t}(U) \cap U \neq \emptyset$.

For the converse, suppose that α is not continuous. In this case, using the fact that α is an additive group homomorphism, it is elementary to show that for any $T > 0$ there are arbitrarily small $t \in \mathbb{R}$ such that $|\alpha(t)| > T$. Suppose that U is bounded away from 0 and 1. Since ψ^t has no fixed points in $(0, 1)$, for T large enough, $\psi^s(U)$ will either be contained in a small neighborhood of 0 or a small neighborhood of 1 whenever $|s| > T$. We can ensure that this neighborhood is disjoint from U by taking T large. In particular, when $|\alpha(t)| > T$ we will have $\psi^{\alpha(t)}(U) \cap U = \emptyset$. □

The next step in the proof of Proposition 3.3 is to detect intersections of sets algebraically (i.e. through the algebraic structure of the diffeomorphism group), using the results of §2. The following notation will be useful.

Notation 3.5. For a group G of diffeomorphisms of a manifold M and a subset $U \subset M$, let

$$G^U := \{g \in G : g(x) = x \text{ for all } x \in U\}.$$

The link between set intersections, subgroups, and centralizers comes from an elementary corollary of Proposition 2.2.

COROLLARY 3.6. *Let $U \subset I$ and $V \subset I$ be closed subsets properly contained in I . The group $\langle G^U, G^V \rangle$ generated by G^U and G^V has non-abelian centralizer if and only if $U \cap V$ contains an open set.*

Proof. The condition that U and V are closed and properly contained in I implies that G^U and G^V are non-trivial and in this case non-abelian. If $\langle G^U, G^V \rangle$ has non-abelian centralizer, then $\text{Fix}(\langle G^U, G^V \rangle)$ contains an open subset of I by Proposition 2.2. This subset must be fixed pointwise by both G^U and G^V , so is a subset of $U \cap V$.

Conversely, if $U \cap V$ contains an open set J , then the group of all diffeomorphisms supported on J commutes with G^U and G^V , so commutes with $\langle G^U, G^V \rangle$. □

Now we can begin the real work of the proof of Proposition 3.3, using our continuity criterion. Let $U \subset I$ be an open set bounded away from 0 and 1. Since ϕ^t is continuous in t , there is some $\delta > 0$ such that for all $t < \delta$ we have $\phi^t(U) \cap U \neq \emptyset$. By Corollary 3.6, this is equivalent to the statement that $\langle G^U, G^{\phi^t(U)} \rangle$ has non-abelian centralizer. We want to translate this into a statement about $\Phi(G^U)$ and $\psi^{\alpha(t)}$.

To do this, note first that

$$\begin{aligned} G^{\phi^t(U)} &:= \{g \in G : g(x) = x \text{ for all } x \in \phi^t(U)\} \\ &= \{\phi^t g \phi^{-t} : g(x) = x \text{ for all } x \in U\} \\ &= \phi^t(G^U)\phi^{-t}, \end{aligned} \tag{1}$$

and more generally, for any group of diffeomorphisms H and any diffeomorphism g , we have

$$g \text{Fix}(H) = \text{Fix}(gHg^{-1}). \tag{2}$$




FIGURE 2. Continuity of α detected by intersections of open sets.

Now consider $\Phi(G^U)$. Since G^U has non-abelian centralizer, $\Phi(G^U)$ has non-abelian centralizer as well. By Proposition 2.2, $\text{Fix } \Phi(G^U)$ contains an open set. Using (1), we have

$$\Phi(G^{\phi^t(U)}) = \Phi(\phi^t)\Phi(G^U)\Phi(\phi^{-t}) = \psi^{\alpha(t)}(\Phi(G^U))\psi^{-\alpha(t)},$$

and, letting U' denote $\text{Fix } \Phi(G^U)$, it follows from (2) that

$$\text{Fix}(\Phi(G^{\phi^t(U)})) = \psi^{\alpha(t)}(U'). \tag{3}$$

For any $t < \delta$, consider the subgroup

$$H_t := \langle \Phi(G^U), \Phi(G^{\phi^t(U)}) \rangle = \langle \Phi(G^U), \psi^{\alpha(t)}(\Phi(G^U))\psi^{-\alpha(t)} \rangle.$$

Using (3), we have

$$\text{Fix}(H_t) = \text{Fix}(\Phi(G^U)) \cap \text{Fix}(\psi^{\alpha(t)}(\Phi(G^U))\psi^{-\alpha(t)}) = U' \cap \psi^{\alpha(t)}(U').$$

H_t also has non-abelian centralizer, so by Proposition 2.2, $\text{Fix}(H_t) = U' \cap \psi^{\alpha(t)}(U')$ contains an open set. If $V = \text{int}(U')$, we know that $V \cap \psi^{\alpha(t)}(V)$ contains an open set as well. (A picture of the situation so far is given in Figure 3.) If we knew further that V was bounded away from 0 and 1, then we could apply Lemma 3.4 to V and conclude that α is continuous.

This would finish our proof. Thus, it remains only to show that V is bounded away from 0 and 1. This is not hard to see, but uses a well-known theorem about diffeomorphism groups called the *fragmentation property*.

THEOREM 3.7. (Fragmentation property) *Let M be a manifold and \mathcal{B} an open cover of M . Then the group $\text{Diff}^r(M)_0$ is generated by the set $\{g \in \text{Diff}^r(M)_0 : \text{supp}(g) \in B \text{ for some } B \in \mathcal{B}\}$.*

Recall that $\text{supp}(g)$ is the *support* of g , the closure of $M \setminus \text{Fix}(g)$. A proof of Theorem 3.7 can be found in [2, Ch. 2].

Since $\{\phi^t\}$ is a flow, for large enough s we have $\phi^s(\bar{U}) \cap \bar{U} = \emptyset$. It follows from the fragmentation property that G^U and $G^{\phi^s(U)}$ together generate G , so $\Phi(G^U)$ and $\Phi(G^{\phi^s(U)})$ generate $\Phi(G)$. In particular, any point in I that is fixed by both $\Phi(G^U)$ and $\Phi(G^{\phi^s(U)})$ is fixed by every element of $\Phi(G)$.

To show now that $\text{int}(U')$ is bounded away from 0 and 1, assume for contradiction that $U' := \text{Fix}(\Phi(G^U))$ contains a neighborhood of 0 (the case where it contains a

neighborhood of 1 is identical). Then $\psi^{\alpha(s)}(U')$ contains a neighborhood of 0 as well, but $\psi^{\alpha(s)}(U') = \text{Fix}(\Phi(G^{\phi^s(U)}))$. By the previous paragraph, we conclude that $\Phi(G)$ pointwise fixes a neighborhood of 0, contradicting the fact that ψ had no fixed points in $(0, 1)$. This concludes the proof of Proposition 3.3. \square

Our next goal is to remove the assumption that our flows had no fixed points and work with \mathbb{R} instead of I . There is some technical work to do here to keep track of supports, but the key ideas are really contained in the proof of Proposition 3.3.

PROPOSITION 3.8. *Let $r \geq 3, p \geq 2$ and let $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_c^p(\mathbb{R})$ be a homomorphism. For any $z \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that any flow $\{\phi^t\} \subset \text{Diff}_c^r(\mathbb{R})$ without fixed points in $(-n, n)$ has $\Phi(\{\phi_t\})$ continuous in t on some neighborhood containing z .*

Remark 3.9. It will follow from Theorem 1.5 that for any flow $\{\phi^t\}$, the image $\Phi(\{\phi^t\})$ will be continuous in t everywhere, but we will not see this until we finish the whole proof of the theorem.

Proof of Proposition 3.8. Let G denote $\text{Diff}_c^r(\mathbb{R})$. We start by taking some steps to make the set-up in this situation as close as possible to the hypotheses of Proposition 3.3.

First, we may assume without loss of generality that $\text{Fix}(\Phi(G)) = \emptyset$. This is because each connected component C of $\mathbb{R} \setminus \text{Fix}(\Phi(G))$ is $\Phi(G)$ -invariant, so we can consider the restriction of elements in the image of Φ to C , giving a homomorphism $\text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_c^p(C) \cong \text{Diff}_c^p(\mathbb{R})$. Since G is perfect, this homomorphism is still injective.

Let $\{\phi^t\} \subset G$ be a flow, and consider a connected component of $\mathbb{R} \setminus \text{Fix}(\{\phi^t\})$. Identify this interval with $(0, 1)$ and consider its half-closure to be $[0, 1) = I$. As in the proof of Proposition 3.3, let $U \subset I$ be an open interval bounded away from 0 and 1.

Choose any $s \neq 0$ and let $\psi = \Phi(\phi^s)$. Since ψ is compactly supported, it has non-empty fixed set in \mathbb{R} . Let J be a connected component of $\mathbb{R} \setminus \text{Fix}(\psi)$. Since ϕ^t and ϕ^s commute, J is $\{\phi^t\}$ -invariant and we can apply Szekeres’s theorem to \bar{J} and conclude as before that there is some flow $\{\psi^t\}$ on J and an additive group homomorphism $\alpha_J : \mathbb{R} \rightarrow \mathbb{R}$ such that $\Phi(\phi_t)|_J = \psi^{\alpha_J(t)}$. If $\text{Fix}(\Phi(G^U)) \cap J$ contains an open set, then the argument from Proposition 3.3 applies verbatim to show that α_J is continuous, and so $\Phi(\{\phi^t\})$ is continuous on J .

Thus, the next lemma will finish the proof of Proposition 3.8.

LEMMA 3.10. *Let $G = \text{Diff}_c^r(\mathbb{R})$ and suppose that $\Phi : G \rightarrow \text{Diff}_c^p(\mathbb{R})$ satisfies $\text{Fix}(\Phi(G)) = \emptyset$. Then:*

- (a) *for any $z \in \mathbb{R}$, there is a bounded set $U \subset \mathbb{R}$ such that z is in the interior of $\text{Fix}(\Phi(G^U))$; and*
- (b) *any flow $\{\phi^t\} \subset G$ with sufficiently large support has $z \notin \text{Fix}(\Phi(\phi^s))$ for some s .*

Given this lemma, we know that any flow with sufficiently large support will have a connected component J of $\mathbb{R} \setminus \text{Fix}(\Phi(\phi^s))$ with z contained inside an open set in $\text{Fix}(\Phi(G^U)) \cap J$, so the argument we just gave shows that $\Phi(\{\phi^t\})$ is continuous on J .

Proof of Lemma 3.10. Let $z \in \mathbb{R}$. Suppose for contradiction that there is no bounded, open interval U such that $z \in \text{int}(\text{Fix}(\Phi(G^U)))$. Let $G_U := \{g \in G : g(x) = x \text{ for all } x \notin U\}$.




FIGURE 3. If $\phi^s(\bar{U}) \cap \bar{U} = \emptyset$, then z is not fixed by $\Phi(\phi^s)$.

By the fragmentation property (Theorem 3.7), the subgroups G_U generate G as U ranges over open intervals of \mathbb{R} .

Now for any open set U , the subgroup G_U commutes with G^U . Also, G_U is perfect and G^U non-abelian, so by Corollary 2.10, $\mathbb{R} \setminus \text{Fix}(\Phi(G^U)) \subset \text{Fix}(\Phi(G_U))$. It follows that $z \in \text{Fix}(\Phi(G_U))$ for every U . Since the $\Phi(G_U)$ generate $\Phi(G)$, we have $z \in \text{Fix}(\Phi(G))$, a contradiction. This proves (a).

To prove (b), take a bounded set U such that z is in the interior of $\text{Fix}(\Phi(G^U))$. We claim that any flow $\{\phi^t\} \subset G$ with support containing U satisfies $z \notin \text{Fix}(\Phi(\phi^s))$ for some s . To show this, suppose that $\{\phi^t\}$ is such a flow. We know that there is some s such that $\phi^s(\bar{U}) \cap \bar{U} = \emptyset$, so ϕ^s and G^U generate G (using fragmentation again). In particular, if $z \in \text{Fix}(\Phi(G^U)) \cap \text{Fix}(\Phi(\phi^s))$, then $z \in \text{Fix}(\Phi(G))$, which we assumed to be empty. So $z \notin \text{Fix}(\Phi(\phi^s))$ and this is what we wanted to show.

This completes the proof of the lemma and the proof of Proposition 3.8. □

To extend Proposition 3.8 to homomorphisms $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_+^p(S^1)$, we need a lemma. This is the diffeomorphism group analog to the fact that any injection $\mathbb{R} \hookrightarrow S^1$ must miss a point.

LEMMA 3.11. *Let $r, p \geq 2$ and let $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_+^p(S^1)$ be a homomorphism. Then $\text{Fix}(\Phi(\text{Diff}_c^r(\mathbb{R}))) \neq \emptyset$.*

Proof. Let G_n be the group of diffeomorphisms with support contained in $(-n, n)$. Then $G_n \subset G_{n+1}$ and

$$G = \bigcup_{n \in \mathbb{N}} G_n.$$

It follows that $\Phi(G_n) \subset \Phi(G_{n+1})$, so

$$\text{Fix}(\Phi(G_n)) \supset \text{Fix}(\Phi(G_{n+1})). \tag{4}$$

Moreover, since

$$\Phi(G) = \bigcup_{n \in \mathbb{N}} \Phi(G_n)$$

we have also

$$\text{Fix}(\Phi(G)) = \bigcap_{n \in \mathbb{N}} \text{Fix}(\Phi(G_n)). \tag{5}$$

Now each G_n commutes with the subgroup H_n of diffeomorphisms fixing $[-n, n]$ pointwise and the subgroups $\Phi(G_n)$ and $\Phi(H_n)$ satisfy the hypotheses of Proposition 2.12. In particular, if we let G'_n be the group of elements of $\Phi(G_n)$ with rotation number zero, this is a non-trivial *subgroup* (by Remark 2.13) so by simplicity of G_n , we have $G'_n = G_n$.

It now follows from Proposition 2.12 that $\text{Fix}(\Phi(G_n)) \neq \emptyset$. Since fixed sets are closed, it follows from (4) and (5) that $\text{Fix}(\Phi(G)) \neq \emptyset$. \square

The following corollary of Proposition 3.8 is now immediate.

COROLLARY 3.12. *Let $r \geq 3$, $p \geq 2$, and suppose that $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_+^p(S^1)$ is a homomorphism. For any $z \in S^1 \setminus \text{Fix}(\Phi(G))$, there is an $n \in \mathbb{N}$ such that if $\{\phi^t\} \subset \text{Diff}_c^r(\mathbb{R})$ is a flow without fixed points in $(-n, n)$, then $\Phi(\{\phi_t\})$ is continuous in t on a neighborhood containing z .*

Proof. By Lemma 3.11, there is some point $x \in \text{Fix}(\Phi(\text{Diff}_c^r(\mathbb{R})))$. Moreover, the derivative of every element $\Phi(g) \in \Phi(\text{Diff}_c^r(\mathbb{R}))$ at x is zero, for $g \mapsto D\Phi(g)(x)$ is a homomorphism to \mathbb{R} , and since $\text{Diff}_c^r(\mathbb{R})$ is perfect, it must be trivial. The same applies to higher order derivatives. Cutting S^1 at x , we can consider Φ to be a homomorphism into the subgroup of diffeomorphisms in $\text{Diff}_+^p(S^1 \setminus \{x\})$ where all derivatives vanish at the endpoints. This is isomorphic to a subgroup of $\text{Diff}_c^p(\mathbb{R})$, and Proposition 3.8 now applies. \square

A similar conclusion holds for homomorphisms $\Phi : \text{Diff}_+^r(S^1) \rightarrow \text{Diff}_+^p(S^1)$. Since we will not use this in the following, we state it informally and only sketch the proof, leaving the details to the reader.

COROLLARY 3.13. *Let $r \geq 3$, and suppose that $\Phi : \text{Diff}_+^r(S^1) \rightarrow \text{Diff}_+^p(S^1)$ is a homomorphism. Then for any $z \in S^1$, any flow $\{\phi^t\} \subset \text{Diff}_+^r(S^1)$ with sufficiently large support has $\Phi(\{\phi^t\})$ continuous at z .*

Proof sketch. There are two classes of \mathbb{R} -subgroups of $\text{Diff}_+(S^1)$: those with non-empty fix set and those conjugate to the rotation subgroup. If $\text{Fix}(\{\phi^t\}) \neq \emptyset$, we can cut S^1 at a point $x \in \text{Fix}(\{\phi^t\})$ and apply Corollary 3.12 with the group of diffeomorphisms of $S^1 \setminus \{x\} \cong \mathbb{R}$ as the domain. Otherwise, $\{\phi^t\}$ is conjugate to the rotation subgroup, and we can use torsion elements and Hölder–Kopell theory to show that $\Phi(\{\phi^t\})$ is conjugate to the rotation subgroup as well. Continuity in t can be demonstrated for small t , since for small t we can choose x and $y \in S^1$ and write rotation by t as $f^t g^t$, where f^t is a flow supported on $S^1 \setminus \{x\}$ and g^t a flow supported on $S^1 \setminus \{y\}$. We may also choose f^t and g^t to have arbitrarily large supports. Corollary 3.12 then implies that $\Phi(\{f^t\})$ and $\Phi(\{g^t\})$ are continuous at every point, hence so is $\Phi(\{\phi^t\})$ —at least for small t . But since $\Phi(\{\phi^t\})$ is an \mathbb{R} -subgroup, it must be continuous for all t . \square

4. Proof of Theorem 1.5

We start by proving Theorem 1.5 for diffeomorphisms of \mathbb{R} . Namely, we show the following.

THEOREM 4.1. *Let $r \geq 3$, $p \geq 2$, and let $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_c^p(\mathbb{R})$ be non-trivial. Then Φ is topologically diagonal. If $r = s$, then Φ is C^r -topologically diagonal.*

The proof of this theorem contains the essence of the proof of Theorem 1.5. The only difference in the statement of Theorem 1.5 is that $\text{Diff}_c^r(\mathbb{R})$ and/or $\text{Diff}_c^p(\mathbb{R})$ can be replaced with $\text{Diff}_+^r(S^1)$ (and/or $\text{Diff}_+^p(S^1)$). We deal with this easily at the end of this section.

Proof of Theorem 4.1. Let G denote the group $\text{Diff}_c^r(\mathbb{R})$ and, as before, for a subset $U \subset \mathbb{R}$ let G^U denote the subgroup $\{g \in G : gx = x \text{ for all } x \in U\}$.

Let $\Phi : G \rightarrow G$ be a non-trivial homomorphism. Assume first for simplicity that $\text{Fix}(\Phi(G)) = \emptyset$. In this case, we need only construct one map $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\Phi(g) = fgf^{-1}$ for all $g \in G$. To construct f , we use Proposition 3.8 (continuity of flows) along with Proposition 2.2 to show that particular subgroups of point stabilizers in G map under Φ to subgroups of point stabilizers. This lets us build a point-to-point $\mathbb{R} \rightarrow \mathbb{R}$ map out of the data of Φ .

For $x \in \text{int}(I)$, let

$$G^x := \langle G^{(-\infty, x]}, G^{[x, \infty)} \rangle.$$

Since $G^{(-\infty, x]}$ and $G^{[x, \infty)}$ commute, we have

$$G^x \cong G^{(-\infty, x]} \times G^{[x, \infty)}. \tag{6}$$

Note that the set G^x is contained in, but not equal to, the point stabilizer G_x —we will need to use the fact that G^x decomposes as a direct product so the full point stabilizer will not work. However, the fragmentation property still implies that

$$\text{if } g(x) \neq x \text{ then } \langle G^x, g \rangle = G. \tag{7}$$

Now let $A_x := \text{Fix}(\Phi(G^x))$. The product structure of G^x in (6) means that

$$\Phi(G^x) = \Phi(G^{(-\infty, x]}) \times \Phi(G^{[x, \infty)}).$$

Corollary 2.9 shows that two non-abelian, commuting subgroups must have a common fixed point. In particular, $\Phi(G^{(-\infty, x]})$ and $\Phi(G^{[x, \infty)})$ have a common fixed point, so $A_x = \text{Fix}(\Phi(G^x)) \neq \emptyset$. Eventually we will see that A_x consists of a *single* fixed point and $f(x) = A_x$ will be our map.

LEMMA 4.2. (Easy properties of A_x) *Let $\Phi : G \rightarrow G$ be a homomorphism and assume that $\text{Fix}(\Phi(G)) = \emptyset$. Let $A_x = \text{Fix}(\Phi(G^x))$ as above. Then:*

- (1) $A_x \neq \emptyset$;
- (2) $A_x \cap A_y = \emptyset$ whenever $x \neq y$;
- (3) if $g \in G$ and $g(x) = y$, then $\Phi(g)(A_x) = A_y$.

Proof. (1) follows from our discussion above. To prove (2), note that, by the fragmentation property (Theorem 3.7), if $x \neq y$ then G^x and G^y together generate G . Now if some point $z \in \mathbb{R}$ is fixed by both $\Phi(G^x)$ and $\Phi(G^y)$, then z is fixed by every element of $\Phi(G) = \langle \Phi(G^x), \Phi(G^y) \rangle$, contradicting our assumption that $\text{Fix}(\Phi(G)) = \emptyset$. This shows that $A_x \cap A_y = \emptyset$.

The proof of (3) is elementary: if g is a diffeomorphism with $g(x) = y$, then

$$G^y = gG^xg^{-1},$$

and so

$$A_y = \text{Fix}(\Phi(gG^xg^{-1})) = \Phi(g)(\text{Fix}(G^x)) = \Phi(g)(A_x). \quad \square$$




FIGURE 4. The definition of $f : \mathbb{R} \rightarrow \mathbb{R}$.

Using Lemma 4.2, we now show that A_x consists of isolated points. Let $z \in A_x$ and let $\{\phi^t\} \subset G$ be a flow such that $\phi^t(x) \neq x$ for $t \neq 0$, and such that $\Phi(\phi^t)$ is continuous in t at z . Such a flow exists by Proposition 3.8; in fact any flow ϕ^t with large enough support will work. Let J be the connected component of $\mathbb{R} \setminus \text{Fix}(\Phi(\phi^t))$ containing z . Proposition 3.8 says that there is a flow ψ^t on J such that $\Phi(\phi^t)|_J = \psi^{\lambda t}$. If $\psi^{\lambda t}(z) = z$ for $t \neq 0$, then z is fixed by both $\psi^{\lambda t}$ and $\Phi(G^x)$. It follows from (7) above that z is then fixed by all of $\Phi(G)$. This contradicts our assumption. Thus, z is not fixed by any such flow.

This shows that z must be an isolated point of A_x ; if not, there is some $z' \in A_x \cap J$ and some small $t \neq 0$ such that $\psi^{\lambda t}(z) = z'$. But property (3) of Lemma 4.2 says that $z' \in A_{\phi^t(x)}$, and property (1) says that $A_{\phi^t(x)} \cap A_x = \emptyset$.

We now build a continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$. For $y \in \mathbb{R}$ define $f(y)$ as follows. Take any flow $\{\phi^t\} \subset G$ such that $\phi^s(x) = y$ for some s and such that $\Phi(\phi^t)$ is continuous in t at z . Then set $f(y) = \Phi(\phi^s)(z)$. See Figure 4. We will show that f is:

- (a) well defined;
- (b) continuous and injective;
- (c) G -equivariant, meaning that $fg(y) = \Phi(g)(f(y))$ for all $g \in G$ and $y \in \mathbb{R}$;
- (d) surjective;
- (e) C^r in the case where $r = p$.

These will all follow from the continuity of flows at z and the easy properties of the sets A_x . In particular, we will use the fact that for any flow $\{\phi^t\}$ used in the definition of $f(y)$, we have

$$\Phi(\phi^s)(z) \in \Phi(\phi^s)(A_x) = A_y.$$

Well defined. Let $\{\phi^t\}$ and $\{\eta^t\}$ be two flows such that $\phi^s(x) = \eta^r(x) = y$ and such that both $\Phi(\phi^t)$ and $\Phi(\eta^t)$ are continuous in t at z . Suppose for contradiction that $\Phi(\phi^s)(z) \neq \Phi(\eta^r)(z)$. Then $\Phi(\phi^s)(z) = \Phi(\eta^q)(z)$ for some $q \neq r$. But then $\eta^q(x) = y'$ for some $y' \neq y$. It follows that $\Phi(\eta^q)(z) \in A_{y'} \cap A_y$, a contradiction.

Continuity and injectivity. Continuity is immediate from the definition of f since $\Phi(\phi^t)$ is continuous on the connected component of its support that contains z . Injectivity follows from continuity and the fact that $A_{y'} \cap A_y = \emptyset$ for $y \neq y'$.

G-equivariance. By construction, the image of f contains exactly one point of A_y for each $y \in \mathbb{R}$. Given $y \in \mathbb{R}$ and $g \in G$, let $\{\phi^t\}$ be a flow such that $\phi^1(x) = y$ and $\phi^s(x) = g(y)$ for some s . We may also choose $\{\phi^t\}$ to have as large support as we want, so as to ensure it is continuous at z . Then

$$f(y) = \Phi(\phi^1)(z)$$

and

$$\Phi(g)f(y) = \Phi(g)\Phi(\phi^1)(z) = \Phi(g\phi^1)(z) \in A_{g(y)}.$$

We also have

$$f(g(y)) = f(\phi^s(x)) = \Phi(\phi^s)(z) \in A_{g(y)},$$

so it follows that $fg(y) = \Phi(g)(f(y))$.

Surjectivity. The G -equivariance of f implies that the image of f is $\Phi(G)$ -invariant. Since we assumed that $\text{Fix}(\Phi(G)) = \emptyset$, the image of f must be \mathbb{R} .

It remains to show that f is C^r under the assumption that $p = r$. To do this, we need a generalization of the following theorem of Takens.

THEOREM 4.3. (Takens [17]) *Let $f : M_1 \rightarrow M_2$ be a bijection between two smooth manifolds M_1 and M_2 with the property that $g : M_1 \rightarrow M_2$ is a C^∞ -diffeomorphism if and only if fgf^{-1} is a C^∞ -diffeomorphism. Then f is also a C^∞ -diffeomorphism.*

In our case, $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies the property that for each $g \in G$ the conjugate $fgf^{-1} = \Phi(g)$ is C^r . We also know that $\Phi(G)$ acts transitively on the image of f . Takens’s original proof can be easily adapted to this case to show that f is of class C^r . We omit the details. The reader may also consult the main theorem of Rybicki in [15] for a similar (but more complicated to state) theorem that applies directly to our situation.

Thus, we have shown that Theorem 4.1 holds provided that $\text{Fix}(\Phi(G)) = \emptyset$. If instead $\text{Fix}(\Phi(G)) \neq \emptyset$, we can pick any connected component J of $\mathbb{R} \setminus \text{Fix}(\Phi(G))$ and run the proof above for the induced group homomorphism $\Phi_J : G \rightarrow \text{Diff}^r(J)$ given by $\Phi_J(g) = \Phi(g)|_J$. What we get is a homeomorphism (or C^r -diffeomorphism in the $p = r$ case) $f : \mathbb{R} \rightarrow J$ such that $f(g(x)) = \Phi(g)(f(x))$ for all $g \in G$. Doing this separately on each connected component defines the topologically diagonal embedding.

This completes the proof of Theorem 4.1. □

Using the fact that the image of our homomorphism is $\text{Diff}_c^p(\mathbb{R})$, that is, all diffeomorphisms in the image have compact support, we may conclude something stronger about the topologically diagonal embedding.

PROPOSITION 4.4. *Suppose that $p, r \geq 2$ and $\Phi : \text{Diff}_c^p(\mathbb{R}) \rightarrow \text{Diff}_c^p(\mathbb{R})$ is a topologically diagonal embedding. Then there are only finitely many connected components of $\mathbb{R} \setminus \text{Fix}(\Phi(\text{Diff}_c^p(\mathbb{R})))$.*

Proof. This will follow from the fact that $\Phi(G)$ acts by C^2 -diffeomorphisms along with the ‘diagonality’ of the action of $\Phi(G)$. That the embedding is topologically diagonal means exactly that for any two connected components J_1 and J_2 of $\mathbb{R} \setminus \text{Fix}(\Phi(G))$ with embeddings $f_i : \mathbb{R} \rightarrow J_i$, we have

$$(f_2 f_1^{-1}) \circ \Phi(g)|_{J_1} \circ (f_1 f_2^{-1}) = \Phi(g)|_{J_2}$$

for each $g \in G$. In other words, $f_2 f_1^{-1}$ conjugates the action of G on J_1 with the action of G on J_2 . We also know that there is some compact subset of \mathbb{R} that contains all the connected components J_i that are bounded. Moreover, Takens’s theorem (Theorem 4.3) implies that each $f_2 f_1^{-1}$ is of class C^s , where $s = \min(p, r) \geq 2$. One can show that such

an action can be made C^1 , but it will necessarily have unbounded derivatives. See, for example, [14] for more details. \square

4.1. *Finishing the proof of Theorem 1.5.* We conclude by stating the necessary modifications to replace either copy of \mathbb{R} with S^1 in the statement of Theorem 4.1 and therefore prove Theorem 1.5. There are three cases to consider.

- (1) Homomorphisms $\Phi : \text{Diff}_+^r(S^1) \rightarrow \text{Diff}_c^p(\mathbb{R})$. There are no non-trivial such homomorphisms since $\text{Diff}_+^r(S^1)$ is simple and has torsion.
- (2) Homomorphisms $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_+^p(S^1)$. For this, the proof of Theorem 4.1 above works verbatim and shows that this embedding is topologically diagonal. Note, in particular, that there must be some point in S^1 fixed by every diffeomorphism in the image of Φ .
- (3) Homomorphisms $\Phi : \text{Diff}_+^r(S^1) \rightarrow \text{Diff}_+^p(S^1)$. $\text{Diff}_+^r(S^1)$ contains many copies of $\text{Diff}_c^r(\mathbb{R})$ (as in Example 1.1(c)) since there are many embeddings of \mathbb{R} in S^1 . For any such copy, consider the restriction of Φ to the $\text{Diff}_c^r(\mathbb{R})$ subgroup, a homomorphism $\text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_+^p(S^1)$. We know already that this must be topologically diagonal. The fact that this is true for *any* copy means that the embeddings are compatible, in the sense that if f_i and f_j are any two maps $\mathbb{R} \rightarrow S^1$ coming from any two embeddings of copies of $\text{Diff}_c^r(\mathbb{R})$, then f_i and f_j agree on their common domain. It follows that the f_i patch together to define a globally defined, continuous map $f : S^1 \rightarrow S^1$, which is locally a homeomorphism. The fact that f is equivariant with respect to the action of $\text{Diff}_+^r(S^1)$ and that $\text{Diff}_+^r(S^1)$ contains finite order rotations means that f must be a global homeomorphism. Again, we can use Takens's theorem to show that f is C^r provided that $p = r$. It is also possible to show that Φ is topologically diagonal in this case by using flows on S^1 and Corollary 3.13.

5. Application: dimension as an algebraic invariant

Recall the statement of Theorem 1.6.

THEOREM 1.6. *Let $r \geq 3$, $p \geq 2$, let M_1 be any manifold and M_2 a 1-manifold. Suppose that $\Phi : \text{Diff}_c^r(M_1) \rightarrow \text{Diff}_c^p(M_2)$ is an injective homomorphism. Then $\dim(M_1) = 1$ and Φ is topologically diagonal. If $r = p$, then Φ is C^r -topologically diagonal.*

To prove this it will be enough to show that $\dim(M_1) = 1$, since in that case it follows from Theorem 1.5 that Φ is topologically diagonal.

We start with an easy corollary of Szekeres's theorem.

COROLLARY 5.1. *Let $r \geq 2$ and let g be a C^r -diffeomorphism of $I = [0, 1)$ with no interior fixed points. Then g^2 and g have the same centralizer in $\text{Diff}_+^r(I)$.*

Proof. By uniqueness of the flow in Szekeres's theorem, $C_{\text{Diff}_+^r(I)}(g) = C_{\text{Diff}_+^r(I)}(g^2)$. Since

$$C_{\text{Diff}_+^r(I)}(g) = C_{\text{Diff}_+^1(I)}(g) \cap \text{Diff}^r(I),$$

we have also

$$C_{\text{Diff}_+^r(I)}(g) = C_{\text{Diff}_+^r(I)}(g^2). \quad \square$$

With a little more work, we can show that Corollary 5.1 is also true for compactly supported diffeomorphisms of \mathbb{R} .

LEMMA 5.2. *Let $r \geq 2$ and let $g \in \text{Diff}_c^r(\mathbb{R})$. Then $C_{\text{Diff}_c^r(\mathbb{R})}(g) = C_{\text{Diff}_c^r(\mathbb{R})}(g^2)$.*

Proof. That $C(g) \subset C(g^2)$ is immediate. We prove that the reverse containment also holds.

Let $g \in \text{Diff}_c^r(\mathbb{R})$ and suppose that $r \geq 2$. Then $\text{Fix}(g) \neq \emptyset$ and for any connected component J of $\mathbb{R} \setminus \text{Fix}(g)$ we can apply Szekeres's theorem to $g|_J$. Note that since g is orientation-preserving, $\text{Fix}(g) = \text{Fix}(g^2)$.

Consider a connected component J of $\mathbb{R} \setminus \text{Fix}(g)$. By Corollary 5.1, any diffeomorphism that leaves J invariant and commutes with g^2 on J also commutes with g on J . A C^r -diffeomorphism h that commutes with g^2 but does *not* leave J invariant must map J to another connected component of $\mathbb{R} \setminus (\text{Fix}(g))$, say J' . Moreover, h must conjugate the action of g^2 on J to the action of g^2 on J' . By uniqueness of the flow from Szekeres's theorem, if g_J^t is the flow on J that commutes with $g^2|_J$ then $hg_J^t h^{-1}$ is the flow on J' that commutes with $g^2|_{J'}$. It follows that h conjugates the action of $g = (g^2)^{1/2}$ on J to the action of g on J' . In other words, for any $x \in J$, we have $hg(x) = gh(x)$, that is, h commutes with g on J . Since this is true for any component of $\mathbb{R} \setminus \text{Fix}(g)$, it follows that h commutes with g . \square

Now we can easily prove Theorem 1.6 if $M_2 = \mathbb{R}$.

Proof of Theorem 1.6, $M_2 = \mathbb{R}$ case. Suppose that M_1 is a manifold of dimension $n \geq 2$ and that Φ is an injective homomorphism $\text{Diff}_c^r(M_1)_0 \rightarrow \text{Diff}_c^p(\mathbb{R})$. Let B^n be an open ball in M_1 and consider the induced injective homomorphism $\text{Diff}_c^r(B^n) \rightarrow \text{Diff}_c^r(\mathbb{R})$ given by restricting the domain of Φ to elements supported in B^n . We claim that no such homomorphism exists.

To see this, we will construct a diffeomorphism $g \in \text{Diff}_c^r(B^n)$ such that

$$C(g^2)_{\text{Diff}_c^r(B^n)} \neq C(g)_{\text{Diff}_c^r(B^n)},$$

that is, such that there is some $h \in \text{Diff}_c^r(B^n) \subset \text{Diff}_c^r(M_1)$, where $[h, g^2] = 1$ but $[h, g] \neq 1$. In this case we will have that $[\Phi(h), \Phi(g^2)] = 1$ but $[\Phi(h), \Phi(g)] \neq 1$ in $\text{Diff}_c^r(\mathbb{R})$, contradicting Lemma 5.2.

To construct g , choose any diffeomorphism that acts as rotation by π on a small cylinder $B^2 \times [0, 1]^{n-2}$ inside B^n , and has trivial centralizer outside $D^2 \times [0, 1]^{n-2}$. Precisely, we want the restriction of g to $B^2 \times [0, 1]^{n-2}$ to be given by $(re^{i\theta}, x) \mapsto (re^{i(\pi+\theta)}, x)$. Now let h be any diffeomorphism supported inside $D^2 \times [0, 1]^{n-2}$ that does not commute with g . By construction, the restriction of g^2 to $D^2 \times [0, 1]^{n-2}$ is trivial, so h commutes with g^2 , as desired. \square

Unfortunately, there *are* elements $g \in \text{Diff}^r(S^1)$ such that $C(g) \neq C(g^2)$ —finite order rotations are one example, but there are infinite order such elements as well. Thus, our proof above will not work if $M_2 = S^1$. To prove Theorem 1.6 when $M_2 = S^1$, we will use the description of $\text{Diff}_c^r(\mathbb{R})$ actions from Theorem 1.5.

Proof of Theorem 1.6, $M_2 = S^1$ case. Suppose that M_1 is a manifold of dimension $n \geq 2$ and that Φ is an injective homomorphism $\text{Diff}_c^r(M_1)_0 \rightarrow \text{Diff}_c^p(S^1)$. We want to produce a contradiction. We claim that it is enough to show that there is no injective homomorphism




FIGURE 5. $\text{Diff}_c^r(\mathbb{R})$ embeds in $\text{Diff}_c^r(B)$ by acting on the radii of an annulus.

$\text{Diff}_c^r(B^2) \rightarrow \text{Diff}_+^p(S^1)$. (Recall that B^2 is the open two-dimensional disc.) This is because of the following easy lemma.

LEMMA 5.3. $\text{Diff}_c^r(B^2)$ embeds as a subgroup of $\text{Diff}_c^r(M_1)$ for any manifold M_1 of dimension at least two.

Proof of Lemma 5.3. This is trivial if $\dim(M_1) = 2$, since B^2 embeds in M_1 as an open submanifold. If $\dim(M_1) = n > 2$, we can embed $\text{Diff}_c^r(B^2)$ in $\text{Diff}_c^r(B^2 \times S^{n-2})$ via $g \cdot (b, s) = (g(b), s)$. Then take any embedding of $B^2 \times S^{n-2}$ in M_1 as an open submanifold. This defines an embedding $\text{Diff}_c^r(B^2 \times S^{n-2}) \hookrightarrow \text{Diff}_c^r(M_1)$. \square

Thus, we have reduced the proof of Theorem 1.6 to the following lemma.

LEMMA 5.4. There is no injective homomorphism $\text{Diff}_c^r(B^2) \rightarrow \text{Diff}_+^p(S^1)$.

Proof of Lemma 5.4. First, we show that $\text{Diff}_c^r(B^2)$ has a subgroup H isomorphic to $\text{Diff}_c^r(\mathbb{R})$. To do this, let $\tau : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_c^r(\mathbb{R} \times S^1)$ be the map given by $\tau(g)(x, s) \mapsto (g(x), s)$, and let $f : (\mathbb{R} \times S^1) \rightarrow B^2$ be a C^r embedding. Then $f_*\tau : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}_c^r(B^2)$ is injective with image $H \cong \text{Diff}_c^r(\mathbb{R})$. See Figure 5.

Suppose for contradiction that $\Phi : \text{Diff}_c^r(B^2) \rightarrow \text{Diff}_+^p(S^1)$ is injective. By Theorem 1.5, $\Phi(H)$ is a topologically diagonal embedding, and by Proposition 4.4, $\text{Fix}(\Phi(H))$ has finitely many connected components. Let N be the number of components of $\text{Fix}(\Phi(H))$. Note that if ϕ is any diffeomorphism of S^1 that permutes the components of $S^1 \setminus \text{Fix}(\Phi(H))$, then $\phi^{N!}$ leaves each component of $S^1 \setminus \text{Fix}(\Phi(H))$ invariant.

We will find a diffeomorphism $g \in \text{Diff}_c^r(B^2)$ with support disjoint from $\text{supp}(H)$, and such that the centralizer of $g^{2(N!)}$ is not equal to the centralizer of $g^{N!}$. To construct such a g , we may use a diffeomorphism that acts as rotation by $\pi/N!$ on a small cylinder disjoint from $\text{supp}(H)$, similar to the construction in the $M_2 = \mathbb{R}$ case above. Let g' be a diffeomorphism supported on the cylinder that does not commute with $g^{N!}$. As in our construction for the $M_2 = \mathbb{R}$ case, g' commutes with H and with $g^{2(N!)}$ but not with $g^{N!}$.

Since g and H commute, $\Phi(g)$ commutes with $\Phi(H)$, and $\Phi(g)^{N!}$ leaves invariant each subinterval or isolated point of $\text{Fix}(\Phi(H))$. Moreover, since $\Phi(g)^{N!}$ commutes with $\Phi(H)$ and H is non-abelian, $\Phi(g)^{N!}$ must be supported on the subintervals fixed by $\Phi(H)$ (we are using the proof of Proposition 2.2 again here). In particular, we can regard $\Phi(g)^{N!}$ and $\Phi(g)^{2N!}$ as diffeomorphisms of an interval, both with the same support.

Finally, consider $\Phi(g')$. Since $\Phi(g')$ commutes with $\Phi(H)$, we also have that $\Phi(g')^{N!}$ leaves invariant each isolated point or subinterval of $\text{Fix}(\Phi(H))$. In particular, it fixes some point and we may regard $\Phi(g')^{N!}$ as a diffeomorphism of the interval. Then $\Phi(g')^{N!}$ commutes with $\Phi(g)^{2N!}$ but not with $\Phi(g)^{N!}$, contradicting Lemma 5.2.

This completes the proof of Lemma 5.4 and the proof of Theorem 1.6. \square

6. Compact support, simplicity, and smoothness

We remark on some of the hypotheses of our theorems and ask about generalizations.

6.1. *Compact support.* As mentioned in the introduction, the group $\text{Diff}_c^r(M)$ is simple for $r \neq \dim(M) + 1$. We believe that it should be possible to prove an analog of Theorem 1.5 for $\text{Diff}^r(\mathbb{R})_0$ rather than $\text{Diff}_c^r(\mathbb{R})$ using the following theorem of Schweitzer, which classifies all normal subgroups of $\text{Diff}^r(\mathbb{R})_0$.

THEOREM 6.1. (Schweitzer [16]) *Let $r \geq 3$. There are only three proper normal subgroups of $\text{Diff}^r(\mathbb{R})_0$:*

- (a) *the group of diffeomorphisms coinciding with the identity in a neighborhood of ∞ ;*
- (b) *those coinciding with the identity in a neighborhood of $-\infty$;*
- (c) $\text{Diff}_c^r(\mathbb{R})$.

Remark 6.2. This is actually a special case of a much more general theorem which classifies normal subgroups of $\text{Diff}^r(\mathbb{R}^n)_0$, for any $r \geq 1$, such that $r \neq n + 1$.

Thus, any non-injective homomorphism $\Phi : \text{Diff}^r(\mathbb{R})_0 \rightarrow \text{Diff}^p(\mathbb{R})_0$ has kernel equal to one of the subgroups above. We conjecture that no such homomorphism exists.

Note also that we did not make very heavy use of the fact that the *target* group was a group of compactly supported diffeomorphisms (or even orientation-preserving diffeomorphisms). It may be possible to adapt our proofs for homomorphisms $\Phi : \text{Diff}_c^r(\mathbb{R}) \rightarrow \text{Diff}^r(\mathbb{R})$, starting with a modified version of Proposition 2.2.

6.2. *Manifolds with boundary.* Our results are not true for manifolds with boundary. For example, $\text{Diff}^r[0, 1]$ admits non-trivial homomorphisms to \mathbb{R} given by taking the derivative at 0 or at 1. These can be used to build homomorphisms of $\text{Diff}^r[0, 1]$ into $\text{Diff}^r[0, 1]$ or $\text{Diff}_c^r(\mathbb{R})$ that are not topologically diagonal.

However, I do not know whether it is possible to construct any counterexamples without using derivatives.

Question 6.3. Let G^∞ be the group of C^∞ diffeomorphisms of the interval that are infinitely tangent to the identity at 0 and at 1. Do there exist homomorphisms $\Phi : G^\infty \rightarrow \text{Diff}_c^r(\mathbb{R})$ that are not topologically diagonal?

6.3. *Smoothness.* There are also counterexamples to some of our tools when we remove assumptions on smoothness. In §2 on commuting subgroups, we assumed all diffeomorphisms were of class at least C^2 . Although C^{1+bv} would have sufficed, Kopell's lemma and Denjoy's theorem do not hold for C^1 -diffeomorphisms. Namely, there are fixed-point-free C^1 -diffeomorphisms of the interval commuting with C^1 -diffeomorphisms that have fixed points, as well as C^1 -diffeomorphisms of the circle with irrational rotation number that are not conjugate to irrational rotations (see [14]). Note also that the group $\text{Diff}^{1+bv}(S^1)$ is known to be *not* simple (see [11]) and whether the group $\text{Diff}^2(S^1)$ is simple is unknown! In particular, $\text{Diff}^{1+bv}(S^1)$ admits non-topologically diagonal and even non-continuous homomorphisms to other groups of diffeomorphisms, and it is possible that $\text{Diff}^2(S^1)$ does as well.

However, working in the category of *homeomorphisms* rather than diffeomorphisms, Militon has recently obtained results in the same spirit as ours, using different techniques. See [12, 13].

6.4. *Higher dimensions.* We conclude with a final remark on one of the difficulties in extending the results of this paper to manifolds of dimension $n > 1$. Our methods here relied heavily on properties of commuting subgroups, namely, the Hölder–Kopell theory and Proposition 2.2. The following example shows that Proposition 2.2 does not hold in higher dimensions.

Example 6.4. (Large centralizers in higher dimensions) Consider the following two subgroups of $\text{Diff}^r(\mathbb{R}^2)$:

$$G := \{f \in \text{Diff}^r(\mathbb{R}^2) : f(x, y) = (f_1(x), y) \text{ for some } f_1 \in \text{Diff}^r(\mathbb{R})\};$$

$$H := \{f \in \text{Diff}^r(\mathbb{R}^2) : f(x, y) = (x, f_2(y)) \text{ for some } f_2 \in \text{Diff}^r(\mathbb{R})\}.$$

These commute, that is, H is a subset of the centralizer of G , they are both 'large' in the sense that they are infinite-dimensional, non-abelian, etc. and yet neither one fixes any non-empty subset of \mathbb{R}^2 .

Similar subgroups can be found in $\text{Diff}^r(\mathbb{T}^2)$, in $\text{Diff}^r(\mathbb{R}^n)$ and $\text{Diff}^r(\mathbb{T}^n)$ for the n -dimensional torus. However, we note that the groups G and H above are *not* groups of compactly supported diffeomorphisms.

Question 6.5. Do there exist examples of large (for any reasonable definition of 'large') commuting subgroups in $\text{Diff}_c^r(\mathbb{R}^n)$ that do not fix any open set? Do there exist examples in $\text{Diff}_c^r(M^n)$ for all n -manifolds?

A negative answer to this question would be the first step towards finding an algebraic–topological correspondence for manifolds of higher dimension, analogous to the correspondence we found here for 1-manifolds.

Acknowledgements. The author wishes to thank Benson Farb, Amie Wilkinson and Bena Tshishiku for many helpful conversations regarding this work, and John Franks and Emmanuel Militon for their comments and careful reading. Thanks also to the anonymous referee for suggestions to correct and clarify work in §2.

REFERENCES

- [1] A. Banyaga. On isomorphic classical diffeomorphism groups I. *Proc. Amer. Math. Soc.* **98** (1986), 113–118.
- [2] A. Banyaga. *The Structure of Classical Diffeomorphism Groups (Mathematics and Its Applications)*. Kluwer Academic Publishers, Dordrecht, 1997.
- [3] B. Farb and J. Franks. Groups of homeomorphisms of one-manifolds, I: Actions of nonlinear groups. *Preprint*, 2001, arXiv:math/0107085v2.
- [4] B. Farb and J. Franks. Group actions on 1-manifolds, II: Extensions of Hölder’s theorem. *Trans. Amer. Math. Soc.* **355**(11) (2003), 4385–4396.
- [5] R. Filipkiewicz. Isomorphisms between diffeomorphism groups. *Ergod. Th. & Dynam. Sys.* **2** (1982), 159–171.
- [6] E. Ghys. Prolongements des difféomorphismes de la sphère. *L’Enseign. Math.* **37** (1991), 45–59.
- [7] O. Hölder. Die Axiome der Quantität und die Lehre vom Mass. *Ber. Verh. Sachs. Ges. Wiss. Leipzig, Math. Phys. Cl* **53** (1901), 1–64.
- [8] N. Kopell. Commuting diffeomorphisms. *Global Analysis (Proceedings of Symposia in Pure Mathematics, XIV)*. Eds. S.-S. Chern and S. Smale. Providence, RI, 1968, pp. 165–184.
- [9] J. Mather. Commutators of diffeomorphisms. *Comment. Math. Helv.* **49** (1974), 512–528.
- [10] J. Mather. Commutators of diffeomorphisms: II. *Comment. Math. Helv.* **50** (1975), 33–40.
- [11] J. Mather. Commutators of diffeomorphisms, III: A group which is not perfect. *Comment. Math. Helv.* **60** (1985), 122–124.
- [12] E. Militon. Continuous actions of the group of homeomorphisms of the circle on surfaces. *Preprint*, 2012, arXiv:1211.0846v2.
- [13] E. Militon. Actions of groups of homeomorphisms on one-manifolds. *Preprint*, 2013, arXiv:1302.3737.
- [14] A. Navas. *Groups of Circle Diffeomorphisms*. University of Chicago Press, Chicago, 2011.
- [15] T. Rybicki. Isomorphisms between groups of diffeomorphisms. *Proc. Amer. Math. Soc.* **123**(1) (1995), 303–310.
- [16] P. Schweitzer. Normal subgroups of diffeomorphism and homeomorphism groups of \mathbb{R}^n and other open manifolds. *Preprint*, 2012, arXiv:0911.4835v3.
- [17] F. Takens. Characterization of a differentiable structure by its group of diffeomorphisms. *Bol. Soc. Brasil. Mat.* **10** (1979), 17–25.