

# Longitudinal epigenetic variation of DNA methyltransferase genes is associated with vulnerability to post-traumatic stress disorder

L. Sipahi<sup>1</sup>, D. E. Wildman<sup>1,2</sup>, A. E. Aiello<sup>3</sup>, K. C. Koenen<sup>4</sup>, S. Galea<sup>4</sup>, A. Abbas<sup>1</sup> and M. Uddin<sup>1,5\*</sup>

<sup>1</sup>Center for Molecular Medicine and Genetics, Wayne State University School of Medicine, Detroit, MI, USA

<sup>2</sup>Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI, USA

<sup>3</sup>Department of Epidemiology, University of North Carolina Gillings School of Global Public Health, Chapel Hill, NC, USA

<sup>4</sup>Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA

<sup>5</sup>Department of Psychiatry and Behavioral Neurosciences, Wayne State University School of Medicine, Detroit, MI, USA

**Background.** Epigenetic differences exist between trauma-exposed individuals with and without post-traumatic stress disorder (PTSD). It is unclear whether these epigenetic differences pre-exist, or arise following, trauma and PTSD onset.

**Method.** In pre- and post-trauma samples from a subset of Detroit Neighborhood Health Study participants, DNA methylation (DNAm) was measured at DNA methyltransferase 1 (*DNMT1*), *DNMT3A*, *DNMT3B* and *DNMT3L*. Pre-trauma DNAm differences and changes in DNAm from pre- to post-trauma were assessed between and within PTSD cases ( $n=30$ ) and age-, gender- and trauma exposure-matched controls ( $n=30$ ). Pre-trauma DNAm was tested for association with post-trauma symptom severity (PTSS) change. Potential functional consequences of DNAm differences were explored via bioinformatic search for putative transcription factor binding sites (TFBS).

**Results.** *DNMT1* DNAm increased following trauma in PTSD cases ( $p=0.001$ ), but not controls ( $p=0.067$ ). *DNMT3A* and *DNMT3B* DNAm increased following trauma in both cases (*DNMT3A*:  $p=0.009$ ; *DNMT3B*:  $p<0.001$ ) and controls (*DNMT3A*:  $p=0.002$ ; *DNMT3B*:  $p<0.001$ ). In cases only, pre-trauma DNAm was lower at a *DNMT3B* CpG site that overlaps with a TFBS involved in epigenetic regulation ( $p=0.001$ ); lower pre-trauma *DNMT3B* DNAm at this site was predictive of worsening of PTSS post-trauma ( $p=0.034$ ). Some effects were attenuated following correction for multiple hypothesis testing.

**Conclusions.** DNAm among trauma-exposed individuals shows both longitudinal changes and pre-existing epigenetic states that differentiate individuals who are resilient *versus* susceptible to PTSD. These distinctive DNAm differences within *DNMT* loci may contribute to genome-wide epigenetic profiles of PTSD.

Received 4 September 2013; Revised 28 February 2014; Accepted 28 March 2014; First published online 25 April 2014

**Key words:** DNA methylation, epidemiology, epigenetics, longitudinal studies, post-traumatic stress disorder.

## Introduction

Post-traumatic stress disorder (PTSD) is a prevalent and debilitating mental health disorder that may arise following exposure to a potentially traumatic event (APA, 2013). While the lifetime prevalence of traumatic exposure is 50–90% (Kessler *et al.* 1995), PTSD in the general US population is estimated to be only 6.8% (Kessler & Wang, 2008). Although the majority of persons exposed to trauma display resiliency (Kessler *et al.* 1995; Breslau *et al.* 1998; Acierio *et al.* 2007; Kessler & Wang, 2008), the molecular

underpinnings of risk remain poorly characterized. The identification of risk markers, and particularly biomarkers, that distinguish between persons at high and low risk of developing PTSD following trauma exposure has been identified as a priority research goal by the Institute of Medicine (2012), Department of Defense (CDMRP, Department of Defense, 2010) and the National Institute of Mental Health (2008). Ideally, the ability to identify persons at high risk of developing PTSD would enable providers to target evidence-based interventions to high-risk groups (Andrews & Neises, 2012). The identification of robust predictive biomarkers may also improve our understanding of the pathophysiology of PTSD and lead to more effective pharmacological interventions.

Although much work has been done to identify social and environmental factors that contribute to PTSD risk (e.g. Kulka *et al.* 1990; Breslau *et al.* 1991,

\* Address for correspondence: M. Uddin, Ph.D., Center for Molecular Medicine and Genetics, Wayne State University School of Medicine, 3309 Scott Hall, 540 E. Canfield Avenue, Detroit, MI 48201, USA.

(Email: monica.uddin@wayne.edu)

2004; Brewin et al. 2000; Koenen et al. 2003; DiGrande et al. 2008; Galea et al. 2008; Kun et al. 2009), the biological underpinning of differential PTSD risk and resiliency remains to be more fully elucidated. Twin studies have demonstrated heritability and genetic contribution to PTSD risk (True et al. 1993; Koenen et al. 2002; Stein et al. 2002) and targeted gene and Genome-wide Association Study (GWAS) approaches have identified both genetic risk loci (Lu et al. 2008; Ressler et al. 2011; Chang et al. 2012; Logue et al. 2013) and important gene × environment interactions (Binder et al. 2008; Xie et al. 2010; Uddin et al. 2013) that contribute to risk for the disorder; nevertheless, a substantial proportion of biologically mediated variance in PTSD risk has yet to be explained.

Epigenetic variability is considered a plausible and increasingly empirically supported contributor to the etiology of phenotypes with marked genetic and environmental influences (Meaney, 2010), including certain psychopathologies (Toyokawa et al. 2012). Indeed, recent advances have revealed that PTSD risk and resiliency are associated with differential epigenetic variation (El-Sayed et al. 2012). Epigenetic mechanisms—including histone modifications, non-protein coding RNAs, and, most notably, DNA methylation (DNAm)—affect gene expression and cellular phenotype without altering the underlying DNA sequence (Feinberg, 2008; Meaney, 2010). DNAm is stably heritable across mitotic replications, but is modifiable throughout the life course in response to lived experiences and environmental exposures (Bird, 2002). In primordial mammalian germ cells, global DNAm is removed (with the exception of imprinted loci) (Reik et al. 2001), with new patterns established by *de novo* DNA methyltransferases DNMT3A, DNMT3B and DNMT3L following fertilization (Bourc'his et al. 2001; Bourc'his & Bestor, 2004; Kaneda et al. 2004; Kato et al. 2007; Ooi et al. 2007). These reprogrammed DNAm patterns are largely maintained throughout mitotic DNA replication by the action of the maintenance methyltransferase, DNMT1 (Li et al. 1992; Seisenberger et al. 2013).

Although influenced by other variables, global DNAm patterns are largely established and maintained by the activity of the DNA methyltransferases DNMT1, DNMT3A, DNMT3B and DNMT3L (Feng & Fan, 2009). Gene expression evidence suggests that these DNMTs may be active throughout the life course (Robertson et al. 1999; Feng et al. 2005; Siegmund et al. 2007), including in brain tissue (Goto et al. 1994; Veldic et al. 2004; Feng et al. 2005) and in association with mental disorders (Veldic et al. 2004, 2005). In addition, protein-level expression of DNMT1 (Inano et al. 2000; Veldic et al. 2005) and DNMT3A (Feng et al. 2005) has been demonstrated in the mouse and human

brain. With respect to PTSD, recent work confirms that DNMT activity plays a role in mediating risk for PTSD-related phenotypes, including fear conditioning and memory consolidation (Miller & Sweatt, 2007; Feng et al. 2010). Together, these findings suggest that DNAm and DNMTs represent promising targets for the identification of epigenetic underpinnings of differential PTSD risk and resiliency.

Studies of epigenetic variation have provided important insights into PTSD risk, but have been largely limited by cross-sectional analyses of post-trauma samples. Most notably, epidemiological cohorts from Detroit (Uddin et al. 2010) and Atlanta (Smith et al. 2011) have been the basis of research that has demonstrated cross-sectional differential DNAm that distinguishes between trauma-exposed individuals with *versus* without PTSD. *DNMT3B* and *DNMT3L* were among the differentially methylated loci identified in the Detroit study (Uddin et al. 2010). More recently, longitudinal DNAm data among PTSD cases and controls have been reported, including studies using samples from a cohort of US military personnel deployed to Iraq and Afghanistan (Rusiecki et al. 2012, 2013). To further elucidate whether differential DNAm between trauma-exposed controls and PTSD cases represent pre-existing susceptibility/resiliency factors or downstream biomarkers of PTSD, additional longitudinal analyses are required. Finally, while the identification of epigenetic variation associated with mental health outcomes is important, work must begin to test the putative functionality of mental health-associated differential DNAm. For example, the identification of transcription factor binding sites (TFBS) that overlap with differentially methylated CpG sites and to which transcription factor binding may be disrupted offer one possibility of supporting DNAm functionality (Weaver et al. 2004).

Here, we analysed DNAm from individuals pre- and post-trauma to identify differences that characterize individuals who are susceptible *versus* resilient to PTSD following trauma. To assess potential functional consequences of examined DNAm differences, we then performed a bioinformatic search for the presence of putative TFBS (Weaver et al. 2004). Results from this work suggest that PTSD-relevant DNAm differences in DNMT loci may exist both prior to and following trauma, with implications for future targeted interventions.

## Method

### Subjects

Samples are from a subset of participants from the Detroit Neighborhood Health Study (DNHS), a

longitudinal, community-representative cohort of adult residents in Detroit, MI, USA. The current study draws on peripheral blood samples and survey data obtained at two time points from 60 DNHS participants; 46 were female and 14 male; 46 were African-American and 12 were Caucasian, and two were Hispanic. The average age was 55.1 years. PTSD diagnosis was assessed via structured interview administered via telephone (Breslau *et al.* 1998). PTSD symptoms were assessed in reference to both the traumatic event that the participant regarded as their worst and one randomly selected traumatic event from the remaining traumas that the participant experienced. Lifetime PTSD cases met all six Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) criteria (APA, 1994) in reference to either the worst or random traumatic event. The diagnostic interview showed good validity against the Clinician Administered PTSD Scale (Blake *et al.* 1995) as described elsewhere (Uddin *et al.* 2010). The Institutional Review Board of the University of Michigan reviewed and approved the study protocol. Incident cases ( $n=30$ ) of PTSD were identified in either waves 2, 3 or 4 of DNHS data collection among individuals for whom blood samples were available at both the wave of first PTSD diagnosis and the immediately previous, pre-incident trauma wave. Non-PTSD controls ( $n=30$ ) were matched to cases on the basis of age, sex and number of traumatic event types. DNA samples were isolated from both pre- and post-trauma time points for both cases and controls. The time between pre- and post-trauma time points was approximately 1 year. Cases and controls had no history of PTSD prior to the post-trauma wave.

#### **Methylation quantification by targeted bisulfite pyrosequencing**

##### *DNA isolation*

DNA was isolated from whole blood acquired via venepuncture when available from DNHS participants selected for inclusion in this study. Blood spots were used as an alternative source of whole blood-derived DNA when venepuncture samples were unavailable. The exact tissue type was shared between matched case-control pairs in all instances. Venepuncture- and bloodspot-derived whole blood represents the same tissue and therefore should not differ with respect to DNAm, as confirmed by numerous studies to date (Wong *et al.* 2008; Aberg *et al.* 2013; Hollegaard *et al.* 2013).

*Whole blood.* DNA was isolated from whole blood using the QIAamp DNA Blood Mini Kit (Qiagen, USA) and the QuickGene DNA Whole blood Kit S

(Lifesciences, FujiFilm, Japan) using the manufacturers' recommended protocols.

*Blood spots.* DNA was isolated using the QIAamp DNA Micro kit (Qiagen, USA) using the manufacturer's recommended protocol. For each sample, one 6 mm punch was taken from dried blood spots using a disposable, sterile biopsy punch (Miltex, USA) within a sterile field and placed immediately into a sterile 1.7 ml microcentrifuge tube. New gloves, biopsy punches and sterile fields were utilized for each sample. Negative controls in the form of blank extractions were included with all DNA isolations.

##### *Bisulfite conversion*

For each sample, about 750 ng of DNA was bisulfite converted using the EpiTect Bisulfite Kit (Qiagen, USA) using the manufacturer's recommended protocol. Negative controls in the form of bisulfite conversion of water were included with each bisulfite conversion.

##### *Pyrosequencing*

Assays to assess the methylation levels of CpG sites found in the *DNMT1*, *DNMT3A*, and *DNMT3L* and *DNMT3B* (see below for assay-specific details) were custom designed using the Pyromark Q24 Assay Design Software 2.0 (Qiagen, USA). Targeted CpG sites were selected based on prior evidence (Uddin *et al.* 2010) of involvement in epigenetic regulation of PTSD risk (*DNMT3B*, *DNMT3L*) and to investigate whether longitudinal, PTSD-associated DNAm differences exist across DNMT genes more broadly (*DNMT1*, *DNMT3A*, *DNMT3B* and *DNMT3L*). Because the *DNMT3B* target CpG is located in a CpG island, our designed assay captures DNAm at 12 CpG sites in an approximately 70 base pair region of exon 1 (see *DNMT3B* assay section below for details). Single CpG sites were assessed at *DNMT1*, *DNMT3A* and *DNMT3L* loci (see individual assay section below for details); these CpG sites did not fall into CpG islands. *DNMT1*, *DNMT3A* and *DNMT3L* CpG sites and 2 *DNMT3B* CpG sites assessed are also found on the HM27 and HM450 K methylation bead chips from Illumina (see below for actual HG19 nucleotide location). The capacity for each assay to capture DNAm levels ranging from 0 to 100% was validated using commercially available demethylated and highly methylated DNA at dilutions of 1:0 (unmethylated), 3:1, 1:1, 1:3 and 0:1 (highly methylated). Polymerase chain reaction (PCR) amplification of target sequences was performed on 20 ng of bisulfite-converted DNA samples using the PyroMark PCR kit (Qiagen, USA).

Bisulfite-converted, PCR-amplified DNA was pyrosequenced on the Pyromark Q24 Pyrosequencer (Qiagen, USA) using the manufacturer's recommended protocol and default settings. All methylation analyses were conducted in triplicate with appropriate negative controls included at each of the following steps: DNA isolation, bisulfite conversion, PCR amplification and pyrosequencing reaction.

Details of each custom assay are listed below.

#### *DNMT1*

PCR forward primer: TTTTTTAGGTGTGA  
TGGGGATAAAG

PCR reverse primer (biotinylated): CAAAAA  
CTCTCACAAACCCTTAAA

PCR program (50 cycles):

Initial	15 min at 95 °C
Denaturation	30 s at 94 °C
Annealing	30 s at 58 °C
Extension	30 s at 72 °C
Final	10 min at 72 °C
Hold	4 °C

Sequencing primer: GTGATGGGGAT  
AAAGT

Target sequence: AGCGAGAAGCCCC  
AAGGGTTGTGAGA (**CpG target in bold**; hg19: chr19:10,305,909-10,305,936)

#### *DNMT3A*:

PCR forward primer: GGTGGGAGGTTGAAT  
GAAATGA

PCR reverse primer (biotinylated): AATACCC  
AACCCCAAATCCTAC

PCR program (50 cycles):

Initial	15 min at 95 °C
Denaturation	30 s at 94 °C
Annealing	30 s at 58 °C
Extension	30 s at 72 °C
Final	10 min at 72 °C
Hold	4 °C

Sequencing primer: AGTTGGAAGATTT  
TGTG

Target sequence: TGTGCCTACACACCG  
CCCTCACCCCTTCACYGTGGGGGCTG  
TTCTCCTCCCATGGAGYGCTCAGG  
GCTCTAGGTTCTGACTTGGGGCACC  
TCTGTCTAATCCACCAGCACAGCCA  
CTCACTATGTGCTCATCTCACTCTCC  
AGCAGCYGCTGTAGGACTTGGGGCT  
GGGCACC (**CpG target in bold**; hg19:  
chr2:25,565,782-25,565,959)

#### *DNMT3B*:

PCR forward primer: GGGGTTAAGTGGTTTAA  
GTAAAT

PCR reverse primer (biotinylated): CCTCCAAAA  
ATCCCTAAAAAATCT CTCC

PCR program (45 cycles):

Initial	15 min at 95 °C
Denaturation	30 s at 94 °C
Annealing	30 s at 52 °C
Extension	30 s at 72 °C
Final	10 min at 72 °C
Hold	4 °C

Sequencing primer: GTTAAGTGGTTTAA  
GTAAATTTAG

Target sequence: CTCGGCGATCGGCGC  
CGGAGATTCGCGAGCCCAGCGCCCTG  
CACGGCCGCCAGCCGGCCTCCCGCCA  
GCCAGCCCCGACCCGCGGCTCCGCCG  
CCCAGCCGCGCCCCAGCCAGCCCTGC  
GGCAGGTGAGCGCCCCGGGGCCC  
(**CpG targets in bold**; hg19: chr20:31,  
350,382-31,350,523)

#### *DNMT3L*:

PCR forward primer: AGTTTTTTTTATTGGGGT  
AGTTAGG

PCR reverse primer (biotinylated): CTAAAA  
ACCAAAAAACCACATTTTAT TCA

PCR program (45 cycles):

Initial	15 min at 95 °C
Denaturation	30 s at 94 °C
Annealing	30 s at 50 °C
Extension	30 s at 72 °C
Final	10 min at 72 °C
Hold	4 °C

Sequencing primer: GATTTAGGGATAGAGAG  
GG

Target sequence: GCGGTAGGGAGTGGGAAAT  
CTGAATAA (**CpG target in bold**; hg19:  
chr21:45,683,527-45,683,553)

To demonstrate the ability of our assays to resolve DNAm differences as small as reported, we computed intraclass correlation coefficients (ICCs) between triplicate replicates for each assay. Average within-sample coefficient of variation was computed using a two-way mixed model, using an absolute agreement definition (Shrout & Fleiss, 1979), as implemented in SPSS (IBM, USA). ICCs for the 15 total CpG sites assayed ranged from 0.703 to 0.937, with a mean ICC of 0.855 (s.d.=0.066). This strongly supports the conclusion that these assays are capable of consistently resolving small DNAm differences.

#### *TFBS prediction*

Putative TFBS were identified that overlap target CpG sites using the MatInspector (Cartharius et al. 2005) tool from Genomatix, with default parameters. Input

**Table 1.** Demographic and pre-trauma characteristics of 30 PTSD case-control pairs

	Controls (n=30)		PTSD (n=30)		t (df)	p
Age, years	55.37	(12.97)	53.71	(12.94)	-0.47 (29)	0.638
Female, n (%)	23	(76.7)	23	(76.7)	N.A. (1)	1.000
African-American, n (%)	22	(73.3)	24	(80)	0.01 (1)	0.938
Lifetime traumas	3.80	(3.83)	4.43	(3.70)	-0.84 (29)	0.407
Assaultive violence	0.87	(1.33)	1.07	(1.55)	-0.55 (29)	0.589
Other injury or shocking experience	0.83	(1.32)	1.27	(1.48)	-1.51 (29)	0.141
Learning about traumas to others	1.20	(1.50)	1.17	(1.37)	0.12 (29)	0.909
Sudden death	0.77	(0.43)	0.70	(0.47)	0.63 (29)	0.536
Other event	0.13	(0.35)	0.23	(0.43)	-1.36 (29)	0.184
Pre-trauma symptom severity	26.90	(10.96)	39.33	(16.20)	-1.12 (29)	0.275
Intrusion	9.58	(7.90)	11.65	(6.08)	-0.99 (29)	0.333
Avoidance	11.65	(11.04)	14.69	(6.97)	-1.14 (29)	0.267
Hyperarousal	8.60	(7.98)	11.24	(5.11)	-1.24 (29)	0.226
Post-trauma symptom severity	25.41	(7.01)	54.20	(11.59)	11.4 (29)	<0.001

Data are given as mean (standard deviation) unless otherwise indicated.

PTSD, Post-traumatic stress disorder; df, degrees of freedom; N.A., not applicable.

sequence included 200 bp up- and downstream of the CpG site. Only putative TFBS that directly overlapped CpG sites of interest were retained.

### Statistical analyses

Statistical testing was performed using SPSS Statistics for Windows, version 21.0 (IBM Corp., USA). DNAm at *DNMT3B* CpG sites was treated on a regional and an individual CpG site basis, similar to previous work (Rusiecki *et al.* 2013). Regional values were calculated as the mean of 12 CpG sites. Paired-sample *t* tests were used to test for differences in pre-trauma DNAm between cases and controls and to test for differences between pre- and post-trauma time points within cases and controls. Linear regression was used to test whether pre-trauma DNAm levels are predictive of post-trauma symptom severity (PTSS) changes. PTSS change was calculated as the difference between PTSD and pre-trauma symptom severity. Analyses included severity scores of individual symptom criteria (hyperarousal, avoidance or intrusion symptoms) as well as a total severity score that is inclusive of each symptom subdomain. Regression models were adjusted for age, gender and pre-trauma symptom severity. The contribution of pre-trauma DNAm to PTSS change was tested via the change in  $R^2$  values comparing full with reduced models. We present primary results uncorrected for multiple testing as is consistent with the current state of the science of DNAm variation in association with psychiatric endpoints (Perroud *et al.* 2011, 2013; Unternaehrer *et al.* 2012; Rusiecki *et al.* 2013). In addition, to assess the extent


to which our results may be attenuated by multiple hypothesis testing correction, we calculated stringent Bonferroni-corrected significance values (Dunn, 1961) as well as false discovery rate (FDR) Q values (Benjamini & Hochberg, 1995). FDR has recently been utilized to correct multiple hypothesis testing in studies utilizing DNAm data, with user-defined Q values ranging from 0.05 to 0.2 (Provencal *et al.* 2013; Zhao *et al.* 2013).

### Results

PTSD cases and controls did not differ in age, gender, ethnicity or pre-trauma symptom severity, including individual symptoms of intrusion, avoidance and hyperarousal (Table 1).

#### Pre-trauma DNAm variation is associated with PTSD

PTSD-associated DNAm variation may both exist before trauma and be associated with post-trauma PTSD outcome. To test for pre-existing protective/risk factors, pre-trauma DNAm at *DNMT1*, *DNMT3A*, *DNMT3B* and *DNMT3L* loci was compared between trauma-exposed individuals with versus without PTSD. Pre-trauma DNAm was higher in controls compared with cases at a single *DNMT3B* CpG site (CpG 9) [Fig. 1;  $t=2.250$ , degrees of freedom (df)=29,  $p=0.032$ ]; no difference in pre-trauma *DNMT3B* regional DNAm mean was observed ( $t=1.538$ , df=29,  $p=0.135$ ). We observed no pre-trauma differences between cases and controls at *DNMT1* ( $t=0.582$ , df=29,



**Fig. 1.** Pre-trauma DNA methyltransferase 3B (*DNMT3B*) DNA methylation (DNAm) is significantly higher in trauma-exposed controls compared with post-traumatic stress disorder (PTSD) cases at CpG 9. Pre-trauma DNAm did not differ between cases and controls at the other 11 *DNMT3B* CpG sites assessed. Values are means, with standard errors represented by vertical bars. Differences between controls and cases were tested by paired-sample *t* tests ( $n=60$ ; 30 cases and 30 matched controls).

\* Mean values were significantly different ( $p < 0.05$ ).

$p=0.565$ ), *DNMT3A* ( $t=0.579$ ,  $df=29$ ,  $p=0.567$ ) and *DNMT3L* ( $t=1.386$ ,  $df=29$ ,  $p=0.176$ ) loci.

#### Pre-trauma DNAm variation predicts post-trauma changes in trauma symptom severity

To explore whether this PTSD-associated pre-trauma DNAm is predictive of trauma response, we performed linear regression analyses with pre-trauma DNAm of *DNMT3B* at CpG 9 and PTSS change as predictor and outcome variables, respectively. Controlling for age, gender and pre-trauma symptom severity, pre-trauma DNAm of CpG 9 (Fig. 2; unstandardized  $B=-2.318$ ,  $s.e.=1.25$ ,  $p=0.034$ ) predicted PTSS change. In this model, only pre-trauma symptom severity and pre-trauma DNAm were significant predictor variables. *DNMT3B* CpG 9 DNAm explained approximately 6.8% of the variance in PTSS change, as revealed by a comparison of the full and reduced models. The full model that included *DNMT3B* CpG 9 DNAm, age, gender and pre-trauma symptom severity explained approximately 24% of the variance in post-trauma PTSS change (adjusted  $R^2=0.242$ ,  $p=0.005$ ).

Because the relationship between pre-trauma DNAm and post-trauma changes in PTSS may be driven by distinct symptom subdomains (hyperarousal, avoidance and intrusion), we regressed separately each subdomain symptom severity change onto pre-trauma DNAm, controlling for age, gender and


pre-trauma symptom severity of the relevant subdomain. Pre-trauma DNAm of *DNMT3B* CpG 9 (hyperarousal:  $p=0.249$ ; avoidance:  $p=0.137$ ; intrusion:  $p=0.071$ ) did not predict change in subdomain symptom severity.

#### Trauma induces PTSD-associated DNAm modifications

DNAm differences may arise following trauma and be associated with PTSD development. To test this, we compared pre-trauma DNAm with post-trauma DNAm within PTSD cases and within trauma-exposed, healthy controls. Both PTSD-associated and PTSD-independent changes in DNAm following trauma were observed at *DNMT* loci (Fig. 3). *DNMT1* DNAm increased (Fig. 3a;  $t=3.887$ ,  $df=29$ ,  $p=0.001$ ) following trauma in the PTSD group, but not in the control group ( $t=1.903$ ,  $df=29$ ,  $p=0.067$ ). At *DNMT3A* (Fig. 3b) and *DNMT3B* (Fig. 3c) loci, DNAm increased following trauma in both PTSD case (*DNMT3A*:  $t=2.806$ ,  $df=29$ ,  $p=0.009$ ; *DNMT3B*:  $t=4.286$ ,  $df=29$ ,  $p < 0.001$ ) and control (*DNMT3A*:  $t=3.421$ ,  $df=29$ ,  $p=0.002$ ; *DNMT3B*:  $t=3.938$ ,  $df=29$ ,  $p < 0.001$ ) groups. No change was observed in *DNMT3L* (Fig. 3d) DNAm in either cases ( $t=1.551$ ,  $df=29$ ,  $p=0.132$ ) or controls ( $t=1.146$ ,  $df=29$ ,  $p=0.261$ ). Table 2 presents a summary including uncorrected  $p$  values, Bonferroni-corrected  $p$  values and FDR values, as well as accompanying effect sizes, of our results described above.

#### TFBS prediction

DNAm is associated with gene expression. One mechanism by which increased DNAm can lead to decreased gene expression is by affecting the binding of *trans*-activating factors to *cis*-regulatory elements. To contextualize our DNAm findings, we used bioinformatic methods to identify putative TFBS that overlap CpG sites showing PTSD-associated DNAm differences. In total, we identified 24 putative TFBS, including 2, 3, 14 and 5 that overlap *DNMT1*, *DNMT3A*, *DNMT3B* and *DNMT3L* CpG target sites, respectively (Table 3). Notable among these 24 TFBS are those that overlap with CpG sites at which we identified PTSD-associated differential methylation (two overlap the *DNMT1* CpG; three overlap *DNMT3B* CpG 9). Binding sites for heat shock factor 1 and E2F-4/DP-2 heterodimeric complex were identified to overlap with the *DNMT1* CpG site at which an increase in DNAm was observed in PTSD cases, but not controls. Overlapping with *DNMT3B* CpG site 9, at which decreased pre-trauma DNAm was associated with PTSD development and predictive of worsening of PTSS, we identified binding sites for human motif



**Fig. 2.** Linear regression model of post-trauma symptom severity (PTSS) change and pre-trauma DNA methyltransferase 3B (*DNMT3B*) CpG 9 DNA methylation (DNAm), adjusting for age, gender and pre-trauma symptom severity ( $n=60$ ). Only pre-trauma symptom severity and DNAm were significant variables in this model. Values in error bar plots are means, with 95% confidence intervals represented by vertical bars. Differences between post-traumatic stress disorder (PTSD cases and trauma-exposed controls) were tested by paired-sample  $t$  tests ( $n=60$ ; 30 PTSD cases and 30 matched controls). \* Mean values were significantly different ( $p<0.05$ ).


ten element, ZF5 POZ domain zinc finger, and the insulator protein CTCF.

## Discussion

Our data represent preliminary findings suggesting that pre-trauma DNAm states and post-trauma DNAm modifications differ between those who develop PTSD following trauma and those who display resiliency. While baseline symptom severity did not differ between cases and controls, baseline DNAm at a *DNMT3B* CpG site was higher in resilient individuals compared with those who eventually developed PTSD. Additionally, longitudinal change in DNAm at a *DNMT1* CpG site was associated with PTSD, with an increase in DNAm being observed in those with PTSD but not controls. Finally, increases in DNAm were observed following trauma at *DNMT3A* and *DNMT3B* loci that were independent of PTSD outcome, being observed in both PTSD cases and trauma-exposed controls. Although some of these results were attenuated following correction for multiple hypothesis testing, our findings suggest that

epigenetic variation plays a complex regulatory role in PTSD risk and etiology.

One way in which DNAm may regulate gene transcription is by altering the strength and occupancy of transcription factor binding (Weaver *et al.* 2004). To provide insight into potential functional consequences of the observed PTSD-associated differences, we conducted a secondary analysis of TFBS overlapping the distinguishing CpG sites. Among the sites identified was a binding site for CTCF, a transcription factor known to be involved in chromatin remodeling (Barkess & West, 2012). We identified this binding site overlapping with *DNMT3B* CpG site 9, at which higher DNAm was identified as a protective/risk factor for PTSD and symptom severity change following trauma exposure. Differential methylation at this site is particularly compelling as a determinant of PTSD risk, given that DNAm at CTCF binding sites has been shown to significantly affect CTCF occupancy (Wang *et al.* 2012) and downstream levels of gene transcription (Renaud *et al.* 2007). Due to the nature of our samples, we are unable to test directly whether DNAm at these identified TFBS influences gene expression. Where available, we have utilized Encyclopedia of



**Fig. 3.** Longitudinal DNA methylation modifications of DNA methyltransferase (*DNMT*) loci in response to trauma in post-traumatic stress disorder (PTSD) cases and trauma-exposed controls. *DNMT3B* (region) represents the mean of 12 CpG sites. Values are means, with standard errors represented by vertical bars. Differences between PTSD cases and trauma-exposed controls were tested by paired-sample *t* tests ( $n=60$ ; 30 PTSD cases and 30 matched controls). Mean values were significantly different: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

DNA Elements (ENCODE) Project data (ENCODE Project Consortium, 2011) to provide evidence for or against transcription factor binding at the PTSD-associated sites in blood-derived cell types. Among the TFBS identified that overlap PTSD-associated CpG sites (*DNMT1* and *DNMT3B* CpG 9), ENCODE data include binding of CTCF and E2F4. ENCODE data support the binding of CTCF to *DNMT3B* in blood tissue (specifically B-lymphocyte cell lines: GM12864 and GM12874), but do not support the binding of E2F4 to *DNMT1*. This supports the potential functionality of observed DNAm differences at *DNMT3B* CpG 9 in pre-trauma samples in cases versus controls.

*DNMTs* have been previously implicated in PTSD, anxiety and fear conditioning. In suicide completers relative to controls, *DNMT3B* was up-regulated in the frontopolar cortex, hypothalamus and dorsal vagal complex, and down-regulated, along with *DNMT1*, in the hippocampus (Poulter et al. 2008). Additionally, *de novo* methyltransferases have been shown to be up-regulated during contextual fear conditioning, also in the hippocampus (Miller & Sweatt, 2007); *DNMTs* are required for fear conditioning and memory consolidation as demonstrated, respectively, by

administration of *DNMT* inhibitors (Miller & Sweatt, 2007) and the creation of mice with the combined knockout of *DNMT1* and *DNMT3A* (Feng et al. 2010). Our results thus add to the growing evidence implicating *DNMTs* in phenotypes of relevance to PTSD, and of psychiatric phenotypes more broadly.

The expression of *DNMTs* at the mRNA (Goto et al. 1994; Veldic et al. 2004; Kang et al. 2011; Sterner et al. 2012) and protein (Inano et al. 2000; Feng et al. 2005; Veldic et al. 2005) levels in post-mitotic neurons of the central nervous system suggests that they are involved in methyltransferase activity that persists into adulthood and that is unrelated to DNA replication (Goto et al. 1994). Indeed, previous work has identified *DNMT1* protein expression in multiple brain regions in rodents (e.g. cortex, cerebellum; Inano et al. 2000), as well as in specific cortical regions in adult humans (e.g. Brodmann area 9; Veldic et al. 2005). Furthermore, recent work suggests that our epigenetic findings in peripheral blood may be relevant to brain tissue: environmental exposures such as trauma have been shown to induce parallel epigenetic modifications in peripheral blood and the brain (McGowan et al. 2011; Klengel et al. 2013). Although the current study, based on living participants drawn from a



**Table 2.** Observed and corrected significance values of tests<sup>a</sup>

Test	Mean difference <sup>b</sup>	(S.E.)	Observed $p^c$	Rank	Bonferroni threshold	FDR thresholds
Pre- v. post-trauma <i>DNMT3B</i> in cases	1.51	(0.35)	0.000*	1	0.002*	0.002*
Pre- v. post-trauma <i>DNMT3B</i> in controls	1.51	(0.38)	0.000*	2	0.002*	0.003*
Pre- v. post-trauma <i>DNMT1</i> in cases	0.84	(0.22)	0.001*	3	0.002*	0.005*
Pre- v. post-trauma <i>DNMT3A</i> in controls	3.38	(0.99)	0.002*	4	0.002*	0.006*
Pre- v. post-trauma <i>DNMT3A</i> in cases	2.41	(0.86)	0.009*	5	0.002	0.008
Cases v. controls <i>DNMT3B</i> (CpG 9) pre-trauma	0.77	(0.34)	0.032*	6	0.002	0.009
CpG 9 regression analysis (all symptoms)	B=-2.318	(1.251)	0.034*	7	0.002	0.011
Cases v. controls <i>DNMT3B</i> (CpG 2) pre-trauma	0.19	(0.11)	0.057	8	0.002	0.013
Pre- v. post-trauma <i>DNMT1</i> in controls	0.56	(0.30)	0.067	9	0.002	0.014
CpG 9 regression analysis (intrusion symptoms)	B=-0.851	(0.458)	0.071	10	0.002	0.016
Cases v. controls <i>DNMT3B</i> (CpG 12) pre-trauma	0.44	(0.24)	0.122	11	0.002	0.017
Cases v. controls <i>DNMT3B</i> (CpG 10) pre-trauma	0.28	(0.19)	0.127	12	0.002	0.019
Pre- v. post-trauma <i>DNMT3L</i> in cases	0.74	(0.47)	0.132	13	0.002	0.020
Cases v. controls <i>DNMT3B</i> pre-trauma	0.27	(0.17)	0.135	14	0.002	0.022
CpG 9 regression analysis (avoidance symptoms)	B=-0.821	(0.541)	0.137	15	0.002	0.023
Cases v. controls <i>DNMT3B</i> (CpG 1) pre-trauma	0.35	(0.23)	0.151	16	0.002	0.025
Cases v. controls <i>DNMT3B</i> (CpG 8) pre-trauma	0.50	(0.33)	0.158	17	0.002	0.027
Cases v. controls <i>DNMT3L</i> pre-trauma	0.91	(0.65)	0.176	18	0.002	0.028
Cases v. controls <i>DNMT3B</i> (CpG 5) pre-trauma	0.25	(0.19)	0.221	19	0.002	0.030
CpG 9 regression analysis (hyperarousal symptoms)	B=-0.466	(0.399)	0.249	20	0.002	0.031
Pre- v. post-trauma <i>DNMT3L</i> in controls	0.36	(0.32)	0.261	21	0.002	0.033
Cases v. controls <i>DNMT3L</i> post-trauma	0.58	(0.56)	0.304	22	0.002	0.034
Cases v. controls <i>DNMT1</i> post-trauma	-0.19	(0.19)	0.323	23	0.002	0.036
Cases v. controls <i>DNMT3B</i> post-trauma	0.18	(0.22)	0.351	24	0.002	0.038
Cases v. controls <i>DNMT3A</i> post-trauma	0.56	(0.66)	0.356	25	0.002	0.039
Cases v. controls <i>DNMT3B</i> (CpG 7) pre-trauma	0.11	(0.12)	0.487	26	0.002	0.041
Cases v. controls <i>DNMT1</i> pre-trauma	0.09	(0.20)	0.565	27	0.002	0.042
Cases v. controls <i>DNMT3A</i> pre-trauma	-0.41	(0.73)	0.567	28	0.002	0.044
Cases v. controls <i>DNMT3B</i> (CpG 6) pre-trauma	0.11	(0.17)	0.603	29	0.002	0.045
Cases v. controls <i>DNMT3B</i> (CpG 3) pre-trauma	0.06	(0.15)	0.711	30	0.002	0.047
Cases v. controls <i>DNMT3B</i> (CpG 11) pre-trauma	0.11	(0.39)	0.773	31	0.002	0.048
Cases v. controls <i>DNMT3B</i> (CpG 4) pre-trauma	0.03	(0.14)	0.881	32	0.002	0.050

S.E., Standard error; FDR, false discovery rate; DNMT, DNA methyltransferase.

<sup>a</sup> Corrected significance thresholds at  $p < 0.05$  are listed using two controlling procedures: Bonferroni and FDR using the procedure of Benjamini & Hochberg (1995).

<sup>b</sup> In percent DNA methylation. For regression analyses, 'B' represents unstandardized  $\beta$  values.

<sup>c</sup> The list of observed  $p$  values is sorted from smallest to largest (indicated by the rank column).

\* Values meet significance at  $p < 0.05$  for the various correction procedures.

population-based cohort, precludes such work, future research is needed to address whether the epigenetic determinants of risk observed here in peripheral blood-derived DNA are also found in brain-derived DNA.

Importantly, this study adds to emerging work utilizing a longitudinal study design capable of measuring biological markers prior to disease onset as well as change between pre-disease and post-disease time points (Rusiecki *et al.* 2012, 2013; Perroud *et al.* 2013; Nieratschker *et al.* 2014). Existing longitudinal studies have documented the importance of DNAm to mental health disorder risk, including differential change in

DNAm of *BDNF* among individuals with *versus* without borderline personality disorder (Perroud *et al.* 2013), increased *DAT* (*SLC6A3*) DNAm with age that may be driven by alcohol dependence (Nieratschker *et al.* 2014), and increasing *SERT* DNAm associated with bullying (Ouellet-Morin *et al.* 2013). Most relevant to the present study is work by Rusiecki *et al.* (2012) which provides evidence for increased global DNAm in controls, but not cases, following trauma exposure, suggesting that resiliency is associated with increased global DNAm, potentially mediated by increased activity and expression of DNMTs. Indeed, our data

**Table 3.** Putative transcription factor binding sites overlap DNMT CpG sites of interest<sup>a</sup>

Gene	Matrix family	Matrix information	Core similarity	DNMT3B CpG overlap	
DNMT1	V\$HEAT	Heat shock factor 1	1.000		
	V\$E2FF	E2F-4/DP-2 heterodimeric complex	0.847	–	
DNMT3A	V\$SP1F	TGFbeta-inducible early gene (TIEG)/early growth response gene alpha (EGRalpha)	0.750	–	
	V\$SP1F	Stimulating protein 1, ubiquitous zinc finger transcription factor	1.000	–	
	V\$EGRF	EGR1, early growth response 1	0.802	–	
	V\$GCMF	Glial cells missing homolog 1, chorion-specific transcription factor GCMa	1.000	–	
	V\$KLFS	Kidney-enriched kruppel-like factor, KLF15	1.000	–	
DNMT3B	O\$MTEN	Human motif ten element	0.839	1, 2, 3, 4, 5	
	V\$PAX5	PAX5 paired domain protein	0.789	1, 2, 3, 4, 5, 6, 7	
	V\$E2FF	E2F transcription factor 3 (secondary DNA binding preference)	1.000	2, 3, 4, 5	
	V\$E2FF	E2F transcription factor 3 (secondary DNA binding preference)	1.000	2, 3, 4, 5, 6, 7	
	V\$ETSF	Ets variant 4	1.000	3, 4, 5, 6, 7	
	O\$MTEN	Human motif ten element	0.961	7, 8, 9	
	V\$ZF5F	ZF5 POZ domain zinc finger, zinc finger protein 161 (secondary DNA binding preference)	0.775	8, 9, 10	
	V\$CTCF	Insulator protein CTCF (CCCTC-binding factor)	0.818	9, 10, 11, 12	
	V\$HDBP	Huntington's disease gene regulatory region-binding protein 1 and 2 (SLC2A4 regulator and papillomavirus binding factor)	1.000	10, 11, 12	
	V\$EGRF	Collagen krox protein (zinc finger protein 67-zfp67)	1.000	11, 12	
	V\$PLAG	Pleomorphic adenoma gene (PLAG) 1, a developmentally regulated C2H2 zinc finger protein	0.958	12	
	V\$ZF02	Transcriptional repressor, binds to elements found predominantly in genes that participate in lipid metabolism	0.776	12	
	DNMT3L	V\$KLFS	Basic transcription element (BTE) binding protein, BTEB3, FKLF-2	1.000	–
		V\$SP1F	Stimulating protein 1, ubiquitous zinc finger transcription factor	1.000	–
V\$MYBL		C-Myb, important in hematopoiesis, cellular equivalent to avian myoblastosis virus oncogene v-myb	0.797	–	
V\$GLIF		Zinc finger transcription factor, Zic family member 2 (odd-paired homolog, Drosophila)	1.000	–	
V\$CP2F		LBP-1c (leader-binding protein-1c), LSF (late SV40 factor, CP2, SEF (SAA3 enhancer factor)	0.875	–	

DNMT, DNA methyltransferase.

<sup>a</sup> V\$ matrix families indicate Genomatix-annotated transcription factor binding site matrix families. DNMT3B CpG sites are described in the Method section.

presented here are consistent with this scenario, as DNAm of *DNMT1* was observed to increase following trauma in cases, but not controls. In contrast, however, we observed an increase in *DNMT3B* DNAm following trauma in both cases and controls, and a pre-trauma association between higher DNAm pre-trauma and resiliency post-trauma. The presence of a CTCF binding site opens the possibility that increased DNAm at this locus is associated with increased gene expression because CTCF can act as either a transcriptional activator or repressor (Phillips & Corces, 2009), with strength of DNA binding inversely correlated with local DNAm (Barkess & West, 2012). If binding of CTCF to the *DNMT3B* locus results in transcriptional repression, then increased DNAm, and concurrent decreased CTCF binding, would be associated with increased, not decreased, gene expression. If true, this would put these findings in line with the previously published, longitudinal, trauma-associated epigenetic data: decreased DNAm in pre-trauma PTSD cases would result in tighter CTCF binding and reduced *DNMT3B* transcription and lower global DNAm levels, as reported by Ruisecki *et al.* (2012). Although DNMT1 is typically thought to maintain DNAm in adult tissues, evidence suggests that DNMT1 and DNMT3B cooperatively maintain DNAm, with one or the other, but not both, required for global DNAm (Rhee *et al.* 2002). More broadly, our data add to the emerging evidence that longitudinal DNAm changes may contribute to the etiology of mental illness and can be taken as a proof of principle that locus-specific epigenetic variability both pre-exists and arises following disease onset in biologically meaningful ways.

While our study is one of the first of its kind to compare pre- and post-trauma DNAm levels with regard to the development of PTSD, there is a minimum of four study limitations that should be kept in mind when interpreting our results. First, it is important to recognize that the epidemiological nature of our cohort precludes sample collection with a well-controlled experimental time course; times between pre-trauma data collection, trauma exposure and post-trauma data collection differed between each test subject. As such, we are unable to resolve whether observed PTSD-associated post-trauma DNAm changes precede PTSD development (i.e. occurred within the first 4 weeks following trauma). As DNMTs are involved in the global regulation of DNAm, it is tempting to conclude from our data that observed changes in *DNMT* DNAm are an upstream process of PTSD development, thereby having the potential to help explain differences in DNAm epigenome-wide reported elsewhere (Uddin *et al.* 2010; Smith *et al.* 2011; Rusiecki *et al.* 2012). However, it is also possible that

the observed PTSD-associated DNAm changes are downstream effects of PTSD development, with no or little involvement in epigenetic modifications across the epigenome. Second, the nature of the epidemiological samples collected precluded the assessment of pre- and post-trauma gene expression differences and changes; furthermore, we did not collect data on blood cell composition. Third, the DNAm differences and effect sizes reported here are small; however, they are consistent with published work showing functional effects of DNAm variation (Tyrka *et al.* 2012). High ICCs between experimental replicates for each of our assays increase confidence of the validity of observed DNAm differences. Indeed, our sample size and observed effect sizes are consistent with published work in the field (Perroud *et al.* 2011; Byrne *et al.* 2013). Fourth, our primary results presented in this work are not corrected for multiple testing. Although this is consistent with the current state of the science of DNAm variation in association with psychiatric endpoints (Perroud *et al.* 2011; Unternaehrer *et al.* 2012; Perroud *et al.* 2013; Rusiecki *et al.* 2013), we do report corrected results (Table 2) to assess the degree to which our findings might be attenuated by multiple hypothesis test correction. Accepting a stringent FDR of 0.05 requires that we reject several findings reported as significant in our study, notably pre-trauma DNAm differences between cases and controls at *DNMT3B* CpG 9. However, it also means that a significant association between DNAm and PTSD emerges as a result of correction, as a significant change in DNAm at *DNMT3A* following trauma is only seen in controls at this stringent FDR cut-off and would therefore be suggestive of a resiliency-associated change in DNAm (Table 2). While we have chosen to utilize a stringent FDR cut-off of 0.05, other DNAm analyses have accepted a cut-off as high as 0.20 (Provencal *et al.* 2013). Overall, we stress the preliminary nature of these findings – both uncorrected and corrected for multiple hypothesis testing – and the importance of replication in an independent cohort.

Individuals exposed to trauma differ in their risk for subsequent PTSD. Our data suggest that variation in pre-trauma DNAm and post-trauma DNAm change may be part of the molecular underpinnings of PTSD risk and resiliency. Future research is needed to determine if the DNAm variation observed here is associated with functional changes that affect the long-term biology of individuals exposed to trauma. The identification of risk markers, including epigenetic markers, is an important step to understanding the biological underpinnings of PTSD risk and may lead to the development of tools to identify those individuals most at risk of developing PTSD as well as to develop evidence-based interventions.

## Acknowledgements

We thank Dr Chet Sherwood and Amy Bauernfeind from The George Washington University for contributions to this work. We thank the many Detroit residents who chose to participate in the DNHS.

This research was funded by National Institutes of Health grants (3R01DA022720-05, 3R01DA022720 and 1RC1MH088283-01) and a National Science Foundation grant (BCS-0827546). L.S. was funded by a Graduate Research Assistantship from the Wayne State University Office of the Vice President for Research, and a Grant-in-Aid of Research from Sigma Xi. K.C.K. is funded by R01MH093612, RC4MH092707, P51RR000165, U01OH010416 and U01OH010407.

## Declaration of Interest

S.G. is funded in part by a grant from Merck Pharmaceuticals for work unrelated to this project. D.E.W. is funded in part from Lung, LLC for work unrelated to this project and also reports receiving honoraria from Elsevier, Inc. A.E.A. consults for SCA Tork for work unrelated to this study.

## References

- Aberg KA, Xie LY, Nerella S, Copeland WE, Costello EJ, van den Oord EJ (2013). High quality methylome-wide investigations through next-generation sequencing of DNA from a single archived dry blood spot. *Epigenetics* **8**, 542–547.
- Acierno R, Ruggiero KJ, Galea S, Resnick HS, Koenen K, Roitzsch J, de Arellano M, Boyle J, Kilpatrick DG (2007). Psychological sequelae resulting from the 2004 Florida hurricanes: implications for postdisaster intervention. *American Journal of Public Health* **97** (Suppl. 1), S103–S108.
- Andrews JA, Neises KD (2012). Cells, biomarkers, and post-traumatic stress disorder: evidence for peripheral involvement in a central disease. *Journal of Neurochemistry* **120**, 26–36.
- APA (1994). *Diagnostic and Statistical Manual of Mental Disorders, 4th edn* (DSM-IV). American Psychiatric Association: Washington, DC.
- APA (2013). *Diagnostic and Statistical Manual of Mental Disorders, 5th edn*. (DSM-5). American Psychiatric Publishing: Arlington, VA.
- Barkess G, West AG (2012). Chromatin insulator elements: establishing barriers to set heterochromatin boundaries. *Epigenomics* **4**, 67–80.
- Benjamini Y, Hochberg Y (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B* **57**, 289–300.
- Binder EB, Bradley RG, Liu W, Epstein MP, Deveau TC, Mercer KB, Tang Y, Gillespie CF, Heim CM, Nemeroff CB, Schwartz AC, Cubells JF, Ressler KJ (2008). Association of FKBP5 polymorphisms and childhood abuse with risk of posttraumatic stress disorder symptoms in adults. *Journal of the American Medical Association* **299**, 1291–1305.
- Bird A (2002). DNA methylation patterns and epigenetic memory. *Genes and Development* **16**, 6–21.
- Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Gusman FD, Charney DS, Keane TM (1995). The development of a clinician-administered PTSD scale. *Journal of Traumatic Stress* **8**, 75–90.
- Bourc'his D, Bestor TH (2004). Meiotic catastrophe and retrotransposon reactivation in male germ cells lacking Dnmt3L. *Nature* **431**, 96–99.
- Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH (2001). Dnmt3L and the establishment of maternal genomic imprints. *Science* **294**, 2536–2539.
- Breslau N, Davis GC, Andreski P, Peterson E (1991). Traumatic events and posttraumatic stress disorder in an urban population of young adults. *Archives of General Psychiatry* **48**, 216–222.
- Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P (1998). Trauma and posttraumatic stress disorder in the community: the 1996 Detroit Area Survey of Trauma. *Archives of General Psychiatry* **55**, 626–632.
- Breslau N, Wilcox HC, Storr CL, Lucia VC, Anthony JC (2004). Trauma exposure and posttraumatic stress disorder: a study of youths in urban America. *Journal of Urban Health* **81**, 530–544.
- Brewin CR, Andrews B, Valentine JD (2000). Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *Journal of Consulting and Clinical Psychology* **68**, 317–336.
- Byrne EM, Carrillo-Roa T, Henders AK, Bowdler L, McRae AF, Heath AC, Martin NG, Montgomery GW, Krause L, Wray NR (2013). Monozygotic twins affected with major depressive disorder have greater variance in methylation than their unaffected co-twin. *Translational Psychiatry* **3**, e269.
- Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, Frisch M, Bayerlein M, Werner T (2005). MatInspector and beyond: promoter analysis based on transcription factor binding sites. *Bioinformatics* **21**, 2933–2942.
- Chang SC, Koenen KC, Galea S, Aiello AE, Soliven R, Wildman DE, Uddin M (2012). Molecular variation at the SLC6A3 locus predicts lifetime risk of PTSD in the Detroit Neighborhood Health Study. *PLOS ONE* **7**, e39184.
- Congressionally Directed Medical Research Programs Department of Defense (2010). *CDMRP Department of Defense Psychological Health/Traumatic Brain Injury Research Program*.
- DiGrande L, Perrin MA, Thorpe LE, Thalji L, Murphy J, Wu D, Farfel M, Brackbill RM (2008). Posttraumatic stress symptoms, PTSD, and risk factors among lower Manhattan residents 2–3 years after the September 11, 2001 terrorist attacks. *Journal of Traumatic Stress* **21**, 264–273.

- Dunn OJ (1961). Multiple comparisons among means. *Journal of the American Statistical Association* **56**, 52–64.
- El-Sayed AM, Haloossim MR, Galea S, Koenen KC (2012). Epigenetic modifications associated with suicide and common mood and anxiety disorders: a systematic review of the literature. *Biology of Mood and Anxiety Disorders* **2**, 10.
- ENCODE Project Consortium (2011). A user's guide to the encyclopedia of DNA elements (ENCODE). *PLoS Biology* **9**, e1001046.
- Feinberg AP (2008). Epigenetics at the epicenter of modern medicine. *Journal of the American Medical Association* **299**, 1345–1350.
- Feng J, Chang H, Li E, Fan G (2005). Dynamic expression of *de novo* DNA methyltransferases Dnmt3a and Dnmt3b in the central nervous system. *Journal of Neuroscience Research* **79**, 734–746.
- Feng J, Fan G (2009). The role of DNA methylation in the central nervous system and neuropsychiatric disorders. *International Review of Neurobiology* **89**, 67–84.
- Feng J, Zhou Y, Campbell SL, Le T, Li E, Sweatt JD, Silva AJ, Fan G (2010). Dnmt1 and Dnmt3a maintain DNA methylation and regulate synaptic function in adult forebrain neurons. *Nature Neuroscience* **13**, 423–430.
- Galea S, Ahern J, Tracy M, Hubbard A, Cerda M, Goldmann E, Vlahov D (2008). Longitudinal determinants of posttraumatic stress in a population-based cohort study. *Epidemiology* **19**, 47–54.
- Goto K, Numata M, Komura JI, Ono T, Bestor TH, Kondo H (1994). Expression of DNA methyltransferase gene in mature and immature neurons as well as proliferating cells in mice. *Differentiation* **56**, 39–44.
- Hollegaard MV, Grauholm J, Norgaard-Pedersen B, Hougaard DM (2013). DNA methylome profiling using neonatal dried blood spot samples: a proof-of-principle study. *Molecular Genetics and Metabolism* **108**, 225–231.
- Inano K, Suetake I, Ueda T, Miyake Y, Nakamura M, Okada M, Tajima S (2000). Maintenance-type DNA methyltransferase is highly expressed in post-mitotic neurons and localized in the cytoplasmic compartment. *Journal of Biochemistry* **128**, 315–321.
- Institute of Medicine (2012). *Treatment for Posttraumatic Stress Disorder in Military and Veteran Populations: Initial Assessment*. National Academy of Sciences: Washington, DC.
- Kaneda M, Okano M, Hata K, Sado T, Tsujimoto N, Li E, Sasaki H (2004). Essential role for *de novo* DNA methyltransferase Dnmt3a in paternal and maternal imprinting. *Nature* **429**, 900–903.
- Kang HJ, Kawasawa YI, Cheng F, Zhu Y, Xu X, Li M, Sousa AM, Pletikos M, Meyer KA, Sedmak G, Guennel T, Shin Y, Johnson MB, Krsnik Z, Mayer S, Fertuzinhos S, Umlauf S, Lisgo SN, Vortmeyer A, Weinberger DR, Mane S, Hyde TM, Huttner A, Reimers M, Kleinman JE, Sestan N (2011). Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483–489.
- Kato Y, Kaneda M, Hata K, Kumaki K, Hisano M, Kohara Y, Okano M, Li E, Nozaki M, Sasaki H (2007). Role of the Dnmt3 family in *de novo* methylation of imprinted and repetitive sequences during male germ cell development in the mouse. *Human Molecular Genetics* **16**, 2272–2280.
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry* **52**, 1048–1060.
- Kessler RC, Wang PS (2008). The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annual Review of Public Health* **29**, 115–129.
- Klengel T, Mehta D, Anacker C, Rex-Haffner M, Pruessner JC, Pariante CM, Pace TW, Mercer KB, Mayberg HS, Bradley B, Nemeroff CB, Holsboer F, Heim CM, Ressler KJ, Rein T, Binder EB (2013). Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nature Neuroscience* **16**, 33–41.
- Koenen KC, Harley R, Lyons MJ, Wolfe J, Simpson JC, Goldberg J, Eisen SA, Tsuang M (2002). A twin registry study of familial and individual risk factors for trauma exposure and posttraumatic stress disorder. *Journal of Nervous and Mental Disease* **190**, 209–218.
- Koenen KC, Stellman JM, Stellman SD, Sommer JF Jr. (2003). Risk factors for course of posttraumatic stress disorder among Vietnam veterans: a 14-year follow-up of American legionnaires. *Journal of Consulting and Clinical Psychology* **71**, 980–986.
- Kulka R, Schlenger W, Fairbank J, Hough R, Jordan BK, Marmar C, Weiss D, Grady D (1990). *Trauma and the Vietnam War Generation: Report of the Findings from the National Vietnam Veterans Readjustment Study*. Brunner/Mazel: New York.
- Kun P, Han S, Chen X, Yao L (2009). Prevalence and risk factors for posttraumatic stress disorder: a cross-sectional study among survivors of the Wenchuan 2008 earthquake in China. *Depression and Anxiety* **26**, 1134–1140.
- Li E, Bestor TH, Jaenisch R (1992). Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* **69**, 915–926.
- Logue MW, Baldwin C, Guffanti G, Melista E, Wolf EJ, Reardon AF, Uddin M, Wildman D, Galea S, Koenen KC, Miller MW (2013). A genome-wide association study of post-traumatic stress disorder identifies the retinoid-related orphan receptor alpha (RORA) gene as a significant risk locus. *Molecular Psychiatry* **18**, 937–942.
- Lu AT, Ogdie MN, Jarvelin MR, Moilanen IK, Loo SK, McCracken JT, McGough JJ, Yang MH, Peltonen L, Nelson SF, Cantor RM, Smalley SL (2008). Association of the cannabinoid receptor gene (CNR1) with ADHD and post-traumatic stress disorder. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics Part B* **147B**, 1488–1494.
- McGowan PO, Suderman M, Sasaki A, Huang TC, Hallett M, Meaney MJ, Szyf M (2011). Broad epigenetic signature of maternal care in the brain of adult rats. *PLoS ONE* **6**, e14739.
- Meaney MJ (2010). Epigenetics and the biological definition of gene × environment interactions. *Child Development* **81**, 41–79.

- Miller CA, Sweatt JD (2007). Covalent modification of DNA regulates memory formation. *Neuron* **53**, 857–869.
- Nieratschker V, Grosshans M, Frank J, Strohmaier J, von der Goltz C, El-Maarri O, Witt SH, Cichon S, Nöthen MM, Kiefer F, Rietschel M (2014). Epigenetic alteration of the dopamine transporter gene in alcohol-dependent patients is associated with age. *Addiction Biology* **19**, 305–311.
- National Institute of Mental Health (2008). *National Institute of Mental Health Strategic Plan*. National Institutes of Health: Bethesda.
- Ooi SK, Qiu C, Bernstein E, Li K, Jia D, Yang Z, Erdjument-Bromage H, Tempst P, Lin SP, Allis CD, Cheng X, Bestor TH (2007). DNMT3L connects unmethylated lysine 4 of histone H3 to *de novo* methylation of DNA. *Nature* **448**, 714–717.
- Ouellet-Morin I, Wong CC, Danese A, Pariante CM, Papadopoulos AS, Mill J, Arseneault L (2013). Increased serotonin transporter gene (SERT) DNA methylation is associated with bullying victimization and blunted cortisol response to stress in childhood: a longitudinal study of discordant monozygotic twins. *Psychological Medicine* **43**, 1813–1823.
- Perroud N, Paoloni-Giacobino A, Prada P, Olie E, Salzmann A, Nicastro R, Guillaume S, Mouthon D, Stouder C, Dieben K, Huguelet P, Courtet P, Malafosse A (2011). Increased methylation of glucocorticoid receptor gene (*NR3C1*) in adults with a history of childhood maltreatment: a link with the severity and type of trauma. *Translational Psychiatry* **1**, e59.
- Perroud N, Salzmann A, Prada P, Nicastro R, Hoeppli ME, Furrer S, Ardu S, Krejci I, Karege F, Malafosse A (2013). Response to psychotherapy in borderline personality disorder and methylation status of the *BDNF* gene. *Translational Psychiatry* **3**, e207.
- Phillips JE, Corces VG (2009). CTCF: master weaver of the genome. *Cell* **137**, 1194–1211.
- Poulter MO, Du L, Weaver IC, Palkovits M, Faludi G, Merali Z, Szyf M, Anisman H (2008). GABAA receptor promoter hypermethylation in suicide brain: implications for the involvement of epigenetic processes. *Biological Psychiatry* **64**, 645–652.
- Provencal N, Suderman MJ, Caramaschi D, Wang D, Hallett M, Vitaro F, Tremblay RE, Szyf M (2013). Differential DNA methylation regions in cytokine and transcription factor genomic loci associate with childhood physical aggression. *PLOS ONE* **8**, e71691.
- Reik W, Dean W, Walter J (2001). Epigenetic reprogramming in mammalian development. *Science* **293**, 1089–1093.
- Renaud S, Loukinov D, Abdullaev Z, Guilleret I, Bosman FT, Lobanekov V, Benhattar J (2007). Dual role of DNA methylation inside and outside of CTCF-binding regions in the transcriptional regulation of the telomerase hTERT gene. *Nucleic Acids Research* **35**, 1245–1256.
- Ressler KJ, Mercer KB, Bradley B, Jovanovic T, Mahan A, Kerley K, Norrholm SD, Kilaru V, Smith AK, Myers AJ, Ramirez M, Engel A, Hammack SE, Toufexis D, Braas KM, Binder EB, May V (2011). Post-traumatic stress disorder is associated with PACAP and the PAC1 receptor. *Nature* **470**, 492–497.
- Rhee I, Bachman KE, Park BH, Jair KW, Yen RW, Schuebel KE, Cui H, Feinberg AP, Lengauer C, Kinzler KW, Baylin SB, Vogelstein B (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* **416**, 552–556.
- Robertson KD, Uzvolgyi E, Liang G, Talmadge C, Sumegi J, Gonzales FA, Jones PA (1999). The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Research* **27**, 2291–2298.
- Rusiecki JA, Byrne C, Galdzicki Z, Srikantan V, Chen L, Poulin M, Yan L, Baccarelli A (2013). PTSD and DNA methylation in select immune function gene promoter regions: a repeated measures case–control study of U.S. military service members. *Frontiers in Psychiatry* **4**, 56.
- Rusiecki JA, Chen L, Srikantan V, Zhang L, Yan L, Polin ML, Baccarelli A (2012). DNA methylation in repetitive elements and post-traumatic stress disorder: a case–control study of US military service members. *Epigenomics* **4**, 29–40.
- Seisenberger S, Peat JR, Hore TA, Santos F, Dean W, Reik W (2013). Reprogramming DNA methylation in the mammalian life cycle: building and breaking epigenetic barriers. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368**, 20110330.
- Shrout PE, Fleiss JL (1979). Intraclass correlations: uses in assessing rater reliability. *Psychological Bulletin* **86**, 420–428.
- Siegmund KD, Connor CM, Campan M, Long TI, Weisenberger DJ, Biniszkiwicz D, Jaenisch R, Laird PW, Akbarian S (2007). DNA methylation in the human cerebral cortex is dynamically regulated throughout the life span and involves differentiated neurons. *PLoS ONE* **2**, e895.
- Smith AK, Conneely KN, Kilaru V, Mercer KB, Weiss TE, Bradley B, Tang Y, Gillespie CF, Cubells JF, Ressler KJ (2011). Differential immune system DNA methylation and cytokine regulation in post-traumatic stress disorder. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* **156**, 700–708.
- Stein MB, Jang KL, Taylor S, Vernon PA, Livesley WJ (2002). Genetic and environmental influences on trauma exposure and posttraumatic stress disorder symptoms: a twin study. *American Journal of Psychiatry* **159**, 1675–1681.
- Sterner KN, Weckle A, Chugani HT, Tarca AL, Sherwood CC, Hof PR, Kuzawa CW, Boddy AM, Abbas A, Raaum RL, Gregoire L, Lipovich L, Grossman LI, Uddin M, Goodman M, Wildman DE (2012). Dynamic gene expression in the human cerebral cortex distinguishes children from adults. *PLOS ONE* **7**, e37714.
- Toyokawa S, Uddin M, Koenen KC, Galea S (2012). How does the social environment ‘get into the mind’? Epigenetics at the intersection of social and psychiatric epidemiology. *Social Science and Medicine* **74**, 67–74.

- True WR, Rice J, Eisen SA, Heath AC, Goldberg J, Lyons MJ, Nowak J** (1993). A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Archives of General Psychiatry* **50**, 257–264.
- Tyrka AR, Price LH, Marsit C, Walters OC, Carpenter LL** (2012). Childhood adversity and epigenetic modulation of the leukocyte glucocorticoid receptor: preliminary findings in healthy adults. *PLoS ONE* **7**, e30148.
- Uddin M, Aiello AE, Wildman DE, Koenen KC, Pawelec G, de Los Santos R, Goldmann E, Galea S** (2010). Epigenetic and immune function profiles associated with posttraumatic stress disorder. *Proceedings of the National Academy of Sciences USA* **107**, 9470–9475.
- Uddin M, Chang SC, Zhang C, Ressler K, Mercer KB, Galea S, Keyes KM, McLaughlin KA, Wildman DE, Aiello AE, Koenen KC** (2013). Adcyap1r1 genotype, posttraumatic stress disorder, and depression among women exposed to childhood maltreatment. *Depression and Anxiety* **30**, 251–258.
- Unternaehrer E, Luers P, Mill J, Dempster E, Meyer AH, Staehli S, Lieb R, Hellhammer DH, Meinlschmidt G** (2012). Dynamic changes in DNA methylation of stress-associated genes (*OXR*, *BDNF*) after acute psychosocial stress. *Translational Psychiatry* **2**, e150.
- Veldic M, Caruncho HJ, Liu WS, Davis J, Satta R, Grayson DR, Guidotti A, Costa E** (2004). DNA-methyltransferase 1 mRNA is selectively overexpressed in telencephalic GABAergic interneurons of schizophrenia brains. *Proceedings of the National Academy of Sciences USA* **101**, 348–353.
- Veldic M, Guidotti A, Maloku E, Davis JM, Costa E** (2005). In psychosis, cortical interneurons overexpress DNA-methyltransferase 1. *Proceedings of the National Academy of Sciences USA* **102**, 2152–2157.
- Wang H, Maurano MT, Qu H, Varley KE, Gertz J, Pauli F, Lee K, Canfield T, Weaver M, Sandstrom R, Thurman RE, Kaul R, Myers RM, Stamatoyannopoulos JA** (2012). Widespread plasticity in CTCF occupancy linked to DNA methylation. *Genome Research* **22**, 1680–1688.
- Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M, Meaney MJ** (2004). Epigenetic programming by maternal behavior. *Nature Neuroscience* **7**, 847–854.
- Wong N, Morley R, Saffery R, Craig J** (2008). Archived Guthrie blood spots as a novel source for quantitative DNA methylation analysis. *Biotechniques* **45**, 423–424, 426, 428 passim.
- Xie P, Kranzler HR, Poling J, Stein MB, Anton RF, Farrer LA, Gelernter J** (2010). Interaction of FKBP5 with childhood adversity on risk for post-traumatic stress disorder. *Neuropsychopharmacology* **35**, 1684–1692.
- Zhao J, Goldberg J, Bremner JD, Vaccarino V** (2013). Association between promoter methylation of serotonin transporter gene and depressive symptoms: a monozygotic twin study. *Psychosomatic Medicine* **75**, 523–529.