

Original Article

Cite this article: Fielding-Smith SF, Greenwood KE, Wichers M, Peters E, Hayward M (2022). Associations between responses to voices, distress and appraisals during daily life: an ecological validation of the cognitive behavioural model. *Psychological Medicine* 52, 538–547. <https://doi.org/10.1017/S0033291720002238>

Received: 17 January 2020

Revised: 6 June 2020

Accepted: 7 June 2020

First published online: 10 July 2020


Key words:

Appraisals; auditory verbal hallucinations; cognitive model; experience sampling; distress; responses; voice hearing

Author for correspondence:

Mark Hayward,
E-mail: m.i.hayward@sussex.ac.uk

Associations between responses to voices, distress and appraisals during daily life: an ecological validation of the cognitive behavioural model

Sarah F. Fielding-Smith¹, Kathryn E. Greenwood^{1,2}, Marieke Wichers³, Emmanuelle Peters^{4,5} and Mark Hayward^{1,2} 

¹School of Psychology, University of Sussex, Brighton, UK; ²Research & Development Department, Sussex Partnership NHS Foundation Trust, Hove, UK; ³University of Groningen, University Medical Centre of Groningen, Department of Psychiatry, Interdisciplinary Center Psychopathology and Emotion Regulation (ICPE), University Medical Centre of Groningen, Groningen, the Netherlands; ⁴Department of Psychology, Institute of Psychiatry, Psychology & Neuroscience, Kings College London, London, UK and ⁵South London and Maudsley NHS Foundation Trust, Bethlem Royal Hospital, Monks Orchard Road, Beckenham, Kent, BR3 3BX, UK

Abstract

Background. Cognitive models propose that behavioural responses to voices maintain distress by preventing disconfirmation of negative beliefs about voices. We used Experience Sampling Methodology (ESM) to examine the hypothesized maintenance role of behavioural responses during daily life.

Method. Thirty-one outpatients with frequent voices completed a smartphone-based ESM questionnaire 10 times a day over 9 days, assessing voice-related distress; resistance and compliance responses to voices; voice characteristics (intensity and negative content); appraisals of voice dominance, uncontrollability and intrusiveness.

Results. In line with predictions, behavioural responses were associated with voice appraisals (dominance and uncontrollability), but not voice characteristics. Greater resistance and compliance were reported in moments of increased voice distress, but these associations did not persist after controlling for concurrent voice appraisals and characteristics. Voice distress was predicted by appraisals, and, unexpectedly, also by voice characteristics. As predicted, compliance and resistance were related to increases in distress at subsequent timepoints, whilst antecedent voice appraisals and characteristics had no such effect. Compliance, but not resistance, additionally predicted subsequent increases in voice uncontrollability. In both cases, the reverse models showed no association, indicating directional effects of responses on subsequent distress, and of compliance on uncontrollability appraisals.

Conclusions. These results provide support for the cognitive model by suggesting that momentary behavioural and emotional responses to voices are associated with concurrent negative voice appraisals. Findings suggest that behavioural responses may be driven by voice appraisals, rather than directly by distress, and may in turn maintain voice appraisals and associated distress during the course of daily life.

Introduction

Voice hearing experiences, or auditory verbal hallucinations, are typically defined in relation to their perceptual characteristics (David, 2004). However, such definitions belie the fact that voice hearers are typically not passive bystanders of these experiences (Beavan, 2011; Nayani & David, 1996). Voice hearers commonly report being drawn in to reacting or responding to their voices (Thomas, Morris, Shawyer, & Farhall, 2013), either via direct and reciprocal acts of communication (Hayward et al., 2011), or via the use of actions to mitigate their activation or negative impact (Farhall, Greenwood, & Jackson, 2007).

The cognitive model proposes that these behavioural responses are driven primarily by the beliefs a person holds about their voices (Chadwick & Birchwood, 1994). Voices are typically perceived as powerful, intrusive beings with malevolent intent towards the hearer or others, over whom the hearer has little control or ability to escape. They are suggested to evoke innate evolved defences of fight, flight or submission, similar to those observed in real-world social interactions (Gilbert et al., 2001), as a means of mitigating perceived threat (Morrison, 1998). A number of studies have demonstrated that efforts to resist voices (by arguing back, avoiding cues that trigger voices, or employing distraction tactics), and attempts to appease the perceived agent of the voices by complying with voice commands, are common responses to voices (Chaix et al., 2014; Hacker, Birchwood, Tudway, Meaden, & Amphlett, 2008;

Howard, Forsyth, Spencer, Young, & Turkington, 2013). Furthermore, both resistance and compliance are predicted by perceptions of voice dominance (Birchwood et al., 2004; Gilbert et al., 2001; Hayward et al., 2008; Reynold & Scragg, 2010) and intrusiveness (Hayward et al., 2008; Mackinnon, Copolov, & Trauer, 2004), whilst resistance (to command hallucinations, and voices more generally) is additionally associated with perceived voice malevolence (Birchwood et al., 2004; Hayward, 2003; Peters, Williams, Cooke, & Kuipers, 2012b; van der Gaag, Hageman, & Birchwood, 2003; Vaughan & Fowler, 2004).

Behavioural responses to voices are a central target of cognitive behaviour therapy for psychosis (CBTp) (Morrison & Barratt, 2010), due to their potentially immediate distressing or harmful effects. These harmful effects are most evident in the case of compliance with voice commands, which can pose significant risks of danger to self and others, with an estimated 30% of hearers reporting at least partial compliance with harmful voice commands (Shawyer, Mackinnon, Farhall, Trauer, & Copolov, 2003). Cognitive models further suggest that, rather than simply representing a consequence of voice beliefs, behavioural responses in turn play a role in maintaining voices and associated distress and disability (Chadwick & Birchwood, 1994; Morrison, 1998). In the case of resistance, Morrison (1998) suggested that efforts to resist voices may be counterproductive, serving to increase voice frequency in a manner similar to the demonstrated effects of thought-suppression on the frequency of intrusive thoughts (Abramowitz, Tolin, & Street, 2001). Morrison also proposed that resistance and compliance should be conceptualized as 'safety behaviours' (Salkovskis, 1991); compensatory actions that afford short-term relief, but contribute to the longer-term maintenance of voice distress, by preventing opportunities for disconfirmation of negative voice beliefs (Michail & Birchwood, 2010).

It has indeed been demonstrated that whilst voice hearers typically perceive their responses as being effective in reducing the sense of immediate threat from voices (Hacker et al., 2008), there is a positive association between levels of voice-related distress and resistance, avoidance (Farhall & Gehrke, 1997; Hayward et al., 2008; Vaughan & Fowler, 2004) and compliance (Hacker et al., 2008). Hacker et al. (2008) demonstrated that the positive association between safety behaviour use and distress is mediated by perceived voice omnipotence, consistent with the notion that safety-seeking behaviours exert their effect on distress by preventing disconfirmation of threat. Furthermore, randomized controlled trials (RCTs) of cognitive therapy for command hallucinations (COMMAND), which aim to change omnipotence beliefs through the use of behavioural experiments to test the consequences of resisting commands (Meaden, Keen, Aston, Barton, & Bucci, 2013), have demonstrated reductions in both compliance behaviours, and beliefs about the perceived power of voices (Birchwood et al., 2014; Trower et al., 2004).

However, whilst the initial COMMAND pilot RCT reported reductions in voice-related distress (Trower et al., 2004), this finding was not replicated in the full-scale trial (Birchwood et al., 2014), suggesting that other factors are involved in maintaining distress, in addition to behaviours and appraisals. Furthermore, to date, the majority of research exploring this issue has been cross-sectional in nature. Whilst findings are consistent with the interpretation that resistance and compliance responses may contribute to or maintain voice-related distress, the opposite inference cannot be ruled out; distressing voices may lead to the hearer persisting with ineffective responses (Farhall et al., 2007;

Hacker et al., 2008). Furthermore, these studies have relied on 'retrospective' accounts of voice hearers, and so it remains unknown how different response styles might impact on voice-related distress during the course of daily life.

Experience Sampling Method (ESM) may shed further light on the role of behavioural responses in maintaining voice distress and associated appraisals. ESM involves assessing constructs of interest using questions delivered by paper or electronic means at semi-random intervals during participants' daily life. This provides a rich, ecologically valid dataset within which to examine the relationships between variables as they fluctuate over time. ESM has previously been used to assess relevant aspects of the cognitive behavioural model of voice hearing. Peters et al. (2012b) demonstrated significant associations between momentary levels of voice distress, voice intensity and concurrent appraisals of voice power and uncontrollability, whilst Hartley, Haddock, Vasconcelos e Sa, Emsley, and Barrowclough (2015) used a 'micro-longitudinal' approach to demonstrate that momentary increases in voice intensity and distress are predicted by antecedent attempts to control or suppress thoughts. Most recently, So et al. (2020) demonstrated how negative affect and voice hearing experiences formed a feedback loop that maintained voices, with these associations being exacerbated by appraisals of voices. The authors found no associations between 'resistance' and 'engagement' responses (on the BAVQ-R; Chadwick, Lees, & Birchwood, 2000); however, the measurement of appraisals and responses was conducted in a cross-sectional manner. To date, no study has used ESM to assess the dynamic relationships between voice appraisals, responses and distress.

The present study aimed to test the predictions of the cognitive model that: (a) behavioural and emotional responses to voices during daily life are driven primarily by beliefs about voices (e.g. omnipotence and malevolence), rather than voice characteristics (e.g. negative content or intensity) and (b) behavioural responses serve to maintain or exacerbate negative voice appraisals and distress from moment-to-moment. These predictions were tested by assessing both momentary and micro-longitudinal relationships between variables.

Four predictions were made. First, we predicted that momentary resistance and compliance responses to voices would be more closely associated with concurrent voice appraisals, rather than voice characteristics. Specifically, in line with past cross-sectional research, it was expected that resistance and compliance responses would be most closely related to appraisals of voice dominance and intrusiveness, with uncontrollability uniquely predicting compliance.

Second, we predicted that momentary voice distress would be related to concurrent resistance and compliance responses (indicating greater use of these behaviours at times of distress), but that these associations would not persist after controlling for concurrent voice appraisals (i.e. suggesting, in line with the cognitive model, that responses are driven by voice appraisals, rather than by distress).

Third, in line with their hypothesized role as safety behaviours, we predicted that resistance and compliance behaviours would be associated with increases in voice distress from moment-to-moment.

Finally, based on the suggestion that responses serve to maintain distress by reinforcing negative voice appraisals, we predicted that resistance and compliance behaviours would be associated with moment-to-moment increases in appraisals of voice dominance, intrusiveness and uncontrollability.

Method

Sample

Thirty-five participants were recruited from mental health services across Sussex, UK. Inclusion criteria were: aged 18 or over; currently treated as an outpatient of mental health services; currently experiencing frequent auditory verbal hallucinations [score of 2 ('at least once a day') or above on the frequency item of the Psychotic Symptoms Rating Scale – Auditory Hallucinations (PSYRATS-AH; Haddock, McCarron, Tarrier, & Faragher, 1999)]; adequate command of the English language. Exclusion criteria were: unable to provide fully informed written consent; symptoms precipitated by an organic cause; evidence of primary substance dependence; previously received 16 sessions or more of NICE-adherent CBTp. Ethical approval was obtained from the Camberwell St Giles National Research Ethics Committee (REC reference: 14/LO/0475).

Data collection

Basic sample characteristics

Data on age, gender, ethnicity, level of education and employment status were collected using a modified version of the Medical Research Council socio-demographic schedule (Mallet, 1997). DSM-IV diagnoses were determined based on a structured examination of case records using the OPERational CRITeria+ (OPCRIT+) system (Rucker *et al.*, 2011). Data on psychotropic medication use were collected using a medication checklist, which was completed based on a close examination of clinical documentation.

ESM measures

Items assessing voice characteristics, voice appraisals, behavioural responses and emotional response were included in the ESM questionnaire and were rated on 1–7 Likert scales (1: not at all; 7: very much). A detailed description of the ESM items is shown in Table 1. Extensive work was undertaken on the item development, including literature and scale reviews for constructs of interest, patient consultation and piloting. Psychometric evaluation of the ESM items indicated a sufficient degree of reliability (split week reliabilities ranging from 0.67 to 0.94; see Table 3) and convergent and divergent validity.

ESM procedure

All participants were provided with a smartphone pre-loaded with the movisensXS ESM app (<https://xs.movisens.com/>), via which the ESM measure was administered 10 times per day. We used a time-based design with stratified random sampling (i.e. with ESM assessments scheduled at random within set blocks of time) (Myin-Germeys *et al.*, 2009; Palmier-Claus *et al.*, 2011; Stone, Shiffman, Atienza, & Nebeling, 2007). On each day over nine consecutive days, the smartphone emitted 10 'beep' signals at semi-random moments within set blocks of time. During an initial briefing session, participants were trained in the use of the smartphone and practising its usage by going through a practice questionnaire. In this session, participants were given instructions about the forthcoming ESM assessment; they were informed that each time the device emitted the beep signal, they should stop their activity and respond to a comprehensive diary questionnaire assessing voice phenomenology and social interactions in daily life. During the assessment period, which was selected to start at any day of the week at the discretion of

the participants (to optimize compliance and achieve sufficient spread of week and weekend days in our sample), the ESM questionnaire was available to participants for the duration of 15 min after emission of the beep signal. Participants were contacted twice during the assessment period to assess their adherence to instructions, identify any problems and help participants overcome any potential barriers. At the end of the assessment period, participants' reactivity to, and compliance with, the method was examined in a debriefing session. Participants were required to provide valid responses to at least one-third of the emitted beeps (i.e. 30 data points) to be included in the analysis, in line with previous ESM research.

Statistical analysis

ESM data have a multilevel structure, such that multiple observations (level-1) are nested within participants (level-2). Linear mixed models were therefore used to control for within-participant clustering of multiple observations using the MIXED module in STATA 14.0 (StataCorp, 2015). Restricted maximum likelihood estimation of these models allows for the use of all available data under the relatively unrestrictive assumption that data are missing at random (Mallinckrodt, Clark, & David, 2001). Where model assumptions were violated, standard errors of the final models were estimated using robust maximum likelihood methods. Effect sizes from predictors in the multilevel model were expressed as β , representing the unstandardized fixed regression coefficient. This can be interpreted in the same way as unstandardized B estimates in unilevel regression analysis.

The improved fit of complex models above baseline models was evaluated using Akaike's and Schwarz's Bayesian information criteria (Burnham, Anderson, & Anderson, 2004). In all models, dependent variables were entered un-centred, and all predictor variables were entered group (i.e. person) mean centred, in order to control for between-person differences in experience intensity (Nezlek, 2012). Intercepts and slopes were modelled as random effects, wherever this resulted in an improved model fit (i.e. indicating a significant random effect). Fixed and random linear effects of time (i.e. measurement occasion) on the dependent variables were explored and controlled for when necessary (Bolger & Laurenceau, 2013). In all models, an independent random-effects covariance matrix was specified to allow for distinct variances of all random effects. Given the possibility of serial autocorrelation between residual errors in ESM data (Bolger & Laurenceau, 2013), in all analyses described, we explored whether model fit was improved by modelling the residual error structure using an autoregressive process of order 1 (Walls, Höppner, & Goodwin, 2007). For these analyses, Bonferroni-adjusted Wald χ^2 tests, based on the multilevel regression models, were used to test the equality of fixed effects in order to assess the relative contribution of each significant predictor on the outcome variable.

Results

Basic sample characteristics and ESM item descriptives

A total of 35 participants were assessed with the ESM during the study period. Of these, 31 participants (88.5%) provided ≥ 30 valid responses and were included in the analyses. Demographic and clinical information for included participants is summarized in Table 2.

ESM data were provided on 1682 occasions, of which voices were reported at 1094 moments (65% of measurement occasions).

Table 1. ESM measures

Domain	ESM measure
Voice characteristics	Voice characteristics were assessed using two ESM items prefaced by the phrase 'Right before the beep...': <i>Voice intensity</i> : '...I could hear a voice or voices that other people couldn't hear' <i>Negative voice content</i> : 'the voice ^a was saying negative things'
Voice appraisals	Voice appraisals were assessed using three ESM items prefaced by the phrase 'Right before the beep...': <i>Voice dominance</i> : '...I felt inferior to the voice ^a ' <i>Voice uncontrollability</i> : '...I felt that the voice ^a was out of my control' <i>Voice intrusiveness</i> : '...I felt that the voice ^a was intruding on my personal space'
Behavioural responses	Behavioural responses to voices were assessed using two ESM items prefaced by the phrase 'Right before the beep...': <i>Resistance</i> : '...I was trying to ignore the voice ^a or stop it from talking' <i>Compliance</i> : '...I was doing what the voice ^a was telling me to do'
Emotional response	The emotional response was assessed using one ESM item prefaced by the phrase 'Right before the beep...': <i>Voice-related distress</i> : '...the voice ^a was upsetting me'

^aOn occasions when more than one voice was reported, all voice-relevant items referred to the 'voices' rather than the 'voice', with appropriate associated grammar (e.g. the voices were as opposed to was; them as opposed to it, etc.).

Descriptive statistics for all ESM constructs are displayed in Table 3. All participants reported voice-hearing episodes, with a mean of 35.3 voice reports (range 2–69) per participant. Thirty participants (96.8%) reported attempting to resist their voices at least once over the course of the 9-day assessment period. Across these individuals, some degree of resistance (i.e. score >1) was reported on 88.4% of measurement occasions during which voices were experienced. Twenty-four participants (77.4%) reported complying with voice demands at least once. On average, these individuals reported complying with their voices to some degree (score >1) on 58.5% of occasions when voices were reported. There were no significant changes in compliance ($B = 0.06$, $z = 1.61$, $p = 0.11$, 95% CI -0.01 to 0.01) or resistance ($B = -0.01$, $z = -1.35$, $p = 0.18$, 95% CI -0.01 to 0.01) behaviours over the course of the 9 days.

Variability was assessed using intraclass correlation (ICC), which provides an index of the percentage of between-person variability relative to the total variability and can thus be used to assess the degree to which items vary between persons, or from moment-to-moment within persons. Standard deviations (s.d.) were also calculated at both the within- and between-person levels, providing a further indication of the variation residing at each level of analysis. A summary of item means, ICCs and within- and between-person s.d. are displayed in Table 3.

ICC values indicate a significant level of clustering for each of the items, indicating substantial between-person variation (i.e. individual differences) in these mean levels. In particular, these analyses indicate high between-person variation in voice intensity and distress, perceived voice dominance and uncontrollability and degree of resistance to voices, indicating heterogeneity in both the experience of voice hearing, and in the psychological mechanisms underlying these experiences.

The values of within-person s.d. also reveal substantial within-person (i.e. temporal) variation in various constructs related to the experience of distressing voices, including voice characteristics (voice intensity and negative content), voice appraisals (perceived voice dominance, uncontrollability and intrusiveness), behavioural responses (resistance and compliance) and emotional response (voice-related distress). Of these constructs, perceived voice dominance demonstrated the least within-person variation,

suggesting that this experience might be somewhat more 'trait-like' in nature, demonstrating greater variation between individuals than within individuals. Variables demonstrating particularly high within-person variability included voice intensity, distress, perceived voice intrusiveness and compliance with voices.

Are behavioural responses to voices best predicted by concurrent voice characteristics or voice appraisals?

Predictors of momentary responses to voices were explored using two multilevel models, with resistance and compliance responses as the outcome variables, and voice characteristics (intensity; negative content) and appraisals (voice dominance, uncontrollability and intrusiveness) as predictors. Table 4 reports the results of these analyses.

In line with predictions, momentary reports of voice resistance and compliance responses were significantly associated with concurrent voice appraisals, but not voice characteristics (although note that the momentary association between negative voice content and resistance responses approached significance; $p = 0.05$). Also supporting predictions, momentary compliance behaviours were associated with appraisals of voice dominance and uncontrollability, with the results indicating that, on average, a unit increase in perceived voice dominance was accompanied by a 0.16-unit increase in voice compliance. A Wald χ^2 test indicated that there was no significant difference in the magnitude of the effects of perceived voice dominance and uncontrollability on compliance behaviours [$\chi^2(1) = 0.27$, $p = 0.60$]. Unexpectedly, perceived voice uncontrollability was the only significant predictor of momentary resistance to voices, whilst voice intrusiveness was not significantly associated with either compliance or resistance behaviours.

Are behavioural responses related to momentary levels of voice distress?

A multilevel model with voice distress as the outcome variable, and resistance and compliance responses as predictors, indicated that momentary distress was significantly associated with both resistance ($B = 0.23$, $z = 3.55$, $p < 0.001$, 95% CI 0.10 – 0.35) and

Table 2. Demographic and clinical characteristics ($N = 31$)

Mean age (s.d.)	41.9 (11.4)
Gender, n (%)	
Male	11 (35.5)
Female	18 (58.1)
Other	2 (6.5)
Ethnicity, n (%)	
White British	27 (87.1)
White Other	1 (3.2)
Other	3 (9.7)
Place of birth	
UK-born	28 (90.3)
Non-UK-born	3 (9.7)
Level of education, n (%)	
School	7 (22.6)
Further	17 (54.8)
Higher	7 (22.6)
Employment	
Unemployed	14 (45.2)
Other	17 (54.8)
OPCRIT+ DSM-IV diagnosis, n (%)	
Schizophrenia	12 (38.7)
Schizoaffective disorder	2 (6.5)
Other psychotic disorder	3 (9.7)
Borderline personality disorder	10 (32.3)
Depression with psychotic features	3 (9.7)
Bipolar disorder	1 (3.2)
Psychotropic medication, n (%)	
Antipsychotic	28 (90.3)
Antidepressant	21 (67.7)
Other	10 (32.3)

compliance ($B = 0.18$, $z = 3.48$, $p < 0.001$, 95% CI 0.09–0.29) responses. However, in line with predictions, these effects did not persist once voice characteristics and appraisals were added to the model (Table 5), indicating that there is no direct relationship between distress and resistance/compliance responses.

As predicted by the cognitive model, voice distress was significantly associated with appraisals of voice dominance, uncontrollability and intrusiveness. However, unexpectedly, both voice intensity and negative voice content made significant independent contributions to the prediction of momentary voice distress.

Are behavioural responses related to subsequent increases in voice distress?

Next, reports of behavioural responses at the previous ESM measurement occasion (time $t-1$) were entered as predictor variables in a multilevel regression analysis assessing current voice distress (time t) as the dependent variable. This analysis controlled for

the effects of voice appraisals, characteristics and distress at $t-1$. Table 6 reports the results of this analysis, including all model covariates.

As predicted, the results indicate that both resistance and compliance behaviours are associated with increases in voice-related distress at subsequent measurement occasions, even after controlling for antecedent effects of voice characteristics, appraisals and voice-related distress. A unit increase in voice compliance or resistance at time $t-1$ were associated with a 0.1-unit increase in voice-related distress at time t , indicating that these responses might serve to maintain or exacerbate voice-related distress. A Wald χ^2 test indicated that there was no significant difference in the magnitude of the effects of resistance and compliance on subsequent distress [$\chi^2(1) = 0.04$, $p = 0.85$].

Running the reverse models indicated that levels of voice distress reported at $t-1$ did not significantly predict compliance ($B = 0.07$, $z = 1.15$, $p = 0.25$, 95% CI -0.05 to 0.18) or resistance ($B = -0.03$, $z = -0.66$, $p = 0.51$, 95% CI -0.12 to 0.06) at time t , indicating directional effects of these behavioural responses on subsequent distress.

Are behavioural responses related to subsequent increases in negative voice appraisals?

Finally, a series of multilevel analyses were performed to determine whether behavioural responses at time $t-1$ predict subsequent increases in voice appraisals. Here, voice dominance, uncontrollability and intrusiveness at time t were the outcome variables, whilst behavioural responses at the previous time point ($t-1$) were the predictor variables. Voice characteristics, appraisals and associated distress at time $t-1$ were controlled for in these analyses. Table 7 reports the results of these analyses including all model covariates.

Resisting voices at time $t-1$ did not independently predict changes in voice appraisals at time t , although the associations between resistance and subsequent increases in perceived intrusiveness approached significance ($B = 0.10$, $z = 1.80$, $p = 0.07$, 95% CI -0.01 to 0.21).

On the other hand, compliance with voices at time $t-1$ was significantly associated with increases in perceived uncontrollability of voices, at time t . Running the reverse model indicated that compliance at time t was not significantly predicted by voice uncontrollability ($B = 0.06$, $z = 0.91$, $p = 0.36$, 95% CI -0.07 to 0.20) at time $t-1$, suggesting directional effects of compliance on this outcome.

Discussion

In support of the cognitive model, findings suggest that both momentary behavioural and emotional responses to voices are associated with concurrent negative voice appraisals. Whilst voice distress was associated with both resistance and compliance responses, these effects did not persist after controlling for concurrent voice appraisals, suggesting that these behaviours are not direct responses to (or causes of) momentary voice distress, but to beliefs about voices. Furthermore, in line with the hypothesized role of behavioural responses in the maintenance of voice distress, the 'micro-longitudinal' analyses indicated that both resistance and compliance behaviours were associated with increases in voice-related distress at subsequent measurement occasions, even after controlling for antecedent effects of voice characteristics, appraisals and voice-related distress. Compliance

Table 3. Descriptive statistics for ESM items assessing voice characteristics, appraisals and responses

	<i>M</i>	ICC	s.d. (within-person)	s.d. (between-person)	Split-week reliability (<i>r</i>) ^a
Intensity	4.02	0.49	1.83	1.85	0.94***
Negative content	5.91	0.52	1.23	1.37	0.93***
Voice dominance	3.94	0.73	1.10	1.91	0.84***
Uncontrollability	4.88	0.66	1.23	1.90	0.89***
Intrusiveness	4.78	0.46	1.33	1.32	0.67***
Resistance	4.85 ^b	0.66	1.21 ^b	1.64 ^b	0.93***
Compliance	2.71 ^b	0.48	1.46 ^b	1.33 ^b	0.83***
Distress	4.32	0.51	1.30	1.51	0.83***

^aThe split-week reliability (the ESM equivalent of test-retest reliability) was calculated as the correlation between mean within-person item scores from the first half (days 1–4) and the second half (days 5–9) of the sampling period.

^bCalculated across participants who reported this response (score >1) on at least one occasion.

Table 4. Momentary associations between behavioural responses, voice characteristics and appraisals

Outcome variables ^a	Predictor variables				
	Voice characteristics		Voice appraisals		
	Intensity <i>B</i> ^b (s.e.)	Negative content <i>B</i> ^b (s.e.)	Dominance <i>B</i> ^b (s.e.)	Uncontrollability <i>B</i> ^b (s.e.)	Intrusiveness <i>B</i> ^b (s.e.)
Resistance (<i>N</i> = 30)	−0.01 (0.04)	0.13 (0.07)	−0.03 (0.06)	0.11 (0.05)*	0.07 (0.06)
Compliance (<i>N</i> = 24)	0.03 (0.04)	−0.02 (0.04)	0.16 (0.08)*	0.21 (0.06)**	0.03 (0.04)

^aIncludes only participants who reported response with score >1 on at least one occasion; see text for details.

^bThe *B* is the unstandardized fixed regression coefficient of the predictor in the multi-level model. Robust standard errors are reported for all coefficients.

p* < 0.05, *p* < 0.01, ****p* < 0.001 (significant findings are shown in bold).

Table 5. Momentary associations between voice-related distress and behavioural responses, voice characteristics and appraisals

Outcome variable ^a	Predictor variables						
	Behavioural responses (<i>t</i>)		Voice characteristics (<i>t</i>)		Voice appraisals (<i>t</i>)		
	Resistance <i>B</i> ^b (s.e.)	Compliance <i>B</i> ^b (s.e.)	Intensity <i>B</i> ^b (s.e.)	Negative content <i>B</i> ^b (s.e.)	Dominance <i>B</i> ^b (s.e.)	Uncontrollability <i>B</i> ^b (s.e.)	Intrusiveness <i>B</i> ^b (s.e.)
Distress (<i>t</i>)	0.04 (0.04)	0.03 (0.02)	0.18 (0.04)***	0.08 (0.04)*	0.09 (0.04)*	0.17 (0.05)**	0.31 (0.06)***

^aAnalysis excluded participants who did not report compliance or resistance responses. Thus, this analysis is based on 922 observations from 24 participants.

^bThe *B* is the unstandardized fixed regression coefficient of the predictor in the multi-level model. Robust standard errors are reported for all coefficients.

p* < 0.05, *p* < 0.01, ****p* < 0.001 (significant findings are shown in bold).

Table 6. Time-lagged associations between voice-related distress at time *t* and behavioural responses at time *t*−1

Outcome variable ^a	Predictor variables		Model covariates					
	Behavioural responses (<i>t</i> −1)		Voice characteristics (<i>t</i> −1)		Voice appraisals (<i>t</i> −1)			Emotional response (<i>t</i> −1)
	Resistance <i>B</i> ^b (s.e.)	Compliance <i>B</i> ^b (s.e.)	Intensity <i>B</i> ^b (s.e.)	Negative content <i>B</i> ^b (s.e.)	Dominance <i>B</i> ^b (s.e.)	Uncontrollability <i>B</i> ^b (s.e.)	Intrusiveness <i>B</i> ^b (s.e.)	Distress <i>B</i> ^b (s.e.)
Distress (<i>t</i>)	0.10 (0.05)*	0.11 (0.03)**	−0.04 (0.05)	−0.05 (0.05)	0.03 (0.05)	−0.06 (0.05)	0.07 (0.05)	0.27 (0.08)**

^aAnalysis excluded participants who did not report compliance or resistance responses. Thus, this analysis is based on 519 time-lagged observations from 24 participants.

^bThe *B* is the unstandardized fixed regression coefficient of the predictor in the multi-level model. Robust standard errors are reported for all coefficients.

p* < 0.05, *p* < 0.01, ****p* < 0.001 (significant findings are shown in bold).

Table 7. Time-lagged associations between voice appraisals at time *t* and behavioural responses at time *t*-1

Outcome variable ^a	Predictor variables			Model covariates					
	Behavioural responses (<i>t</i> -1)		Compliance B^b (S.E.)	Voice characteristics (<i>t</i> -1)			Voice appraisals (<i>t</i> -1)		Emotional response (<i>t</i> -1) Distress B^b (S.E.)
	Resistance B^b (S.E.)	Intensity B^b (S.E.)		Negative content B^b (S.E.)	Dominance B^b (S.E.)	Uncontrollability B^b (S.E.)	Intrusiveness B^b (S.E.)		
Dominance (<i>t</i>)	0.01 (0.04)	0.02 (0.05)	0.03 (0.03)	-0.06 (0.05)	0.16 (0.08)	0.08 (0.05)	-0.01 (0.04)	0.05 (0.07)	
Uncontrollability (<i>t</i>)	0.01 (0.05)	0.01 (0.06)	0.08 (0.04)*	-0.05 (0.06)	0.01 (0.08)	0.14 (0.07)*	0.08 (0.05)	0.04 (0.08)	
Intrusiveness (<i>t</i>)	0.10 (0.06)	-0.09 (0.07)	0.06 (0.04)	0.02 (0.07)	0.08 (0.06)	0.01 (0.06)	0.20(0.08)**	0.16(0.07)*	

^aAnalyses excluded participants who did not report compliance or resistance responses. Thus, these analyses are based on 517-639 time-lagged observations from 24 participants.

^bThe B is the unstandardized fixed regression coefficient of the predictor in the multi-level models. Robust standard errors are reported for all coefficients.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (significant findings are shown in bold).

was additionally associated with increases in appraisals of voice uncontrollability over time, suggesting a mechanism via which responses may serve to maintain voice distress.

Considering first the results of the momentary analyses, the demonstrated associations between negative voice appraisals and both voice distress and behavioural responses are consistent with the possibility of a mediating role of voice appraisals in both behavioural and emotional responses to voices (Chadwick & Birchwood, 1994). In line with past cross-sectional and ESM research, voice distress was associated with concurrent appraisals of voice dominance, uncontrollability and intrusiveness (Beavan & Read, 2010; Birchwood, Meaden, Trower, Gilbert, & Plaistow, 2000; Hayward et al., 2008; Peters et al., 2012a; So et al., 2020), whilst both compliance and resistance were associated with appraisals of voice uncontrollability, with compliance additionally being associated with appraisals of voice dominance. The finding that associations between voice distress and behavioural responses did not persist after controlling for concurrent voice appraisals parallel those of Hacker et al. (2008), who demonstrated that the observed cross-sectional association between safety behaviour use and voice distress was mediated by appraisals of voice omnipotence, suggesting that safety behaviours reflect attempts to mitigate perceived threat, rather than to reduce distress *per se*. Findings from the current study suggest that similar mechanisms may be at play during daily life.

The demonstrated association between compliance and perceived voice dominance is consistent with a wealth of cross-sectional findings implicating perceived voice rank (Reynold & Scragg, 2010) and omnipotence (Bucci et al., 2013; Fox, Gray, & Lewis, 2004; Reynold & Scragg, 2010) as predictors of compliance with voice commands. However, whilst it was predicted, based on past cross-sectional research (Birchwood et al., 2004; Gilbert et al., 2001; Hayward et al., 2008), that voice dominance would also be associated with resistance, this was not born out in the findings.

Interestingly, some studies have failed to demonstrate an association between voice omnipotence (a construct closely related to voice dominance) and resistance, after controlling for the perceived malevolent intent of voices (Peters et al., 2012b; van der Gaag et al., 2003). Indeed, social relating theories (Benjamin, 1989; Hayward et al., 2011; Thomas, McLeod, & Brewin, 2009) propose that voices perceived as dominant will elicit complementary submissive responses, such as compliance, whilst resistance is more likely to be elicited by voices that are perceived as intrusive or hostile. Whilst no evidence was found of an association between voice intrusiveness and resistance responses, the finding that dominance uniquely predicted compliance, and not resistance, is in line with this suggestion. Based on their findings, Peters et al. (2012b) suggested that voice malevolence might be more critical in eliciting resistance than voice power/dominance. This notion is supported by findings from the command hallucination literature, where resistance to commands is best predicted by perceived voice malevolence, and with compliance being more likely when voices are perceived as powerful (Barrowcliff & Haddock, 2006; Bucci et al., 2013). However, since there was no attempt to assess perceived voice malevolence, this possibility could not be tested.

Findings from the current study additionally highlight the importance of appraisals of voice uncontrollability in both resistance and compliance responses. To our knowledge, this construct has not previously been assessed as a predictor of voice compliance or resistance, but research has demonstrated cross-sectional associations between voice-related distress and both perceived

voice uncontrollability (Beavan & Read, 2010; Peters et al., 2012a), and metacognitive beliefs about the uncontrollability of voices and their associated danger (Morrison, Nothard, Bove, & Wells, 2004; Varese et al., 2016). It has been suggested that perceived loss of control may elicit maladaptive attempts to control or suppress voices (Varese et al., 2016), or alternatively, reinforce appraisals of voice power, eliciting submissive responses such as compliance (Benjamin, 1989; Thomas et al., 2009). Whilst the findings from the current study are consistent with these suggestions, it is of course equally possible that appraisals of uncontrollability may stem from failed attempts to resist voices or their commands.

Using a micro-longitudinal approach, we found evidence that resistance and compliance responses played a role in maintaining or exacerbating voice distress and negative appraisals over time in daily life. Furthermore, these associations appear to be directional; antecedent distress did not predict increases in resistance or compliance responses. Similar findings have previously been demonstrated with regard to the role of attempts to control or suppress thoughts on subsequent voice distress (Hartley et al., 2015); results from the current study suggest that this effect applies to attempts to control or resist voices.

Some support was also found for the notion that behavioural responses might maintain distress via their effect on reinforcing and/or preventing disconfirmation of negative voice appraisals (Michail & Birchwood, 2010; Morrison, 1998). Compliance with voices was associated with subsequent increases in appraisals of voice uncontrollability, whilst the time-lagged association between resistance and perceived voice intrusiveness approached significance. This dynamic association between compliance and voice uncontrollability is particularly interesting in light of the findings of momentary associations between uncontrollability appraisals and both voice distress and compliance, suggesting a mechanism via which compliance may serve to exacerbate voice distress, and prompt further compliance, over time, by reinforcing appraisals of voice uncontrollability.

An unexpected finding concerned the associations between momentary voice distress and both voice intensity and negative content, even after controlling for voice appraisals. Whilst this lies in contrast to previous cross-sectional research (van der Gaag et al., 2003), similar findings were obtained in previous ESM studies (Peters et al., 2012a; So et al., 2020), suggesting that the influence of voice characteristics on voice-related distress as experienced during daily life may have been underestimated. This finding echoes suggestions that exploration of voice content may be a crucial component of both understanding and reducing the distress experienced by some in relation to their voices (Beavan & Read, 2010; Larøi et al., 2018; Longden, Corstens, Escher, & Romme, 2012; Romme & Escher, 2000).

The findings of this study should be interpreted in light of several limitations. First, the findings are consistent with a mediating role of voice appraisals in the relationship between behavioural responses and voice distress, both in the moment and over time, but it was not possible to perform formal tests of mediation whilst controlling for necessary covariates due to model non-convergence. Future research should use a multilevel mediation approach (Bauer, Preacher, & Gil, 2006; Bolger & Laurenceau, 2013) within the context of a larger ESM study to test whether: (i) the observed momentary association between behavioural responses and voice distress is mediated by appraisals of voice dominance and uncontrollability; and (ii) the observed micro-longitudinal relationship between compliance and

subsequent increase in voice distress is mediated by appraisals of voice uncontrollability.

Second, there was no assessment of appraisals of voice malevolence. Whilst an item assessing voice benevolence was included, it was rarely endorsed and demonstrated unacceptably low within-person variability for use (in reverse-coded form) within the present analyses.

Third, the magnitude of the effects was generally small, although not negligible. Small effect sizes are common in ESM studies, but it is the cumulative impact of these effects that can be substantial given the frequency with which they are observed during daily life (Myin-Germeys et al., 2003).

Finally, despite conducting extensive work on item development, the psychometric evaluation of the ESM items did not include an assessment of their construct validity. This should be a focus of future research.

The findings from the current study provide ecological validation for a role of compliance and resistance responses in the maintenance of voice distress and negative voice appraisals during daily life. In addition to providing support for cognitive models, these findings have implications for psychological interventions for distressing voices, supporting the notion that behaviour change should remain a central goal of therapy. However, the results have particular implications for therapies incorporating behavioural experiments encouraging attempts to resist command hallucinations; the findings highlight the importance of differentiating between resistance to voice commands and resisting voice experiences more generally. In this respect, interventions incorporating acceptance and mindfulness approaches (e.g. Chadwick et al., 2016), targeting coping behaviours (Bell et al., 2019; Hayward, Edgecumbe, Jones, Berry, & Strauss, 2018) or interpersonal relationships with voices (Craig et al., 2018; Hayward, Jones, Bogen-Johnston, Thomas, & Strauss, 2017), may offer hearers an alternative way of responding to their voices. Furthermore, whilst the results support the focus of cognitive interventions on re-evaluating appraisals of voice power/dominance and uncontrollability, they highlight the importance of a parallel therapeutic focus on exploring and responding to negative voice content (Larøi et al., 2019).

References

- Abramowitz, J. S., Tolin, D. F., & Street, G. P. (2001). Paradoxical effects of thought suppression: A meta-analysis of controlled studies. *Clinical Psychology Review*, 21(5), 683–703. [http://doi.org/10.1016/S0272-7358\(00\)00057-X](http://doi.org/10.1016/S0272-7358(00)00057-X).
- Barrowcliff, A. L., & Haddock, G. (2006). The relationship between command hallucinations and factors of compliance: A critical review of the literature. *Journal of Forensic Psychiatry & Psychology*, 17(2), 266–298. <http://doi.org/10.1080/14789940500485078>.
- Bauer, D. J., Preacher, K. J., & Gil, K. M. (2006). Conceptualizing and testing random indirect effects and moderated mediation in multilevel models: New procedures and recommendations. *Psychological Methods*, 11(2), 142–163. <http://doi.org/10.1037/1082-989X.11.2.142>.
- Beavan, V. (2011). Towards a definition of 'hearing voices': A phenomenological approach. *Psychosis*, 3(1), 63–73. <http://doi.org/10.1080/17522431003615622>.
- Beavan, V., & Read, J. (2010). Hearing voices and listening to what they say: The importance of voice content in understanding and working with distressing voices. *The Journal of Nervous and Mental Disease*, 198(3), 201–205. <http://doi.org/10.1097/NMD.0b013e3181d14612>.
- Bell, I. H., Fielding-Smith, S. F., Hayward, M., Rossell, S. L., Lim, M. H., Farhall, J., & Thomas, N. (2019). Pilot randomised controlled trial of a brief coping-focused intervention for hearing voices blended with smartphone-based ecological momentary assessment and intervention (SAVVy): Feasibility,

- acceptability and preliminary clinical outcomes. *Schizophrenia Research*, 216, 479–487. <https://doi.org/10.1016/j.schres.2019.10.026>.
- Benjamin, L. S. (1989). Is chronicity a function of the relationship between the person and the auditory hallucination? *Schizophrenia Bulletin*, 15(2), 291–310. <http://doi.org/10.1093/schbul/15.2.291>.
- Birchwood, M., Gilbert, P., Gilbert, J., Trower, P., Meaden, A., Hay, J., ... Miles, J. N. V. (2004). Interpersonal and role-related schema influence the relationship with the dominant 'voice' in schizophrenia: A comparison of three models. *Psychological Medicine*, 34(8), 1571–1580. <http://doi.org/10.1017/S0033291704002636>.
- Birchwood, M., Meaden, A., Trower, P., Gilbert, P., & Plaistow, J. (2000). The power and omnipotence of voices: Subordination and entrapment by voices and significant others. *Psychological Medicine*, 30(2), 337–344. <http://doi.org/10.1017/S0033291799001828>.
- Birchwood, M., Michail, M., Meaden, A., Tarrrier, N., Lewis, S. W., Wykes, T., ... Peters, E. R. (2014). Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): A randomised controlled trial. *The Lancet Psychiatry*, 1(1), 23–33. [http://doi.org/10.1016/S2215-0366\(14\)70247-0](http://doi.org/10.1016/S2215-0366(14)70247-0).
- Bolger, N., & Laurenceau, P.-J. (2013). *Intensive longitudinal methods: An introduction to diary and experience sampling research*. New York: The Guildford Press.
- Bucci, S., Birchwood, M., Twist, L., Tarrrier, N., Emsley, R., & Haddock, G. (2013). Predicting compliance with command hallucinations: Anger, impulsivity and appraisals of voices' power and intent. *Schizophrenia Research*, 147(1), 163–168. <http://doi.org/10.1016/j.schres.2013.02.037>.
- Burnham, K. P., Anderson, D. R., & Anderson, R. P. (2004). Multimodel inference: Understanding AIC and BIC in model selection. *Sociological Methods & Research*, 33(2), 261–304. <https://doi.org/10.1177/0049124104268644>.
- Chadwick, P., & Birchwood, M. (1994). The omnipotence of voices: A cognitive approach to hallucinations. *British Journal of Psychiatry*, 164, 190–201. Retrieved from <http://bjp.rcpsych.org/content/164/2/190.short>.
- Chadwick, P., Lees, S., & Birchwood, M. (2000). The revised beliefs about voices questionnaire (BAVQ-R). *British Journal of Psychiatry*, 177(SEPT), 229–232. <http://doi.org/10.1192/bjp.177.3.229>.
- Chadwick, P., Strauss, C., Jones, A.-M., Kingdon, D., Ellet, L., Dannahy, L., & Hayward, M. (2016). Group mindfulness-based intervention for distressing voices: A pragmatic randomised controlled trial. *Schizophrenia Research*, 175, 168–175.
- Chaix, J., Ma, E., Nguyen, A., Ortiz Collado, M. A., Rexhaj, S., & Favrod, J. (2014). Safety-seeking behaviours and verbal auditory hallucinations in schizophrenia. *Psychiatry Research*, 220(1–2), 158–162. <http://doi.org/10.1016/j.psychres.2014.08.041>.
- Craig, T. K. J., Rus-Calafell, M., Ward, T., Leff, J. P., Huckvale, M., Howarth, E., ... Garety, P. A. (2018). AVATAR therapy for auditory verbal hallucinations in people with psychosis: A single-blind, randomised controlled trial. *The Lancet Psychiatry*, 5, 31–40. [http://dx.doi.org/10.1016/S2215-0366\(17\)30427-3](http://dx.doi.org/10.1016/S2215-0366(17)30427-3).
- David, A. S. (2004). The cognitive neuropsychiatry of auditory verbal hallucinations: An overview. *Cognitive Neuropsychiatry*, 9(1–2), 107–123. <http://doi.org/10.1080/13546800344000183>.
- Farhall, J., & Gehrke, M. (1997). Coping with hallucinations: Exploring stress and coping framework. *The British Journal of Clinical Psychology/The British Psychological Society*, 36(Pt 2), 259–261.
- Farhall, J., Greenwood, K. M., & Jackson, H. J. (2007). Coping with hallucinated voices in schizophrenia: A review of self-initiated strategies and therapeutic interventions. *Clinical Psychology Review*, 27(4), 476–493. <http://doi.org/10.1016/j.cpr.2006.12.002>.
- Fox, J. R., Gray, N. S., & Lewis, H. (2004). Factors determining compliance with command hallucinations with violent content: The role of social rank, perceived power of the voice and voice malevolence. *Journal of Forensic Psychiatry & Psychology*, 15(3), 511–531. <http://doi.org/10.1080/1478994042000226741>.
- Gilbert, P., Birchwood, M., Gilbert, J., Trower, P., Hay, J., Murray, B., ... Miles, J. N. V. (2001). An exploration of evolved mental mechanisms for dominant and subordinate behaviour in relation to auditory hallucinations in schizophrenia and critical thoughts in depression. *Psychological Medicine*, 31(6), 1117–1127. <http://doi.org/10.1017/S0033291701004093>.
- Hacker, D., Birchwood, M., Tudway, J., Meaden, A., & Amplett, C. (2008). Acting on voices: Omnipotence, sources of threat, and safety-seeking behaviours. *British Journal of Clinical Psychology*, 47(2), 201–213. <http://doi.org/10.1348/014466507X249093>.
- Haddock, G., McCarron, J., Tarrrier, N., & Faragher, E. B. (1999). Scales to measure dimensions of hallucinations and delusions: The psychotic symptom rating scales (PSYRATS). *Psychological Medicine*, 29(4), 879–889. <http://doi.org/10.1017/S0033291799008661>.
- Hartley, S., Haddock, G., Vasconcelos e Sa, D., Emsley, R., & Barrowclough, C. (2015). The influence of thought control on the experience of persecutory delusions and auditory hallucinations in daily life. *Behaviour Research and Therapy*, 65, 1–4. <http://doi.org/10.1016/j.brat.2014.12.002>.
- Hayward, M. (2003). Interpersonal relating and voice hearing: To what extent does relating to the voice reflect social relating? *Psychology and Psychotherapy*, 76(Pt 4), 369–383. <http://doi.org/10.1348/147608303770584737>.
- Hayward, M., Berry, K., & Ashton, A. (2011). Applying interpersonal theories to the understanding of and therapy for auditory hallucinations. *Clinical Psychology Review*, 31, 1313–1323.
- Hayward, M., Denney, J., Vaughan, S., & Fowler, D. (2008). The Voice and You: Development and psychometric evaluation of a measure of relationships with voices. *Clinical Psychology and Psychotherapy*, 15, 45–52.
- Hayward, M., Edgcombe, R., Jones, A.-M., Berry, C., & Strauss, C. (2018). Brief coping strategy enhancement for distressing voices: An evaluation in routine clinical practice. *Behavioural and Cognitive Psychotherapy*, 46, 226–237. <http://doi.org/10.1017/S1352465817000388>.
- Hayward, M., Jones, A.-M., Bogen-Johnston, L., Thomas, N., & Strauss, C. (2017). Relating therapy for distressing auditory hallucinations: A pilot randomized controlled trial. *Schizophrenia Research*, 183, 137–142.
- Howard, A., Forsyth, A., Spencer, H., Young, E. W., & Turkington, D. (2013). Do voice hearers naturally use focusing and metacognitive coping techniques? *Psychosis*, 5(2), 119–126. <http://doi.org/10.1080/17522439.2012.668926>.
- Larøi, F., Thomas, N., Aleman, A., Fernyhough, C., Wilkinson, S., Deamer, F., & McCarthy-Jones, S. (2018). The ice in voices: Understanding negative content in auditory verbal hallucinations. *Clinical Psychology Review*, 67, 1–10.
- Larøi, F., Thomas, N., Aleman, A., Fernyhough, C., Wilkinson, S., Deamer, F., & McCarthy-Jones, S. (2019). The ice in voices: Understanding negative content in auditory-verbal hallucinations. *Clinical Psychology Review*, 67, 1–10.
- Longden, E., Corstens, D., Escher, S., & Romme, M. A. J. (2012). Voice hearing in a biographical context: A model for formulating the relationship between voices and life history. *Psychosis*, 4(3), 224–234. <http://doi.org/10.1080/17522439.2011.596566>.
- Mackinnon, A., Copolov, D. L., & Trauer, T. (2004). Factors associated with compliance and resistance to command hallucinations. *The Journal of Nervous and Mental Disease*, 192(5), 357–362. <http://doi.org/10.1097/01.nmd.0000126728.70060.fa>.
- Mallet, R. (1997). *Sociodemographic schedule*. London, UK: Section of Social Psychiatry, Institute of Psychiatry.
- Mallinckrodt, C. H., Clark, W. S., & David, S. R. (2001). Accounting for drop-out bias using mixed-effects models. *Journal of Biopharmaceutical Statistics*, 11(1–2), 9–21.
- Meaden, A., Keen, N., Aston, R., Barton, K., & Bucci, S. (2013). *Cognitive therapy for command hallucinations: An advanced practical companion*. Hove: Routledge.
- Michail, M., & Birchwood, M. (2010). Command hallucinations: Theory and psychological interventions. In R. Hagen, D. Turkington & T. Berge (Eds.), *CBT for psychosis: A symptom-based approach* (pp. 99–115). New York, NY: Routledge.
- Morrison, A. P. (1998). A cognitive analysis of the maintenance of auditory hallucinations: Are voices to schizophrenia what bodily sensations are to panic? *Behavioural and Cognitive Psychotherapy*, 26(1973), 289–302. Retrieved from http://journals.cambridge.org/abstract_S1352465898264010.
- Morrison, A. P., & Barratt, S. (2010). What are the components of CBT for psychosis? A Delphi study. *Schizophrenia Bulletin*, 36(1), 136–142. <http://doi.org/10.1093/schbul/sbp118>.
- Morrison, A. P., Nothard, S., Bowe, S. E., & Wells, A. (2004). Interpretations of voices in patients with hallucinations and non-patient controls: A

- comparison and predictors of distress in patients. *Behaviour Research and Therapy*, 42(11), 1315–1323. <http://doi.org/10.1016/j.brat.2003.08.009>.
- Myin-Germeys, I., Oorschot, M., Collip, D., Lataster, J., Delespaul, P., & van Os, J. (2009). Experience sampling research in psychopathology: Opening the black box of daily life. *Psychological Medicine*, 39(9), 1533–1547. <http://doi.org/10.1017/S0033291708004947>.
- Myin-Germeys, I., Peeters, F., Havermans, R., Nicolson, N. A., DeVries, M. W., Delespaul, P., & van Os, J. (2003). Emotional reactivity to daily life stress in psychosis and affective disorder: An experience sampling study. *Acta Psychiatrica Scandinavica*, 107(2), 124–131. <https://doi.org/10.1034/j.1600-0447.2003.02025.x>.
- Nayani, T. H., & David, A. S. (1996). The auditory hallucination: A phenomenological survey. *Psychological Medicine*, 26(1), 177. <http://doi.org/10.1017/S003329170003381X>.
- Nezlek, J. B. (2012). *Diary methods for social and personality psychology*. London: Sage.
- Palmier-Claus, J. E., Myin-Germeys, I., Barkus, E., Bentley, L., Udachina, A., Delespaul, P., ... Dunn, G. (2011). Experience sampling research in individuals with mental illness: Reflections and guidance. *Acta Psychiatrica Scandinavica*, 123(1), 12–20. <http://doi.org/10.1111/j.1600-0447.2010.01596.x>.
- Peters, E. R., Lataster, T., Greenwood, K. E., Kuipers, E., Scott, J., Williams, S. L., ... Myin-Germeys, I. (2012a). Appraisals, psychotic symptoms and affect in daily life. *Psychological Medicine*, 42(5), 1013–1023. <http://doi.org/10.1017/S0033291711001802>.
- Peters, E. R., Williams, S. L., Cooke, M. A., & Kuipers, E. (2012b). It's not what you hear, it's the way you think about it: Appraisals as determinants of affect and behaviour in voice hearers. *Psychological Medicine*, 42(7), 1507–1514. <http://doi.org/10.1017/S0033291711002650>.
- Reynold, N., & Scragg, P. (2010). Compliance with command hallucinations: The role of power in relation to the voice, and social rank in relation to the voice and others. *Journal of Forensic Psychiatry & Psychology*, 21(1), 121–138. <http://doi.org/10.1080/14789940903194111>.
- Romme, M. A. J., & Escher, S. (2000). *Making sense of voices*. London, UK: Mind.
- Rucker, J., Newman, S., Gray, J., Gunasinghe, C., Broadbent, M., Brittain, P., ... McGuffin, P. (2011). OPCRIT + : An electronic system for psychiatric diagnosis and data collection in clinical and research settings. *British Journal of Psychiatry*, 199(2), 151–155. <http://doi.org/10.1192/bjp.bp.110.082925>.
- Salkovskis, P. M. (1991). The importance of behaviour in the maintenance of anxiety and panic: A cognitive account. *Behavioural Psychotherapy*, 19(1), 6. <http://doi.org/10.1017/S0141347300011472>.
- Shawyer, F., Mackinnon, A., Farhall, J., Trauer, T., & Copolov, D. (2003). Command hallucinations and violence: Implications for detention and treatment. *Psychiatry, Psychology and Law*, 10(1), 97–107. <http://doi.org/10.1375/pplt.2003.10.1.97>.
- So, S. -W., Chung, L. -H., Tse, C. -Y., Chan, S. -M., Chong, G., Hung, K. -Y., & Sommer, I. E. C. (2020). Moment-to-moment dynamics between auditory verbal hallucinations and negative affect and the role of beliefs about voices. *Psychological Medicine*, 1–7. <https://doi.org/10.1017/S0033291719003611>.
- StataCorp (2015). *Stata Statistical Software: Release 14*. 2015. College Station, TX: Stata Corporation. <http://doi.org/10.2307/2234838>.
- Stone, A. A., Shiffman, S., Atienza, A., & Nebeling, L. (2007). *The science of real-time data capture: Self-reports in health research*. New York: Oxford University Press.
- Thomas, N., McLeod, H. J., & Brewin, C. R. (2009). Interpersonal complementarity in responses to auditory hallucinations in psychosis. *The British Journal of Clinical Psychology/the British Psychological Society*, 48(Pt 4), 411–424. <http://doi.org/10.1348/014466509X411937>.
- Thomas, N., Morris, E. M. J., Shawyer, F., & Farhall, J. (2013). Acceptance and commitment therapy for voices. In E. M. J. Morris, L. C. Johns & J. Oliver (Eds.), *Acceptance and commitment therapy and mindfulness for psychosis*. Chichester, England: Wiley, pp. 95–111.
- Trower, P., Birchwood, M., Meaden, A., Byrne, S., Nelson, A., & Ross, K. (2004). Cognitive therapy for command hallucinations: Randomised controlled trial. *British Journal of Psychiatry*, 184(APR), 312–320. <http://doi.org/10.1192/bjp.184.4.312>.
- van der Gaag, M., Hageman, M. C., & Birchwood, M. (2003). Evidence for a cognitive model of auditory hallucinations. *The Journal of Nervous and Mental Disease*, 191(8), 542–545. <http://doi.org/10.1097/01.nmd.0000082183.95853.ec>.
- Varese, F., Morrison, A. P., Beck, R., Heffernan, S., Law, H., & Bentall, R. P. (2016). Experiential avoidance and appraisals of voices as predictors of voice-related distress. *British Journal of Clinical Psychology*, 55(3), 320–331. <http://doi.org/10.1111/bjc.12102>.
- Vaughan, S., & Fowler, D. G. (2004). The distress experienced by voice hearers is associated with the perceived relationship between the voice hearer and the voice. *The British Journal of Clinical Psychology/The British Psychological Society*, 43(Pt 2), 143–153. <http://doi.org/10.1348/014466504323088024>.
- Walls, T. A., Höppner, B. B., & Goodwin, M. S. (2007). Statistical issues in intensive longitudinal data analysis. In A.A. Stone, S. Shiffman, A.A. Atienza & L. Nebeling (Eds.), *The science of real-time data capture: Self-reports in health research* (pp. 338–360). Oxford University Press.