


Evolutionary and taphonomic implications of a new species of *Amphoracrinus* from the early Viséan of Kentucky

William I. Ausich,¹  Steven C. Koenig,^{2,3} Alan Goldstein,⁴ and Gretel Monreal²

¹Department of Geological Sciences, 155 South Oval Mall, The Ohio State University, Columbus, Ohio 43210, USA <ausich.1@osu.edu>

²Department of Cardiovascular and Thoracic Surgery, University of Louisville, Louisville, Kentucky 40202, USA <gretel.monreal@louisville.edu>

³Department of Bioengineering, University of Louisville, Louisville, Kentucky 40292, USA <steven.koenig@louisville.edu>

⁴Falls of the Ohio State Park, 201 W. Riverside Drive, Clarksville, Indiana 47129-3148, USA <agoldstein@dnr.in.gov>

Abstract.—The youngest species of *Amphoracrinus*, *A. tenax* new species, is described from the Muldraugh Member of the Borden Formation (early Viséan) of north-central Kentucky. With this new occurrence, both the oldest and youngest named species of *Amphoracrinus* are from North America. Numerous Tournaisian and Viséan crinoid faunas are documented in the United States, but only four are known to contain *Amphoracrinus*. Morphological analysis indicates that *A. tenax* is more closely aligned with species from China than with species from Western Europe or other species from North America, where *Amphoracrinus* was most diverse and abundant, which has implications for understanding paleogeographic dispersal. The holotype of *A. tenax* was partially disarticulated on the seafloor before burial, and final burial occurred early during disarticulation. The relative state of disarticulation from pinnules to columns suggests that plates bound only with ligaments disarticulated as a function of surface area of ligaments binding an articulation.

UUID: <http://zoobank.org/c7faf06e-bdd1-43a2-8c10-1364a0aeae0d>

Introduction

An *Amphoracrinus* Austin, 1848 specimen from England was the first illustrated Paleozoic crinoid (Lister, 1673; Ausich and Kammer, 2006). *Amphoracrinus* is now known from Western Europe, the United States, and China (Tables 1 and 2). The genus had a duration of as much as 20 million years from the Famennian through the early Viséan (Mississippian). With the addition of *Amphoracrinus tenax* new species (Muldraugh Member, Borden Formation, early Viséan), 17 named species (one with a named variation) are currently assigned to this genus (Wright, 1943, 1955; Chen and Yao, 1993; Ausich and Sevastopulo, 2001; Donovan et al., 2006; Ausich and Kammer, 2006, 2008a; Webster et al., 2009) (Table 1).

Despite its relatively short temporal duration, *Amphoracrinus* spread across much of the globe in the northern tropics and subtropics, in both western and eastern Laurussia and on the Sibumasu Block, which was part of northern Gondwana during the Mississippian (present-day southern China). The most common occurrences of *Amphoracrinus* are on the Tournaisian mixed carbonate and siliciclastic ramp of southern Ireland and south-western United Kingdom (Hook Head Formation, Ausich and Sevastopulo, 1994, 2001) and in facies associated with Tournaisian and Viséan Waulsortian buildups in Ireland, England, and Wales (Wright, 1955).

The Mississippian, recognized as the “Age of Crinoids” (Kammer and Ausich, 2006; Ausich et al., 2020a), is known for diverse and abundant crinoid occurrences, yet

Amphoracrinus is absent from all but four North American crinoid faunas. Surprisingly, with the description of *A. tenax* n. sp. (Fig. 1), both the oldest named species, *A. viminalis* (Hall, 1863) (Meadville Shale Member, Cuyahoga Formation, early Tournaisian, Kinderhookian, late Hastarian of Ohio), and the youngest species, *A. tenax* n. sp. (Muldraugh Member, Borden Formation, early Viséan, late Osagean, Arundian of Kentucky), occur in the United States. The taphonomic implications of *A. tenax* are also discussed in this study

Location and stratigraphy

This new specimen was collected from the Muldraugh Member of the Borden Formation in north-central Kentucky. This location differs from the Muldraugh location (a quarry) described previously by Ausich et al. (2000) and Kammer et al. (2007). The specimen was collected from a road cut on Kentucky route 313 in Hardin County, approximately 7.3 km west of I-65 (exit 102), coordinates 37°48'23.492"N, 85°49'29.513"W (UTM 16N 591623 4184859) (Fig. 2). The road-cut stratigraphy includes (in ascending order) the New Providence Shale Member of the Borden Formation, the Muldraugh Member of the Borden Formation, and the Harrodsburg Limestone (Kepferle, 1977) that resulted from the time-transgressive, basin-filling progradation of the Borden Delta (e.g., Lane and DuBar, 1983; Richardson and Ausich, 2004). The Muldraugh Member has much lenticular bedding (Ausich et al., 2000), and this new specimen was collected from loose matrix on the surface of a

Table 1. Species of *Amphoracrinus* listed by country occurrence with appropriate international ages as well as regional stages.

Species	Country	International Series	European Regional Stages	North American Stages
<i>A. atlas</i>	UK	Tournaisian-Viséan	lower to upper Chadian	
<i>A. bollandensis</i>	UK	Tournaisian	lower to upper Chadian	
<i>A. compressus</i>	UK	Tournaisian	lower Chadian	
<i>A. gilbertsoni</i>	UK	Tournaisian-Viséan	lower to upper Chadian	
<i>A. granulatus</i>	UK	Tournaisian	Ivorian	
<i>A. portlocki</i>	UK	Tournaisian	lower Chadian	
<i>A. rotundus</i>	UK	Tournaisian	lower Chadian	
<i>A. rugosus</i>	UK	unknown		
<i>A. turgidus</i>	UK	Tournaisian-Viséan	lower to upper Chadian	
<i>A. crassus</i>	Ireland	Tournaisian	Ivorian	
<i>A. gigas</i>	Ireland	Tournaisian-Viséan	lower to upper Chadian	
<i>A. gilbertsoni</i>	Ireland	Viséan	lower to upper Chadian	
<i>A. gilbertsoni</i> var. <i>triangularis</i>	Ireland	Viséan	upper Chadian	
<i>A. granulatus</i>	Ireland	Tournaisian	Ivorian	
<i>A. portlocki</i>	Ireland	Tournaisian	lower Chadian	
<i>A. gigas</i>	Belgium	Tournaisian	Ivorian	
<i>A. rupinus</i>	USA	Viséan	lower Chadian	middle Osagean
<i>A. tenax</i> n. sp.	USA	Viséan	Arundian	upper Osagean
<i>A. viminalis</i>	USA	Tournaisian	upper Hasterian	upper Kinderhookian
<i>A. cheni</i>	China	Tournaisian		
<i>A. pseudoturgidus</i>	China	Tournaisian		
<i>A.?</i> <i>atlas</i>	China	Tournaisian		
<i>A.?</i> <i>bollandensis</i>	China	Tournaisian		
<i>A.?</i> <i>gilbertsoni</i>	China	Tournaisian		
<i>Amphoracrinus</i> sp.	China	Famennian		

Table 2. Chronostratigraphy of the Mississippian. Correlation of International series, European and North American regional stages, and time bins of Ausich and Kammer (2006) and Kammer and Ausich (2007).

System	International Series	Mississippian Time Bins	European Regional Stages	North American Regional Stages
Mississippian	Serpukhovian	10	Pendelian	Chesterian
	Viséan	9	Brigantian	
		8	Asbian	Meramecian
		7	Holkerian	
		6	Arundian	
		5	upper Chadian	Osagean
	4	lower Chadian		
	Tournaisian	3	Ivorian	Kinderhookian
		2	upper Hasterian	
		1	lower Hasterian	

lenticular bed approximately 28 m above the New Providence Shale Member (Fig. 2).

The Muldraugh Member was a mixed carbonate-siliclastic, tempestite-dominated ramp environment and the shallow-water equivalent of the Fort Payne Formation in Kentucky and Tennessee (Nicoll and Rexroad, 1975; Peterson and Kepferle, 1970; Kammer and Ausich, 1987; Ausich and Meyer, 1990; Sable and Dever, 1990; Ausich et al., 2000). The Muldraugh Member consists of argillaceous, coarse-grained packstones, yellow-gray dolomitic siltstone, silty dolomite, crinoidal limestone, shale, chert, and geodes (Sable et al., 1966; Kepferle, 1977; Sable and Dever, 1990; Ausich et al., 2000). In the

Muldraugh Member, advanced cladids, followed by camerates, dominated in both species richness and the number of specimens (Ausich et al., 2000). Other contemporaneous deltaic localities with a comparably diverse crinoid fauna include the New Providence Shale Member (prodeltaic environments) in southern Indiana and north-central Kentucky (Ausich et al., 1979; Kammer, 1984), the Edwardsville Formation (delta platform environments) in central and southern Indiana (Lane, 1963, 1973; Van Sant and Lane, 1964; Ausich and Lane, 1982), and the Fort Payne Formation of south-central Kentucky (Krivicich et al., 2014 and references cited therein). In other locations, the Muldraugh Member contains a diverse assemblage, including



Figure 1. *Amphoracrinus tenax* n. sp. holotype from the Muldraugh Member (early Viséan) of the Borden Formation, north-central Kentucky. Posterior side of entire specimen. USNM PAL 771320, holotype.

brachiopods, bryozoans, corals (primarily rugosans), conulariids, other echinoderms, gastropods (abundant platyceratids), sponges (abundant hexactinellids), arthropods, trace fossils, and vertebrates (Chondrichthyes teeth and dorsal fin spines). However, in the bed containing *A. tenax*, the only other skeletal elements are crinoid columnals and pluricolumnals.

Methods

Table 1 is a list of taxa considered in this study, including their distribution and age. Note that Ausich and Kammer (2008a) designated *A. blairi* Miller and Gurley, 1896a and *A. jessieae* Miller and Gurley, 1896b as nomen dubia; *Amphoracrinus rouchi* Delpy, 1941 was transferred to *Ectocrinus* by Webster et al. (2004); and *A. atlas* (M'Coy, 1849), *A. bollandensis* Wright, 1955, and *A. gilbertsoni* (Miller in Phillips, 1836) are only questionably recognized in China (Webster et al., 2009). In addition, Webster and Waters (2009) described two poorly preserved Devonian crinoids from northwestern China as *Amphoracrinus* sp. These specimens are from the Hongguleng Formation (Famennian). Unfortunately, both specimens are too poorly preserved to code their morphological characters.

Cluster analyses were conducted using PAST 4.04 (Hammer and Harper, 2006; Hammer et al., 2006). The paired group (UPGMA) algorithm, the Euclidean similarity index, and the no constraints setting were employed to generate dendrograms with the characters listed in Supplemental Tables 1, 2.

Parsimony analyses were performed using PAUP 4.0a168 (Swofford, 2015). Analyses were run using the criterion of maximum parsimony and heuristic searches with random addition repeated 1,000 times. All characters were treated as equally weighted and unordered. Branch swapping was with the tree bisection–reconnection algorithm. Strict consensus, 50% majority rule, and Adams consensus trees were evaluated, the latter to identify potential “wildcard taxa” for possible elimination (Wiley and Lieberman, 2011). The consistency index (ci), retention index (ri), and rescaled index (rc) were calculated, and new trees were generated using rescaled consistency indices. Bayesian, tip-dating method was not attempted because there is insufficient temporal resolution for this approach. Combined, the outgroup taxa span much of the Devonian, but all ingroup species belong to either one or both of two time bins (Tournaisian and Viséan). Tree support was evaluated with bootstrap values, jackknife values, and Bremer support.

Three outgroup taxa are designated. All are from the Periechocrinidae, which were probably ancestors of the Amphoracrinidae (Ausich and Kammer, 2008a, 2008b). Outgroup taxa are *Athabascacrinus orientale* (Waters et al., 2003) (Famennian, China), *Pyxidocrinus collensis* Breimer, 1962 (Emsian, Spain), and *Megistocrinus depressus* (Hall, 1862) (Eifelian-Givetian, United States). Numerous morphological characters were coded; however, based on results of PAUP, only 16 were informative for 14 species of *Amphoracrinus* and three outgroup taxa (Table 1, Supplementary Tables 2, 3). The initial parsimony analysis yielded 1,460 trees with a length of 55. An Adams consensus tree yielded more than one polytomy, and the following five wildcard species were identified and eliminated from further analyses: *A. atlas*, *A. bollandensis*, *A. portlocki* Wright, 1955, *A. rotundus* Wright, 1955, and *A. turgidus* Wright, 1943.



Figure 2. Road cut in Hardin County, north-central Kentucky, where *A. tenax* n. sp. was collected from the Muldraugh Member of the Borden Formation. Lithostratigraphy noted on right; arrow indicates its location on the bench.

With these wildcard taxa eliminated, the resulting analysis yielded 91 trees of length 45. However, as noted, this tree lacked any bootstrap, jackknife, or Bremer support. Next, trees were generated using rescaled consistency indices (rc), resulting in 12 trees of length 26.64722 (Fig. 3.2) that also lacked any bootstrap, jackknife, or Bremer support.

Repository and institutional abbreviation.—The holotype of *A. tenax* n. sp. is deposited in the Smithsonian National Museum of Natural History (USNM), USNM PAL 771320.

Systematic paleontology

The superfamilial classification used here follows Cole (2017), Wright (2017), and Wright et al. (2017); family-level classifications follow Moore and Teichert (1978). Morphologic terminology follows Webster (1974), Ubaghs (1978b), and Ausich et al. (2020b). The plating of interrays is given by the number of plates in each range, from proximalmost plate to the last range before the tegmen. In the posterior interray, the pramanal is indicated by “P,” and the first interradial plate in regular interrays is indicated by “1.” A “?” indicates that more-distal plating is unknown (Ausich, 2021a). Abbreviations used in designating measurements include the following: CrH, crown height; CaH, calyx height; CaW, calyx width; CoH, column height. An * indicates a measurement was incomplete.

Class Crinoidea Miller, 1821
 Subclass Camerata Wachsmuth and Springer, 1885
 Infraclass Eucamerata Cole, 2017
 Order Monobathrida Moore and Laudon, 1943
 Suborder Composocrinina Ubaghs, 1978a
 Superfamily Periechocinacea Bronn, 1849
 Family Amphoracrinidae Bather, 1899
 Genus *Amphoracrinus* Austin, 1848

Type species.—*Amphoracrinus gilbertsoni* Miller in Phillips, 1836.

Included species.—*A. atlas* (M’Coy, 1849); *A. bollandensis* Wright, 1955; *A. cheni* Webster et al., 2009; *A. compressus* Wright, 1943; *A. crassus* (Austin and Austin, 1843); *A. gigas* Austin in Wright, 1955; *A. gilbertsoni* (Miller in Phillips, 1836); *A. granulatus* (Austin and Austin, 1843); *A. portlocki* Wright, 1955; *A. pseudoturgidus* Chen and Yao, 1993; *A. rotundus* Wright, 1955; *A. rugosus* (Rofe, 1865); *A. rupinus* Webster and Lane, 1987; *A. tenax* n. sp.; *A. turgidus* Wright, 1943; and *A. viminalis* (Hall, 1863).

Amphoracrinus tenax new species
 Figures 1, 4

Type.—The holotype of *A. tenax* n. sp. is designated USNM PAL 771320.




Figure 3. Results from morphological analyses of *Amphoracrinus*. (1) Dendrogram from cluster analysis of morphological characters of *Amphoracrinus* species. (2) A 50% majority-rule tree of nine species of *Amphoracrinus* with three outgroup taxa. Values at nodes indicate the percentage of most parsimonious trees that recovered that relationship (data in Supplemental Tables 1, 2, 3).




Figure 4. Holotype of *A. tenax* n. sp. USNM PAL 771320. (1, 2) Well-articulated posterior side of the crown; (3) proximal portion of column, compare with Figure 1; (4) portion of middle column, compare with Figure 1; (5) poorly articulated anterior side of crown. (2–5) Coated with ammonium chloride.

Diagnosis.—Calyx low cone shape; calyx plates convex, smooth; basal circler 20% of calyx height, ridge at base of calyx absent; radial plates as high as wide, unequal in size, smaller than primanal; three interradsial plates in regular interrays with distalmost range adjacent to primibrachials, regular interrays depressed; first fixed primibrachial tetragonal, as high as wide; CD interray very much wider than other regular interrays, three plates above primanal; first primibrachial smaller than radial plate and primaxil, more than five CD-interray plates in calyx; two fixed primibrachials, 15–20 free arms project upward and branch.

Occurrence.—Muldraugh Member of the Borden Formation, Hardin County, north-central Kentucky, 37°48'23.492"N, 85°49'29.513"W, UTM 16N 591623 4184859.

Description.—Calyx medium sized, low cone shape (Fig. 4.1), ~0.52 times higher than wide; arms grouped and interrays moderately depressed; calyx plates low convex shape, smooth plate sculpturing. Basal circler high, visible in side view, ~20% of calyx height (Fig. 4.2); three basal plates, unequal in size, wider than high, much smaller than radial plates. Radial circler ~35% of calyx height; radial plates five, hexagonal, as high as wide. Regular interrays narrowly in contact with tegmen, all plates convex as noted; first interradsial hexagonal, as high as wide, smaller than radials and first primibrachial; plating in second range unknown. Primanal heptagonal, higher than wide, largest plate in calyx, interrupts the radial circler; plating in CD interray P–3–5–? (Fig. 4.2), plates have impressed sutures; second range of interradsial plates in posterior with subtle plate plications that are ridges that connect with like ridges on adjoining plates (radials, primanal, first and second primibrachials, adjacent plates in the first and second ranges of interradsial plates, and some plates of the third range of posterior interray plates) (Fig. 4.2). CD interray in contact with tegmen. First primibrachial tetragonal or pentagonal, approximately as high as wide, convex, smaller than radial plates and primaxil; second primibrachial axillary, approximately twice as wide as high. Intrabrachial plates between adjacent half-rays absent. Tegmen not visible.

Three or four free arms per ray (C ray four arms; A and D rays three arms). Free arms begin with first tertibrachial, first tertibrachial either uniserial or biserial (Fig. 4.2); all subsequent brachials biserial. Free arms robust, aborally rounded; directed obliquely upward, narrow distally, and abaxially incurved at distal end of crown. Each free arm may have as many as four branches (Fig. 4.2), with branching pattern variable above the first branch (may be either endotomous or exotomous) (Fig. 4.2). Pinnules slender. Column circular (Figs. 1, 4); in proximal column heteromorphic (N212 pattern), latus of nodals very convex and latus of internodals progressively less convex (Fig. 4.3); mesistele with very subtle N212 pattern and some places approach homeomorphy, latus of columnals convex (Fig. 4.4).

Etymology.—The name, *tenax* (m), is Latin for steadfast or tenacious.

Measurements.—CrH, 90.0; CaH, 25.0*; CaW, 12.0; CoH, 230*.

Remarks.—USNM PAL 771320 is the most complete known specimen of this genus. Most species are known from only the theca (calyx and tegmen). Because the arms are largely complete, they hide the tegmen, so comparison to other species is limited to characters of the calyx.

Amphoracrinus tenax n. sp. differs from other North American species because *A. rupinus* has a very low cone shape, coarsely granulose plate sculpturing, a continuous ridge around the base of the calyx, radial plates wider than high and the largest plates in the calyx, most-distal plates in calyx interrays that abut secundibrachials, first primibrachial wider than high, CD interray wider than other interrays, three or four extra plates in the CD interray, and 10 free arms. By contrast, *A. viminalis* has a very low bowl-shaped calyx, coarsely granulose plate sculpturing, a broken ridge around the base of the calyx, radial plates wider than high and the largest plates in the calyx, distal-most interradsials in calyx that abut secundibrachials, first primibrachial wider than high, CD interray wider than other interrays, two plates above the primanal, and 10 free arms (see Supplemental Table 5 for comparison with other species).

Results and discussion

Paleogeography and evolution.—*Amphoracrinus viminalis*, the oldest named species, is from northeastern Ohio from the Meadville Shale Member of the Cuyahoga Formation and dates to the late Kinderhookian (Tournaisian, late Hastarian) (Table 1). The Meadville Shale was deposited in a shallow marine deltaic tempestite shelf setting that resulted in excellent crinoid preservation (Ausich and Roeser, 2012; Kammer and Roeser, 2012). *A. rupinus* (Webster and Lane, 1987) is from southern Nevada (Meadow Valley Range) from the Anchor Limestone of the Monte Cristo Group (early Osagean, Tournaisian, Ivorian), where monobathrid camerates were the dominant element of the crinoid fauna (Webster and Lane, 1987). *A. rupinus* was also documented in New Mexico from the Nunn Member of the Lake Valley Formation (early Osagean/late Tournaisian) (Rhenberg and Kammer, 2013). The Nunn Member in general was a narrow and relatively deepwater low-energy carbonate ramp with Waulsortian mounds; however, the specific location where *Amphoracrinus* occurred was from a shallower shelf subenvironment comparable to the Burlington Shelf (Rhenberg and Kammer, 2013). *Amphoracrinus tenax* n. sp. is now the youngest (early Viséan, Arundian) described species of *Amphoracrinus*.

Amphoracrinus is best known from Western Europe, where it occurs from the Tournaisian (Ivorian) to the early Viséan (upper Chadian) in England, Wales, Ireland, and Belgium. In Ireland, *Amphoracrinus* is present in the Hook Head Formation (Tournaisian, Ivorian), particularly the shallow-water “Michelinia Beds” (Ausich and Sevastopulo, 2001), which was a mixed carbonate and siliclastic tempestite shelf-ramp (Ausich and Sevastopulo, 1994). It is also present in the equivalent ramp setting in southwestern United Kingdom. *Amphoracrinus* was a dominant faunal element in the Tournaisian (lower Chadian) to the Viséan (upper Chadian) Waulsortian-related facies in England, Wales, Ireland, and Belgium (Wright, 1955; Ausich and Sevastopulo, 1994, 2001; Ausich and Kammer, 2006, 2008a; Donovan et al., 2006).

As noted, the oldest occurrence of *Amphoracrinus* is from the Famennian of China (Webster and Waters, 2009). Chen and Yao (1993) described *A. atlas*, *A. bollandensis*, *A. gigus*, *A. gilbertsoni*, and *A. pseudoturgidus* from the Tournaisian of China. These species assignments were revisited by Webster et al. (2009), who described *A. cheni* and *A. pseudoturgidus*; but they recognized the occurrences *A. atlas?* and *A. gilbertsoni?* as questionable. Both the Famennian and Tournaisian occurrences in China are interpreted to have been from shallow-water habitats with no clear evidence of association with carbonate buildups.

Cluster analysis was performed on morphological characters (Supplemental Tables 1, 2) to identify any morphological similarities among *Amphoracrinus* species within and between geographic areas. In the resultant dendrogram, the three North American species are not grouped together. *Amphoracrinus tenax* n. sp. and *A. pseudoturgidus* form a group at the base of the dendrogram, and *A. crassus*, *A. rupinus*, and *A. viminalis* form a group (Fig. 3.1). *Amphoracrinus bollandensis*, *A. gigas*, and *A. cheni* group together, and the remaining Western European species (with *A. rupinus* and *A. viminalis*) group together in various clusters.

Phylogenetic analyses were also performed. Whereas the phylogenetic tree in Figure 3.2 cannot be demonstrated to be a robust phylogenetic solution, it displays interesting relationships, with European, North American, and Chinese species intermixed. For example, *A. tenax*, *A. cheni*, and *A. pseudoturgidus* (Tournaisian, China) are ladderized at the base of the ingroup (Fig. 3.2). By contrast, the two other North American species (*A. viminalis*, Tournaisian, and *A. rupinus*, Viséan) are in a clade with all European species. *Amphoracrinus rupinus* and *A. crassus* are in the crownward clade and are on a polytomy. *A. viminalis* is in a basal position of the crownward group. Thus, the three North American species are in reverse stratigraphic order and are associated with species of contrasting paleogeographies. Interpretation of the phylogenetic analysis (Fig. 3.2) is regarded as tentative, but the repeated recovery of relationships at the base of the tree suggest that these taxa have close phylogenetic and paleogeographic relationships. The older species of North American *Amphoracrinus* (*A. rupinus* and *A. viminalis*) are most closely associated with European species, whereas the youngest North American species (*A. tenax* n. sp.) is more closely associated with species from China, which may suggest either separate paleogeographic origins or multiple migration events for North American members of this genus. These relationships will not be resolved until characters of the tegmen, arms, and column are known for most species, which is unlikely. At least tegmen characters for *A. tenax* may become known with additional specimens, which is significant because tegmen characters are highly variable and phylogenetically important (Kammer et al., 2013; Ausich and Kammer, 2016).

Ausich and Kammer (2013) evaluated the temporal and paleogeographic patterns among Mississippian crinoids in North America and Western Europe. On the basis of analyses of first occurrences of genera, paleobiogeography through the history of a genus, and biodiversity changes through the Mississippian, Ausich and Kammer (2013) concluded that the evolutionary center of monobathrid camerate crinoids, disparid crinoids, and cyathocrin crinoids was predominantly in North America, with subsequent migration to Western Europe. A

major factor driving migration was interpreted to be sea-level rise following the late Kinderhookian glaciation (Walker et al., 2002; Caputo et al., 2008; Kammer and Matchen, 2008). Waters et al. (2008) and Webster et al. (2009) concluded that most Chinese Mississippian crinoids migrated from Western Europe. Thus, it follows that the Tournaisian crinoid faunas from China were originally seeded by migration from North America and Europe as a result of sea-level rise following late Kinderhookian glaciations.

With *Amphoracrinus* sp. (Webster and Waters, 2009) (Famennian) assigned to *Amphoracrinus*, a more-nuanced paleogeographic history must have existed. Origination of *Amphoracrinus* in northwestern China during the Famennian would be consistent with the overall conclusions of several studies that indicate that Famennian blastoids and crinoids in China have close affinities to early Mississippian crinoids in North America (Lane et al., 1997; Waters et al., 2003, 2008; Webster and Waters, 2009). This would suggest that during the latest Devonian and earliest Mississippian, migration routes were more freely open than traditionally understood between China, North America, and Western Europe, including westward migration.

A potential history of *Amphoracrinus* is that this genus originated during the Late Devonian of China, where it thrived in shallow-water, nonbuildup-associated habitats. It then migrated westward, with the first preserved occurrence in western Laurussia (North America) in a siliciclastic deltaic setting (*A. viminalis*) and the first occurrence in central Laurussia (Western Europe) in a mixed carbonate–siliciclastic setting (*A. crassus*, *A. granulatus*, and *A. gigas*) (Table 1). Younger species in China and North America persisted in mixed carbonate–siliciclastic settings, whereas in Western Europe, *Amphoracrinus* thrived in facies associated with Waulsortian mudmounds as well as other settings. As noted, the morphological and phylogenetic analyses suggest that migratory pathways were open throughout the Famennian–Viséan interval.

To better resolve the paleogeographic history of *Amphoracrinus*, a more-detailed understanding of *Amphoracrinus* sp. from the Famennian of China is required, as well as more occurrences of the genus from both the late Devonian and Tournaisian. Analyses of multiple clades through this interval are also needed to more fully understand this important episode in crinoid evolutionary history.

Taphonomy.—*Amphoracrinus tenax* n. sp. is an illustrative example of the earliest phases of disarticulation of a robust camerate crinoid. Preservation of two sides of this specimen are strikingly different (Fig. 4.2, 4.5). This preservational contrast is well known for many clades with multielement skeletons (e.g., see Brett and Baird, 1986 and Meyer and Milsom, 2001 for discussions on this phenomena). This style of preservation has been noted for Paleozoic (Meyer et al., 1981; Brett and Baird, 1986) and Mesozoic (Springer, 1901; Struve, 1957; Seilacher et al., 1968; Hagdom, 1978, 1998; Neugebauer, 1978; Simms, 1986; Milsom et al., 1994; Hess, 1999) crinoids.

Brett and Baird (1986) discussed this differential preservation in the Silurian camerate crinoid *Eucalyptocrinites* Goldfuss, 1831. In this Silurian crinoid, arms were preserved mostly intact on the lower surface, but arms were absent from the upper surface. In both cases, calyx plates were present but displaced due to compaction.

The *A. tenax* (Figs. 1, 4) holotype USNM PAL 771320 is important for understanding the taphonomy of the crown and the significant height of the column. Further, this allows consideration of relative disarticulation of nearly the entire skeleton of a crinoid inferred to have had only ligamentary articulations. The CD interray is very well preserved, whereas the opposite side (A ray) is in a much more advanced stage of disarticulation. The A ray was the stratigraphically up side when this specimen was found (Fig. 4.1, 4.2). On the A-ray side, the radial plates and fixed brachials remain firmly sutured, but the interrarial plates are slightly askew (Fig. 4.5). Every free arm is disarticulated from the calyx. Most of the preserved arms are in approximately the correct position, but the biserial brachials are beginning to separate along some arm segments. In other places, brachials are completely disarticulated but remain as part of the specimen. Pinnules are in the correct position and articulated, in their correct position but slightly askew, or completely disarticulated. By contrast, the CD interray, the stratigraphically down side of the specimen, was buried in the sediment. Calyx plates are sutured; all free arms are in place, with three having minor displacement (Fig. 4.1, 4.2). Proximal arms are mostly intact, but in the middle and distal parts of the arms some disarticulation is present. However, the distal, narrower, incurled arms are intact. A few pinnules are intact, but they are mostly disarticulated. Pinnules may be approximately in place or completely disarticulated. The proximal and middle column are completely articulated, but the distal column is absent (Figs. 1, 4.3, 4.4)

Actualistic studies of disarticulation and preservation have been conducted with articuliform crinoids that have muscles in their arms, and muscles are the most rapidly decaying tissue (Meyer, 1971; Liddell, 1975; Meyer and Oji, 1992). By contrast, Mississippian camerate crinoids are inferred to have had only ligamentary connective tissues, on the basis of both brachial articulations and disarticulation patterns (e.g., Lane and Burke, 1976; Ausich and Baumiller, 1993).

A possible scenario for the preservation of USNM PAL 771320 is that the crown and 230 mm of the column broke away from the more-distal column, presumably by a storm or gravity flow (Taylor and Brett, 1996). The crinoid was transported some unknown distance and was only partially buried. The enclosing matrix is uniformly fine grained, so if it was displaced by a storm or gravity flow it was likely transported in the upper, finer-grained portion of the resulting deposit, as discussed by Ausich and Meyer (1990), Taylor and Brett (1996), and Ausich (2021b). The upper, more-exposed surface decayed and disarticulated more rapidly, and the lower buried surface may have decayed more slowly. The specimen was eventually permanently buried before it was completely disarticulated. Current action did not disperse skeletal elements, and it does not appear to have been significantly attacked by scavengers. However, note the lack of plates on the lower arms (left side of Fig. 4.1, 4.2 and the opposite side of Fig. 4.5). It is possible that after burial, this specimen was disturbed by a burrowing organism (Maples and Archer, 1989) (other trace fossils occur in this unit). Thus, although buried and intact, all connective tissues had decayed. Finally, the specimen was compacted and lithified, yielding its present condition (Figs. 1, 4).

The poor articulation on the upper surface was either a reflection of differential disarticulation on the seafloor or a reflection of

differential decay of connective tissues and disarticulation manifested by compaction. Overall, the pinnules are the elements displaying the most disarticulation, followed by the arms, calyx, and column. Camerate calyx plates are traditionally considered to have been cemented to one another more than in other clades during life (e.g., Ubahgs, 1978a); however, their more-secure articulations may have been due to stereom interlocking (Gorzelaek et al., 2012; Ausich, 2021b). The difference in disarticulation propensity of pinnules, arms, and column may be the result of surface area of ligamentary articulations, with the smaller pinnular facets more likely than the larger brachial facets to disarticulate. The largest ligamentary articulations between columnals are all intact. This result is consistent with studies by Allison (1990) and Breton (1997), who concluded that disarticulation rates among asteroid clades were also correlated with plate size and robustness.

During decay of a dead extant crinoid, Meyer and Oji (1992) convincingly demonstrated that some sort of a microbial sheath was responsible for the excellent preservation of slabs of *Uintacrinus socialis* Grinnell, 1876, which includes numerous specimens with the down sides very well preserved and the up-side plates disarticulated. Crinoid preservation enhancement by microbial sheaths has been hypothesized in other examples of exceptional preservation (e.g., Seilacher et al., 1985). It is not possible to determine whether a microbial sheath enhanced the preservation of this Muldraugh Member crinoid because, if present, a microbial sheath would not be likely to have been preserved in sediments that lack any evidence of having been anoxic.

Acknowledgments

We thank D. Cooper and B. Cooper of Trilobites of America for preparation of this specimen. G. Monreal is supported in part by a gift from R.M. Prizant to the Legacy Foundation of Kentucky. We also thank G.D. Sevastopulo and J.-P. Lin for discussion of stratigraphy and paleogeography in Ireland and China; J.A. Waters for discussion of paleoenvironmental occurrences in China; D.F. Wright for discussion of numerical analyses; and S.R. Cole, T.W. Kammer, and S. Zamora for reviews that improved this manuscript.

Data availability statement

Data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.jq2bvq889>.

References

- Allison, P.A., 1990, Variation in rates of decay and disarticulation of Echinodermata: implications for the application of actualistic data: PALAIOS, v. 5, p. 432–440.
- Ausich, W.I., 2021a, Glossary of patterns, abbreviations, and symbols to designate crinoid morphology, in Ausich, W.I., ed., Treatise Online no. 145, Part T, Revised, Chapter 6, 5 p., journals.ku.edu/treatiseonline/article/view/15161
- Ausich, W.I., 2021b, Disarticulation and preservation of fossil echinoderms: recognition of ecological-time information in the echinoderm fossil record: Elements of Paleontology, 56 p., <https://doi.org/10.1017/9781108893374>
- Ausich, W.I., and Baumiller, T.K., 1993, Taphonomic method for determining muscular articulations in fossil crinoids: PALAIOS, v. 8, p. 477–484.
- Ausich, W.I., and Kammer, T.W., 2006, Stratigraphical and geographical distribution of Mississippian (lower Carboniferous) Crinoidea from England and Wales: Proceedings of the Yorkshire Geological Society, v. 56, p. 91–109.

- Ausich, W.I., and Kammer, T.W., 2008a, Generic concepts in the Amphoracrinidae Bather, 1899 (class Crinoidea) and evaluation of generic assignments of North American species: *Journal of Paleontology*, v. 82, p. 1139–1149.
- Ausich, W.I., and Kammer, T.W., 2008b, Evolution and extinction of a Paleozoic crinoid clade: phylogenetics, paleogeography, and environmental distribution of the periechocrinids, in Ausich, W.I., and Webster, G.D., eds, *Echinoderm Paleobiology*: Bloomington, Indiana University Press, p. 144–171.
- Ausich, W.I., and Kammer, T.W., 2013, Mississippian crinoid biodiversity, biogeography, and macroevolution: *Palaeontology*, v. 56, p. 727–740.
- Ausich, W.I., and Kammer, T.W., 2016, Exaptation of pelmatozoan oral surfaces: constructional pathways in tegmen evolution: *Journal of Paleontology*, v. 90, p. 689–720.
- Ausich, W.I., and Lane, N.G., 1982, Crinoids from the Edwardsville Formation (Lower Mississippian) of southern Indiana: *Journal of Paleontology*, v. 56, p. 1343–1361.
- Ausich, W.I., and Meyer, D.L., 1990, Origin and composition of carbonate build-ups and associated facies in the Fort Payne Formation (Lower Mississippian, south-central Kentucky): an integrated sedimentologic and paleoecologic analysis: *Geological Society of America Bulletin*, v. 102, p. 129–146.
- Ausich, W.I., and Roeser, E.W., 2012, Camerate and disparid crinoids from the late Kinderhookian Meadville Shale Cuyahoga Formation from Ohio: *Journal of Paleontology*, v. 86, p. 488–507.
- Ausich, W.I., and Sevastopulo, G.D., 1994, Taphonomy of lower Carboniferous crinoids from the Hook Head Formation, Co. Wexford, Ireland: *Lethaia*, v. 27, p. 245–256.
- Ausich, W.I., and Sevastopulo, G.D., 2001, The lower Carboniferous (Tournaisian) crinoids from Hook Head, County Wexford, Ireland: *Monograph of the Palaeontographical Society*, p. 617, 136 p.
- Ausich, W.I., Kammer, T.W., Lane, N.G., 1979, Fossil communities of the Borden (Mississippi) delta in Indiana and northern Kentucky: *Journal of Paleontology*, v. 53, p. 1182–1196.
- Ausich, W.I., Goldstein, A., and Yates, R., 2000, Crinoids from the Muldraugh Member of the Borden Formation in north-central Kentucky (Echinodermata, Lower Mississippian): *Journal of Paleontology*, v. 74, p. 1072–1082.
- Ausich, W.I., Kammer, T.W., and Mirantsev, G.V., 2020a, Carboniferous crinoids, in Lucas, S., Schneider, S.G., Wang, J.W., and Nikolaeva, S., eds., *The Carboniferous Timescale*: Geological Society of London Special Publications 512, <https://doi.org/10.1144/SP512-2020-71>
- Ausich, W.I., Wright, D.F., Cole, S.R., and Sevastopulo, G.D., 2020b, Homology of posterior interarray plates in crinoids: a review and new perspectives from phylogenetics, the fossil record, and development: *Palaeontology*, v. 63, p. 552–545.
- Austin, T., 1848, Observations on the Cystidea of M. von Buch, and the Crinoidea generally: *Geological Society of London, Quarterly Journal*, v. 4, p. 291–294.
- Austin, T., and Austin, T., 1843–1847, A Monograph on Recent and Fossil Crinoidea, with Figures and Descriptions of Some Recent and Fossil Allied Genera, v. 1, 2, p. 1–32, pl. 1–4, frontispiece (1843); v. 3, p. 33–48, pl. 5, 6 (1844); v. 4, p. 49–64, pl. 7, 8 (1845); v. 5, p. 65–80, pl. 9, 10 (1846); v. 6–8, p. 81–128, pl. 11–16 (1847); 128 p., 16 pl.
- Bather, F.A., 1899, A phylogenetic classification of the Pelmatozoa: Report of the British Association for the Advancement of Science 68th Meeting, p. 916–923.
- Breimer, A., 1962, A monograph on Spanish Paleozoic Crinoidea: *Overdruk uit Leidse Geologische Mededelingen*, v. 27, 190 p.
- Breton, G., 1997, Deux étoiles de mer du Bajocien du nord-est du bassin de Paris (France): leur allies actuels sond des fossils vivants: *Bulletin trimestriel de la Société Géologique de Normandie et des Amis du Museum du Havre*, v. 84, p. 23–34.
- Brett, C.E., and Baird, G.C., 1986, Comparative taphonomy: a key to paleo-environmental interpretation based on fossil preservation: *PALAIOS*, v. 1, p. 207–227.
- Bronn, H.G., 1849, Index palaeontologicus, unter Mitwirkung der Herren Prof. H.R. Göppert und H. von Meyer, *Handbuch einer Geschichte der Natur*, 5, Abt. 1, A: Nomenclator Palaeontologicus, 1381 p.
- Caputo, M.V., de Melo, J.H.G., Streeb, M., and Isbell, J.L., 2008, Late Devonian and early Carboniferous glacial records of South America, in Fielding, C.R., Frank, T.D., and Isbell, J.L., eds., *Resolving the Late Paleozoic Ice Age in Time and Space*: Geological Society of America Special Paper 441, p. 161–173.
- Chen, Z.-T., and Yao, J.-H., 1993, Palaeozoic Echinoderm Fossils of Western Yunnan, China: Beijing, Geological Publishing House, 102 p.
- Cole, S.R., 2017, Phylogeny and morphological evolution of the Ordovician Camerata (Class Crinoidea, Phylum Echinodermata): *Journal of Paleontology*, v. 91, p. 815–828.
- Delpey, G., 1941, Crinoïdes du viséan supérieur dans le pays de Skouva, au Maroc: *Bulletin de la Société des Sciences Naturelles du Maroc Rabat*, v. 19, p. 214–220.
- Donovan, S.K., Lewis, D.N., and Kabra, P., 2006, A dense epizoobiontic infestation of a lower Carboniferous crinoid (*Amphoracrinus gilbertsoni* (Phillips)) by *Oichnus paraboloides* Bromley: *Ichnos*, v. 13, p. 43–45.
- Goldfuss, G.A., 1826–1844, *Petrefacta Germaniae*, tam ea, Quae in Museo Universitatis Regiae Borussicae Fridericiae Wilhelmae Rhenanae, serventur, quam alia quaecunque in Museis Hoeninghusiano Muensteriano aliisque, extant, iconibus et descriptionibus illustrata. -- Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angränzende Länder, unter Mitwirkung des Herrn Grafen Georg zu Münster, herausgegeben von August Goldfuss, v. 1 (1826–1833), Divisio prima. Zoophytorum reliquiae, p. 1–114; Divisio secunda. Radiariorum reliquiae, p. 115–221 [Echinodermata]; Divisio tertia. Annulatorum reliquiae, p. 222–242; v. 2 (1834–1840), Divisio quarta. Molluscorum acephalicorum reliquiae. I. Bivalvia, p. 65–286; II. Brachiopoda, p. 287–303; III. (1841–1844), Divisio quinta. Molluscorum gasteropodum reliquiae, p. 1–121; atlas of plates, 1–199, Düsseldorf, Arnz & Co. v. 1, p. 1–76 (1826), p. 77–164 (1829), p. 165–240 (1831), p. 241–252 (1833); v. 2, p. 1–68 (1833), p. 69–140 (1836), p. 141–224 (1837), p. 225–312 (1840); v. 3, p. 1–128 (1844).
- Gozelak, P., Stolarski, J., Mazur, M., and Meibom, A., 2012, Micro- to nanostructure and geochemistry of extant crinoidal echinoderm skeletons: *Geobiology*, 15 p., <https://doi.org/10.1111/gbi.12012>
- Grinnell, G.B., 1876, On a new crinoid form the Cretaceous formation of the West: *American Journal of Science*, v. 3, p. 81–83.
- Hagdorn, H., 1978, Muschel/Krinoiden-Bioherme im Oberen Muschelkalk (mol. I, Anis) von Craillsheim und Schwäbisch Hall (Süd-westdeutschland): *Zeitschrift für Hobbypaläontologie Abhandlungen*, v. 156, p. 31–86.
- Hagdorn, H., 1998, *Traumatocrinus*, eine long verkannte Seelilie aus der Triassic: *Zeitschrift für Hobbypaläontologie Abhandlungen*, v. 15, p. 269–275.
- Hall, J., 1862, Preliminary notice of some of the species of Crinoidea known in the Upper Helderberg and Hamilton groups of New York: *New York State Cabinet of Natural History 15th Annual Report*, p. 87–125.
- Hall, J., 1863, Preliminary notice, of some species of Crinoidea from the Waverly Sandstone series of Summit Co., Ohio, supposed to be of the age of the Chemung Group of New York: *Preprint of Seventeenth Annual Report of the Regents of the University of the state of New-York, on the Condition of the State Cabinet of Natural History, and the Historical and Antiquarian Collection annexed thereto, State of New York in Senate Document 189*, Albany, Comstock and Cassiday Printers, p. 50–60.
- Hammer, Ø., and Harper, D.A.T., 2006, *Paleontological Data Analysis*: Oxford, Blackwell, 352 p.
- Hammer, Ø., Harper, D.A.T., and Ryan, P.D., 2006, PAST: Paleontological Statistics Software Package for Education and Data Analysis: *Palaeontologica Electronica*, v. 4, 9 p., https://palaeo-electronica.org/2001_1/past/issue1_01.htm
- Hess, H., 1999, Middle Jurassic of northern Switzerland, in Hess, H., Ausich, W.I., Brett, C.E., and Simms, M.J., eds., *Fossil Crinoids*: Cambridge, Cambridge University Press, p. 203–215.
- Kammer, T.W., 1984, Crinoids from the New Providence Shale Member of the Borden Formation (Mississippian) in Kentucky and Indiana: *Journal of Paleontology*, v. 58, p. 115–130.
- Kammer, T.W., and Ausich, W.I., 1987, Aerosol suspension feeding and current velocities: distributional controls for late Osagean crinoids: *Paleobiology*, v. 13, p. 379–395.
- Kammer, T.W., and Ausich, W.I., 2006, The “Age of Crinoids”: a Mississippian biodiversity spike coincident with widespread carbonate ramps: *PALAIOS*, v. 21, p. 238–248.
- Kammer, W. W., and Ausich, W. I., 2007, New cladid and flexible crinoids from the Mississippian (Tournaisian, Ivorian) of England and Wales: *Palaeontology*, v. 50, p. 1039–1050.
- Kammer, T.W., and Matchen, D.L., 2008, Evidence for eustasy at the Kinderhookian–Osagean (Mississippian) boundary in the United States: response to late Tournaisian glaciation? in Fielding, C.R., Frank, T.D., and Isbell, J.L., eds., *Resolving the Late Paleozoic Ice Age in Time and Space*: Geological Society of America Special Paper 441, p. 261–274.
- Kammer, T.W., and Roeser, E.W., 2012, Cladid crinoids from the late Kinderhookian Meadville Shale Cuyahoga Formation of Ohio: *Journal of Paleontology*, v. 86, p. 470–487.
- Kammer, T.W., Ausich, W.I., and Goldstein, A., 2007, *Gilmocrinus kentuckyensis* n. sp. from the late Osagean (Mississippian) Muldraugh Member of the Borden Formation in Kentucky: a European immigrant originally derived from North America?: *Journal of Paleontology*, v. 81, p. 209–212.
- Kammer, T.W., Sumrall, C.D., Zamora, S., Ausich, W.I., and Deline, B., 2013, Oral region homologies in Paleozoic crinoids and other plesiomorphic pentaradiate echinoderms. *PLoS ONE*, v. 8, n. 11, 16 p., <https://dx.plos.org/10.1371/journal.pone.0077989>
- Kepferle, R.C., 1977, Stratigraphy, Petrology, and Depositional Environment of the Kenwood Siltstone Member, Borden Formation (Mississippian), Kentucky and Indiana: U.S. Geological Survey Professional Paper 1007, 49 p.
- Krivicich, E.B., Ausich, W.I., and Meyer, D.L., 2014, Crinoid assemblages from the Fort Payne Formation (late Osagean, early Viséan, Mississippian)

- from Kentucky, Tennessee, and Alabama: *Journal of Paleontology*, v. 88, p. 1154–1162.
- Lane, N.G., 1963, The Berkeley crinoid collection from Crawfordsville, Indiana: *Journal of Paleontology*, v. 3, p. 1001–1008.
- Lane, N.G., 1973, Paleontology and paleoecology of the Crawfordsville fossil site (upper Osagian, Indiana): *California University Publications in the Geological Sciences*, v. 99, 141 p.
- Lane, N.G., and Burke, J.J., 1976, Arm movement and feeding mode of inadunate crinoids with biserial muscular arm articulations: *Paleobiology*, v. 2, p. 202–208.
- Lane, N.G., and DuBar, J.R., 1983, Progradation of the Borden delta: new evidence from crinoids: *Journal of Paleontology*, v. 57, p. 112–123.
- Lane, N.G., Waters, J.A., and Maples, C.G., 1997, Echinoderm faunas of the Hongguleleng Formation, Late Devonian (Famennian), Xinjiang-Uygur Autonomous Region, People's Republic of China: *Journal of Paleontology*, v. 71, no. S47, p. 1–43.
- Liddell, W.D., 1975, Recent crinoid biostratigraphy: *Geological Society of America, Abstracts with Programs*, v. 7, p. 1169.
- Lister, M., 1673, A description of certain stones figured like plants, and by some observing men esteemed to be plants petrified: *Philosophical Transactions of the Royal Society of London*, v. 100, p. 6181–6191.
- Maples, C.G., and Archer, A.W., 1989, Paleoecological and sedimentological significance of bioturbated crinoid calyces: *PALAIOS*, v. 4, p. 379–383.
- M'Coy, F., 1849, On some new Paleozoic Echinodermata: *Annals and Magazine of Natural History*, ser. 2, v. 3, p. 244–254.
- Meyer, D.L., 1971, Post-mortem disarticulation of Recent crinoids and ophiuroids under natural conditions: *Geological Society of America Abstracts with Programs*, v. 3, p. 645.
- Meyer, D.L., and Milsom, C.V., 2001, Microbial sealing in the biostratigraphy of *Urintacrinus* Lagerstätten in the Upper Cretaceous of Kansas and Colorado, USA: *PALAIOS*, v. 16, p. 535–546.
- Meyer, D.L., and Oji, T., 1992, Experimental taphonomy of a Recent stalked crinoid: implications for the crinoid fossil record: *Geological Society of America Abstracts with Programs*, v. 24, p. 55.
- Meyer, D.L., Tobin, R.C., Pryor, W.A., Harrison, W.B., and Osgood, R.G., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnati Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, in Roberts, T.G., ed., *Geological Society of America, Cincinnati 1981, Field Trip Guidebooks*, v. 1: American Geological Institute, p. 31–71.
- Miller, J.S., 1821, A Natural History of the Crinoidea or Lily-shaped Animals; with Observations on the Genera *Asteria*, *Euryale*, *Comatula* and *Marsupites*: Bristol, Bryan & Co., 150 p.
- Miller, S.A., and Gurley, W.F.E., 1896a, Description of new and remarkable fossils from the Palaeozoic rocks of the Mississippi Valley: *Illinois State Museum Bulletin* 8, p. 1–65.
- Miller, S.A., and Gurley, W.F.E., 1896b, New species of Echinodermata and a new crustacean from the Palaeozoic rocks: *Illinois State Museum Bulletin* 10, p. 1–91.
- Milsom, C.V., Simms, M.J., and Gale, A.S., 1994, Phylogeny and palaeobiology of *Marsupites* and *Urintacrinus*: *Palaeontology*, v. 37, p. 595–607.
- Moore, R.C., and Laudon, L.R., 1943, Evolution and classification of Paleozoic crinoids: *Geological Society of America Special Paper* 46, 154 p.
- Moore, R.C., and Teichert, C., eds., 1978, *Treatise on Invertebrate Paleontology, Part T, Echinodermata 2*: Boulder, Colorado, and Lawrence, Kansas, Geological Society of America and University of Kansas Press, 1027 p.
- Neugebauer, J., 1978, Micritization of crinoids by diagenetic dissolution: *Sedimentology*, v. 25, p. 267–283.
- Nicoll, R.S., and Rexroad, C.B., 1975, Stratigraphy and conodont paleontology of the Sanders Group (Mississippian) in Indiana and adjacent Kentucky: *Indiana Geological Survey Bulletin* 51, 36 p.
- Peterson, W.L., and Kepferle, R.C., 1970, Deltaic deposits of the Borden Formation in central Kentucky: *U.S. Geological Survey Professional Paper* 700-D, p. D49–D54.
- Phillips, J., 1836, *Illustrations of the Geology of Yorkshire, or a Description of the Strata and Organic Remains. Pt. 2, The Mountain Limestone Districts* (second edition): London, John Murray, p. 203–308.
- Rhenberg, E.C., and Kammer, T.W., 2013, Camerate crinoids from the Nunn Member (Tournaian, Osagean) of the Lower Mississippian Lake Valley Formation, New Mexico: *Journal of Paleontology*, v. 87, p. 312–340.
- Richardson, J.G., and Ausich, W.I., 2004, Miospore biostratigraphy of the Borden delta (Lower Mississippian; Osagean) in Kentucky and Indiana, USA: *Palynology*, v. 28, p. 159–174.
- Rofe, J., 1865, Notes on some Echinodermata from the Mountain Limestone, etc.: *Geological Magazine*, v. 2, p. 245–252.
- Sable, E.G., and Dever, G.R., Jr., 1990, Mississippian rocks in Kentucky: *U.S. Geological Survey Professional Paper* 1503, 125 p.
- Sable, E.G., Kepferle, R.C., and Peterson, W.L., 1966, Harrodsburg Limestone in Kentucky. *Contributions to Stratigraphy: Geological Survey Bulletin* 1224-I, 12 p.
- Seilacher, A., Drozdowski, G., and Haude, R., 1968, Form and function of the stem in a pseudoplanktonic crinoid (*Seirocrinus*): *Palaeontology*, v. 11, p. 275–282.
- Seilacher, A., Reif, W.E., and Westphal, F., 1985, Sedimentological, ecological and temporal patterns of fossil Lagerstätten: *Philosophical Transactions of the Royal Society of London B*, v. 311, p. 5–23.
- Simms, M.J., 1986, Contrasting lifestyles of Lower Jurassic crinoids: a comparison of benthic and pseudopelagic Isocrinida: *Palaeontology*, v. 29, p. 475–493.
- Springer, F., 1901, *Urintacrinus*: its structure and relations: *Memoirs of the Museum of Comparative Zoology, Harvard University*, v. 25, p. 1–89.
- Struve, W., 1957, Ein Massengrab kreidezeitlicher Seelilien: die *Urintacrinus*-Platte des Senckenberg-Museums: *Natur und Volk*, v. 87, p. 361–373.
- Swofford, D.L., 2015, PAUP*: Phylogenetic analysis using parsimony (and other methods), version 4.0a142. <http://phylosolutions.com/paup-test>.
- Taylor, W., and Brett, C.E., 1996, Taphonomy and paleoecology of echinoderm Laterstätten from the Silurian (Wenlockian) Rochester Shale: *PALAIOS*, v. 11, p. 118–140.
- Ubaghs, G., 1978a, Camerates, in Moore, R.C., and Teichert, C., eds., *Treatise on Invertebrate Paleontology, Part T, Echinodermata 2*: Geological Society of America and University of Kansas Press, p. T409–T519.
- Ubaghs, G., 1978b, Skeletal morphology of fossil crinoids, in Moore, R.C., and Teichert, C., eds., *Treatise on Invertebrate Paleontology, Part T, Echinodermata 2*: Boulder, Colorado, and Lawrence, Kansas, Geological Society of America and University of Kansas Press, p. T58–T216.
- Van Sant, J.F., and Lane, N.G., 1964, *Crawfordsville (Indiana) crinoid studies*: University of Kansas Paleontological Contributions, Article 35, Echinodermata 7, 136 p.
- Wachsmuth, C., and Springer, F., 1880–1886, *Revision of the Palaeocrinoidea, Part 2, Family Sphaeroidocrinidae, with the sub-families Platycrinidae, Rhodocrinidae, and Actinocrinidae; Part 3, Section 1, Discussion of the classification and relations of the brachiote crinoids, and conclusions of the generic descriptions*: *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1881, p. 177–411 (Part 2); 1885, p. 225–364 (Part 3).
- Walker, L.J., Wilkinson, B.H., and Ivany, L.C., 2002, Continental drift and Phanerozoic carbonate accumulation in shallow-shelf and deep-marine settings: *Journal of Geology*, v. 110, p. 75–87.
- Waters, J.A., Maples, C.G., Lane, N.G., Marcus, S., Liao, Z-T., Liu, L., Hou, H-F., and Wang, J-X., 2003, A quadrupling of Famennian pelmatozoan diversity: new Late Devonian blastoids and crinoids from northwest China: *Journal of Paleontology*, v. 77, p. 922–948.
- Waters, J.A., Marcus, S.A., Maples, C.G., Lane, N.G., Hou, H., Liao, Z-T., Wang, J-X., and Liu, L., 2008, Overview of Paleozoic stemmed echinoderms from China, in Ausich, W.I., and Webster, G.D., eds., *Echinoderm Paleobiology*: Bloomington, Indiana University Press, p. 346–367.
- Webster, G.D., 1974, Crinoid pluricolumnal noditaxis patterns: *Journal of Paleontology*, v. 48, p. 1283–1288.
- Webster, G.D., and Lane, N.G., 1987, Crinoids from the Anchor Limestone (Lower Mississippian) of the Monte Cristo Group southern Nevada: *University of Kansas Paleontological Contributions Paper* 119, 55 p.
- Webster, G.D., and Waters, J.A., 2009, Late Devonian echinoderms from the Hongguleleng Formation of northwestern China, in Königshof, P., ed., *Devonian Change: Case Studies in Palaeogeography and Palaeoecology*: The Geological Society of London Special Publications, v. 314, p. 263–287.
- Webster, G.D., Maples, C.G., Sevastopulo, G.D., Frest, T., and Waters, J.A., 2004, Carboniferous (Viséan–Moscowian) echinoderms from the Béchar Basin area of western Algeria: *Bulletins of American Paleontology*, no. 368, 98 p.
- Webster, G.D., Waters, J.A., and Chen X., 2009, Revision of the Chen and Yao Devonian to Permian crinoids from Western Yunnan: *Palaeobiodiversity and Palaeoenvironments*, v. 89, p. 119–160.
- Wiley, E.O., and Lieberman, B.S., 2011, *Phylogenetics: Theory and Practice of Phylogenetic Systematics*: Wiley-Blackwell, 432 p.
- Wright, D.F., 2017, Bayesian estimation of fossil phylogenies and the evolution of early to middle Paleozoic crinoids (Echinodermata): *Journal of Paleontology*, v. 91, p. 799–814.
- Wright, D.F., Ausich, W.I., Cole, S.R., Peter, M.E., and Rhenberg, E.C., 2017, Phylogenetic taxonomy and classification of the Crinoidea (Echinodermata): *Journal of Paleontology*, v. 91, p. 829–846.
- Wright, J., 1943, *Pimlicocrinus* gen. nov. and two new species of *Amphoracrinus* from the Carboniferous Limestone: *Geological Magazine*, v. 80, p. 81–94.
- Wright, J., 1950–1960, *The British Carboniferous Crinoidea*. *Palaeontographical Society Monograph*, v. 1, no. 1, p. 1–24, pl. 1–7, 1950; v. 1, no. 2, p. 25–46, pl. 8–12, 1951a; v. 1, no. 3, p. 47–102, pl. 13–31, 1951b; v. 1, no. 4, p. 103–148, pl. 32–40, 1952a; v. 1, no. 5, p. 149–190, pl. 41–47, 1954a; v. 2, no. 1, p. 191–254, pl. 48–63, 1955a; v. 2, no. 2, p. 255–272, pl. 64–67, 1955b; v. 2, no. 3, p. 273–306, pl. 68–75, 1956b; v. 2, no. 4, p. 307–328, pl. 76–81, 1958; v. 2, no. 5, p. 329–347, pl. A, B, 1960.

Accepted: 10 May 2021