

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. *Tables* Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLES

Affective problems and decline in cognitive state in older adults: a systematic review and meta-analysis

John A, Patel U, Rusted J, Richards M & Gaysina D 353

Effectiveness of psychotherapy for treatment-resistant depression: a meta-analysis and meta-regression

van Bronswijk S, Moopen N, Beijers L, Ruhe HG & Peeters F 366

INVITED REVIEW

Can we predict the direction of sudden shifts in symptoms? Transdiagnostic implications from a complex systems perspective on psychopathology

Wichers M, Schreuder MJ, Goekoop R & Groen RN 380

ORIGINAL ARTICLES

Links between psychotic and neurotic symptoms in the general population: an analysis of longitudinal British National Survey data using Directed Acyclic Graphs

Kuipers J, Moffa G, Kuipers E, Freeman D, Bebbington P 388

Diagnostic accuracy of dopaminergic imaging in prodromal dementia with Lewy bodies

Thomas AJ, Donaghy P, Roberts G, Colloby SJ, Barnett NA, Petrides G, Lloyd J, Olsen K, Taylor J-P, McKeith I & O'Brien JT 396

Short sleep duration and poor sleep quality predict next-day suicidal ideation: an ecological momentary assessment study

Littlewood DL, Kyle SD, Carter L-A, Peters S, Pratt D & Gooding P 403

Structural covariance and cortical reorganisation in schizophrenia: a MRI-based morphometric study

Palaniyappan L, Hodgson O, Balain V, Iwabuchi S, Gowland P & Liddle P 412

Common genetic contributions to high-risk trauma exposure and self-injurious thoughts and behaviors

Richmond-Rakerd LS, Trull TJ, Gizer IR, McLaughlin K, Scheiderer EM, Nelson EC, Agrawal A, Lynskey MT, Madden PAF, Heath AC, Statham DJ & Martin NG 421

Exercise effects on cognitive functioning in young adults with first-episode psychosis: FitForLife

Hallgren M, Skott M, Ekblom Ö, Firth J, Schembri A & Forsell Y 431

Measuring disorganized speech in schizophrenia: automated analysis explains variance in cognitive deficits beyond clinician-rated scales

Minor KS, Willits JA, Marggraf MP, Jones MN & Lysaker PH 440

Tree analysis modeling of the associations between PHQ-9 depressive symptoms and doctor diagnosis of depression in primary care

Chin W-Y, Wan EYF, Dowrick C, Arroll B & Lam CLK 449

Being left-behind, mental disorder, and elderly suicide in rural China: a case-control psychological autopsy study

Zhou L, Wang G, Jia C & Ma Z 458

Long-term outcomes of acute treatment with cognitive therapy v. interpersonal psychotherapy for adult depression: follow-up of a randomized controlled trial

Lemmens LHJM, van Bronswijk SC, Peeters F, Arntz A, Hollon SD & Huibers MJH 465

Neighborhood alcohol outlet density and genetic influences on alcohol use: evidence for gene-environment interaction

Slutske WS, Deutsch AR & Piasecki TM 474

Clinical implications of the proposed ICD-11 PTSD diagnostic criteria

Barbano AC, van der Mei WF, Bryant RA, Delahanty DL, deRoos-Cassini TA, Matsuoka YJ, Olf M, Qi W, Ratanatharathorn A, Schnyder U, Seedat S, Kessler RC, Koenen KC & Shalev AY 483

Efficacy of digital CBT for insomnia to reduce depression across demographic groups: a randomized trial

Cheng P, Luiik AI, Fellman-Couture C, Peterson E, Joseph CLM, Tallent G, Tran KM, Ahmedani BK, Roehrs T, Roth T & Drake CL 491

Does online insomnia treatment reduce depressive symptoms? A randomized controlled trial in individuals with both insomnia and depressive symptoms

van der Zweerde T, van Straten A, Effting M, Kyle SD & Lancee J 501

Disrupted rich club organization and structural brain connectome in unmedicated bipolar disorder

Wang Y, Deng F, Jia Y, Wang J, Zhong S, Huang H, Chen L, Chen G, Hu H, Huang L & Huang R 510

Prevalence and determinants of cognitive impairment in the euthymic phase of bipolar disorders: results from the FACE-BD cohort

Roux P, Etain B, Cannavo A-S, Aubin V, Auizerate B, Azorin J-M, Bellivier F, Belzeaux R, Bougerol T, Cussac I, Courtet P, Kahn J-P, Leboyer M, M'Bailara K, Payet MP, Olié E, the FondaMental Advanced Centers of Expertise in Bipolar Disorders (FACE-BD) Collaborators, Henry C & Passerieux C 519

Corrigendum

528