

Assessment of genetic divergence in diploid cotton (*Gossypium arboreum* L.) germplasm using fibre quality traits

A. Manivannan^{1*}  and V. N. Waghmare²

¹ICAR-Central Institute for Cotton Research, Regional Station, Coimbatore 641003, India and ²ICAR-Central Institute for Cotton Research, Nagpur 441108, India

Received 22 May 2020; Accepted 3 October 2020 – First published online 4 November 2020

Abstract

Cotton is one of the most important crops among natural fibres. Fibre quality determines the spinning ability, which is negatively correlated with yield and yield-contributing traits. Limited efforts have been made to improve fibre quality and yield in diploid cotton. Therefore, screening a large panel of germplasm lines can help identify genotypes with better fibre quality and yield. We evaluated 712 desi cotton genotypes for fibre quality traits. The genotypes showed a significant difference for all the traits, suggesting considerable variability for fibre quality improvement. Fibre length and strength showed high phenotypic and genotypic coefficients of variation. Heritability was high for fibre strength, length, and elongation. Fibre length and strength were positively correlated; however, micronaire was negatively correlated with these two traits. Superior accessions were identified for fibre length (11), strength (20), uniformity (7), and elongation (25) among genotypes. Most of the desi lines (71%) had medium micronaire values. Twenty accessions identified were ideal for spinning, showing the fibre strength-to-length ratio of one. Cluster analysis based on Euclidean distance grouped all 712 accessions into four major clusters. Principal component analysis biplot revealed that accessions AC3418, 360-SP1, AC3522B, Kanpur A, Gao16CB-9, and AC3370 were genetically diverse. The superior accessions for fibre quality identified in this study are potential lines for the diploid cotton improvement programme.

Keywords: characterization, cotton, diploid, fibre quality, germplasm, *Gossypium arboreum*

Introduction

The cotton genus (*Gossypium* L.) has 50 species; among them, *Gossypium arboreum* L. is a diploid originating from the Indian Subcontinent (Li *et al.*, 2014). Diploid cotton grows in marginal environments under rain-fed situation and can sustain climate changes (Wendel *et al.*, 2012). It exhibits resistance to various pests and diseases (Miyazaki *et al.*, 2014). Despite all the virtues of diploid cotton, its fibre is short, poor in strength, and coarse and has a non-spinning ability, which are great impediments for the widespread use of diploid cotton in textiles in comparison with tetraploid cotton (Romeu-Dalmau *et al.*, 2015).

The quality of cotton is determined by its fibre length, strength, micronaire, uniformity, and elongation. The length determines the spinning efficiency as lengthier fibre produces long yarns (Koebernick *et al.*, 2019). Strength is necessary to maintain cotton's natural qualities after fabric processing. Micronaire measures the maturity and fineness of cotton. A high micronaire is linked with fibre coarseness (Han *et al.*, 1998). Uniformity is associated with spinning property, which determines the efficiency of fabric production (Hequet *et al.*, 2006). Elongation measures the elasticity of fibres, as breakage causes inefficiency in yarn manufacturing and affects the end product quality (Ruan *et al.*, 2004). All these traits are highly influenced by the environments, and desi cotton grows in various marginal environments, making its fibre vulnerable. After the Bt cotton regime, desi cotton gained importance for its

*Corresponding author. E-mail: manivannan461@gmail.com

resilience to climatic change. Hence, the improvement of fibre properties of desi cotton is important for fulfilling the growing requirements of the textile industry (Chandra and Sreenivasan, 2011). Breeding for high-yield cotton with good fibre quality is a challenging task because of the strong negative correlation of quality with yield (Yu *et al.*, 2013). Such negative linkages between fibre yield and quality can be reversed by adapting diverse breeding strategies (Campbell and Jones, 2005). Thus, the simultaneous improvement of yield and quality is a major challenge for the cotton breeding programme (Wang *et al.*, 2016). Large-scale screening of the germplasm is expected to identify genotypes with superior fibre properties, which in turn could be used for improving yield (Chen *et al.*, 2018). In this study, we analysed the genetic diversity and population structure of 712 *G. arboreum* accessions collected from different parts of India as a germplasm collection using six fibre quality traits. Our objectives were to estimate the levels of genetic diversity and characterize the *G. arboreum* germplasm for identifying the superior fibre quality genotypes for further breeding.

Materials and methods

A set of 712 *G. arboreum* accessions belonging to three races, namely, *indicum*, *cernuum*, and *bengalense*, were grown in augmented design at ICAR-Central Institute for Cotton Research (CICR), Regional Station, Coimbatore, Tamil Nadu, India (11.0168°N, 76.9558°E) during the 2016 *kharif* crop season (online Supplementary Table S1). The germplasm accessions were procured from the Cotton Gene Bank of ICAR-CICR, Nagpur, Maharashtra, India. They were raised in a 3 m row with a spacing of 60 cm × 30 cm. All agronomical practices were followed as per ICAR recommendations of packages of practice for cotton. Fully burst bolls were picked from 10 plants per entry and sundried well for further processing. Seed cotton of all individual plants in each entry was ginned with a cloy gin in the laboratory. Lint was conditioned by placing at 65% humidity and 18–20°C in an air-conditioned room using a humidifier before fibre testing. Lint samples were used to measure the following fibre properties: 2.5% span length (mm), length uniformity ratio (UR), fibre strength (g/tex), fibre elongation, and micronaire by using a High Volume Instrumentation (HVI-900-SA; Uster). Fibre analysis was performed under ICC mode at ICAR-Central Institute for Research on Cotton Technology, Regional Unit, Coimbatore.

Analysis of variance (ANOVA) was performed to ascertain the genetic variation present among the genotypes. The genetic parameters were studied by calculating the genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) (Burton and Davane, 1952),

heritability (Hanson *et al.*, 1956), and genetic advance as mean (GAM) (Johnson *et al.*, 1955) for all the characteristics. Pearson's correlation coefficient arrived for all fibre traits and a correlation matrix was formed for comparing various traits. Principal component analysis (PCA) was performed to identify the key variant contributing traits. The factors of these traits were used to determine the contribution of each factor towards variation. Standardized values were used to perform PCA. A scree plot was drawn using eigenvalues, which helped in visually accessing the factors representing most of the variability. Dissimilarity matrices were derived based on Euclidean distance. Highly similar and less dissimilar entries were identified using this dissimilarity matrix. This matrix was employed for constructing hierarchical cluster analysis using pooled genotyping data for genotype grouping (Ward, 1963). ANOVA, GCV, PCV, and correlation were computed using the STATGRAPHICS, Centurion XVI (StatPoint Technologies Inc., Warrenton, VA, USA), while PCA was performed using Past3 (Hammer *et al.*, 2001) and dissimilarity matrix was derived using DARwin 5 (Perrier and Jacquemod-Collet, 2006).

Results

Analysis of components of variance

Analysis of the quantitative traits showed that genotypes had a wide range of variations (Table 1). Fibre staple length (SL) varied from 19 to 29.3 mm (mean: 24.1 mm). UR ranged from 45 to 60 (mean: 51.7). Micronaire ranged from 2.7 to 6.8 (mean: 5.0). Fibre strength varied from 13.6 to 27.6 g/tex (mean: 19.8 g/tex). Elongation ranged from 5 to 7.7 (mean: 5.8). Phenotypic variance (PV) and genotypic variance (GV) showed wide variations. PV ranged from 0.1 for elongation to 3.1 for fibre strength, while GV ranged from 0.09 for elongation to 2.3 for fibre strength. Similarly, the environmental variance (EV) was negligible (nearly zero) for elongation to 2.2 for uniformity. Fibre strength and length had high PCV and GCV values. The remaining traits exhibited medium to low values. In general, estimates of both PCV and GCV showed a wide range of variations (3 and 1% in UR to 10.1 and 8.4% in fibre length). The PCV values were slightly higher than their corresponding GCV values for all the traits (Table 1).

Estimates of heritability in a broad sense (h^2) and genetic advance (GA)

The estimate of heritability in a broad sense (b^2) showed differences ranging from 12% for uniformity to 75.7% for strength. Fibre SL (68.8%), strength (75.7%), and elongation (56.8%) showed a higher level of heritability. Similarly,

Table 1. Mean, range, and variance components of *G. arboreum* germplasm for fibre traits

| Characters | Mean | Range | PV | EV | GV | PCV | GCV | h^2 | GA | GA % mean |
|------------|------|-----------|-----|------|------|------|-----|-------|-----|-----------|
| SL | 24.1 | 19.0–29.3 | 0.3 | 0.1 | 0.2 | 10.1 | 8.4 | 68.8 | 1.4 | 28.3 |
| UR | 51.7 | 45–60 | 2.4 | 2.2 | 0.3 | 3.0 | 1.0 | 12.0 | 0.6 | 1.1 |
| MC | 5.0 | 2.7–6.8 | 1.7 | 1.2 | 0.5 | 5.3 | 2.8 | 28.0 | 0.7 | 3.1 |
| ST | 19.8 | 13.6–27.6 | 3.1 | 0.8 | 2.3 | 8.9 | 7.7 | 75.7 | 6.7 | 34.0 |
| EL | 5.8 | 5.0–7.7 | 0.1 | 0.01 | 0.09 | 4.1 | 3.1 | 56.8 | 0.8 | 14.1 |

SL, staple length; UR, uniformity ratio; MC, micronaire; ST, strength; EL, elongation percentage.

estimates of GA varied from 0.6 (uniformity) to 6.7 (strength), whereas estimates of GA as a per cent of traits mean varied from 1.1 (uniformity) to 34 (strength).

Principal component analysis

Five principal components were identified and contributed to variations. The first principal component (PC 1) accounted for 56.12%, second (PC 2) for 29.41%, third (PC 3) for 10.68%, fourth (PC 4) for 3.16%, and fifth (PC 5) for 0.63% of total variation (online Supplementary Table S2). Eigenvectors were calculated to quantify the principal components for fibre traits (online Supplementary Table S3). The first two principal component axes jointly accounted for 85.53% of the total variation. PC 1 was associated with strength (0.700), SL (0.511), and elongation (0.014). PC 2 was associated with UR (0.774), strength (0.622), and elongation (0.016). PC 3 correlated with SL (0.849), UR (0.394), micronaire (0.118), and elongation (0.013). PC 4 was associated with micronaire (0.977), strength (0.112), and elongation (0.162), while PC 5 was only associated with elongation (0.986). According to all the principal components, maximum variation was recorded for strength (54.37%), moderate variability was observed for SL (34.28%), and low variability for elongation (2.53%), and micronaire had the least variability (Table 2).

Biplot analysis

A PC biplot was drawn using PC 1 and PC 2 factor scores, and a clear pattern of grouping between the genotypes was observed in the factor plane (online Supplementary Fig. S1). The gap within traits concerning PC 1 and PC 2 depicted the contribution of these traits in creating genotypic variation. Each trait was depicted as a vector on the biplot and the length of the vector was proportionate to the ability to distinguish genotypes. However, the overall biplot diagram demonstrated that fibre strength, SL, and uniformity contributed significantly towards germplasm diversity, while micronaire and elongation were different for all characteristics. Genotypes AC3418, 360-SP1, AC3522B,

Table 2. Variability by different fibre traits of *G. arboreum* accessions

| Characters | Variability contribution (%) |
|------------|------------------------------|
| SL | 34.28 |
| UR | 0.91 |
| Mic | 0.98 |
| ST | 54.37 |
| EL | 2.53 |

SL, staple length; UR, uniformity ratio; MC, micronaire; ST, strength; EL, elongation percentage.

Arboreum (Kanpur A), Gao16CB-9, and AC3370 occupied the convex of the hull. These genotypes were entirely distinct from each other, and they were occupying the extreme corner of the vector plane (Fig. 1).

Association analysis

Pearson's correlation coefficient was calculated among the fibre traits. Among the inter-correlation coefficients, four were significant. The highest significant positive correlation was observed between strength and length (0.498). A significant correlation was observed between elongation and micronaire (0.214). SL and uniformity were significantly negatively correlated with each other (−0.479); strength and micronaire were also significantly negative correlated (−0.304). All other inter-correlations were non-significant (online Supplementary Table S3).

Cluster analysis

The factors corresponding to four PCs were subjected to cluster analysis based on Euclidean distances and grouped by an unweighted paired group method using arithmetic average by using DARwin 5. The dendrogram depicted four distinct clusters of 306, 232, 46, and 128 accessions in each accession (Fig. 2).




Fig. 2. Dendrogram based on unweighted pair group method with arithmetic mean (UPGMA) of qualitative characteristics.

Table 3. Superior *G. arboreum* accessions for fibre traits

| Genotype | SL | Genotype | UR | Genotype | ST | Genotype | EL |
|----------------------------|------|---|------|-----------|------|--|------|
| 360-SP1 | 29.3 | Gao 16 CB-9 | 60 | AC 3522 B | 27.6 | GDH 149 (Sel.) | 7.7 |
| Shamali | 28.2 | Obtusifolium-B-Indica | 58 | AC 3451 | 26.6 | AC 3370 | 7.4 |
| Desi-103 | 28.1 | 360, GDH 149 (Sel.) | 57 | AC 3284 | 25.4 | Obtusifolium-B-Indica, Gao 16 CB-9 | 7.0 |
| 5974, 6582 | 28.0 | Arboreum (Kanpur A), Chinese broad lobe, PBN 48 | 56 | AC 3234 | 25.0 | N 31-24 | 6.8 |
| 30,859, AK606-SP1, AC 3695 | 27.9 | | | AH 71 | 24.7 | Desi 87, AC 622-SP2 | 6.5 |
| Sarguja-NL-WF, 30,843 | 27.8 | | | SC 97 | 24.3 | 79/BH 113 | 6.4 |
| 30,839 | 27.4 | | | 6920 | 24.0 | AH 71, 6187, 30,809, AC 3547 | 6.3 |
| | | | | | | 6920, 564, 6164, AK 592, 30,841-SP1, 1173 WR, 30,861, 30,827, 30,828, 2927, AKA 603, 30,844-SP1, H 502 | 6.2 |
| Population mean | 24.1 | | 51.7 | | 19.8 | | 5.8 |
| CD 5% | 3.2 | | 4.3 | | 2.5 | | 0.45 |

SL, staple length; UR, uniformity ratio; ST, strength; EL, elongation percentage.

Table 4. Strength-to-length ratio among the selected *G. arboreum* accessions

| Genotype | ST | SL | ST/SL |
|---------------------|------|------|-------|
| AC 3522 B | 27.6 | 22.5 | 1.23 |
| AC 3451 | 26.6 | 23.2 | 1.15 |
| Arboreum (Kanpur A) | 23.2 | 22.8 | 1.02 |
| AC 3284 | 25.4 | 25.4 | 1.00 |
| AKA 14 | 23.0 | 23.2 | 0.99 |
| AH 71 | 24.7 | 25.0 | 0.99 |
| H 575 | 22.6 | 23.0 | 0.98 |
| AC 3234 | 25.0 | 25.5 | 0.98 |
| AC 3289 | 22.6 | 23.1 | 0.98 |
| Desi-1 | 23.4 | 24.0 | 0.98 |
| PS-135 | 23.0 | 23.6 | 0.97 |
| SC 97 | 24.3 | 25.0 | 0.97 |
| 79/BH-97 | 23.7 | 24.5 | 0.97 |
| H 52-473 | 20.7 | 21.4 | 0.97 |
| CC-1-1-3 | 23.2 | 24.0 | 0.97 |
| AC 3368 | 21.7 | 22.5 | 0.96 |
| 7763 | 22.8 | 23.7 | 0.96 |
| 8410-2 | 23.9 | 24.9 | 0.96 |
| PBN 6977 × AKH4 | 21.9 | 22.9 | 0.96 |
| 30,819-SP1 | 23.2 | 24.4 | 0.95 |

ST, strength; SL, staple length; ST/SL, strength-to-length ratio.

the population. When high heritability is combined with high GA, researchers should follow selection strategies in the next generation of crops for improving these traits (Rahman, 2016). High to medium values of heritability estimates were found associated with moderate GAM in fibre elongation. This trait could be used in the early generation, but it would be more effective if used in the late generation (Kumar *et al.*, 2017).

Effective grouping of genotypes could be done with the multivariate statistics of PCA (Pearson's 1901). PCA is the widely used multivariate statistical analysis based on the principle of data reduction. Among the several observed variables in a dataset, PCA identifies the significant variables contributing to the majority of variability. PCA simplifies multidimensionality into lower dimensions. The linear transformation technique is used to transform the variability into new coordinate systems, in which higher variability occupies the first principal coordinate and it moves in descending orders. Rathinavel (2018) applied PCA in 108 extant varieties of cotton using fibre length, strength, fineness, and uniformity and grouped the varieties into five major clusters. Since the biological explanation of principal components is subtle, the best way is to find the degree of influence of each variable weight on each of the components. Primary variability was governed by fibre

strength, length, and elongation. Hence, PCA is important in identifying traits contributing most to the variation among genotypes (online Supplementary Fig. S1). Biplot diagram depicted that fibre strength, SL, and uniformity contributed significantly towards germplasm diversity; however, micronaire and elongation contributed less towards diversity. Highly diverse accession for all the traits plotted in the distance from each other. These genotypes, namely, AC3418, 360-SP1, AC3522B, Arboreum (Kanpur A), Gao16CB-9, and AC3370 were plotted distal among all the accessions (Fig. 2). AC 3522B was highly superior to other genotypes, as it had a fibre length of 22.5 mm, a strength of 27.6 g/tex, and a strength-to-length ratio of 1.2. Similarly, Arboreum (Kanpur A) had a fibre length of 22.8 mm and a strength of 23.2 g/tex. These genotypes could be employed for improving fibre length by using them directly for selection or as potential donor parents in a crossing programme. The association between different traits is the most important criterion to demonstrate cotton improvement strategies (Ali *et al.*, 2009). In our study, cotton fibre length and strength positively correlated with each other; however, micronaire negatively correlated with both traits. These results were similar to those of Karademir *et al.* (2011) and Clement *et al.* (2012), as the authors observed such an inverse relationship among these traits. In textile industries, fibre length is an important factor for yarn strength and processing performance (Chaudhary, 2000). The short fibre length of *G. arboreum* affects the efficiency of high-speed spinning machines. Genotypes 360-SP1, Shamali, Desi-103, 5974, 6582, 30859, AK606-SP1, AC 3695, Sarguja-NL-WF, 30843, and 30839 showed longer fibres; these genotypes would be efficient donors for fibre improvement in desi cotton. Fibre uniformity is important for determining the utility of lint for various purposes, and it directly influences yarn strength, elongation, and twist (Parsi *et al.*, 2016). Higher uniformity affects the production of uniform yarn in size and strength, resulting in lesser fibre wastage. Lower uniformity results in the production of short fibre content (Ibrahim, 2019). CB-9, Obtusifolium-B-Indica, 360, GDH 149 (Sel.), Arboreum (Kanpur A), Chinese broad lobe, and PBN 48 were superior for the UR. Fibre strength is critical to the processing of fibre into yarn, and it consequently affects the end product quality. The breaking strength plays a major role in determining yarn strength (Chandra and Sreenivasan, 2011). Among 712 accessions, 54 exhibited higher strength over the mean population mean. Four accessions, AC 3522 B(27.6), AC 3451(26.6), AC 3284(25.4), and AC 3234(25.0), had a strength of >25 g/tex, which is on par with the strength of tetraploid cotton (*Gossypium hirsutum*). These high strength lines can be crossed with high length materials, and improved lines for both strength and length can be obtained, and such lines by nature would be spinnable.

Fibre fineness, an important genetic character after fibre length, determines the texture of cotton fibre as fine or coarse. Micronaire measures both the fineness and maturity of cotton. Micronaire values <3.5 indicate immature fibres, which are weak and prone to break while spinning, have poor dye absorption and create fibre entanglements (neps) that affect fabric texture and appearance (Hanan *et al.*, 2017). Micronaire values >5.0 indicate coarse fibres, which cannot be spun as yarn. Micronaire is not always the ideal measure of fineness as it is a product of fibre maturity (which is the result of secondary cell wall thickening) and linear density. Generally, low micronaire values indicate immature fibres. High micronaire values are associated with coarseness. Hence, finer fibres allow more fibres per cross-section of yarn to improve yarn strength, resulting in finer yarn (Han *et al.*, 1998). Among the accessions, 76 had a micronaire value <4.5, 511 had 4.5–5.5, and 88 had >5.5. Fibre elongation (breaking elongation) directly affects yarn toughness. Twenty-five accessions had elongation percentage of >6%, among them GDH 149 (Sel.), AC 3370, Obtusifolium-B-Indica, and Gao16CB-9 had elongation percentage of >7%. The relation between strength and length is expressed as the strength-to-length ratio; a ratio of one is considered as an ideal parameter for spinning. According to the strength-to-length ratio, 20 genotypes, AC 3522 B, AC 3451, Arboreum (Kanpur A), AC 3284, AKA 14, AH 71, H 575, AC 3234, AC 3289, Desi-1, PS-135, SC 97, 79/BH-97, H 52-473, CC-1-1-3, AC 3368, 7763, 8410-2, PBN 6977 × AKH4-SP1, and 30819-SP1, had a ratio of one and are suitable for spinning, since diploid cotton *per se* has poor spinnability. Spinnable *G. arboreum* cotton is fetching a huge demand in the market, especially for medical textile production (Kranti, 2015).

Conclusion

We found that the magnitude of variation for fibre quality traits exists among the *G. arboreum* accessions in the germplasm. Multivariate analysis is a useful statistical method to evaluate germplasm, which plays an important role in characterizing genotypes in terms of their discriminatory ability to separate each variable in the single-trait analysis. Association analysis revealed that cotton fibre length and strength are positively correlated with each other, but micronaire is negatively associated with these two traits. These traits have to be emphasized in breeding programmes for further enhancement. Accessions AC3418, 360-SP1, AC3522B, Arboreum (Kanpur A), Gao16CB-9, and AC3370 are highly diverse for these fibre traits. These genotypes could serve as parents for harnessing in hybridization to exploit heterosis. Accessions AC 3522 B, AC 3451, Arboreum (Kanpur A), AC 3284, AKA 14, AH 71, H 575, AC

3234, AC 3289, Desi-1, PS-135, SC 97, 79/BH-97, H 52-473, CC-1-1-3, AC 3368, 7763, 8410-2, PBN 6977 × AKH4-SP1, and 30819-SP1 have higher strength and length and are more amenable for spinning because of their fibre strength-to-length ratio. These accessions can be utilized in diploid cotton breeding to improve fibre quality traits.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262120000374>.

Acknowledgements

This research was supported by ICAR-Central Institute for Cotton Research, Regional Station, Coimbatore. The authors are thankful to ICAR-Central Institute for Cotton Research, Nagpur for providing germplasm for this work.

References

- Ali MA, Khan IA and Nawab NN (2009) Estimation of genetic divergence and linkage for fibre quality traits in upland cotton. *Journal of Agricultural Research* 47: 229–236.
- Burton GW and Davane EM (1952) Estimating heritability in fall fescue (*Festuca Arundinacea* L.) from replicated clonal material. *Agronomy Journal* 45: 478–481.
- Campbell BT and Jones MA (2005) Assessment of genotype × environment interactions for yield and fiber quality in cotton performance trials. *Euphytica* 144: 69–78.
- Chandra M and Sreenivasan S (2011) Studies on improved *Gossypium arboreum* cotton: part I – fibre quality parameters. *Indian Journal of Fibre and Textile Research* 36: 24–34.
- Chaudhary R (2000) Fibre length: achievements and new challenges. *The International Cotton Advisory Committee Recorder* 18: 3.
- Chen Y, Liu G, Ma H, Song Z, Zhang C, Zhang J, Zhang J, Wang F and Zhang J (2018) Identification of introgressed alleles conferring high fiber quality derived from *Gossypium barbadense* L. in secondary mapping populations of *G. hirsutum* L. *Frontiers in Plant Science* 9: 1023.
- Clement JD, Constable GA, Stiller WN and Liu SM (2012) Negative associations still exist between yield and fibre quality in cotton breeding programs in Australia and USA. *Field Crops Research* 128: 1–7.
- Gadissa F, Tesfaye K, Dagne K and Geleta M (2020) Morphological traits based genetic diversity assessment of Ethiopian potato (*Plectranthus edulis* (Vatke) Agnew) populations from Ethiopia. *Genetic Resources and Crop Evolution* 67: 809–829.
- Hammer O, Harper DAT and Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 1–9.
- Han YJ, Cho YJ, Lambert WE and Bragg CK (1998) Identification and measurement of convolutions in cotton fibre using image analysis. *Artificial Intelligence Review* 12: 201–211.
- Hanan G, Ghith A and Benameu T (2017) Open-end yarn properties prediction using hvi fibre properties and process parameters. *Alexandria Science Exchange Research Journal* 17: 6–11.

- Hanson CH, Robinson HF and Comstock RE (1956) Biometrical studies of yield in segregating populations of Korean lespe-diza. *Agronomy Journal* 48: 268–272.
- Hequet E, Wyatt B and Abidi N (2006) Creation of a set of reference material for cotton fiber maturity measurements. *Textile Research Journal* 76: 576–586.
- Hovav R, Udall JA, Chaudhary B, Hovav E, Flagel L, Hu G and Wendel JF (2008) The evolution of spinnable cotton fiber entailed prolonged development and a novel metabolism. *PLoS Genetics* 4: e25.
- Ibrahim AE (2019) Effect of fiber length and short fiber per cent in cotton on fiber and yarn quality. *Alexandria Science Exchange Journal* 39: 663–667.
- Johnson HW, Robinson HF and Comstock RE (1955) Estimates of genetic and environmental variability in soya bean. *Agronomy Journal* 47: 314–318.
- Karademir C, Karademir E and Gencer O (2011) Yield and fiber quality of f_1 and f_2 generations of cotton (*Gossypium hirsutum* L.) under drought stress conditions. *Bulgarian Journal of Agricultural Science* 17: 795–805.
- Koebnick JC, Liu S, Constable GA and Stiller WN (2019) Parental selection strategy for improving fibre strength and maintaining lint yield in cotton. *Industrial Crops and Products* 129: 585–593.
- Kranti KR (2015) Desi cotton return? *Cotton Statistics and Newsletter* 15: 1–4.
- Kumar SK, Nidagundi JM and Hosamani AC (2017) Association analysis for seed cotton yield and fiber quality improvement in intra specific crosses of cotton (*Gossypium hirsutum* L.). *International Journal of Pure and Applied Bioscience* 5: 282–284.
- Li F, Fan G, Wang K, Sun F, Yuan Y, Song G, Li Q, Ma Z, Lu C, Zou C, Chen W, Liang X, Shang H, Liu W, Shi C, Xiao G, Gou C, Ye W, Xu X, Zhang X, Wei H, Li Z, Hang G, Wang J, Liu K, Kohel RJ, Percy RG, Yu JZ, Zhu YX, Wang J, Yu S (2014) Genome sequence of the cultivated cotton *Gossypium arboreum*. *Nature Genetics* 46: 567–572. <https://doi.org/10.1038/ng.2987>
- McCouch S, Baute GJ, Bradeen J, Bramel P, Bretting PK, Buckler E, Burke JM, Charest D, Cloutier S, Cole G, Dempewolf H, Dingkuhn M, Feuillet C, Gepts P, Grattapaglia D, Guarino L, Jackson S, Knapp S, Langridge P, Lawton-Rauh A, Lijua Q, Lusty C, Michael T, Myles S, Naito K, Nelson RL, Pontarollo R, Richards CM, Rieseberg L, Ross-Ibarra J, Rounsley S, Hamilton RS, Schurr U, Stein N, Tomooka N, van der Knaap E, van Tassel D, Toll J, Valls J, Varshney RK, Ward J, Waugh R, Wenzl P, Zamir D (2013) Agriculture: feeding the future. *Nature* 499: 23–24.
- Miyazaki J, Stiller WN, Truong TT, Xu Q, Hocart CH, Wilson LJ and Wilson IW (2014) Jasmonic acid is associated with resistance to two spotted spider mites in diploid cotton (*Gossypium arboreum*). *Functional Plant Biology* 41: 748–757.
- Parsi RD, Kakde MV, Pawar K and Patil RSP (2016) Influence of fibre length on ring spun yarn quality. *International Journal of Research and Scientific Innovation* 3: 154–156.
- Pearson K (1901) On lines and planes of closest fit to systems of points in space. *Philosophical Magazine* 2: 559–572.
- Perrier X and Jacquemoud-Collet JP (2006) DARwin software. Available at <http://darwin.cirad.fr/>
- Rahman M (2016) Cotton improvement for environmentally stressed economies. *The International Cotton Advisory Committee Recorder* 34: 12.
- Rathinavel K (2018) Principal component analysis with quantitative traits in extant cotton varieties (*Gossypium hirsutum* L.) and parental lines for diversity. *Current Agriculture Research Journal* 6: 54–64.
- Romeu-Dalmau C, Bonsall M, Willis K, Dolan L. (2015) Asiatic cotton can generate similar economic benefits to Bt cotton under rainfed conditions. *Nature Plants* 1: 15072. <https://doi.org/10.1038/nplants.2015.72>
- Ruan YL, Xu SM, White R and Furbank RT (2004) Genotypic and developmental evidence for the role of plasmodesmatal regulation in cotton fibre elongation mediated by callose turnover. *Plant Physiology* 136: 4104–4113.
- Singh B (2001) *Plant Breeding: Principles and Methods*, 6th edn. New Delhi, India: Kalyani Publishers.
- Subramaniam S and Madhav M (1973) Inheritance of short stature in rice. *Madras Agriculture Journal* 60: 1129–1133.
- Wang F, Zhang C, Liu G, Chen Y, Zhang J and Qiao Q (2016) Phenotypic variation analysis and QTL mapping for cotton (*Gossypium hirsutum* L.) fiber quality grown in different cotton-producing regions. *Euphytica* 211: 169–183.
- Ward JH (1963) Hierarchical grouping to optimize an objective function. *The Journal of the American Statistical Association* 58: 236–244.
- Wendel JF, Flagel LE and Adams KL (2012) Jeans, genes, and genomes: cotton as a model for studying polyploidy. In: Soltis P and Soltis D (eds) *Polyploidy and Genome Evolution*, pp. 181–207. Berlin, Heidelberg: Springer.
- Yu J, Zhang K, Li S, Yu S, Zhai H, Wu M, Li X, Fan S, Song M, Yang D, Li Y, Zhang J (2013) Mapping quantitative trait loci for lint yield and fiber quality across environments in a *Gossypium hirsutum* × *Gossypium barbadense* backcross inbred line population. *Theoretical and Applied Genetics* 126: 275–287.