

PARTITIONING THE REAL LINE INTO BOREL SETS

WILL BRIAN

Abstract. For which infinite cardinals κ is there a partition of the real line \mathbb{R} into precisely κ Borel sets? Work of Lusin, Souslin, and Hausdorff shows that \mathbb{R} can be partitioned into \aleph_1 Borel sets. But other than this, we show that the spectrum of possible sizes of partitions of \mathbb{R} into Borel sets can be fairly arbitrary. For example, given any $A \subseteq \omega$ with $0, 1 \in A$, there is a forcing extension in which $A = \{n : \text{there is a partition of } \mathbb{R} \text{ into } \aleph_n \text{ Borel sets}\}$. We also look at the corresponding question for partitions of \mathbb{R} into closed sets. We show that, like with partitions into Borel sets, the set of all uncountable κ such that there is a partition of \mathbb{R} into precisely κ closed sets can be fairly arbitrary.

§1. Introduction. By work of Lusin and Sierpiński in [16], there is an analytic, non-Borel subset of \mathbb{R} , and every analytic and every coanalytic set is the union of \aleph_1 Borel sets. It follows that \mathbb{R} can be partitioned into \aleph_1 Borel sets. Hausdorff later sharpened this in [10], showing that \mathbb{R} , or any other uncountable Polish space, can be written as an increasing union of $\aleph_1 G_\delta$ sets, and therefore can be partitioned into $\aleph_1 F_{\sigma\delta}$ sets. This raises the question:

(Q1) For which uncountable cardinals κ is there a partition of \mathbb{R} into κ nonempty Borel sets?

Let us define the *Borel partition spectrum*, denoted $\text{sp}(\text{Borel})$, to be the answer to this question:

$$\text{sp}(\text{Borel}) = \{|\mathcal{P}| : \mathcal{P} \text{ is a partition of } \mathbb{R} \text{ into uncountably many Borel sets}\}.$$

The main result of this paper shows that the Borel partition spectrum can be fairly arbitrary. (A precise statement of the result can be found in Corollary 3.3.) For example, given any set A of positive integers, there is a forcing extension in which $\text{sp}(\text{Borel}) = \{\aleph_n : n \in A\} \cup \{\aleph_1, \aleph_\omega, \aleph_{\omega+1}\}$.

A related question can be asked for any given class of subsets of \mathbb{R} . As \mathbb{R} is connected, there is no partition of \mathbb{R} into two or more open sets. So the descriptively simplest sets one can ask this for are the closed sets.

(Q2) For which cardinals κ is there a partition of \mathbb{R} into κ closed sets?

Making use of the Baire Category Theorem, Sierpiński proved in [23] that any partition of \mathbb{R} into at least two nonempty closed sets is uncountable. (Using the modern vocabulary, his proof actually shows that any partition of \mathbb{R} into closed sets

Received January 6, 2022.

2020 *Mathematics Subject Classification*. Primary 03E35, Secondary 03E17.

Key words and phrases. partitions, Borel sets, forcing, isomorphism-of-names arguments, cardinal characteristics of the continuum.

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

0022-4812/24/8902-0005

DOI:10.1017/jsl.2023.20



has size $\geq \text{cov}(\mathcal{M})$.) Let us define the *closed partition spectrum*, denoted $\text{sp}^{(\text{closed})}$, to be the answer to (Q2):

$$\text{sp}^{(\text{closed})} = \{|\mathcal{P}| : \mathcal{P} \text{ is a partition of } \mathbb{R} \text{ into at least two closed sets}\}.$$

We show that, like the Borel partition spectrum, $\text{sp}^{(\text{closed})}$ can be fairly arbitrary—even more so, in fact, since the closed partition spectrum need not contain \aleph_1 . For example, given any nonempty $A \subseteq \omega \setminus \{0\}$, there is a forcing extension in which $\text{sp}^{(\text{closed})} = \{\aleph_n : n \in A\} \cup \{\aleph_\omega, \aleph_{\omega+1}\}$.

These results about $\text{sp}^{(\text{Borel})}$ and $\text{sp}^{(\text{closed})}$ are encompassed in a single theorem, Theorem 3.2. The proof identifies a notion of forcing that adds partitions of \mathbb{R} into closed sets having certain prescribed sizes, while avoiding partitions of \mathbb{R} into Borel sets with other sizes. The second of these tasks is the more difficult. It is accomplished via an isomorphism-of-names argument, similar to the folklore proof (found, e.g., in [3, Theorem 3.1]) that, after many mutually generic Cohen reals are added to a model of CH, there are no MAD families of size strictly between \aleph_1 and \mathfrak{c} .

Let us point out that analogues of (Q1) and (Q2) have been asked and answered (or partly answered) concerning other extremal families. For example, consider $\text{sp}^{(\text{MAD})} = \{\kappa \geq \aleph_0 : \text{there is a MAD family of size } \kappa\}$. Hechler showed in [11] that $\text{sp}^{(\text{MAD})}$ can include any prescribed set of cardinals, and Blass showed in [2] how to exclude certain cardinals from $\text{sp}^{(\text{MAD})}$. Shelah and Spinas proved the strongest results in [22], showing that $\text{sp}^{(\text{MAD})}$ can be rather arbitrary, especially on the regular cardinals. Similarly, the spectrum of possible sizes of maximal independent families, $\text{sp}^{(\text{mif})}$, was investigated recently by Fischer and Shelah in [7, 8]. Like with $\text{sp}^{(\text{MAD})}$, they proved that $\text{sp}^{(\text{mif})}$ can be fairly arbitrary, especially on the regular cardinals. Similar work concerning maximal cofinitary groups was done by Fischer in [6]. Ultimately, all these proofs share the same core idea: variations on the isomorphism-of-names argument mentioned above. In every case, the key to making this kind of argument work is to find an automorphism-rich poset that can be used to add extremal families of prescribed sizes. One notable exception to this rule is Shelah’s analysis of the set of possible sizes of ultrafilter bases in [21], where he proves, from large cardinal hypotheses, that this spectrum can exhibit fairly chaotic behavior.

§2. $\text{sp}^{(\text{Borel})}$ and $\text{sp}^{(\text{closed})}$ do not depend on \mathbb{R} . We begin this section by observing that the definition of $\text{sp}^{(\text{Borel})}$ does not depend on \mathbb{R} , and remains unchanged when \mathbb{R} is replaced by any other uncountable Polish space:

THEOREM 2.1. *If X is any uncountable Polish space, then*

$$\text{sp}^{(\text{Borel})} = \{\kappa > \aleph_0 : \text{there is a partition of } X \text{ into } \kappa \text{ Borel sets}\}.$$

PROOF. By a theorem of Kuratowski (see [14, Theorem 15.6]), any two uncountable Polish spaces are Borel isomorphic: in other words, there is a bijection $f : \mathbb{R} \rightarrow X$ such that $A \subseteq \mathbb{R}$ is Borel if and only if $f[A]$ is Borel. Thus if \mathcal{P} is any partition of \mathbb{R} into Borel sets, then $\{f[B] : B \in \mathcal{P}\}$ is a partition of X into Borel sets, and if \mathcal{Q} is any partition of X into Borel sets, then $\{f^{-1}[B] : B \in \mathcal{Q}\}$ is a partition of \mathbb{R} into Borel sets. ◻

It turns out that the same is true for $\mathfrak{sp}(\text{closed})$: if X is any uncountable Polish space, then

$$\mathfrak{sp}(\text{closed}) = \{\kappa > \aleph_0 : \text{there is a partition of } X \text{ into } \kappa \text{ closed sets}\}.$$

A related theorem is proved by Miller in [17, Theorem 3]: \mathbb{R} can be partitioned into \aleph_1 closed sets if and only if some uncountable Polish space can be, if and only if every uncountable Polish space can be. We wish to prove the same, but with an arbitrary uncountable cardinal κ in place of \aleph_1 . Miller's proof does not readily adapt to this task, because it uses in an essential way the fact that \aleph_1 is the smallest uncountable cardinal. So we take a different approach. First we need a few lemmas.

LEMMA 2.2. *For any uncountable cardinal κ , the following are equivalent:*

- (1) *There is a partition of 2^ω into κ closed sets.*
- (2) *There is a partition of ω^ω into κ compact sets.*
- (3) *For every uncountable Polish space X , there is a partition of X into κ compact sets.*
- (4) *For some uncountable compact Polish space X , there is a partition of X into κF_σ sets.*

PROOF. Throughout the proof, if $X \subseteq Y$ and \mathcal{P} is a partition of Y , then $\mathcal{P} \upharpoonright X = \{K \cap X : K \in \mathcal{P}\}$ denotes the restriction of \mathcal{P} to X . We prove that (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (1).

(1) \Rightarrow (2): Suppose \mathcal{P} is a partition of 2^ω into κ closed sets. We claim first that there is a closed $X \subseteq 2^\omega$ such that $\mathcal{P} \upharpoonright X$ is a partition of X into κ nowhere dense closed sets (nowhere dense in X , that is).

To see this, we define a descending transfinite sequence of closed subsets of 2^ω . Let $X_0 = 2^\omega$, and if α is a limit ordinal, take $X_\alpha = \bigcap_{\xi < \alpha} X_\xi$. At stage α , given X_α , form $X_{\alpha+1}$ by removing any open subset of X_α contained in a single member of \mathcal{P} : $X_{\alpha+1} = X_\alpha \setminus \bigcup \{U : U \text{ is open in } X_\alpha \text{ and } |\mathcal{P} \upharpoonright U| = 1\}$. Because 2^ω is second countable, there is some $\beta < \omega_1$ such that $X_\beta = X_\gamma$ for all $\gamma \geq \beta$. Let $X = X_\beta$. Clearly, $\mathcal{P} \upharpoonright X$ is a partition of X into compact nowhere dense sets. Furthermore, at any stage $\alpha < \beta$ of our recursion, $\{K \in \mathcal{P} : K \cap X_\alpha \neq K \cap X_{\alpha+1}\}$ is countable. Thus $\{K \in \mathcal{P} : K \cap X \neq K\}$ is countable. It follows that $|\mathcal{P} \upharpoonright X| = |\mathcal{P}| = \kappa$.

Now, we claim there is a subspace Y of X such that $Y \approx \omega^\omega$ and $\mathcal{P} \upharpoonright Y$ is a partition of Y into κ compact sets. Fix a countable basis \mathcal{B} for X , and for every $U \in \mathcal{B}$ fix some $K_U \in \mathcal{P}$ such that $K_U \cap U \neq \emptyset$. Let $Y = X \setminus \bigcup \{K_U : U \in \mathcal{B}\}$. Clearly $\mathcal{P} \upharpoonright Y = \mathcal{P} \setminus \{K_U : U \in \mathcal{B}\}$, so \mathcal{P} partitions Y into κ compact sets. The set $\{K_U : U \in \mathcal{B}\}$ is both dense and meager in X . It follows that no relatively (cl)open subset of Y is closed in X , and therefore no (cl)open subset of Y is compact. Furthermore Y is G_δ in X , and therefore Polish. That $Y \approx \omega^\omega$ now follows from the Alexander–Urysohn characterization of ω^ω as the unique nowhere compact, zero-dimensional Polish space (see [14, Theorem 7.7]).

(2) \Rightarrow (3): Suppose \mathcal{P} is a partition of ω^ω into κ compact sets, and let X be any uncountable Polish space. Decompose X into its scattered part and perfect part: i.e., let $X = Y \cup Z$, where Y is countable and Z is closed in X (hence still Polish) and Z has no isolated points. By [14, Exercise 7.15], there is a continuous bijection $f : \omega^\omega \rightarrow Z$. But then $\{f[K] : K \in \mathcal{P}\} \cup \{\{y\} : y \in Y\}$ is a partition of X into κ compact sets.

(3) \Rightarrow (4): This implication is obvious, since “every uncountable Polish space” includes some compact spaces, and compact sets are F_σ .

(4) \Rightarrow (1): Suppose X is an uncountable compact Polish space. By [14, Theorem 7.4], there is a continuous surjection $f : 2^\omega \rightarrow X$. If \mathcal{P} is any partition of X into κF_σ sets, then $\mathcal{Q} = \{f^{-1}[K] : K \in \mathcal{P}\}$ is a partition of 2^ω into κF_σ sets. But every F_σ subset of 2^ω can be partitioned into countably many closed sets. (This observation is attributed to Luzin in [17, Theorem 2].) Thus, by breaking up any non-closed members of \mathcal{Q} into countably many closed pieces, we can refine \mathcal{Q} to obtain a partition of 2^ω into κ closed sets. \dashv

Define the *Sierpiński cardinal* \mathfrak{n} to be the minimum size of a cardinal satisfying the equivalent statements in Lemma 2.2: that is,

$$\begin{aligned} \mathfrak{n} &= \min \{|\mathcal{P}| : \mathcal{P} \text{ is a partition of } 2^\omega \text{ into uncountably many closed sets}\} \\ &= \min \{|\mathcal{P}| : \mathcal{P} \text{ is a partition of } \omega^\omega \text{ into compact sets}\}. \end{aligned}$$

Recall that the dominating number \mathfrak{d} is equal to the smallest covering of ω^ω by compact sets. Hence $\mathfrak{d} \leq \mathfrak{n}$. And clearly $\mathfrak{n} \leq \mathfrak{c}$, because ω^ω can be partitioned into singletons. Thus we may consider \mathfrak{n} to be a cardinal characteristic of the continuum. Quite a bit is known already about this cardinal. The main results are due to Stern [25], Miller [17], Newelski [19], and Spinas [24], who studied this cardinal implicitly without giving it a name, and Hrušák [12, 13], who denotes it \mathfrak{a}_T . The name “Sierpiński cardinal” and the notation \mathfrak{n} were suggested by Banach in [1].

A set S of cardinals is *closed under singular limits* if for every singular cardinal κ , if $S \cap \kappa$ is unbounded in κ , then $\kappa \in S$.

LEMMA 2.3. *The set $S = \{\kappa : \text{there is a partition of } 2^\omega \text{ into } \kappa \text{ closed sets}\}$ is closed under singular limits.*

PROOF. Suppose κ is a singular cardinal and $S \cap \kappa$ is unbounded in κ . Let $\nu = \text{cf}(\kappa) < \kappa$, and let $\langle \alpha_\xi : \xi < \nu \rangle$ be a sequence of cardinals in S increasing up to κ . Fix some $\lambda \in S$ with $\nu \leq \lambda < \kappa$.

Let $\{K_\xi : \xi < \lambda\}$ be a partition of 2^ω into closed sets. Observe that $\{2^\omega \times K_\xi : \xi < \lambda\}$ is a partition of $2^\omega \times 2^\omega \approx 2^\omega$ into λ copies of 2^ω . Thus we may (and do) assume that $K_\xi \approx 2^\omega$ for each $\xi < \lambda$. For each $\xi < \nu$, let \mathcal{P}_ξ be a partition of K_ξ into α_ξ closed sets (using the fact that $\alpha_\xi \in S$). Then $\bigcup_{\xi < \nu} \mathcal{P}_\xi \cup \{K_\xi : \nu \leq \xi < \lambda\}$ is a partition of 2^ω into κ closed sets. \dashv

THEOREM 2.4. *Let κ be an uncountable cardinal. Then all six statements of the following form are equivalent:*

Some/every uncountable Polish space can be partitioned into κ compact/closed/ F_σ sets.

PROOF. It is clear that (every-compact) \Rightarrow (every-closed) \Rightarrow (every- F_σ), and that (some-compact) \Rightarrow (some-closed) \Rightarrow (some- F_σ). Also, Lemma 2.2 implies that (every- F_σ) \Rightarrow (some-compact): because if every uncountable Polish space can be partitioned into κF_σ sets, then in particular some uncountable compact Polish space can be, and by Lemma 2.2 this implies ω^ω can be partitioned into κ compact sets. Thus, to prove the theorem, we need to show (some- F_σ) \Rightarrow (every-compact).

So let X be some uncountable Polish space, and suppose \mathcal{P} is a partition of X into κF_σ sets. If $\kappa = \acute{n}$, then by definition, there is a partition of ω^ω into \acute{n} compact sets, and by Lemma 2.2 every uncountable Polish space can be partitioned into κ compact sets, and we are done. So let us suppose $\kappa > \acute{n}$. We consider two cases.

For the first case, suppose κ is regular. By the definition of \acute{n} and Lemma 2.2, there is a partition \mathcal{Q} of X into \acute{n} compact sets. Because $|\mathcal{P}| = \kappa > \acute{n}$ and κ is regular, there is some $K \in \mathcal{Q}$ such that $|\mathcal{P} \upharpoonright K| = \kappa$. Thus $\mathcal{P} \upharpoonright K$ is a partition of K into κ compact subsets. But K is a compact Polish space, and uncountable because $|K| \geq |\mathcal{P} \upharpoonright K| = \kappa$. Thus some compact uncountable Polish space can be partitioned into κF_σ sets. Invoking Lemma 2.2 again, this implies every uncountable Polish space can be partitioned into κ compact sets.

For the second case, suppose κ is singular. As in the first case, there is a partition \mathcal{Q} of X into \acute{n} compact sets. If $|\mathcal{P} \upharpoonright K| = \kappa$ for some $K \in \mathcal{Q}$, then we may argue as in the first case and conclude that every uncountable Polish space can be partitioned into κ compact sets, and we are done. So let us suppose instead that $|\mathcal{P} \upharpoonright K| < \kappa$ for every $K \in \mathcal{Q}$. Let λ be any cardinal with $\acute{n} < \lambda < \kappa$. Because $|\mathcal{P}| = \kappa > \lambda > \acute{n} = |\mathcal{Q}|$, there is some $K \in \mathcal{Q}$ such that $|\mathcal{P} \upharpoonright K| \geq \lambda$. Let $\mu = |\mathcal{P} \upharpoonright K|$, and note that $\mu < \kappa$ (by the third sentence of this paragraph). Now, as in the previous paragraph, K is a compact uncountable Polish space that can be partitioned into μF_σ sets. By Lemma 2.2, this implies 2^ω can be partitioned into μ closed sets. Because λ was an arbitrary cardinal below κ and $\lambda \leq \mu < \kappa$, this shows that $S = \{\mu : \text{there is a partition of } 2^\omega \text{ into } \mu \text{ closed sets}\}$ is unbounded below κ . By Lemma 2.3, $\kappa \in S$. By Lemma 2.2, this implies every uncountable Polish space can be partitioned into κ compact sets. ⊣

COROLLARY 2.5. *For any uncountable Polish space X ,*

$$\begin{aligned} \mathfrak{sp}(\text{closed}) &= \{\kappa > \aleph_0 : \text{there is a partition of } X \text{ into } \kappa \text{ compact sets}\} \\ &= \{\kappa > \aleph_0 : \text{there is a partition of } X \text{ into } \kappa \text{ closed sets}\} \\ &= \{\kappa > \aleph_0 : \text{there is a partition of } X \text{ into } \kappa F_\sigma \text{ sets}\}. \end{aligned}$$

COROLLARY 2.6. $\min(\mathfrak{sp}(\text{closed})) = \acute{n} \geq \mathfrak{d}$.

We note that the inequality $\acute{n} \geq \mathfrak{d}$ has been observed before, and can be considered folklore. It is (arguably) implicit in [17], and was observed explicitly by Hrušák in [13] and later by Banach in [1]. Anticipating the main theorem in Section 3, note that this inequality gives us an easy way of excluding an initial segment of the uncountable cardinals from $\mathfrak{sp}(\text{closed})$: simply make \mathfrak{d} big.

COROLLARY 2.7. $\mathfrak{sp}(\text{closed})$ is closed under singular limits.

PROOF. This follows from Corollary 2.5 and Lemma 2.3. ⊣

It is worth pointing out that the same result holds for $\mathfrak{sp}(\text{Borel})$, by a very similar argument. The analogous result also holds for the set $\mathfrak{sp}(\text{MAD})$ mentioned in the introduction [11, Theorem 3.1], again by a similar argument.

THEOREM 2.8. $\mathfrak{sp}(\text{Borel})$ is closed under singular limits.

PROOF. Suppose κ is a singular cardinal and $\mathfrak{sp}(\text{Borel}) \cap \kappa$ is unbounded in κ . Let $\nu = \text{cf}(\kappa) < \kappa$, and let $\langle \alpha_\xi : \xi < \nu \rangle$ be a sequence of cardinals in $\mathfrak{sp}(\text{Borel})$ increasing up to κ . Fix some $\lambda \in \mathfrak{sp}(\text{Borel})$ with $\nu \leq \lambda < \kappa$.

Let $\{B_\xi : \xi < \lambda\}$ be a partition of 2^ω into λ Borel sets. Observe that $\{2^\omega \times B_\xi : \xi < \lambda\}$ is a partition of $2^\omega \times 2^\omega \approx 2^\omega$ into λ uncountable Borel sets. Thus we may (and do) assume that B_ξ is uncountable for each $\xi < \lambda$. Every uncountable Borel set contains a closed subspace homeomorphic to 2^ω . For each $\xi < \nu$, fix $C_\xi \subseteq B_\xi$ with $C_\xi \approx 2^\omega$, and let \mathcal{P}_ξ be a partition of C_ξ into α_ξ Borel sets. Then $\bigcup_{\xi < \nu} \mathcal{P}_\xi \cup \{B_\xi \setminus C_\xi : \xi < \nu\} \cup \{B_\xi : \nu \leq \xi < \lambda\}$ is a partition of 2^ω into κ Borel sets. ⊣

We end this section with some terminology regarding trees, and two open questions regarding the Sierpiński cardinal \mathfrak{n} .

Recall that a *subtree* of $2^{<\omega}$ is a subset of $2^{<\omega}$ that is closed under taking initial segments. A subtree T of $2^{<\omega}$ is *pruned* if every node of T has a successor in T . A *branch* through T means a function $b \in 2^\omega$ such that $b \upharpoonright n \in T$ for all $n < \omega$. If T is a subtree of $2^{<\omega}$, we denote by $\llbracket T \rrbracket$ the set of all branches through T . It is not difficult to see that $\llbracket T \rrbracket$ is closed in 2^ω for any subtree T of $2^{<\omega}$; and conversely, for every closed $C \subseteq 2^\omega$ there is a subtree T of $2^{<\omega}$ with $C = \llbracket T \rrbracket$ (for details, see [14, Chapter 2]). Similar terminology is used for subtrees of $\omega^{<\omega}$ or of $2^{<k}$.

Representing closed sets with trees in this way, Theorem 2.4 states that $\kappa \in \mathfrak{sp}(\text{closed})$ if and only if $\kappa > \aleph_0$ and there is a MAD family of κ subtrees of $2^{<\omega}$. Similarly, $\kappa \in \mathfrak{sp}(\text{closed})$ if and only if there is a MAD family of κ finitely branching, pruned subtrees of $\omega^{<\omega}$. These characterizations of $\mathfrak{sp}(\text{closed})$ explain Hrušák’s notation, writing \mathfrak{a}_T for $\min(\mathfrak{sp}(\text{closed}))$. This relationship between \mathfrak{n} and \mathfrak{a} raises the following questions.

QUESTION 2.9. *Is it consistent that \mathfrak{n} has countable cofinality?*

The corresponding question for \mathfrak{a} was solved by Brendle in [4], where he used Shelah’s template forcing technique to obtain a model of $\mathfrak{a} = \aleph_\omega$. It is relatively easy to make \mathfrak{n} singular of uncountable cofinality: e.g., by adding \aleph_{ω_1} Cohen reals to a model of CH, we get a model where $\mathfrak{d} = \mathfrak{n} = \mathfrak{c} = \aleph_{\omega_1}$.

It is also simple to prove the consistency of $\mathfrak{a} < \mathfrak{n}$. This holds, for example, in the Cohen model, where $\aleph_1 = \mathfrak{a} < \mathfrak{d} = \mathfrak{n} = \mathfrak{c}$.

QUESTION 2.10. *Is $\mathfrak{a} \leq \mathfrak{n}$?*

We note that if $\mathfrak{n} < \mathfrak{a}$ is consistent, then proving it is likely very difficult. This is because $\mathfrak{n} < \mathfrak{a}$ implies $\mathfrak{d} < \mathfrak{a}$. The consistency of $\mathfrak{d} < \mathfrak{a}$ was an open question for a long time, solved by Shelah in [20]. Shelah’s technique will not work, however, for obtaining a model of $\mathfrak{n} < \mathfrak{a}$: his technique makes \mathfrak{n} large for the same reasons it makes \mathfrak{a} large. (Roughly, by an “averaging of names” argument, an ultrapower $\mathbb{P}^\kappa/\mathcal{U}$ forces that if $\mathfrak{n} \geq \kappa$ then $\mathfrak{n} = \mathfrak{c}$.)

§3. Forcing an (almost) arbitrary spectrum. Every $\kappa \in \mathfrak{sp}(\text{Borel})$ is an example what Blass calls a “ Δ_1^1 characteristic” in [2]. Blass proves [2, Theorems 8 and 9] that there can be many cardinals between \aleph_1 and \mathfrak{c} that are not Δ_1^1 characteristics, and therefore are not in $\mathfrak{sp}(\text{Borel})$. For example, it is consistent to have the set of Δ_1^1

characteristics be equal to precisely $\{\aleph_n : n \text{ is a power of } 17\} \cup \{\aleph_\omega, \aleph_{\omega+1}\}$. Blass' method does not guarantee that any of these Δ_1^1 characteristics will be in $\mathfrak{sp}(\text{Borel})$, except of course for \aleph_1 and \mathfrak{c} . But his results do show at least that $\mathfrak{sp}(\text{Borel})$ can contain large gaps, and that it is possible to surgically exclude specific cardinals from $\mathfrak{sp}(\text{Borel})$. (We should mention that the consistency of having $\aleph_2 \notin \mathfrak{sp}(\text{Borel})$ predates Blass' work, and is due to Miller [18].)

In the other direction, extending a prior result of Miller's [17, Theorem 4], Miller and the author showed in [5, Theorem 3.12] that:

THEOREM 3.1. *For any $\kappa \geq \mathfrak{c}$ with $\text{cf}(\kappa) > \omega$, there is a ccc forcing extension in which $\mathfrak{c} = \kappa$ and $\mathfrak{sp}(\text{closed}) = [\aleph_1, \kappa]$.*

In particular, for any set C of cardinals, there is a ccc forcing extension in which $C \subseteq \mathfrak{sp}(\text{closed})$. We now sketch a proof of this theorem, in order to introduce some of the ideas used in the proof of the main theorem below, but in a simpler context. This sketch can be skipped by the reader requiring no such introduction.

PROOF SKETCH OF THEOREM 3.1. Given some $X \subseteq 2^\omega$ with empty interior, let \mathbb{T}_X be the poset whose conditions are pairs (t, B) , where

- There is some $k \in \omega$ such that t is a subtree of $2^{<k}$, and t is pruned: i.e., if $\sigma \in t$ with $|\sigma| < k$, then σ has a proper extension in t .
- B is a finite subset of $2^\omega \setminus X$, and $b \upharpoonright k$ is a branch of t for each $b \in B$.

The ordering on \mathbb{T}_X is defined by: (t', B') extends (t, B) if and only if $B' \supseteq B$ and t' is an end extension of t (meaning that $t = t' \cap 2^{<k}$ for some k).

Roughly, a condition (t, B) can be thought of as a finite approximation t to a subtree T of $2^{<\omega}$ that we are trying to build generically, and a promise that some finite set B of reals will be branches of T .

For any $X \subseteq 2^\omega$, \mathbb{T}_X is σ -centered. The poset \mathbb{T}_X generically adds an infinite pruned subtree T of $2^{<\omega}$, defined from a generic filter G on \mathbb{T}_X as $T = \bigcup \{t : (t, B) \in G \text{ for some } B\}$. (Equivalently, T is the evaluation in $V[G]$ of the name $\dot{T} = \{\langle \sigma, q \rangle : q = (t, B) \text{ for some } B, \text{ and } \sigma \in t\}$.) In the extension, $\llbracket T \rrbracket$ is a closed subset of 2^ω disjoint from X .

Let \mathbb{T}_X^ω denote the finite support product of countably many copies of \mathbb{T}_X . For any $X \subseteq 2^\omega$, \mathbb{T}_X^ω is σ -centered. The poset \mathbb{T}_X^ω generically adds countably many infinite pruned subtrees T_0, T_1, T_2, \dots of $2^{<\omega}$, and in the extension,

$$(2^\omega)^V \cap \bigcup_{n < \omega} \llbracket T_n \rrbracket = (2^\omega)^V \setminus X.$$

In other words, \mathbb{T}_X^ω generically adds an F_σ subset of 2^ω that intersects the ground model reals in precisely the complement of X . (Note: this poset may look familiar: it is the one usually used for showing that Martin's Axiom implies every $< \mathfrak{c}$ -sized subset of 2^ω is a Q -set.)

Let C denote the set of cardinals in $[\aleph_1, \kappa]$. We now define a finite support iteration of length ω_1 as follows. At stage 0, force with the poset $\mathbb{Q}_0 = \mathbb{C}_\kappa$ of finite partial functions $\kappa \rightarrow 2$, in order to add a set of κ mutually generic Cohen reals. Let $\{c_\xi : \xi < \kappa\}$ be an enumeration of these Cohen reals in $V^{\mathbb{Q}_0}$, and for each $\mu \in C$, let $\mathcal{P}_0^\mu = \{\{c_\xi\} : \xi < \mu\}$. Note that \mathcal{P}_0^μ is a μ -sized collection of disjoint subsets of \mathbb{R} : we think of \mathcal{P}_0^μ as a first approximation to a μ -sized partition we are trying to build. At a later stage α of the iteration, suppose we have already obtained, for each

$\mu \in C$, a μ -sized collection \mathcal{P}_α^μ of disjoint subsets of \mathbb{R} . In $V^{\mathbb{Q}_\alpha}$, define $X_\alpha^\mu = \bigcup \mathcal{P}_\alpha^\mu$ for each $\mu \in C$, and then obtain $V^{\mathbb{Q}_{\alpha+1}}$ from $V^{\mathbb{Q}_\alpha}$ by forcing with $\prod_{\mu \in C} \mathbb{T}_{X_\alpha^\mu}^\omega$. This adds countably many generic trees $T_{\alpha,0}^\mu, T_{\alpha,1}^\mu, T_{\alpha,2}^\mu, \dots$ for each $\mu \in C$, and in $V^{\mathbb{Q}_{\alpha+1}}$ we define $\mathcal{P}_{\alpha+1}^\mu = \mathcal{P}_\alpha^\mu \cup \{\bigcup_{n \in \omega} \llbracket T_{\alpha,n}^\mu \rrbracket\}$.

At the end of the iteration, in $V^{\mathbb{Q}_{\omega_1}}$, let $\mathcal{P}^\mu = \bigcup_{\alpha < \omega_1} \mathcal{P}_\alpha^\mu$. This is a μ -sized collection of disjoint F_σ subsets of \mathbb{R} . Furthermore, if x is a real in $V^{\mathbb{Q}_{\omega_1}}$, then there is some $\alpha < \omega_1$ with $x \in V^{\mathbb{Q}_\alpha}$. At that stage of the iteration, either $x \in \bigcup \mathcal{P}_\alpha^\mu$, or if not, then $x \in \bigcup_{n \in \omega} \llbracket T_{\mu,n}^\alpha \rrbracket$ because $(2^\omega)^{V^{\mathbb{P}_\alpha}} \cap \bigcup_{n < \omega} \llbracket T_{\mu,n}^\alpha \rrbracket = (2^\omega)^{V^{\mathbb{P}_\alpha}} \setminus X_\alpha^\mu$. Either way, $x \in \bigcup \mathcal{P}_{\alpha+1}^\mu \subseteq \bigcup \mathcal{P}^\mu$. Thus \mathcal{P}^μ is a partition of 2^ω into F_σ sets in $V^{\mathbb{Q}_{\omega_1}}$, and this means $\mu \in \mathfrak{sp}(\text{closed})$ by Corollary 2.5. \dashv

The main theorem is proved with a modification of this poset, with two major changes. First, the set C will not necessarily be an interval of cardinals, but will consist only of those cardinals we wish to add to $\mathfrak{sp}(\text{closed})$. Second, instead of a true iteration, we use a streamlined modification. This modified iteration has actual finite sequences (rather than names for them) for the working parts of the $\mathbb{T}_{X_\alpha^\mu}$, which is helpful for proving that certain permutations of the Cohen reals extend to automorphisms of the entire poset. (And these automorphisms are essential for excluding cardinals $\notin C$ from $\mathfrak{sp}(\text{Borel})$.) The definition of this modified iteration is reminiscent of the template forcing notions in [3], but with a well-ordered template, so that it is essentially an iteration.

THEOREM 3.2. *Let C be a set of uncountable cardinals such that:*

- $\min(C)$ is regular,
- $|C| < \min(C)$,
- C has a maximum with $\text{cf}(\max(C)) > \omega$,
- C is closed under singular limits, and
- if λ is singular and $\lambda \in C$, then $\lambda^+ \in C$.

Assuming GCH holds up to $\max(C)$, there is a ccc forcing extension in which $\mathfrak{c} = \max(C)$, $\mathfrak{sp}(\text{closed}) = C$, and if $\min(C) < \lambda \notin C$ then $\lambda \notin \mathfrak{sp}(\text{Borel})$.

Before proving this theorem, let us deduce some relatively easy corollaries from it, including the results mentioned in the introduction.

COROLLARY 3.3. *Let C be a set of uncountable cardinals such that:*

- $\aleph_1 \in C$,
- C is at most countable,
- C has a maximum with $\text{cf}(\max(C)) > \omega$,
- C is closed under singular limits, and
- if λ is singular and $\lambda \in C$, then $\lambda^+ \in C$.

Assuming GCH holds up to $\max(C)$, there is a ccc forcing extension in which $\mathfrak{c} = \max(C)$ and $\mathfrak{sp}(\text{Borel}) = \mathfrak{sp}(\text{closed}) = C$.

PROOF. This follows immediately from the previous theorem. \dashv

COROLLARY 3.4. *Given any $A \subseteq \omega \setminus \{0\}$, there is a forcing extension in which $\mathfrak{sp}(\text{Borel}) = \{\aleph_n : n \in A\} \cup \{\aleph_1, \aleph_\omega, \aleph_{\omega+1}\}$, and if $A \neq \emptyset$ there is a forcing extension in which $\mathfrak{sp}(\text{closed}) = \{\aleph_n : n \in A\} \cup \{\aleph_\omega, \aleph_{\omega+1}\}$.*

PROOF. Fix some $A \subseteq \omega \setminus \{0\}$. First pass to a forcing extension in which GCH holds up to $\aleph_{\omega+1}$. Then, for the result about $\mathfrak{sp}(\text{closed})$, apply Theorem 3.2 with $C = \{\aleph_n : n \in A\} \cup \{\aleph_\omega, \aleph_{\omega+1}\}$. Or, for the result about $\mathfrak{sp}(\text{Borel})$, apply Corollary 3.3 with $C = \{\aleph_n : n \in A\} \cup \{\aleph_1, \aleph_\omega, \aleph_{\omega+1}\}$. \dashv

COROLLARY 3.5. *Given any finite nonempty $A \subseteq \omega \setminus \{0\}$, there is a forcing extension in which $\mathfrak{sp}(\text{Borel}) = \{\aleph_n : n \in A\} \cup \{\aleph_1\}$, and there is a forcing extension in which $\mathfrak{sp}(\text{closed}) = \{\aleph_n : n \in A\}$.*

PROOF. This is proved in exactly the same way as the previous corollary, but without \aleph_ω and $\aleph_{\omega+1}$ put into C . \dashv

PROOF OF THEOREM 3.2. Let C be a set of uncountable cardinals satisfying the hypotheses listed in the statement of the theorem, and assume GCH holds up to $\max(C)$. Let $\kappa = \min(C)$ and let $\theta = \max(C)$.

For each $\mu \in C$, let $I_\mu = \mu \times \{\mu, \theta^+\}$. (These are merely indexing sets, and for all practical purposes one may think of each I_μ as a set of atoms, or urelements, in the set-theoretic universe. The relevant properties of these I_μ 's are that they are pairwise disjoint, each I_μ has size μ , and the I_μ do not “interact” in any accidental way with any other sets in the proof.)

Let $I = \bigcup_{\mu \in C} I_\mu$, and let \mathbb{P}_0 denote the poset of finite partial functions from $I \times \omega$ to 2. Note that this is equivalent to the standard poset \mathbb{C}_θ for adding θ mutually generic Cohen reals. For each $i \in I$, let c_i denote in $V^{\mathbb{P}_0}$ the Cohen real added by \mathbb{P}_0 in coordinate i . More formally, c_i denotes in $V^{\mathbb{P}_0}$ the evaluation of the name $\dot{c}_i = \{\langle (n, j), p \rangle : p(i, n) = j\}$.

Next we define, recursively, a poset $\langle \mathbb{P}_\alpha, \leq_{\mathbb{P}_\alpha} \rangle$ for each $\alpha \leq \kappa$ with $\alpha > 0$. At every stage of the recursion, \mathbb{P}_α is defined so that \mathbb{P}_β is a sub-poset of \mathbb{P}_α for each $\beta < \alpha$. Conditions in \mathbb{P}_α are finite partial functions on $(I \times \omega) \cup (\alpha \times C \times \omega)$, and the \mathbb{P}_α are defined so that for any $\beta < \alpha$, the restriction of a condition in \mathbb{P}_α to $(I \times \omega) \cup (\beta \times C \times \omega)$ is a condition in \mathbb{P}_β .

At limit stages, we take $\mathbb{P}_\alpha = \bigcup_{\xi < \alpha} \mathbb{P}_\xi$ and $\leq_{\mathbb{P}_\alpha} = \bigcup_{\xi < \alpha} \leq_{\mathbb{P}_\xi}$. In other words, $\langle \mathbb{P}_\alpha, \leq_{\mathbb{P}_\alpha} \rangle$ is the direct limit of $\langle \langle \mathbb{P}_\xi, \leq_{\mathbb{P}_\xi} \rangle : \xi < \alpha \rangle$ for limit α .

At successor stages, suppose \mathbb{P}_ξ is given for every $\xi \leq \beta$, and $\mathbb{P}_\xi \supseteq \mathbb{P}_\zeta$ whenever $\zeta \leq \xi \leq \beta$. Let $\alpha = \beta + 1$. Conditions in \mathbb{P}_α are finite partial functions p on $(I \times \omega) \cup (\alpha \times C \times \omega)$ such that:

- o If $(i, n) \in (I \times \omega) \cap \text{dom}(p)$, then $p(i, n) \in \{0, 1\}$.
- o If $(\gamma, \mu, n) \in \text{dom}(p)$ for some $\gamma \leq \beta$, $\mu \in C$, and $n \in \omega$, then $p(\gamma, \mu, n) = (t, B)$, where:
 - o t is a pruned subtree of $2^{<k}$ for some $k \in \omega$ (in this context, “pruned” means that if $\sigma \in t$ and $|\sigma| < k$, then σ has an extension in t).
 - o B is a finite set of nice \mathbb{P}_γ -names for members of 2^ω such that for every $\dot{b} \in B$,

$$\begin{aligned}
 p \upharpoonright ((I \times \omega) \cup (\gamma \times C \times \omega)) \Vdash_{\mathbb{P}_\gamma} & \dot{b} \upharpoonright j \in t \text{ for every } j < k, \\
 & \dot{b} \neq \dot{c}_i \text{ for every } i \in I_\mu, \text{ and} \\
 & \dot{b} \text{ is not a branch of } \dot{T}_{\mu, m}^\xi \\
 & \text{for any } \xi < \gamma \text{ and } m \in \omega,
 \end{aligned}$$

where \dot{c}_i is the \mathbb{P}_0 -name described above, but interpreted as a \mathbb{P}_γ -name, and where for every $\xi < \gamma$ and $m \in \omega$, $\dot{T}_{\mu,m}^\xi$ is the \mathbb{P}_γ -name for a subtree of $2^{<\omega}$ defined as follows:

$$\dot{T}_{\mu,m}^\xi = \{ \langle \sigma, q \rangle : q(\xi, \mu, m) = (t', B') \text{ for some } B', \text{ and } \sigma \in t' \}.$$

The extension relation $\leq_{\mathbb{P}_\alpha}$ on \mathbb{P}_α is defined as follows: given $p, q \in \mathbb{P}_\alpha$, we write $q \leq_{\mathbb{P}_\alpha} p$ (meaning that q extends p) if and only if:

- o $\text{dom}(q) \supseteq \text{dom}(p)$,
- o $q \upharpoonright (I \times \omega) \supseteq p \upharpoonright (I \times \omega)$, and
- o if $(\gamma, \mu, n) \in \text{dom}(p)$ for some $\gamma \leq \beta$, $\mu \in C$, and $n \in \omega$, and if $p(\gamma, \mu, n) = (t, B)$ and $q(\gamma, \mu, n) = (t', B')$, then $B' \supseteq B$ and t' is an end extension of t (meaning that $t = t' \cap 2^{<k}$ for some k).

Naturally, we abbreviate “ $\leq_{\mathbb{P}_\alpha}$ ” with “ \leq ” in situations where this creates no ambiguity. We also write $p \upharpoonright \alpha$ as an abbreviation for $p \upharpoonright ((I \times \omega) \cup (\alpha \times C \times \omega))$ where doing so creates no ambiguity.

This completes the recursive definition of the \mathbb{P}_α and $\leq_{\mathbb{P}_\alpha}$. It is easy to see that $\leq_{\mathbb{P}_\alpha}$ is a partial order on \mathbb{P}_α for all $\alpha \leq \kappa$, and that \mathbb{P}_β is a sub-poset of \mathbb{P}_α whenever $\beta \leq \alpha \leq \kappa$.

LEMMA 3.6. \mathbb{P}_β is a complete sub-poset of \mathbb{P}_α for all $\beta \leq \alpha \leq \kappa$.

PROOF. Fix $\beta \leq \alpha \leq \kappa$. By definition, the ordering \leq_β on \mathbb{P}_β is equal to the ordering \leq_α on \mathbb{P}_α restricted to \mathbb{P}_β . So to prove the lemma, it suffices to show that every maximal antichain in \mathbb{P}_β is also a maximal antichain in \mathbb{P}_α . Suppose \mathcal{A} is a maximal antichain in \mathbb{P}_β , and let $p \in \mathbb{P}_\alpha$. By the maximality of \mathcal{A} , there is some $q \in \mathcal{A}$ such that q is compatible with $p \upharpoonright \beta$ in \mathbb{P}_β . Let $r \in \mathbb{P}_\beta$ be a common extension of q and $p \upharpoonright \beta$. Define a function s on $(I \times \omega) \cup (\alpha \times C \times \omega)$ by setting $s = r$ on $(I \times \omega) \cup (\beta \times C \times \omega)$ and setting $s = p$ on $(I \times \omega) \cup ((\alpha \setminus \beta) \times C \times \omega)$. Then $s \in \mathbb{P}_\alpha$, and s is a common extension of p and q in \mathbb{P}_α . As p was arbitrary, this shows \mathcal{A} is a maximal antichain in \mathbb{P}_α . \dashv

LEMMA 3.7. \mathbb{P}_κ has the ccc.

PROOF. Let \mathcal{A}_0 be an uncountable subset of \mathbb{P}_κ . By the Δ -system lemma, there is an uncountable $\mathcal{A}_1 \subseteq \mathcal{A}_0$ and a finite $R \subseteq (I \times \omega) \cup (\kappa \times C \times \omega)$ such that for any $p, q \in \mathcal{A}_1$, $\text{dom}(p) \cap \text{dom}(q) = R$. If $i \in R \cap (I \times \omega)$, then $p(i) \in \{0, 1\}$ for every $p \in \mathcal{A}_1$. As there are only finitely many functions $R \cap (I \times \omega) \rightarrow 2$, this implies there is some uncountable $\mathcal{A}_2 \subseteq \mathcal{A}_1$ such that $p \upharpoonright (R \cap (I \times \omega)) = q \upharpoonright (R \cap (I \times \omega))$ for any $p, q \in \mathcal{A}_2$. For every $p \in \mathcal{A}_2$, if $j \in R \setminus (I \times \omega)$, then $p(j) = (t, B)$ for some finite subtree t of $2^{<\omega}$. As there are only countably many finite subtrees of $2^{<\omega}$, there is some uncountable $\mathcal{A}_3 \subseteq \mathcal{A}_2$ such that for each $j \in R \setminus (I \times \omega)$, there is some fixed t_j such that for any $p \in \mathcal{A}_3$, $p(j) = (t_j, B_j^p)$ for some B_j^p . But any two members of \mathcal{A}_3 are compatible: if $p, q \in \mathcal{A}_3$, then

$$r(j) = \begin{cases} p(j), & \text{if } j \in \text{dom}(p) \setminus R, \\ q(j), & \text{if } j \in \text{dom}(q) \setminus R, \\ (t_j, B_j^p \cup B_j^q), & \text{if } j \in \text{dom}(p) \cap \text{dom}(q) = R \end{cases}$$

is a common extension of p and q . Thus \mathbb{P}_κ has no uncountable antichains, and is ccc. (In fact we have shown a bit more: \mathbb{P}_κ has property K .) \dashv

If $\beta \leq \alpha$, then because $\mathbb{P}_\beta \subseteq \mathbb{P}_\alpha$, we may (and do) consider every \mathbb{P}_β -name to be a \mathbb{P}_α -name as well. Let us also set the convention that in $V^{\mathbb{P}_\alpha}$, the evaluation of a \mathbb{P}_α -name is indicated by removing its dot. So, for example, c_i denotes in $V^{\mathbb{P}_\alpha}$ (for any $\alpha \leq \kappa$) the Cohen real added by \mathbb{P}_0 in coordinate i , and $T_{\mu,n}^\zeta$ denotes in $V^{\mathbb{P}_\alpha}$, for any $\alpha \geq \zeta$, the evaluation of the name $\dot{T}_{\mu,n}^\zeta$.

LEMMA 3.8. $C \subseteq \text{sp}(\text{closed})$ in $V^{\mathbb{P}_\kappa}$.

PROOF. Let $\mu \in C$. We claim that in $V^{\mathbb{P}_\kappa}$,

$$\mathcal{P} = \{\{c_i\} : i \in I_\mu\} \cup \{\bigcup_{n \in \omega} \llbracket T_{\mu,n}^\alpha \rrbracket : \alpha < \kappa\}$$

is a partition of 2^ω . Observe that every member of \mathcal{P} is an F_σ subset of 2^ω , and $|\mathcal{P}| = \mu$. (It is clear that each $T_{\mu,n}^\alpha$ is a subtree of $2^{<\omega}$, which means the sets of the form $\bigcup_{n \in \omega} \llbracket T_{\mu,n}^\alpha \rrbracket$ are all F_σ .) Thus if \mathcal{P} is a partition, then $\mu \in \text{sp}(\text{closed})$ by Theorem 2.4.

The members of \mathcal{P} come in two types, so in order to show they are pairwise disjoint, we have three things to prove.

First, if $i, i' \in I_\mu$ and $i \neq i'$, then clearly $c_i \neq c_{i'}$, i.e., $\{c_i\} \cap \{c_{i'}\} = \emptyset$.

Second, fix $i \in I_\mu$, $\alpha < \kappa$, and $n \in \omega$. We claim $\{c_i\} \cap \llbracket T_{\mu,n}^\alpha \rrbracket = \emptyset$, or equivalently, $c_i \notin \llbracket T_{\mu,n}^\alpha \rrbracket$. To see this, fix some $p \in \mathbb{P}_\kappa$: we will find an extension r of p forcing $c_i \notin \llbracket T_{\mu,n}^\alpha \rrbracket$. Extending p to $p \cup \{(\langle \alpha, \mu, n \rangle, (\emptyset, \emptyset))\}$, if necessary, we may (and do) assume $(\alpha, \mu, n) \in \text{dom}(p)$. Let $(t, B) = p(\alpha, \mu, n)$, with t a pruned subtree of $2^{<k}$. From the definition of \mathbb{P}_κ , we know $p \restriction \alpha \Vdash_{\mathbb{P}_\alpha} \dot{b} \neq \dot{c}_i$ for every $\dot{b} \in B$. Using this, and the fact that B is finite, there is an extension q of p in \mathbb{P}_α , and some finite sequence $\sigma \in 2^\ell$ for some $\ell > k$, such that $q \restriction \alpha \Vdash_{\mathbb{P}_\alpha} \dot{c}_i \restriction \ell = \sigma$ but $\dot{b} \restriction \ell \neq \sigma$ for all $\dot{b} \in B$. Let t' be the largest subtree of $2^{<\ell+1}$ that is an end extension of t and that does not contain σ . (In other words, $\tau \in t'$ if and only if $\tau \in 2^{<\ell+1} \setminus \{\sigma\}$ and $\tau \restriction k \in t$. This is a subtree of $2^{<\ell+1}$, and it is an end extension of t because $\ell > k$ and $\sigma \in 2^\ell$.) Define a function r on $(I \times \omega) \cup (\kappa \times C \times \omega)$ by setting

$$r(j) = \begin{cases} q(j), & \text{if } j \in (I \times \omega) \cup (\alpha \times C \times \omega), \\ p(j), & \text{if } j \in (I \times \omega) \cup ((\kappa \setminus \alpha) \times C \times \omega) \text{ but } j \neq (\alpha, \mu, n), \\ (t', B), & \text{if } j = (\alpha, \mu, n). \end{cases}$$

Then $r \in \mathbb{P}_\kappa$, and $r \Vdash \sigma \notin \dot{T}_{\mu,n}^\alpha$ and $\dot{c}_i \restriction \ell = \sigma$, which means $r \Vdash \dot{c}_i \notin \llbracket \dot{T}_{\mu,n}^\alpha \rrbracket$. As p was arbitrary, this shows that $c_i \notin \llbracket T_{\mu,n}^\alpha \rrbracket$ in $V^{\mathbb{P}_\kappa}$.

Third, fix $\beta < \alpha < \kappa$, and $m, n \in \omega$. We claim that $\llbracket T_{\mu,m}^\beta \rrbracket \cap \llbracket T_{\mu,n}^\alpha \rrbracket = \emptyset$. To see this, fix $p \in \mathbb{P}_\kappa$: we will find an extension of p forcing $\llbracket T_{\mu,m}^\beta \rrbracket \cap \llbracket T_{\mu,n}^\alpha \rrbracket = \emptyset$. Extending p to $p \cup \{(\langle \beta, \mu, m \rangle, (\emptyset, \emptyset)), \langle \alpha, \mu, n \rangle, (\emptyset, \emptyset)\}$, if necessary, we may (and do) assume $(\beta, \mu, m), (\alpha, \mu, n) \in \text{dom}(p)$. Let $(t_\beta, B_\beta) = p(\beta, \mu, m)$, where t_β is a pruned subtree of $2^{<k_\beta}$ for some $k_\beta \in \omega$, and let $(t_\alpha, B_\alpha) = p(\alpha, \mu, n)$, where t_α is a pruned subtree of $2^{<k_\alpha}$ for some $k_\alpha \in \omega$. From the definition of \mathbb{P}_κ , we know that $p \restriction ((I \times \omega) \cup (\alpha \times C \times \omega)) \Vdash_{\mathbb{P}_\alpha} \dot{b} \neq \dot{a}$ for every $\dot{b} \in B_\beta$ and $\dot{a} \in B_\alpha$. Using this, and the fact that B_β and B_α are both finite, there is an extension q of p in \mathbb{P}_α , and some $\ell > k_\beta, k_\alpha$, such

that q “decides” all the \dot{b} and \dot{a} up to ℓ , and in such a way that witnesses $\dot{b} \neq \dot{a}$ for all $\dot{b} \in B_\beta$ and $\dot{a} \in B_\alpha$. More precisely: for each $\dot{b} \in B_\beta$, there is a particular branch $c(\dot{b})$ of $2^{<\ell+1}$ such that $q \Vdash_{\mathbb{P}_\alpha} \text{“}\dot{b} \upharpoonright \ell = c(\dot{b})\text{”}$; and similarly, for each $\dot{a} \in B_\alpha$, there is a particular branch $c(\dot{a})$ of $2^{<\ell+1}$ such that $q \Vdash_{\mathbb{P}_\alpha} \text{“}\dot{a} \upharpoonright \ell = c(\dot{a})\text{”}$; and furthermore, $c(\dot{b}) \upharpoonright \ell \neq c(\dot{a}) \upharpoonright \ell$ for all $\dot{b} \in B_\beta$ and $\dot{a} \in B_\alpha$. Let t'_β be the largest end extension of t_β that is a pruned subtree of $2^{<\ell}$, and let t''_β be the end extension of t'_β to a pruned subtree of $2^{<\ell+1}$ containing on level ℓ the nodes:

- $\sigma \frown 0$ if $\sigma \in t'_\beta, \sigma = c(\dot{a})$ for some $\dot{a} \in B_\alpha$, and $c(\dot{a})(\ell) = 1$,
- $\sigma \frown 1$ if $\sigma \in t'_\beta, \sigma = c(\dot{a})$ for some $\dot{a} \in B_\alpha$, and $c(\dot{a})(\ell) = 0$,
- $\sigma \frown 0$ if $\sigma \in t'_\beta$ and $\sigma \neq c(\dot{a})$ for any $\dot{a} \in B_\alpha$.

Similarly, let t'_α be the largest end extension of t_α that is a pruned subtree of $2^{<\ell}$, and let t''_α be the end extension of t'_α to a pruned subtree of $2^{<\ell+1}$ containing on level ℓ the nodes:

- $\sigma \frown 0$ if $\sigma \in t'_\alpha, \sigma = c(\dot{b})$ for some $\dot{b} \in B_\beta$, and $c(\dot{b})(\ell) = 1$,
- $\sigma \frown 1$ if $\sigma \in t'_\alpha, \sigma = c(\dot{b})$ for some $\dot{b} \in B_\beta$, and $c(\dot{b})(\ell) = 0$,
- $\sigma \frown 1$ if $\sigma \in t'_\alpha$ and $\sigma \neq c(\dot{b})$ for any $\dot{b} \in B_\beta$.

Observe that t''_β and t''_α have no common nodes on level ℓ . Define a function r on $(I \times \omega) \cup (\kappa \times \mathcal{C} \times \omega)$ by setting

$$r(j) = \begin{cases} q(j), & \text{if } j \in (I \times \omega) \cup (\alpha \times \mathcal{C} \times \omega) \text{ but } j \neq (\beta, \mu, m), \\ p(j), & \text{if } j \in (I \times \omega) \cup ((\kappa \setminus \alpha) \times \mathcal{C} \times \omega) \text{ but } j \neq (\alpha, \mu, n), \\ (t''_\beta, B_\beta), & \text{if } j = (\beta, \mu, m), \\ (t''_\alpha, B_\alpha), & \text{if } j = (\alpha, \mu, n). \end{cases}$$

Then $r \in \mathbb{P}_\kappa$, and $r \Vdash t''_\beta$ is a subtree of $\dot{T}_{\mu,m}^\beta$ and $r \Vdash t''_\alpha$ is a subtree of $\dot{T}_{\mu,n}^\alpha$. Because t''_β and t''_α share no nodes on level ℓ , $r \Vdash \llbracket \dot{T}_{\mu,m}^\beta \rrbracket \cap \llbracket \dot{T}_{\mu,n}^\alpha \rrbracket = \emptyset$. As p was arbitrary, this shows that $\llbracket T_{\mu,m}^\beta \rrbracket \cap \llbracket T_{\mu,n}^\alpha \rrbracket = \emptyset$ in $V^{\mathbb{P}_\kappa}$.

Thus \mathcal{P} is a pairwise disjoint collection of F_σ subsets of 2^ω in $V^{\mathbb{P}_\kappa}$. To finish the proof, we must show also that $\bigcup \mathcal{P} = 2^\omega$.

Fix $x \in 2^\omega$ in $V^{\mathbb{P}_\kappa}$. Because \mathbb{P}_κ has the ccc, and each \mathbb{P}_α is a complete sub-poset of \mathbb{P}_κ , there is some $\alpha < \kappa$ such that $x \in V^{\mathbb{P}_\alpha}$. If $x = c_i$ for some $i \in I_\mu$, or if $x \in \llbracket T_{\mu,n}^\xi \rrbracket$ for some $\xi < \alpha$, then we’re done. If not, let \dot{x} be a nice \mathbb{P}_α -name for x , and fix some $p \in \mathbb{P}_\kappa$ such that $p \Vdash \text{“}\dot{x} \neq \dot{c}_i \text{ for all } i \in I_\mu, \text{ and } \dot{x} \notin \llbracket T_{\mu,n}^\xi \rrbracket \text{ for all } \xi < \alpha \text{ and } n \in \omega.\text{”}$ Because p has finite support, there is some $N \in \omega$ such that $(\alpha, \mu, N) \notin \text{dom}(p)$. But then $q = p \cup \{(\alpha, \mu, N), (\emptyset, \{\dot{x}\})\}$ is in \mathbb{P}_α , and $q \Vdash \dot{x} \in \llbracket T_{\mu,N}^\alpha \rrbracket$. Thus for every condition p forcing $x \notin \bigcup_{i \in I_\mu} \{c_i\} \cup \bigcup_{\xi < \alpha} \bigcup_{n \in \omega} \llbracket T_{\mu,n}^\xi \rrbracket$, there is an extension q of p forcing $x \in \bigcup_{n \in \omega} \llbracket T_{\mu,n}^\alpha \rrbracket$. Hence $x \in \bigcup_{i \in I_\mu} \{c_i\} \cup \bigcup_{\xi \leq \alpha} \bigcup_{n \in \omega} \llbracket T_{\mu,n}^\xi \rrbracket \subseteq \bigcup \mathcal{P}$. As x was arbitrary, $\bigcup \mathcal{P} = 2^\omega$ as claimed. \dashv

LEMMA 3.9. $c = \theta = \max(C)$ in $V^{\mathbb{P}_\kappa}$.

PROOF. A straightforward transfinite induction on α shows that $|\mathbb{P}_\alpha| = \theta$ for all $\alpha \leq \kappa$. (For the base case, clearly $|\mathbb{P}_0| = \theta$, and the limit case is also clear. For the successor case, suppose $|\mathbb{P}_\alpha| = \theta$. Because \mathbb{P}_α has the ccc, there are $\theta^{\aleph_0} = \theta$ nice \mathbb{P}_α -names for reals, and it follows that $|\mathbb{P}_{\alpha+1}| = \theta$.) In particular, $|\mathbb{P}_\kappa| = \theta$, and because \mathbb{P}_κ has the ccc, it follows that there are $\theta^{\aleph_0} = \theta$ nice \mathbb{P}_κ -names for reals. Hence $\mathfrak{c} \leq \theta$ in $V^{\mathbb{P}_\kappa}$.

On the other hand, it is obvious that $\mathfrak{c} \geq \theta$ in $V^{\mathbb{P}_\kappa}$, because of the Cohen reals added by \mathbb{P}_0 . ⊣

LEMMA 3.10. $\text{cov}(\mathcal{M}) = \mathfrak{d} = \mathfrak{u} = \kappa = \min(C)$ in $V^{\mathbb{P}_\kappa}$.

PROOF. By Lemma 3.8 and Corollary 2.6, $\kappa \geq \mathfrak{u} \geq \mathfrak{d} \geq \text{cov}(\mathcal{M})$ in $V^{\mathbb{P}_\kappa}$. So to prove the lemma, it suffices to show that 2^ω cannot be covered with $< \kappa$ meager sets in $V^{\mathbb{P}_\kappa}$.

Suppose $\lambda < \kappa$ and $\{C_\xi : \xi < \lambda\}$ is a collection of closed nowhere dense subsets of 2^ω in $V^{\mathbb{P}_\kappa}$. For each C_ξ , let A_ξ be a Borel code that evaluates to C_ξ . Because \mathbb{P}_κ has the ccc and each \mathbb{P}_α is a complete sub-poset of \mathbb{P}_κ , there is for each ξ some $\alpha < \kappa$ such that $A_\xi \in V^{\mathbb{P}_\alpha}$. Because κ is regular and $\lambda < \kappa$, there is some particular $\alpha < \kappa$ such that $A_\xi \in V^{\mathbb{P}_\alpha}$ for all $\xi < \lambda$. So to prove the lemma, it suffices to show that for each $\alpha < \kappa$ there is some $c \in V^{\mathbb{P}_\kappa}$ not contained in any closed nowhere dense subset of 2^ω coded in $V^{\mathbb{P}_\alpha}$ (i.e., we want c to be Cohen-generic over $V^{\mathbb{P}_\alpha}$).

Fix $\alpha < \kappa$, and define $c : \omega \rightarrow 2$ by setting

$$c(n) = 0 \quad \text{if and only if} \quad \langle 0 \rangle \in T_{\kappa,n}^\alpha.$$

We claim that c is Cohen-generic over $V^{\mathbb{P}_\alpha}$. To see this, let $U \subseteq 2^\omega$ be a dense open set whose Borel code is in $V^{\mathbb{P}_\alpha}$. In particular, $\tilde{U} = \{\tau \in 2^{<\omega} : \llbracket \tau \rrbracket \subseteq U\} \in V^{\mathbb{P}_\alpha}$ (where, abusing notation slightly, $\llbracket \tau \rrbracket$ denotes all those x in 2^ω with $x \upharpoonright \text{dom}(\tau) = \tau$). Now fix $p \in \mathbb{P}_\kappa$. We will find an extension r of p forcing that $c \in U$. Extending p if necessary, we may (and do) assume that there is some $N \in \mathbb{N}$ such that $(\alpha, \mu, n) \in \text{dom}(p)$ for all $n < N$, but $(\alpha, \mu, n) \notin \text{dom}(p)$ for all $n \geq N$. For all $n < N$, let $p(\alpha, \mu, n) = (t_n, B_n)$. Define $\sigma \in 2^N$ by setting

$$\sigma(n) = 0 \quad \text{if and only if} \quad \langle 0 \rangle \in t_n$$

for all $n < N$. Because U is dense in 2^ω , there is some $\tau \in \tilde{U}$ such that $\tau \upharpoonright N = \sigma$. Because $\tilde{U} \in V^{\mathbb{P}_\alpha}$, there is some $q \in \mathbb{P}_\alpha$ with $q \leq_{\mathbb{P}_\alpha} p \upharpoonright \alpha$ forcing that $\tau \in \tilde{U}$. Define $r \in \mathbb{P}_\kappa$ by

$$r(j) = \begin{cases} r(j), & \text{if } j \in (I \times \omega) \cup (\alpha \times C \times \omega), \\ (\langle \tau(n) \rangle, \emptyset), & \text{if } j = (\alpha, \mu, n) \text{ for some } n \in \text{dom}(\tau) \setminus \text{dom}(\sigma), \\ p(j), & \text{otherwise.} \end{cases}$$

Then $r \in \mathbb{P}_\kappa$, and r forces $c \in U$. As U was an arbitrary open dense subset of 2^ω coded in $V^{\mathbb{P}_\alpha}$, this shows c is Cohen-generic over $V^{\mathbb{P}_\alpha}$. ⊣

From the last two lemmas, it follows that $\text{sp}(\text{closed}) \subseteq [\kappa, \theta]$ in $V^{\mathbb{P}_\kappa}$, and if $\lambda > \theta$ then $\lambda \notin \text{sp}(\text{Borel})$. To finish proving the theorem, it remains only to show that if $\lambda \in [\kappa, \theta]$ and $\lambda \notin C$, then $\lambda \notin \text{sp}(\text{Borel})$. This is where the isomorphism-of-names argument comes in. This argument requires a rich supply of automorphisms of \mathbb{P}_κ , which we describe next.

If $\phi : I \rightarrow I$ is a permutation, then for every set x we define $\bar{\phi}(x)$ to be the set obtained from x by replacing every $i \in I$ in the transitive closure of x with $\phi(i)$. This is well defined, because if $i, j \in I$ then i is not in the transitive closure of j . (One may think of I as a set of urelements, ϕ as a permutation of them, and $\bar{\phi}$ as the automorphism of the universe induced by ϕ .) Alternatively, $\bar{\phi}$ is described via well-founded recursion by the relation

$$\bar{\phi}(x) = \{\bar{\phi}(y) : y \in x\}$$

if $x \notin I$, and $\bar{\phi} \upharpoonright I = \phi$. In particular, if $p \in \mathbb{P}_\kappa$, $\text{dom}(\bar{\phi}(p)) = \bar{\phi}(\text{dom}(p))$ and

$$\bar{\phi}(p)(j) = \begin{cases} p(i), & \text{if } i \in (I \times \omega) \cap \text{dom}(p) \text{ and } \bar{\phi}(i) = j, \\ (t, \{\bar{\phi}(\dot{b}) : \dot{b} \in B\}), & \text{if } j \in \text{dom}(p) \setminus (I \times \omega) \text{ and } p(j) = (t, B), \end{cases}$$

and this expression can be taken as a recursive definition of $\bar{\phi}$ on \mathbb{P}_κ , and the natural extension of $\bar{\phi}$ to the class of \mathbb{P}_κ -names.

LEMMA 3.11. *Let ϕ be a permutation of I such that $\phi \upharpoonright I_\mu$ is a permutation of I_μ for all $\mu \in C$. Then $\bar{\phi} \upharpoonright \mathbb{P}_\alpha$ is an automorphism of \mathbb{P}_α for all $\alpha \leq \kappa$.*

PROOF. First, note that for all $p, q \in \mathbb{P}_\alpha$, we have

$$\langle p, q \rangle \in \leq_{\mathbb{P}_\alpha} \quad \text{if and only if} \quad \bar{\phi}(\langle p, q \rangle) = \langle \bar{\phi}(p), \bar{\phi}(q) \rangle \in \bar{\phi}(\leq_{\mathbb{P}_\alpha}).$$

Together with the fact that $\bar{\phi}$ is invertible (with inverse $\bar{\phi}^{-1} = \overline{\phi^{-1}}$), this shows that the image of \mathbb{P}_α under $\bar{\phi}$ is a poset, and is naturally isomorphic to \mathbb{P}_α as witnessed by $\bar{\phi}$. But it is not immediately clear that the image of \mathbb{P}_α under $\bar{\phi}$ is equal to \mathbb{P}_α , which is of course required for our claim that $\bar{\phi} \upharpoonright \mathbb{P}_\alpha$ is an automorphism of \mathbb{P}_α . (In fact, one may show that this would not be true for $\alpha \geq 1$ if ϕ were not required to fix all the I_μ .) This is proved for $\bar{\phi}$ and its inverse, simultaneously, by induction on α .

The base case $\alpha = 0$ is clear. And for limit α , if $\bar{\phi} \upharpoonright \mathbb{P}_\xi$ is an automorphism of \mathbb{P}_ξ , for all $\xi < \alpha$, then $\bar{\phi} \upharpoonright \mathbb{P}_\alpha$ is an automorphism of \mathbb{P}_α because $\langle \mathbb{P}_\alpha, \leq_{\mathbb{P}_\alpha} \rangle$ is the direct limit of $\langle \mathbb{P}_\xi, \leq_{\mathbb{P}_\xi} : \xi < \alpha \rangle$. The same applies to $\bar{\phi}^{-1}$.

For the successor case, fix some $\beta < \kappa$ and suppose $\bar{\phi} \upharpoonright \mathbb{P}_\xi$ is an automorphism of \mathbb{P}_ξ for all $\xi \leq \beta$ (the inductive hypothesis). Let $\alpha = \beta + 1$, and fix $p \in \mathbb{P}_\alpha$. We claim that $\bar{\phi}(p) \in \mathbb{P}_\alpha$. Recall that

$$\bar{\phi}(p)(j) = \begin{cases} p(i), & \text{if } i \in (I \times \omega) \cap \text{dom}(p) \text{ and } \bar{\phi}(i) = j, \\ (t, \{\bar{\phi}(\dot{b}) : \dot{b} \in B\}), & \text{if } j \in \text{dom}(p) \setminus (I \times \omega) \text{ and } p(j) = (t, B). \end{cases}$$

It is clear that $\bar{\phi}(p)$ is a finite partial function on $(I \times \omega) \cup (\alpha \times C \times \omega)$, and that if $i \in (I \times \omega) \cap \text{dom}(\bar{\phi}(p))$, then $\bar{\phi}(p)(i) = p(\phi^{-1}(i)) \in \{0, 1\}$. It remains to check that the last bullet point in the definition of the conditions in \mathbb{P}_α is satisfied by $\bar{\phi}(p)$.

Suppose $(\gamma, \mu, n) \in \text{dom}(\bar{\phi}(p))$ for some $\gamma \leq \beta$, $\mu \in C$, and $n \in \omega$. This means $(\gamma, \mu, n) \in \text{dom}(p)$ as well: let us denote $p(\gamma, \mu, n) = (t, B)$. Then

$$\bar{\phi}(p)(\gamma, \mu, n) = (t, \{\bar{\phi}(\dot{b}) : \dot{b} \in B\}).$$

Because $p \in \mathbb{P}_\alpha$, t is a pruned subtree of $2^{<k}$ for some $k \in \omega$, and (by the inductive hypothesis, that $\bar{\phi} \upharpoonright \mathbb{P}_\gamma$ is an automorphism of \mathbb{P}_γ), $\{\bar{\phi}(\dot{b}) : \dot{b} \in B\}$ is a finite set of nice \mathbb{P}_γ -names for members of 2^ω . Furthermore, by applying the automorphism $\bar{\phi}$ of \mathbb{P}_γ to the displayed statement in the last bullet point in the definition of the conditions in \mathbb{P}_α , we get that for every $\dot{b} \in B$,

$$\begin{aligned} \bar{\phi}(p) \upharpoonright ((I \times \omega) \cup (\gamma \times C \times \omega)) \Vdash_{\mathbb{P}_\gamma} & \quad \bar{\phi}(\dot{b}) \upharpoonright j \in t \text{ for every } j < k, \\ & \quad \bar{\phi}(\dot{b}) \neq \bar{\phi}(\dot{c}_i) \text{ for every } i \in I_\mu, \text{ and} \\ & \quad \bar{\phi}(\dot{b}) \text{ is not a branch of } \bar{\phi}(\dot{T}_{\mu,m}^\xi) \\ & \quad \text{for any } \xi < \gamma \text{ and } m \in \omega. \end{aligned}$$

Considering the second of these three statements, note that $\bar{\phi}(\dot{c}_i) = \dot{c}_{\phi(i)}$ for every $i \in I$. Because $\phi \upharpoonright I_\mu$ is a permutation of I_μ , this means that the assertion “ $\bar{\phi}(\dot{b}) \neq \bar{\phi}(\dot{c}_i)$ for every $i \in I_\mu$ ” is equivalent to the assertion “ $\bar{\phi}(\dot{b}) \neq \dot{c}_i$ for every $i \in I_\mu$.” Thus

$$\bar{\phi}(p) \upharpoonright ((I \times \omega) \cup (\gamma \times C \times \omega)) \Vdash_{\mathbb{P}_\gamma} \quad \bar{\phi}(\dot{b}) \neq \dot{c}_i \text{ for every } i \in I_\mu.$$

Considering the third of these three statements, observe that

$$\begin{aligned} \bar{\phi}(\dot{T}_{\mu,n}^\xi) &= \{\bar{\phi}(\langle \sigma, p \rangle) : p(\xi, \mu, m) = (t, B) \text{ for some } B, \text{ and } \sigma \in t\} \\ &= \{\langle \sigma, \bar{\phi}(p) \rangle : p(\xi, \mu, m) = (t, B) \text{ for some } B, \text{ and } \sigma \in t\} \\ &= \{\langle \sigma, \bar{\phi}(p) \rangle : \bar{\phi}(p)(\xi, \mu, m) = (t, \bar{\phi}(B)) \text{ for some } B, \text{ and } \sigma \in t\} \\ &= \{\langle \sigma, \bar{\phi}(p) \rangle : \bar{\phi}(p)(\xi, \mu, m) = (t, B') \text{ for some } B', \text{ and } \sigma \in t\} \\ &= \{\langle \sigma, q \rangle : q(\xi, \mu, m) = (t, B') \text{ for some } B', \text{ and } \sigma \in t\} \\ &= \dot{T}_{\mu,n}^\xi. \end{aligned}$$

The third equality is true because for any condition p , $p(\xi, \mu, m) = (t, B)$ if and only if $\bar{\phi}(p)(\xi, \mu, m) = (t, \bar{\phi}(B))$. The fourth equality uses the inductive hypothesis, that $\bar{\phi} \upharpoonright \mathbb{P}_\gamma$ is an automorphism: $p(\xi, \mu, m) = (t, \bar{\phi}(B))$ for some B if and only if $p(\xi, \mu, m) = (t, B')$ for some B' , specifically for $B' = \bar{\phi}^{-1}(B)$. The fifth equality also uses the fact that $\bar{\phi} \upharpoonright \mathbb{P}_\gamma$ is an automorphism. Thus

$$\begin{aligned} \bar{\phi}(p) \upharpoonright ((I \times \omega) \cup (\gamma \times C \times \omega)) \Vdash_{\mathbb{P}_\gamma} & \quad \bar{\phi}(\dot{b}) \text{ is not a branch of } \dot{T}_{\mu,m}^\xi \\ & \quad \text{for any } \xi < \gamma \text{ and } m \in \omega. \end{aligned}$$

Putting these together, we obtain

$$\begin{aligned} \bar{\phi}(p) \upharpoonright ((I \times \omega) \cup (\gamma \times C \times \omega)) \Vdash_{\mathbb{P}_\gamma} & \quad \bar{\phi}(\dot{b}) \upharpoonright j \in t \text{ for every } j < k, \\ & \quad \bar{\phi}(\dot{b}) \neq \dot{c}_i \text{ for every } i \in I_\mu, \text{ and} \\ & \quad \bar{\phi}(\dot{b}) \text{ is not a branch of } \dot{T}_{\mu,m}^\xi \\ & \quad \text{for any } \xi < \beta \text{ and } m \in \omega. \end{aligned}$$

In other words, $\bar{\phi}(p)$ satisfies the final bullet point in the definition of the \mathbb{P}_α conditions. Hence $\bar{\phi}(p) \in \mathbb{P}_\alpha$, as claimed.

This shows that (the restriction of) $\bar{\phi}$ is an injective morphism from \mathbb{P}_α to \mathbb{P}_α . To get surjectivity, simply note that the same argument applies to $\bar{\phi}^{-1} = \overline{\phi^{-1}}$ as well. \dashv

From now on, we work in the ground model. Let λ be a cardinal such that $\lambda \in [\kappa, \theta]$ and $\lambda \notin C$. Suppose $\{\dot{B}_\alpha : \alpha < \lambda\}$ is a set of nice \mathbb{P}_κ -names for Borel codes for subsets of \mathbb{R} . (Any standard method of constructing Borel codes can be used for the proof, provided only that the codes are hereditarily countable sets. But for concreteness, let us take a Borel code to be a subset of ω .) We let $B_\alpha \subseteq \omega$ denote the evaluation of the name \dot{B}_α in $V^{\mathbb{P}_\kappa}$, and we let $\tilde{B}_\alpha \subseteq \mathbb{R}$ denote the interpretation of B_α .

We aim to show that $\{\tilde{B}_\alpha : \alpha < \lambda\}$ is not a partition of \mathbb{R} in $V^{\mathbb{P}_\kappa}$. To this end, suppose $q \in \mathbb{P}_\kappa$ and q forces each of the \tilde{B}_α is nonempty, and $\tilde{B}_\alpha \cap \tilde{B}_\beta = \emptyset$ whenever $\alpha \neq \beta$. We will show that q also forces $\bigcup_{\alpha < \lambda} \tilde{B}_\alpha \neq \mathbb{R}$.

Fix a cardinal $\nu \leq \lambda$ such that $\alpha^{\aleph_0} < \nu$ for all cardinals $\alpha < \nu$, and such that C contains no cardinals in the interval $[\nu, \lambda]$. If λ is neither singular nor the successor of a singular cardinal, then we may simply take $\lambda = \nu$. Otherwise, using the last two bullet points in our description of C , there is an infinite interval of cardinals below λ and disjoint from C , and we may take ν to be any successor-of-a-successor cardinal in this interval. In either case, $\alpha^{\aleph_0} < \nu$ for all cardinals $\alpha < \nu$ by the GCH.

Given $\alpha \leq \kappa$ and a condition $p \in \mathbb{P}_\alpha$, for each $\mu \in C$ define

$$\text{hdom}_\mu(p) = \text{TC}(p) \cap I_\mu,$$

where $\text{TC}(p)$ denotes the transitive closure of p . We think of $\text{hdom}_\mu(p)$ as the “hereditary domain” of p on I_μ : all those $i \in I_\mu$ that are used at any stage in building the condition p . A straightforward transfinite induction on α shows that $|\text{hdom}_\mu(p)| \leq \aleph_0$ for every $p \in \mathbb{P}_\alpha$ and every $\mu \in C$. (The base case and the limit case are clear. For the successor case, use the fact that \mathbb{P}_α has the ccc.) Similarly, for each \mathbb{P}_κ -name \dot{x} and each $\mu \in C$, let $\text{hdom}_\mu(\dot{x}) = \text{TC}(\dot{x}) \cap I_\mu$. As before, it is not difficult to see that if \dot{x} is a nice \mathbb{P}_κ -name for a Borel code (or for any hereditarily countable set), then $\text{hdom}_\mu(\dot{x})$ is countable for every $\mu \in C$.

For each $\alpha < \lambda$ and each $\mu \in C$, let $D_\mu^\alpha = \text{hdom}_\mu(\dot{B}_\alpha)$. Note that D_μ^α is countable for each $\alpha < \lambda$ and $\mu \in C$. Expanding some of the D_μ^α if necessary, we may (and do) assume each D_μ^α is countably infinite and includes $\text{hdom}_\mu(q)$. For each $\alpha < \lambda$, let $D^\alpha = \bigcup_{\mu \in C} D_\mu^\alpha$. We note that $|D^\alpha| = \sum_{\mu \in C} |D_\mu^\alpha| = |C| \cdot \aleph_0$ for all $\alpha < \lambda$, and $|C| \cdot \aleph_0 < \min(C) = \kappa$. (Note: the inequality $|C| \cdot \aleph_0 < \min(C)$ uses the second bullet point in our description of C .)

By our choice of ν , together with the aforementioned fact that $|D^\alpha| < \kappa$ for all $\alpha < \lambda$, $\{D^\alpha : \alpha < \nu\}$ meets the conditions of the generalized Δ -system lemma [15, Lemma III.6.15]. Thus there is some $\mathcal{A}_0 \subseteq \nu$ with $|\mathcal{A}_0| = \nu$ such that $\{D^\alpha : \alpha \in \mathcal{A}_0\}$ is a Δ -system with root R .

Let $\mathcal{A}_1 = \{\alpha \in \mathcal{A}_0 : D_\mu^\alpha \setminus R = \emptyset \text{ for all } \mu \in C \text{ with } \mu < \lambda\}$. For all $\mu \in C$, $\{D_\mu^\alpha \setminus R : \alpha \in \mathcal{A}_0\}$ is a ν -size collection of pairwise disjoint subsets of I_μ . If also $\mu < \lambda$, then $\mu < \nu$ and it follows that $D_\mu^\alpha \setminus R = \emptyset$ for all but (at most) μ members of \mathcal{A}_0 . Furthermore, $|\{\mu \in C : \mu < \lambda\}| \leq |C| < \kappa < \nu$. It follows that $|\mathcal{A}_1| = \nu$.

For each $\alpha < \lambda$ and $\mu \in C$, fix a bijection $\phi_\mu^\alpha : D_\mu^\alpha \rightarrow \omega$, in such a way that $\phi_\mu^\alpha \upharpoonright R = \phi_\mu^\beta \upharpoonright R$ for all $\alpha, \beta < \lambda$. For each $\alpha, \beta < \lambda$ and $\mu \in C$, let $\phi_\mu^{\alpha,\beta}$ be the involution of I_μ given by

$$\phi_\mu^{\alpha,\beta}(i) = \begin{cases} (\phi_\mu^\beta)^{-1} \circ \phi_\mu^\alpha(i), & \text{if } i \in D_\mu^\alpha, \\ (\phi_\mu^\alpha)^{-1} \circ \phi_\mu^\beta(i), & \text{if } i \in D_\mu^\beta, \\ i, & \text{otherwise.} \end{cases}$$

(Recall that an ‘‘involution’’ of a set S means a bijection $f : S \rightarrow S$ such that $\psi \circ \psi = \text{id}_S$.) This function is well defined, because $D_\mu^\alpha \cap D_\mu^\beta = R$, and $(\phi_\mu^\beta)^{-1} \circ \phi_\mu^\alpha(i) = i = (\phi_\mu^\alpha)^{-1} \circ \phi_\mu^\beta(i)$ whenever $i \in R$. This gives us a collection $\{\phi_\mu^{\alpha,\beta} : \alpha, \beta < \lambda\}$ of involutions of I_μ such that for any $\alpha, \beta < \lambda$,

- $\phi_\mu^{\alpha,\beta}$ maps D_μ^α onto D_μ^β and D_μ^β onto D_μ^α , but acts as the identity on the rest of I_μ ,
- $\phi_\mu^{\alpha,\beta}$ acts as the identity on $R \cap I_\mu$, and
- $\phi_\mu^{\alpha,\beta} = \phi_\mu^{\gamma,\beta} \circ \phi_\mu^{\alpha,\gamma}$ for any $\gamma < \lambda$.

For each $\alpha, \beta < \lambda$, let $\phi^{\alpha,\beta}$ denote the product map $\bigotimes_{\mu \in C} \phi_\mu^{\alpha,\beta}$ (that is, the map defined by setting $\phi^{\alpha,\beta}(i) = \phi_\mu^{\alpha,\beta}(i)$ whenever $i \in I_\mu$). This is an involution of I , and restricts to $\phi_\mu^{\alpha,\beta}$ on each I_μ . In particular, Lemma 3.11 applies, and $\bar{\phi}^{\alpha,\beta} \upharpoonright \mathbb{P}_\kappa$ is an automorphism of \mathbb{P}_κ for each $\alpha, \beta < \lambda$.

LEMMA 3.12. *There are at most κ nice \mathbb{P}_κ -names \dot{x} for subsets of ω with the property that $\text{hdom}(\dot{x}) \subseteq D^0$.*

PROOF. For each $\alpha \leq \kappa$, let $\mathbb{P}_\alpha \upharpoonright D^0 = \{p \in \mathbb{P}_\alpha : \text{hdom}(p) \subseteq D^0\}$. We prove by transfinite induction on α that for all $\alpha \leq \kappa$, $|\mathbb{P}_\alpha \upharpoonright D^0| \leq \kappa$ and there are $\leq \kappa$ nice $\mathbb{P}_\alpha \upharpoonright D^0$ -names for subsets of ω . Note that a nice $\mathbb{P}_\alpha \upharpoonright D^0$ -name for a subset of ω is the same thing as a nice \mathbb{P}_α -name \dot{x} for a subset of ω with the property that $\text{hdom}(\dot{x}) \subseteq D^0$.

For the base case, $\mathbb{P}_0 \upharpoonright D^0$ is just the set of finite partial functions $D^0 \rightarrow 2$, so $|\mathbb{P}_0 \upharpoonright D^0| = \aleph_0 \leq \kappa$ because D^0 is countable. Because $\mathbb{P}_0 \upharpoonright D^0$ has the ccc, there are \aleph_0 nice $\mathbb{P}_0 \upharpoonright D^0$ -names for subsets of ω .

For the other cases, fix $\alpha \leq \kappa$, and suppose $|\mathbb{P}_\xi \upharpoonright D^0| \leq \kappa$ and there are $\leq \kappa$ nice $\mathbb{P}_\xi \upharpoonright D^0$ -names for subsets of ω , for all $\xi < \alpha$. A condition in $\mathbb{P}_\alpha \upharpoonright D^0$ is defined in the same way as a condition in \mathbb{P}_α , except that we restrict the \dot{b} in that definition to members of $\mathbb{P}_\gamma \upharpoonright D^0$. Using the inductive hypothesis on $\mathbb{P}_\gamma \upharpoonright D^0$ -names, it follows that $|\mathbb{P}_\alpha \upharpoonright D^0| \leq \kappa$. Each nice $\mathbb{P}_\alpha \upharpoonright D^0$ -name \dot{x} for a subset of ω has the form $\dot{x} = \{\langle n, p \rangle : p \in \mathcal{A}_n\}$, where each \mathcal{A}_n is an antichain in $\mathbb{P}_\alpha \upharpoonright D^0$. Because \mathbb{P}_α has the ccc, each \mathcal{A}_n is countable. Thus there are $\kappa^{\aleph_0} = \kappa$ names \dot{x} with this form. \dashv

In particular, there are κ nice \mathbb{P}_κ -names \dot{x} for subsets of ω having the property that $\text{hdom}(\dot{x}) \subseteq D^0$. But for each $\alpha \in \mathcal{A}_1$, $\bar{\phi}^{\alpha,0}(\dot{B}_\alpha)$ is just such a name. By the pigeonhole principle, and the fact that $\kappa < \nu$, this implies that there is some $\mathcal{A}_2 \subseteq \mathcal{A}_1$ with $|\mathcal{A}_2| = \nu$ such that $\bar{\phi}^{\alpha,0}(\dot{B}_\alpha) = \bar{\phi}^{\beta,0}(\dot{B}_\beta)$ whenever $\alpha, \beta \in \mathcal{A}_2$. Reindexing the \dot{B}_α 's if necessary, we may (and do) assume $0 \in \mathcal{A}_2$.

We now proceed to define a new \mathbb{P}_κ -name \dot{B}_λ for a subset of ω .

First, for all $\mu \in C$ with $\mu < \lambda$, let $D_\mu^\lambda = R \cap I_\mu$. (Recall that $D_\mu^\alpha \cap I_\mu \subseteq R$ whenever $\mu \in C$ and $\mu < \lambda$ and $\alpha \in \mathcal{A}_1$. Thus, in this case, $D_\mu^\lambda \cap D_\mu^\alpha = R \cap I_\mu$ for all $\alpha \in \mathcal{A}_1 \supseteq \mathcal{A}_2$.) Next, for each $\mu \in C$ with $\mu > \lambda$, let D_μ^λ be a countable subset of I_μ with $D_\mu^\lambda \cap \bigcup \{D_\mu^\alpha : \alpha < \lambda\} = R \cap I_\mu$, and with $|D_\mu^\lambda \setminus R| = |D_\mu^0 \setminus R|$. Some such set exists because $|\bigcup \{D_\mu^\alpha : \alpha < \lambda\}| = \lambda < \mu = |I_\mu|$, so we may take D_μ^λ to be any subset (of the appropriate size) of $I_\mu \setminus \bigcup \{D_\mu^\alpha : \alpha < \lambda\}$, together with $R \cap I_\mu$.

For each $\mu \in C$, fix a bijection $\phi_\mu^\lambda : D_\mu^\lambda \rightarrow \omega$ such that $\phi_\mu^\lambda \upharpoonright R = \phi_\mu^0 \upharpoonright R$. For each $\alpha < \lambda$ and $\mu \in C$, let $\phi_\mu^{\alpha,\lambda}$ and $\phi_\mu^{\lambda,\alpha}$ be the involutions of I_μ defined just like the $\phi_\mu^{\alpha,\beta}$ above, but using ϕ_μ^λ in place of ϕ_μ^β . This naturally extends the system of involutions described above: for any $\alpha, \beta \leq \lambda$,

- $\phi_\mu^{\alpha,\beta}$ maps D_μ^α onto D_μ^β and D_μ^β onto D_μ^α , but acts as the identity on the rest of I_μ ,
- each $\phi_\mu^{\alpha,\beta}$ acts as the identity on $R \cap I_\mu$, and
- $\phi_\mu^{\alpha,\beta} = \phi_\mu^{\gamma,\beta} \circ \phi_\mu^{\alpha,\gamma}$ for any $\gamma \leq \lambda$.

For each $\alpha, \beta \leq \lambda$, let $\phi^{\alpha,\beta} = \bigotimes_{\mu \in C} \phi_\mu^{\alpha,\beta}$. By Lemma 3.11, $\bar{\phi}^{\alpha,\beta} \upharpoonright \mathbb{P}_\kappa$ is an automorphism of \mathbb{P}_κ for each $\alpha, \beta \leq \lambda$.

Because $\text{hdom}_\mu(q) \subseteq R$ for all μ and $\phi^{0,\lambda}$ fixes R , $\bar{\phi}^{0,\lambda}(q) = q$. Recall

$q \Vdash$ the evaluation B_0 of \dot{B}_0 codes a nonempty Borel subset of \mathbb{R} .

Applying standard facts about automorphisms of forcing posets,

$\bar{\phi}^{0,\lambda}(q) \Vdash$ the evaluation of $\bar{\phi}^{0,\lambda}(\dot{B}_\lambda)$ codes a nonempty Borel subset of \mathbb{R} .

Let B_λ denote the evaluation of \dot{B}_λ in $V^{\mathbb{P}_\kappa}$, and let \tilde{B}_λ be the Borel set that it codes. So, in particular, the displayed statement above implies that $q \Vdash \bar{\phi}^{0,\lambda}(q) \Vdash \tilde{B}_\lambda \neq \emptyset$. To show $q \Vdash \bigcup_{\alpha < \lambda} \tilde{B}_\alpha \neq \mathbb{R}$, it now suffices to show $q \Vdash \tilde{B}_\lambda \cap \tilde{B}_\beta = \emptyset$ for all $\beta < \lambda$.

Fix $\beta < \lambda$. By our choice of the sets D_μ^λ , we have $D_\mu^\lambda \cap D_\mu^\beta \subseteq R$ for all $\mu \in C$, and therefore $D^\lambda \cap D^\beta \subseteq R$. Because $\{D^\alpha : \alpha \in \mathcal{A}_2\}$ is a Δ -system of size v , and because $|D^\beta| \leq |C| \cdot \aleph_0 < \kappa < v$, there is some $\alpha \in \mathcal{A}_2$ with $D^\alpha \cap D^\beta \subseteq R$ (regardless of whether $\beta \in \mathcal{A}_2$). Fix some such α .

Note that $\phi^{\alpha,\lambda}$ sends D^α to D^λ , while acting as the identity on D^β . And because $0, \alpha \in \mathcal{A}_2$, our choice of \mathcal{A}_2 implies $\bar{\phi}^{0,\alpha}(\dot{B}_0) = \dot{B}_\alpha$. It follows that

$$\bar{\phi}^{\alpha,\lambda}(\dot{B}_\alpha) = \bar{\phi}^{\alpha,\lambda} \circ \bar{\phi}^{0,\alpha}(\dot{B}_0) = \bar{\phi}^{0,\lambda}(\dot{B}_0) = \dot{B}_\lambda,$$

while $\bar{\phi}^{\alpha,\lambda}(\dot{B}_\beta) = \dot{B}_\beta$. But

$q \Vdash B_\alpha$ and B_β code disjoint Borel sets.

Applying the automorphism $\bar{\phi}^{\alpha,\lambda}$ of \mathbb{P}_κ ,

$$\bar{\phi}^{\alpha,\lambda}(q) \Vdash B_\lambda \text{ and } B_\beta \text{ code disjoint Borel sets.}$$

Because β was arbitrary, this shows that $q \Vdash \tilde{B}_\lambda \cap \tilde{B}_\beta = \emptyset$ for all $\beta < \lambda$. In other words, $q \Vdash \bigcup_{\alpha < \lambda} \tilde{B}_\alpha \neq \mathbb{R}$. ⊢

Note that Theorem 3.2 only gives us models in which \acute{n} is regular and $|\text{sp}(\text{closed})| \leq \acute{n}$. Both of these are merely artifacts of the proof: neither need be true of $\text{sp}(\text{closed})$

in general. As mentioned near the end of Section 2, it is possible that $\mathfrak{d} = \mathfrak{c} = \kappa$ for some singular cardinal κ of uncountable cofinality, and this makes $\mathfrak{u} = \kappa$ also. And of course, Theorem 3.1 shows it is possible to have $|\mathfrak{sp}(\text{closed})| > \mathfrak{u}$.

Theorem 3.2 also cannot produce a model in which $\mathfrak{sp}(\text{closed}) \cap \kappa$ is unbounded for a regular limit cardinal κ . Theorem 3.1 implies that this is possible, but in such a case automatically gives $\kappa \in \mathfrak{sp}(\text{closed})$. This suggests the following question:

QUESTION 3.13. *Is $\mathfrak{sp}(\text{closed})$ or $\mathfrak{sp}(\text{Borel})$ closed under regular limits?*

We also do not know whether our final condition on C is an artifact of the proof.

QUESTION 3.14. *Is it consistent that $\aleph_\omega \in \mathfrak{sp}(\text{closed})$ but $\aleph_{\omega+1} \notin \mathfrak{sp}(\text{closed})$? What about $\mathfrak{sp}(\text{Borel})$?*

For any pointclass Γ , define

$$\mathfrak{sp}(\Gamma) = \{|\mathcal{P}| > \aleph_0 : \mathcal{P} \text{ is a partition of } \mathbb{R} \text{ into sets in } \Gamma\}.$$

To strengthen Theorem 3.2, we could replace $\mathfrak{sp}(\text{Borel})$ in the conclusion of the theorem with $\mathfrak{sp}(\Gamma)$ for some pointclass Γ strictly containing the Borel sets. The ultimate result in this direction would be to take $\Gamma = \text{OD}(\mathbb{R})$, since this is essentially the largest pointclass of interest from a descriptive point of view.

While we have chosen to focus on $\mathfrak{sp}(\text{Borel})$ and $\mathfrak{sp}(\text{closed})$ until now, let us observe that minor modifications to the proof of Theorem 3.2 will give a proof of the stronger version, where $\mathfrak{sp}(\text{Borel})$ is replaced by $\mathfrak{sp}(\text{OD}(\mathbb{R}))$ in the conclusion. Consequently, Corollaries 3.4 and 3.5 can also be strengthened by replacing $\mathfrak{sp}(\text{Borel})$ with $\mathfrak{sp}(\text{OD}(\mathbb{R}))$. This has the interesting consequence that large cardinals do not imply the existence of a partition of \mathbb{R} into \aleph_2 projective sets. This consequence is already known, by the work of Blass [2] mentioned at the start of this section.

We close with three more open questions.

QUESTION 3.15. *Is $\mathfrak{sp}(\text{Borel}) = \mathfrak{sp}(\text{OD}(\mathbb{R}))$?*

QUESTION 3.16. *Given $\alpha < \omega_1$, is it consistent that $\mathfrak{sp}(\Pi_\alpha^0) \neq \mathfrak{sp}(\Pi_{\alpha+1}^0)$?*

The answer to this question is currently known for $\alpha = 1, 2$. For $\alpha = 1$, Miller proved the consistency of $\mathfrak{sp}(\Pi_1^0) \neq \mathfrak{sp}(\Pi_2^0)$ (i.e., $\mathfrak{sp}(\text{closed}) \neq \mathfrak{sp}(G_\delta)$) by constructing a model with $\aleph_1 = \text{cov}(\mathcal{M}) < \mathfrak{u}$ [17, Theorem 5]. This shows $\mathfrak{sp}(\text{closed}) \neq \mathfrak{sp}(G_\delta)$, because if $\aleph_1 = \text{cov}(\mathcal{M})$, then $\aleph_1 \in \mathfrak{sp}(G_\delta)$. (Proof: if $\{F_\alpha : \alpha < \omega_1\}$ is a collection of closed nowhere dense sets covering \mathbb{R} , then $\{F_\alpha \setminus \bigcup_{\xi < \alpha} F_\xi : \alpha < \omega_1\}$ is a partition of \mathbb{R} into G_δ sets.) For $\alpha = 2$, the consistency of $\mathfrak{sp}(\Pi_2^0) \neq \mathfrak{sp}(\Pi_3^0)$ (i.e., $\mathfrak{sp}(G_\delta) \neq \mathfrak{sp}(F_{\text{od}})$) follows from Hausdorff's theorem that $\aleph_1 \in \mathfrak{sp}(\Pi_3^0)$, together with a result of Fremlin and Shelah [9] stating that $\min(\mathfrak{sp}(\Pi_2^0)) \geq \text{cov}(\mathcal{M})$.

QUESTION 3.17. *Is it consistent that $\mathfrak{sp}(\text{closed}) \neq \mathfrak{sp}(G_\delta)$ and $\text{cov}(\mathcal{M}) > \aleph_1$?*

REFERENCES

[1] T. BANAKH, *Are the Sierpiński cardinal \mathfrak{u} and its measure modification \mathfrak{m} equal to some known small uncountable cardinals?* MathOverflow question 285780 (2017). Available at <https://mathoverflow.net/questions/285780>.
 [2] A. R. BLASS, *Simple cardinal characteristics of the continuum*, *Set Theory of the Reals* (H. Judah, editor), Israel Mathematical Conferences Proceedings, vol. 6, American Mathematical Society, Providence, 1993, pp. 63–90.

- [3] J. BRENDLE, *Mad families and iteration theory*, **Logic and Algebra** (Y. Zhang, editor), Contemporary Mathematics, vol. 302, American Mathematical Society, Providence, 2002, pp. 1–31.
- [4] ———, *The almost-disjointness number may have countable cofinality*. **Transactions of the American Mathematical Society**, vol. 355 (2003), no. 7, pp. 2633–2649.
- [5] W. BRIAN and A. W. MILLER, *Partitions of 2^{ω} and completely ultrametrizable spaces*. **Topology and Its Applications**, vol. 184 (2015), pp. 61–71.
- [6] V. FISCHER, *Maximal cofinitary groups revisited*. **Mathematical Logic Quarterly**, vol. 61 (2015), pp. 367–379.
- [7] V. FISCHER and S. SHELAH, *The spectrum of independence*. **Archive for Mathematical Logic**, vol. 58 (2019), pp. 877–884.
- [8] ———, *The spectrum of independence II*. **Annals of Pure and Applied Logic**, vol. 173 (2022), no. 9, p. 103161.
- [9] D. FREMLIN and S. SHELAH, *On partitions of the real line*. **Israel Journal of Mathematics**, vol. 32 (1979), pp. 299–304.
- [10] F. HAUSDORFF, *Summen von \aleph_1 Mengen*. **Fundamenta Mathematicae**, vol. 26 (1936), no. 1, pp. 241–255.
- [11] S. S. HECHLER, *Short complete nested sequences in $\beta\mathbb{N} \setminus \mathbb{N}$ and small maximal almost-disjoint families*. **General Topology and Its Applications**, vol. 2 (1972), pp. 139–149.
- [12] M. HRUŠÁK, *Another \diamond -like principle*. **Fundamenta Mathematicae**, vol. 167 (2001), no. 3, pp. 277–289.
- [13] ———, *Life in the Sacks model*. **Acta Universitatis Carolinae, Mathematica et Physica**, vol. 42 (2001), no. 2, pp. 43–58.
- [14] A. KECHRIS, **Classical Descriptive Set Theory**, Graduate Texts in Mathematics, vol. 156, Springer, New York, 1995.
- [15] K. KUNEN, **Set Theory: An Introduction to Independence Proofs**, Studies in Logic and the Foundations of Mathematics, vol. 102, Elsevier, Amsterdam, 1980.
- [16] N. LUSIN and W. SIERPIŃSKI, *Sur un ensemble non mesurable B* . **Journal de Mathématiques Pures et Appliquées**, vol. 2 (1923), pp. 53–72.
- [17] A. W. MILLER, *Covering 2^{ω} with ω_1 disjoint closed sets*. **The Kleene Symposium** (J. Barwise, H. J. Keisler, and K. Kunen, editors), North-Holland, Amsterdam, 1980, pp. 415–421.
- [18] ———, *Infinite combinatorics and definability*. **Annals of Pure and Applied Logic**, vol. 41 (1989), pp. 179–203.
- [19] L. NEWELSKI, *On partitions of the real line into compact sets*, this JOURNAL, vol. 52 (1987), pp. 353–359.
- [20] S. SHELAH, *Two cardinal invariants of the continuum ($d < a$) and FS linearly ordered iterated forcing*. **Acta Mathematica**, vol. 192 (2004), pp. 187–223.
- [21] ———, *The character spectrum of $\beta(\mathbb{N})$* . **Topology and Its Applications**, vol. 158 (2011), pp. 2535–2555.
- [22] S. SHELAH and O. SPINAS, *MAD spectra*, this Journal, vol. 80 (2015), no. 3, pp. 901–916.
- [23] W. SIERPIŃSKI, *Un théorème sur les continus*. **Tôhoku Mathematical Journal**, vol. 13 (1918), pp. 300–303.
- [24] O. SPINAS, *Partition numbers*. **Annals of Pure and Applied Logic**, vol. 90 (1997), pp. 243–262.
- [25] J. STERN, *Partitions of the real line into F_{σ} , or G_{δ} subsets*. **Comptes Rendus de l'Académie des Sciences. Paris. Série A**, vol. 284 (1977), 921–922.

DEPARTMENT OF MATHEMATICS AND STATISTICS
 UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE
 9201 UNIVERSITY CITY BOULEVARD
 CHARLOTTE, NC 28223, USA

E-mail: wbrian.math@gmail.com

URL: wrbrian.wordpress.com