

Review

Community-based fish sanctuaries: untapped potential for freshwater fish conservation

SUMAN JUMANI, VANESSA HULL, PARINEETA DANDEKAR and NEETHI MAHESH

Abstract Riverine systems and associated fish populations worldwide are threatened by human impacts, especially in tropical countries with emerging economies. In India, community-based fish sanctuaries are a key mechanism for the conservation of freshwater fish populations, but there are few peer-reviewed studies on this subject. Here we integrate over 35 combined years of field experience with a literature synthesis to define and classify community-based fish sanctuaries. We present a novel, critical analysis of fish sanctuaries as social–ecological systems with a functional characterization based on natural capital, ecosystem services, human well-being, and policy and governance. We find that such sanctuaries are shaped by complex social–ecological processes, including coevolution of religious practices and ecological change, feedback processes created by retaliatory conflicts between river users, and diverse and dynamic governance strategies. These sanctuaries hold great potential for the conservation of rare fish species in India, but are subject to myriad threats at local, regional and global scales. Given the complexity of these social–ecological systems, we outline their conservation potential and highlight directions for future research.

Keywords Community-based conservation, fish conservation zone, participatory management, religion, social–ecological systems, temple sanctuary

Introduction

Industrial and agricultural expansion, rising energy consumption and growing human populations exert severe pressure on freshwater resources (Nilsson, 2005; Grill et al., 2019). Globally, rivers are amongst the most altered ecosystems, affected by > 125,000 large and small dams

(Zarfl et al., 2015; Couto & Olden, 2018), pollution, destruction of riparian habitats, sand mining and destructive fishing practices (Dudgeon, 2002). Freshwater diversity is thus amongst the most threatened in the world, with freshwater fish being particularly vulnerable (Dudgeon, 2002; Dudgeon et al., 2006).

Threats are most pronounced in tropical countries with emerging economies, where dam building overlaps with high freshwater biodiversity (Pandit & Grumbine, 2012; Tockner et al., 2016). In such regions, subsistence communities are directly dependent on the services that aquatic ecosystems provide (Beck et al., 2012). For example, in India, > 10.8 million people depend on inland fisheries for their livelihoods (Planning Commission, 2012). India also ranks third in the world in terms of the numbers of large dams (ICOLD, 2019), with > 6,000 large and small dams, in addition to severe pollution, degradation of riparian vegetation, unsustainable sand mining and harmful fish harvesting practices (Grant et al., 2012; Grumbine & Pandit, 2013; Jumanı et al., 2017; Krishnaswamy et al., 2017).

The Western Ghats mountain range, extending along the west coast of peninsular India, is especially vulnerable. As the most densely populated global biodiversity hotspot (Myers et al., 2000), its catchment provides water for > 400 million people (Molur et al., 2011). This freshwater ecoregion also has high fish species richness and endemism (Abell et al., 2008). Of the 320 fish species recorded in the Western Ghats, 66% are endemic, of which approximately half are threatened or near threatened (Dahanukar & Raghavan, 2013). Millions of people rely on these rivers for their livelihoods and sustenance, with c. 56% of fish species in the region being harvested for consumption (Molur et al., 2011). River-dependent communities use traditional fishing practices and artisanal fishing gear, often made using locally sourced materials (Ramesan, 2006; Kharat et al., 2013; Shaji & Laladhas, 2013). Some communities also associate fishing with important festivals and life events such as marriage or death (Prajith et al., 2016).

Despite their ecological, cultural and socio-economic significance, these rivers continue to be heavily dammed and diverted (Krishnaswamy et al., 2017). Such projects, coupled with pollution, unsustainable fishing and sand-harvesting practices, have severely damaged several river reaches and imperilled the biodiversity that they support (Grant et al., 2012; Atkore et al., 2017; Jumanı et al., 2018). Although > 150 species are threatened in the Western Ghats, existing laws

SUMAN JUMANI (Corresponding author, orcid.org/0000-0002-2292-7996, sumanjumani@ufl.edu) Soil and Water Sciences Department, University of Florida, Gainesville, FL 32611, USA

VANESSA HULL (orcid.org/0000-0002-7530-5672) Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, USA

PARINEETA DANDEKAR South Asia Network for Dams, Rivers and People, Delhi, India

NEETHI MAHESH Wildlife Conservation Society – India, Bangalore, India

Received 11 August 2021. Revision requested 28 October 2021.

Accepted 22 April 2022. First published online 2 September 2022.

This is an Open Access article, distributed under the terms of the Creative Commons-Attribution-ShareAlike licence (<https://creativecommons.org/licenses/by-sa/4.0/>), which permits re-use, distribution, reproduction, transformation, and adaptation in any medium and for any purpose, provided the original work is properly cited.

do not protect freshwater fish species under the Schedules of the Wild Life (Protection) Act, 1972 (Molur et al., 2011). Although fish are included in the definition of ‘wildlife’ under the Act, they are excluded from the definition of ‘wild animals’ (Pinder & Raghavan, 2013). Although the Indian Fisheries Act of 1879 prohibits unsustainable fishing methods, these rules are rarely enforced (Molur et al., 2011). No-fish zones declared under various State Acts are also not well recognized (Vyas et al., 2012), and Forest Department management plans rarely focus on freshwater biodiversity. Almost all of the 981 protected areas in India are set up and managed to conserve terrestrial habitats and offer only incidental protection to rivers and riverine biodiversity within their boundaries (Vasudevan et al., 2006; Abraham & Kelkar, 2012).

One of the few existing models of in situ freshwater fish conservation in India involves community-based fish sanctuaries (Dandekar, 2013). Here we define a community-based fish sanctuary (henceforth fish sanctuary) as a specific waterbody, its associated fish resources and other biotic and abiotic habitat elements that are managed by one or more local communities through formal or informal governance mechanisms. These sanctuaries have existed across India for centuries and usually comprise a river reach (ranging from 50 m to 20 km) or pond protected by local communities from destructive activities such as fishing, sand mining, water abstraction and removal of riparian vegetation.

Although sacred groves (terrestrial community-based protected areas) and sacred waters (rivers of religious significance) have been well studied (Gokhale et al., 1998; Bhagwat & Rutte, 2006), there is little mention of fish sanctuaries in the scientific literature (except Gupta et al., 2016). Large gaps remain in our understanding of the implementation of fish sanctuaries, particularly in terms of critically examining them from a systems perspective.

We conducted a review of the published, peer-reviewed literature compiled from thematic searches on Google Scholar using the keywords ‘fish sanctuary’, ‘community-based conservation’, ‘fish reserves’ and ‘fish god’. We excluded papers pertaining to marine and terrestrial protected areas and those that did not involve community participation. This resulted in a final selection of 19 publications, of which only four pertained to fish sanctuaries in India. Additionally, we also reviewed the grey literature, comprising reports, governmental websites, blogs and news articles. Based on this literature review and primary information gathered over 35 years of combined field experience, we propose a classification of fish sanctuaries and put forth a novel conceptualization of fish sanctuaries as dynamic social-ecological systems. We also synthesize the threats to these systems across scales, their potential role in freshwater fish conservation and future research directions.

Types of fish sanctuaries

Based on the motivation behind their establishment and management, we identify three types of fish sanctuaries in India (Fig. 1, Table 1).

(1) *Temple-based fish sanctuaries* (Plate 1) These are protected river reaches or pools situated adjacent to temples or shrines. Although their numbers are not well established, we have observed > 30 temple-based sanctuaries across six states (Kerala, Karnataka, Maharashtra, Madhya Pradesh, Uttarakhand and Himachal Pradesh), and older records document up to 27 sacred fish ponds in a single district (Simoons, 1974). Temple-based sanctuaries are usually established to worship one or more fish god(s). Fish are considered sacred by various Hindu sects, particularly the *Vaishnavite* (worshippers of Lord Vishnu) and *Saivite* (worshippers of Lord Shiva) groups (Simoons, 1974). The first incarnation of Vishnu is believed to have been a fish who saved humanity from a destructive flood and is worshipped as the *Matsya* or Fish God. The *Vedas*, an ancient Hindu epic, mentions the mahseer fish being used by devotees to appease the souls of their deceased ancestors (Nautiyal, 1989). The Indus script also used the fish as a symbol to represent five important deities (Simoons, 1974). Some of the earliest records of fish sanctuaries date back to c. 295 BCE, whereas more recent records are from British naturalists of colonial India (Simoons, 1974). The iconic mahseer group (species belonging to the genera *Tor*, *Neolissochilus* and *Naziritor*; Eschmeyer & Fricke, 2010; Froese & Pauly, 2010) is revered in particular. These fish have high socio-cultural and religious significance throughout South and Southeast Asia, with references to them existing in various scriptures and sculptures (Nautiyal, 2014; Pinder et al., 2019). In these sanctuaries, fish are protected from hunting and fed by priests and devotees (Table 1), and thus tend to grow large in body size and high in abundance.

(2) *Conservation-based fish reserves* (Plate 2) These reserves are created and managed by local communities to protect freshwater resources and the services that they provide. They may receive governmental or NGO support and are similar to the Freshwater Conservation Reserves found across Southeast Asia (Koning et al., 2020). These fish reserves are gaining popularity in parts of north-east India, where fish diversity is particularly high and local communities heavily rely on fish for sustenance (Dash et al., 2020). For example, during 2012–2018, 54 fish sanctuaries were established in the state of Meghalaya alone (Dash et al., 2020). Fish are not explicitly revered in these reserves, and there may or may not be consistent food provisioning. Some communities offer tourism or catch-and-release angling in these




FIG. 1 Framework to characterize community-based fish sanctuaries in India as a functional social–ecological system (Table 2).

reserves to generate revenue (Baruah & Sarma, 2018; Dash et al., 2020).

(3) *Informal fish sanctuaries* (Plate 3) These are uncommon and documented poorly. They are established through varying levels of protection afforded by one or more individual(s) or landowner(s), usually from a small household, river resort or tourism homestay located near a river. Such protected river reaches are created informally, without a clear intent to form a fish sanctuary. However, the protection they offer against exploitation and habitat destruction, along with food provisioning, leads to congregations of large-sized fish. Hence, they function as protected spaces. Often these are close to homestays or small-scale resorts, where fish congregations serve as tourist attractions.

Fish sanctuaries as social–ecological systems

We conceptualize fish sanctuaries using the social–ecological systems framework, which allows for the integration of social and ecological factors (Fig. 1; Ostrom, 2009). This framework classifies the social–ecological system into four subsystems (resource systems, resource units, governance

system and users) that interact with each other and are set within a larger ecological, social and political setting (Ostrom, 2009; McGinnis & Ostrom, 2014). For fish sanctuaries, we define the resource system as the extent of the sanctuary. The resource units, governance system and user subsystem vary across sanctuary types (Table 2). Additionally, we incorporate a functional characterization of the social–ecological system based on natural capital, ecosystem services and human well-being, which further influence the policy and governance of these systems (adapted from Resilience Alliance, 2010; Kalaba, 2014).

For temple-based fish sanctuaries, the resource units are the revered fish, often belonging to the mahseer group (Nautiyal, 2014). Users include devotees, local communities and tourists; the governance system usually comprises a priest or a committee representing the temple institution. These fish sanctuaries are governed by traditional practices rooted in religious or spiritual beliefs and are further safeguarded by social customs. Devotees are encouraged to feed temple fish with puffed rice or biscuits. These practices have become ritualized as important religious activities, yet potentially they are contentious with conservation goals as these items are not considered by scientists as suitable food sources for wild fish. Scientific knowledge is usually not considered in the governance of fish sanctuaries, and

TABLE 1 Representative examples of community-based fish sanctuaries in India, including sanctuary type, priority fish species protected and key management strategies employed.

Sanctuary type	Fish sanctuary	Reference	State	Year established	River	Priority fish species	Key management strategy
Temple-based	Shishileshwara fish sanctuary	Akshatha (2011)	Karnataka	c. 1300	Netravathi	<i>Tor</i> spp.	Food provisioning, ban on fishing & sand mining
	Walan Kond	Dandekar (2013)	Maharashtra	Unknown	Savitri	<i>Tor khudree</i>	
	Laxman Jhula	P. Dandekar, pers. obs., 2012	Uttarakhand	c. 1927	Ganga	<i>Tor putitora</i>	
Conservation-based	Amlayee River fish sanctuary	Meghalaya State Aquaculture Mission (2014)	Meghalaya	c. 1970	Amlayee	<i>Neolissochilus hexagonolepis</i>	Ban on fishing & sand mining, monitoring, enforcement of fines for offenders
	Khengiang–Yangoulen	Pinto et al. (2021)	Manipur	2020	Tuivang	NA ¹	
	Songkal Wari	Dandekar (2018)	Meghalaya	2003	Simsang	<i>N. hexagonolepis</i>	
Informal	Sitanadi Jungle Camp River of Joy homestay	S. Jumani, pers. obs., 2020	Karnataka	Unknown	Sitanadi, Kapila	NA ¹	Passive protection against fishing, food provisioning

¹NA: information not available.

religious beliefs can contradict scientific facts. For example, priests in the Shishileshwara fish sanctuary assign religious attributes to the fish and believe that mahseer are vegetarian despite their role as top predators in river systems.

Conservation-based fish reserves are examples of local communities self-organizing to manage a common natural resource for sustainable use. Here, the resource units comprise the protected river reach, including the protected fish, water, aquatic biota and adjoining riparian forests. Users include members of the local community, tourists and sometimes anglers (Dash et al., 2020). A locally elected committee comprising representatives from one or more village(s) usually forms the governance system, which may or may not include members from NGOs or local government bodies (Dash et al., 2020). These sanctuaries are primarily managed using the traditional knowledge of local communities, although scientific practices can be incorporated in the management of these sanctuaries when partnering with scientists, NGOs or governmental bodies, as in the case of the Lapalang fish conservation zone in Meghalaya (Pinto et al., 2021). As the fish are not revered, social compliance is achieved through participatory decision-making, penalties for the violation of rules and regular monitoring (Dash et al., 2020).

Informal sanctuaries are variable and their resource units comprise the protected fish populations. Users often include sanctuary owners/guardians and tourists or communities that visit the site. Although they lack a structured

governance system, protection is offered through the presence of one or more vigilant individual(s) who reside(s) by the river. For example, the staff of the Sitanadi Jungle Camp began feeding the fish in the river near their campsite whilst simultaneously warding off illegal fishing activities. The consequent occurrence of high fish densities became a tourist attraction, which further encouraged protection efforts. Such sanctuaries are managed informally, including via food provisioning and the prevention of illegal activities such as destructive fishing practices and sand mining.

Based on the policies and institutions governing fish sanctuaries, their establishment creates natural capital assets, primarily in the form of protected riverine habitats and associated biotic components (Fig. 1, Table 2). Consequently, some level of ecosystem function is preserved, with the extent of this preservation depending on on sanctuary characteristics such as size, location and condition. Natural assets and preserved ecosystem function yield various ecosystem services to users within and beyond the boundary of the social–ecological system. Potential ecosystem services generated include regulatory (water purification, flood control), provisioning (water, food and/or fodder supply), supporting (biodiversity, nutrient cycling) and cultural (spirituality, tourism, recreation) services (Costanza et al., 1997; Millennium Ecosystem Assessment, 2005). These services in turn provide direct benefits and intangible value to local communities, thereby contributing to human well-being (Fig. 1). The extent of ecosystem services



PLATE 1 The Shishileshwara temple fish sanctuary on the Netravathi River in Karnataka, India. Photo: Wikimedia Commons (2007).



PLATE 2 Protector of the Wachi Wari fish sanctuary, a community-based conservation reserve in West Garo hills, Meghalaya, India. Photo: P. Dandekar.

realized is driven further by the policies and institutions governing these sanctuaries (Kalaba, 2014), which vary across sanctuary types.

Threats to fish sanctuaries

As community-based fish sanctuaries are documented poorly and not recognized officially, they are subject to threats (Ramachandra et al., 2013; Gupta et al., 2016), which originate from factors within the boundary of the social–ecological system, within the broader river system and beyond the river system (Fig. 2).

Threats originating within the social–ecological system are linked to the various subsystems and the interactions between them (Jupiter et al., 2017). Often the efficacy of a social–ecological system centred on common-pool resource management relies on the ability of participatory mechanisms to align individual interests with those of the group



PLATE 3 The informal Sitanadi fish sanctuary in Karnataka, India. Photo: S. Machado.

(Ostrom, 2009). The literature on common-pool governance (Ostrom, 2009) suggests that the conservation outcomes of fish sanctuaries are influenced by interactions between subsystems. For example, sanctuaries can be threatened by the collapse of the governance subsystem (e.g. lack of participatory approaches, lack of trust, conflict amongst users, erosion of governance mechanisms), reduced social compliance (because of weakening religious beliefs, improper monitoring, dissatisfaction, changes in local demographics) or the loss of clarity of system boundaries (if boundaries are not well delineated). Such breakdowns in governance have occurred in fish sanctuaries in India. For example, a feud between two villages managing a fish sanctuary led to the collapse of the governance system and social compliance, resulting in the dynamiting and destruction of the Bansamgre fish sanctuary in Meghalaya (Dandekar, 2018). Temple-based sanctuaries are also at risk because of declining religious beliefs, especially in younger generations who are influenced by increased urbanization and modernization (Gupta et al., 2016), or caste-based conflicts. For instance, in 1996 miscreants destroyed the protected fish population at the Shishileshwara fish sanctuary in Karnataka by pouring chemical poisons into the river as an act of sabotage (Akshatha, 2011).

TABLE 2 Components of the social–ecological framework (Fig. 1) across three types of community-based fish sanctuaries in India.

		Temple-based	Conservation-based	Informal
Motivation for establishment		Religious &/or spiritual motivation	Sustainable management of fish populations	Tourism, recreation &/or economic incentives
Social–ecological subsystems	Resource system	Protected river reach	Protected river reach	Protected river reach
	Resource units	Sacred fish	Fish & other aquatic biota, flowing water, riparian edge	Fish community
	Governance system	Priest or temple institution (top–down management)	Elected representatives from one or more village(s); can be associated with NGO/government body	One or more individual(s)
	Users	Temple devotees, local communities, tourists	Local communities, tourists	Owners, tourists, local communities
Functional characterization	Natural capital	Protected ecosystem structure (with respect to stream & riparian edge habitat), riverine biodiversity (with a focus on fish communities) & ecosystem function		
	Ecosystem services	Regulatory (water provisioning & flood control), supporting (biodiversity, nutrient & sediment cycling), provisioning (water, food, gravel, fodder) & cultural (recreational, spiritual, tourism)		
	Human well-being	Benefits: place of worship, recreation, tourism Values: religious, spiritual, cultural, social, economic	Benefits: food & water security, recreation, tourism, education Values: cultural, social, ecological, economic	Benefits: food & water security, recreation, tourism, education Values: cultural, social, economic, ecological
Policy & institutions		Governed by traditional practices; rules set by temple board or priest based on religious beliefs; scientific knowledge not considered	Governed by traditional knowledge & science-based practice through a participatory management approach; allows for collaborations with NGOs/government bodies & capacity development	No clear policy mechanism; managed using traditional knowledge but can incorporate science-based practice



Fig. 2 Threats to community-based fish sanctuaries in India across spatial scales.

Threats at an intermediate scale, from outside the boundary of the social–ecological system but within the river system, are also prevalent. As most fish sanctuaries are not well documented and receive little to no formal protection, they are often ignored in environmental impact assessments and development planning. Consequently, many have been lost or are under threat from hydropower dams, water diversion projects, development

of inland waterways and pollution (Dandekar, 2013; Raghu, 2013; Pinder et al., 2019). Dams can affect sanctuaries through submergence, loss of connectivity, flow regulation, altered riverine habitats and disturbances during their construction and operation (Bhatt & Pandit, 2016; Jumani et al., 2020). For example, the proposed Kukke Stage I small hydropower project threatens the Yenekal and Nakur Gaya temple fish sanctuaries on the Kumaradhara River in Karnataka (Ramachandra et al., 2013). The Ganol and Daribokgre hydropower dams threaten numerous sanctuaries on the Simsang River (Gupta et al., 2016). Similarly, pollution events caused mass fish mortality in two temple sanctuaries on the Indrayani River and one on the Bhima River in Maharashtra (Pinder et al., 2019). The destruction of resource systems and resource units adversely affects natural capital, thereby compromising the entire functional loop of the social–ecological system (Fig. 1). As many fish species in sanctuaries are potamodromous (i.e. those that undertake seasonal migrations to access upstream spawning habitats), fish populations are affected by dams that fragment the river and impede successful migration and reproduction (Nautiyal et al., 2008; Gupta et al., 2016).

At broader scales beyond the river system, threats from global climate change also affect fish sanctuaries. Global warming has caused an increase in extreme climatic events, including more frequent and severe droughts and changes in rainfall patterns and intensity (Perkins et al., 2012; Trenberth et al., 2014). These changes are expected to alter stream flows, increase stream drying, reduce hydrological connectivity and increase maximum water temperatures, which in turn affect endemic fish communities, especially in intermittent streams (Vorosmarty et al., 2000; Meenu et al., 2013). The Himalayas are amongst the hotspots that are most sensitive to global warming (Immerzeel et al., 2010; Hock et al., 2019), with their rivers and fish populations being especially vulnerable (Grumbine & Pandit, 2013; Nautiyal, 2014). Increasing water temperatures have affected the reproduction of fish such as the Indian major carps in the river Ganga (Vass et al., 2009) and the copper mahseer *Neolissochilus hexagonolepis* (a flagship species in many fish reserves) in Meghalaya (Majhi et al., 2013). Additionally, geographical shifts of warm-water species towards colder stretches have also occurred in various streams (Vass et al., 2009; Adve, 2014).

Conservation potential of fish sanctuaries

Well-managed fish sanctuaries have the potential to contribute to the in situ conservation of freshwater fish, other aquatic biota and riparian habitats whilst also providing socio-cultural benefits. Sanctuaries can function as riverine protected areas, protecting not only fish and biotic communities but also entire reaches of riparian habitats, akin to the freshwater conservation zones that have been documented across Southeast Asia (Baird et al., 2005; Koning et al., 2020). By safeguarding against harmful activities, fish sanctuaries have the potential to form refuge habitats for species of high conservation importance such as the black mahseer *Tor khudree* (Least Concern), Malabar mahseer *Tor malabaricus* (Endangered), hump-backed mahseer *Tor remadevii* (Critically Endangered), copper mahseer *N. hexagonolepis* (Near Threatened) and kooral or Curmuca barb *Hypselobarbus curmuca* (Endangered). For example, surveys along the Savitri basin in the Western Ghats recorded the migratory *T. khudree* only in and around the Walan Kond temple-based fish sanctuary (Katwate & Katwate, 2015). As illustrated by studies of similar freshwater conservation zones in Laos (Baird et al., 2005) and Thailand (Koning et al., 2020), fish communities in such sanctuaries can attain higher species richness, density and biomass, and can potentially serve as source populations for adjacent river reaches.

Given that community-based initiatives can achieve in situ conservation whilst circumventing excessive spending and bureaucratic processes (Dandekar, 2018), there is an urgent need to tap into their conservation potential. Temple priests can be powerful allies in efforts to curb prevalent

destructive practices such as illegal sand mining, dynamiting and poison fishing. As temple institutions and leaders are respected by local communities, they can garner local conservation support through communication and education (Sheikh, 2006; Bhatia et al., 2017). Similarly, capacity building with key actors in community-based reserves could help promote best practices. Targeted engagement with landowners near streams that support high freshwater biodiversity can also help to popularize and improve informal fish sanctuaries.

Efforts made by NGOs, scientists and governmental bodies to engage with local communities can also aid in the creation of new fish sanctuaries. For example, the Meghalaya State Aquaculture Mission has been working with local communities to revitalize their traditional fish protection and management practices. Since 2012 they have helped local communities establish > 54 fish sanctuaries in Meghalaya (Meghalaya State Aquaculture Mission, 2014). Although governmental agencies offer infrastructure and logistical support, the local community manages, protects and monitors the sanctuaries (Dandekar, 2018). Such sanctuaries have benefitted local communities through increased capture of food fish, local tourism, generation of employment and improved livelihoods (Meghalaya State Aquaculture Mission, 2014).

Given their socio-ecological importance, sanctuaries should be integrated into broader conservation efforts and recognized in official decision-making processes, impact assessments and development planning, similarly to how protected areas and sacred groves are considered. Consideration should also be given to increasing the recognition of fish sanctuaries by pursuing their official designation, for example as IUCN Category VI protected areas. However, given that sanctuaries are typically small, scattered and vulnerable to threats from within and beyond their boundaries, additional strategies are needed for fish conservation. Multi-pronged approaches that include catchment-scale management and conservation, legislation, education programmes and enforcement will be essential to conserving these systems.

Directions for future research

Community-based fish sanctuaries are complex social-ecological systems that have been documented to some extent in popular articles and historical records, but have been mostly overlooked in the scientific literature (except Gupta et al., 2016). Given their potential for conservation, there is an urgent need to address this lack of scientific knowledge. All of the available information on known sanctuaries needs to be compiled into a publicly available data source, particularly as many such sanctuaries are being destroyed without having been documented.

We highlight the following areas as priorities for further research: (1) Determine the impacts of sanctuaries on fish behaviour and population dynamics, other aquatic biota and riverine habitats. (2) Examine the potential of protected fish communities to serve as source populations within a river system. (3) Determine how factors such as sanctuary size, location, level of protection and local compliance with protective regulations influence the effectiveness of sanctuaries in protecting threatened species. (4) Ascertain the social, ecological and institutional factors that contribute to the various conservation outcomes of these sanctuaries (Gokhale et al., 1998; Ostrom, 2009; Jupiter et al., 2017). (5) Determine the users and governance subsystems across fish sanctuary types, communities and geographical regions and their influence on the efficacy of management techniques, levels of compliance and local attitudes towards these systems. (6) Examine the responses and adaptive capacities of sanctuaries to growing anthropogenic threats. (7) Determine the varied ecosystem services derived from sanctuaries.

Acknowledgements We thank Mahi Puri, Chad Palmer and Woi Sok for comments on manuscript drafts, and the Editor and two anonymous reviewers for their critiques. This research received no specific grant from any funding agency or commercial or not-for-profit sectors.

Author contributions Conceptual development: SJ, with contributions from VH; writing: SJ, with contributions from VH, PD, NM.

Conflicts of interest None.

Ethical standards This research abided by the *Oryx* guidelines on ethical standards.

References

- ABELL, R., THIEME, M.L., REVENGA, C., BRYER, M., KOTTELAT, M., BOGUTSKAYA, N. et al. (2008) Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience*, 58, 403–414.
- ABRAHAM, R.K. & KELKAR, N. (2012) Do terrestrial protected areas conserve freshwater fish diversity? Results from the Western Ghats of India. *Oryx*, 46, 544–553.
- ADVE, N. (2014) Moving home: global warming and the shifts in species' range in India. *Economic and Political Weekly*, 49, 34–38.
- AKSHATHA, M. (2011) This temple has power to attract atheists too. *Deccan Herald*, 11 March 2011. deccanherald.com/content/145072/this-temple-has-power-attract.html [accessed 9 May 2022].
- ATKORE, V., KELKAR, N. & KRISHNASWAMY, J. (2017) Assessing the recovery of fish assemblages downstream of hydrological barriers in India's Western Ghats. *River Research and Applications*, 33, 1026–1035.
- BAIRD, I.G., FLAHERTY, M.S. & BAIRD, I.G. (2005) Mekong river fish conservation zones in southern Laos: assessing effectiveness using local ecological knowledge. *Environmental Management*, 36, 439–454.
- BARUAH, D. & SARMA, D. (2018) Mahseer in recreational fisheries and ecotourism in India. *Aquaculture Asia Magazine*, 22, 3–10.
- BECK, M.W., CLAASSEN, A.H. & HUNDT, P.J. (2012) Environmental and livelihood impacts of dams: common lessons across development gradients that challenge sustainability. *International Journal of River Basin Management*, 10, 73–92.
- BHAGWAT, S.A. & RUTTE, C. (2006) Sacred groves: potential for biodiversity management. *Frontiers in Ecology and the Environment*, 4, 519–524.
- BHATIA, S., REDPATH, S.M., SURYAWANSHI, K. & MISHRA, C. (2017) The relationship between religion and attitudes toward large carnivores in northern India? *Human Dimensions of Wildlife*, 22, 30–42.
- BHATT, J.P. & PANDIT, M.K. (2016) Endangered golden mahseer *Tor putitora* Hamilton: a review of natural history. *Reviews in Fish Biology and Fisheries*, 26, 25–38.
- COSTANZA, R., D'ARGE, R., DE GROOT, R., FARBER, S., GRASSO, M., HANNON, B. et al. (1997) The value of the world's ecosystem services and natural capital. *Nature*, 387, 253–260.
- COUTO, T.B. & OLDEN, J.D. (2018) Global proliferation of small hydropower plants – science and policy. *Frontiers in Ecology and the Environment*, 16, 91–100.
- DAHANUKAR, N. & RAGHAVAN, R. (2013) *Freshwater Fishes of the Western Ghats: Checklist v1.0*, 12 August 2013. [researchgate.net/publication/263086413_Freshwater_fishes_of_the_Western_Ghats_Checklist_v1_0_August_2013](https://www.researchgate.net/publication/263086413_Freshwater_fishes_of_the_Western_Ghats_Checklist_v1_0_August_2013) [accessed 15 May 2021].
- DANDEKAR, P. (2013) Community fish sanctuaries: protecting the fish and their rivers. *South Asia Network on Dams, Rivers and People*, 21 November 2013. sandr.in/2013/11/21/community-fish-sanctuaries-protecting-the-fishes-and-their-rivers [accessed 9 May 2022].
- DANDEKAR, P. (2018) Community-led projects in Meghalaya are helping the rare mahseer thrive. *Scroll.in*, 11 December 2018. scroll.in/article/904599/many-fish-in-the-river-community-led-projects-in-meghalaya-are-helping-the-rare-mahseer-thrive [accessed 22 March 2021].
- DASH, P., TANDEL, R., BARUAH, D. & SARMA, D. (2020) Mahseer sanctuaries of Meghalaya: a conservation and recreational perspective. *Aquaculture Asia Magazine*, 241, 3–7.
- DUDGEON, D. (2002) The most endangered ecosystems in the world? Conservation of riverine biodiversity in Asia. *SIL Proceedings, 1922–2010*, 28, 59–68.
- DUDGEON, D., ARTHINGTON, A.H., GESSNER, M.O., KAWABATA, Z., KNOWLER, D.J., NAIMAN, R.J. et al. (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Review*, 81, 163–182.
- ESCHMEYER, W.N. & FRICKE, R. (2010) *Catalog of Fishes*. Electronic version, 15 January 2010. [research.calacademy.org/ichthyology/catalog/fishcatmain.asp](https://www.research.calacademy.org/ichthyology/catalog/fishcatmain.asp) [accessed 10 May 2010].
- FROESE, R. & PAULY, D. (2010) *FishBase*. Fisheries Centre, University of British Columbia, Vancouver, Canada.
- GOKHALE, Y., VELANKAR, R., CHANDRAN, M.D. & GADGIL, M. (1998) Sacred woods, grasslands and water bodies as self-organized systems of conservation. In *Conserving the Sacred for Biodiversity Management* (eds P.S. Ramakrishnan, K.G. Saxena & U.M. Chandrashekara), pp. 365–396. IBH Publishing Co., Oxford, UK and New Delhi, India.
- GRANT, E.H.C., LYNCH, H.J., MUNEEPEERAKUL, R., ARUNACHALAM, M., RODRIGUEZ-ITURBE, I. & FAGAN, W.F. (2012) Interbasin water transfer, riverine connectivity, and spatial controls on fish biodiversity. *PLOS ONE*, 7, e34170.
- GRILL, G., LEHNER, B., THIEME, M., GEENEN, B., TICKNER, D., ANTONELLI, F. et al. (2019) Mapping the world's free-flowing rivers. *Nature*, 569, 215–221.

- GRUMBINE, R.E. & PANDIT, M.K. (2013) Threats from India's Himalaya dams. *Science*, 339, 36–37.
- GUPTA, N., KANAGAVEL, A., DANDEKAR, P., DAHANUKAR, N., SIVAKUMAR, K., MATHUR, V.B. & RAGHAVAN, R. (2016) God's fishes: religion, culture and freshwater fish conservation in India. *Oryx*, 50, 244–249.
- HOCK, R., RASUL, G., ADLER, C., CÁCERES, B., GRUBER, S., HIRABAYASHI, Y. et al. (2019) High Mountain Areas. In *Special Report on the Ocean and Cryosphere in a Changing Climate*, pp. 131–202. Intergovernmental Panel on Climate Change, Geneva, Switzerland.
- ICOLD (2019) International Commission on Large Dams. [icold-cigb.org/article/GB/world_register/general_synthesis/number-of-dams-by-country-members](https://www.icold.org/article/GB/world_register/general_synthesis/number-of-dams-by-country-members) [accessed 9 June 2019].
- IMMERZEEL, W.W., VAN BEEK, L.P.H. & BIERKENS, M.F.P. (2010) Climate change will affect the Asian water towers. *Science*, 328, 1382–1385.
- JUMANI, S., DEITCH, M.J., KAPLAN, D., ANDERSON, E.P., KRISHNASWAMY, J., LECOUCRS, V. & WHILES, M.R. (2020) River fragmentation and flow alteration metrics: a review of methods and directions for future research. *Environmental Research Letters*, 15, 123009.
- JUMANI, S., RAO, S., KELKAR, N., MACHADO, S., KRISHNASWAMY, J. & VAIDYANATHAN, S. (2018) Fish community responses to stream flow alterations and habitat modifications by small hydropower projects in the Western Ghats biodiversity hotspot, India. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28, 979–993.
- JUMANI, S., RAO, S., MACHADO, S. & PRAKASH, A. (2017) Big concerns with small projects: evaluating the socio-ecological impacts of small hydropower projects in India. *Ambio*, 46, 500–511.
- JUPITER, S.D., EPSTEIN, G., BAN, N.C., MANGUBHAI, S., FOX, M. & COX, M. (2017) A social–ecological systems approach to assessing conservation and fisheries outcomes in Fijian locally managed marine areas. *Society & Natural Resources*, 30, 1096–1111.
- KALABA, F.K. (2014) A conceptual framework for understanding forest socio-ecological systems. *Biodiversity and Conservation*, 23, 3391–3403.
- KATWATE, U. & KATWATE, C. (2015) *Status of Freshwater Fishes in the Sahyadri–Konkan Corridor: Diversity, Distribution and Conservation Assessments in Raigad*. Grant report submitted to CEPF-ATREE Western Ghats Program. Bombay Natural History Society, India. cepf.net/sites/default/files/sg61484-technical-report.pdf [accessed 3 June 2021].
- KHARAT, S., KUMKAR, P. & SONAWANE, K. (2013) Traditional fishing techniques of Adivasi tribes in Tamhini region of Western Ghats. *International Journal of Fisheries and Aquaculture Sciences*, 3, 165–172.
- KONING, A.A., PERALES, K.M., FLUET-CHOUINARD, E. & MCINTYRE, P.B. (2020) A network of grassroots reserves protects tropical river fish diversity. *Nature*, 588, 631–635.
- KRISHNASWAMY, J., KUMAR, M., KELKAR, N., NAIR, T. & ATKORE, V. (2017) Moving from requiem to revival: India's rivers and riverine ecosystems. In *Transcending Boundaries: Reflecting on Twenty Years of Action and Research at ATREE* (eds A.J. Hiremath, N.D. Rai & A. Siddhartha), pp. 94–103. Ashoka Trust for Research in Ecology and the Environment, Bangalore, India.
- MAJHI, S.K., DAS, S.K. & RAJKHOWA, D. (2013) Effects of elevated water temperature on tolerance and stress in chocolate mahseer *Neolissochilus hexagonolepis*: implications for habitat restoration and conservation. *Current Science*, 105, 379–383.
- MCGINNIS, M.D. & OSTROM, E. (2014) Social–ecological system framework: initial changes and continuing challenges. *Ecology and Society*, 19, 30.
- MEENU, R., REHANA, S. & MUJUMDAR, P.P. (2013) Assessment of hydrologic impacts of climate change in Tunga–Bhadra river basin, India with HEC-HMS and SDSM: hydrologic impacts of climate change. *Hydrological Processes*, 27, 1572–1589.
- MILLENNIUM ECOSYSTEM ASSESSMENT (2005) *Ecosystems and Human Well-Being*. Island Press, Washington, DC, USA.
- MOLUR, S., SMITH, K.G. & DANIEL, B.A. (2011) *The Status and Distribution of Freshwater Biodiversity in the Western Ghats, India*. IUCN, Zoo Outreach Organisation, Cambridge, UK and Coimbatore, India.
- MEGHALAYA STATE AQUACULTURE MISSION (2014) *Conversation with People of Meghalaya*. Meghalaya Basin Development Authority, Meghalaya, India.
- MYERS, N., MITTERMEIER, R.A., MITTERMEIER, C.G., DA FONSECA, G.A.B. & KENT, J. (2000) Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.
- NAUTIYAL, P. (1989) Mahseer conservation: problems and prospects. *Journal of Bombay Natural History Society*, 86, 32–36.
- NAUTIYAL, P. (2014) Review of the art and science of Indian mahseer (game fish) from nineteenth to twentieth century: road to extinction or conservation? *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 84, 215–236.
- NAUTIYAL, P., RIZVI, A.F. & DHASMANAA, P. (2008) Life-history traits and decadal trends in the growth parameters of golden mahseer *Tor puititora* (Hamilton 1822) from the Himalayan stretch of the Ganga river system. *Turkish Journal of Fisheries and Aquatic Sciences*, 8, 125–131.
- NILSSON, C. (2005) Fragmentation and flow regulation of the world's large river systems. *Science*, 308, 405–408.
- OSTROM, E. (2009) A general framework for analyzing sustainability of social–ecological systems. *Science*, 325, 419–422.
- PANDIT, M.K. & GRUMBINE, R.E. (2012) Potential effects of ongoing and proposed hydropower development on terrestrial biological diversity in the Indian Himalaya. *Conservation Biology*, 26, 1061–1071.
- PERKINS, S.E., ALEXANDER, L.V. & NAIRN, J.R. (2012) Increasing frequency, intensity and duration of observed global heatwaves and warm spells. *Geophysical Research Letters*, 39, 20714.
- PINDER, A.C. & RAGHAVAN, R. (2013) Conserving the Endangered mahseers (*Tor* spp.) of India: the positive role of recreational fisheries. *Current Science*, 104, 1472–1475.
- PINDER, A.C., BRITTON, J.R., HARRISON, A.J., NAUTIYAL, P., BOWER, S.D., COOKE, S.J. et al. (2019) Mahseer (*Tor* spp.) fishes of the world: status, challenges and opportunities for conservation. *Reviews in Fish Biology and Fisheries*, 29, 417–452.
- PINTO, N., VAIDYANATHAN, S., VARUGHESE, S., KRISHNASWAMY, J., MASSAR, B. & HAOKIP, J.V. (2021) Establishment of community-led fish conservation zones in Meghalaya and Manipur, India. *Oryx*, 55, 493–494.
- PLANNING COMMISSION (ed.) (2012) *Twelfth Five Year Plan (2012–2017)*. SAGE Publications, New Delhi, India and Thousand Oaks, USA.
- PRAJITH, K., REMESAN, M. & EDWIN, L. (2016) Traditional wisdom of fishing techniques and rituals of Kuruman tribe of Wayanad, Western Ghats. *Asian Agri-History*, 20, 119–126.
- RAGHU, S. (2013) The sacred fish tales of Shishila. *The New Indian Express*, 6 October 2013. [newindianexpress.com/magazine/2013/oct/06/The-sacred-fish-tales-of-Shishila-523411.html](https://www.newindianexpress.com/magazine/2013/oct/06/The-sacred-fish-tales-of-Shishila-523411.html) [accessed 9 May 2022].
- RAMACHANDRA, T.V., CHANDRAN, M.D., SHENOY, H.S., RAO, R.G., SHIVAMURTHY, V., MUKRI, V. et al. (2013) *Kumaradhara River Basin, Karnataka Western Ghats: Need for Conservation and Sustainable Use*. wgbis.ces.iisc.ernet.in/biodiversity/pubs/ETR/ETR54/vegetation.htm [accessed 9 May 2022].

- RAMESAN, M.P. (2006) *Studies on inland fishing gears of north Kerala*. PhD thesis. Cochin University of Science and Technology, Cochin, India.
- RESILIENCE ALLIANCE (2010) *Assessing Resilience in Social–Ecological Systems: Workbook for Practitioners*. Version 2. resalliance.org/3871.php [accessed 9 May 2022].
- SHAJI, C.P. & LALADHAS, K.P. (2013) Multifariousness in the structure and fabrication of ‘Koodu’ – a traditional fish trap of Kerala. *Journal of Aquatic Biology & Fisheries*, 1, 77–82.
- SHEIKH, K.M. (2006) Involving religious leaders in conservation education in the Western Karakorum, Pakistan. *Mountain Research and Development*, 26, 319–322.
- SIMOONS, F.J. (1974) Fish as forbidden food: the case of India. *Ecology of Food and Nutrition*, 3, 185–201.
- TOCKNER, K., BERNHARDT, E.S., KOSKA, A. & ZARFL, C. (2016) A global view on future major water engineering projects. In *Society – Water – Technology* (eds R.F. Hüttl, O. Bens, C. Bismuth & S. Hoehstetter), pp. 47–64. Springer International Publishing, Cham, India.
- TRENBERTH, K.E., DAI, A., VAN DER SCHRIER, G., JONES, P.D., BARICHOVICH, J., BRIFFA, K.R. & SHEFFIELD, J. (2014) Global warming and changes in drought. *Nature Climate Change*, 4, 17–22.
- VASS, K.K., DAS, M.K., SRIVASTAVA, P.K. & DEY, S. (2009) Assessing the impact of climate change on inland fisheries in river Ganga and its plains in India. *Aquatic Ecosystem Health & Management*, 12, 138–151.
- VASUDEVAN, K., KUMAR, A. & CHELLAM, R. (2006) Species turnover: the case of stream amphibians of rainforests in the Western Ghats, southern India. *Biodiversity & Conservation*, 15, 3515–3525.
- VOROSMARTY, C.J., GREEN, P., SALISBURY, J. & LAMMERS, R.B. (2000) Global water resources: vulnerability from climate change and population growth. *Science*, 289, 284–288.
- VYAS, V., DAMDE, D. & PARASHAR, V. (2012) Fish biodiversity of Betwa River in Madhya Pradesh, India with special reference to a sacred ghat. *International Journal of Biodiversity and Conservation*, 4, 71–77.
- WIKIMEDIA COMMONS (2007) *File: Shishila Temple Fish Conservation – Feeding Fish*. en.wikipedia.org/wiki/File:Shishila_temple_fish_conservation_-_feeding_fish.JPG [accessed 3 June 2021].
- ZARFL, C., LUMSDON, A.E., BERLEKAMP, J., TYDECKS, L. & TOCKNER, K. (2015) A global boom in hydropower dam construction. *Aquatic Sciences*, 77, 161–170.