

Corrigendum

Cite this article: Fountoulakis KN, Panagiotidis P, Gonda X, Kimiskidis V, Nimatoudis I (2019). Neurological soft signs significantly differentiate schizophrenia patients from healthy controls – CORRIGENDUM. *Acta Neuropsychiatrica* 31:61. doi: 10.1017/neu.2018.37

Corrigendum to “Neurological soft signs significantly differentiate schizophrenia patients from healthy controls”; [*Acta Neuropsychiatrica* 2018;30(2):97–105]

Konstantinos N. Fountoulakis¹, Panagiotis Panagiotidis¹, Xenia Gonda^{2,3,4}, Vasilios Kimiskidis⁵ and Ioannis Nimatoudis¹

¹3rd Department of Psychiatry, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece,

²Department of Psychiatry and Psychotherapy, Semmelweis University, Budapest, Hungary, ³MTA-SE Neuropsychopharmacology and Neurochemistry Research Group, Hungarian Academy of Sciences, Semmelweis University, Budapest, Hungary, ⁴NAP-A-SE New Antidepressant Target Research Group, Semmelweis University, Budapest, Hungary and ⁵Laboratory of Clinical Neurophysiology, School of Medicine, Aristotle University of Thessaloniki, Greece

<https://doi.org/10.1017/neu.2017.29> Published online by Cambridge University Press, 29 November 2017

Table 2 (page 100): The heading ‘Drug free (n = 98) Medicated (n = 35)’ should instead read ‘Medicated (n = 98) Drug free (n = 35)’.

The authors would like to apologise for any inconvenience caused.

Reference

Fountoulakis K, Panagiotidis P, Gonda X, Kimiskidis V and Nimatoudis I. (2018) Neurological soft signs significantly differentiate schizophrenia patients from healthy controls. *Acta Neuropsychiatrica* 30(2), 97–105. doi: 10.1017/neu.2017.29

