

Competency in Chaos: Lifesaving Performance of Care Providers Utilizing a Competency-Based, Multi-Actor Emergency Preparedness Training Curriculum

Lancer A. Scott, MD, FACEP;¹ Derrick A. Swartzentruber, BS;² Christopher Ashby Davis, BS;² P. Tim Maddux, BS;² Jennifer Schnellman, PhD, ELS;³ Amy E. Wahlquist, MS⁴

1. Emergency Medicine, Medical University of South Carolina, Charleston, South Carolina USA
2. College of Medicine, Medical University of South Carolina, Charleston, South Carolina USA
3. Department of Clinical Pharmacy and Outcomes Sciences, Medical University of South Carolina, Charleston, South Carolina USA
4. Division of Biostatistics and Epidemiology, Medical University of South Carolina, Charleston, South Carolina USA

Correspondence:

Lancer A. Scott, MD, FACEP
Division of Emergency Medicine and Pediatric Emergency Medicine
Medical University of South Carolina
MSC 300
115 MUH
169 Ashley Ave.
Charleston, SC 29425 USA
E-mail: scottlan@musc.edu

Conflicts of interest and funding: The authors report no conflicts of interest. This research was partially supported by a grant from the South Carolina Translational Institute.

Keywords: clinical disaster; competency-based training; mass casualty; care provider; simulation

Abbreviations:

CHPTER: Center for Health Professional Training and Emergency Response
ED: emergency department
EPT: emergency preparedness training
IRB: institutional review board
START: Simple Triage and Rapid Treatment
VHA: Veterans Health Administration

Abstract

Objective: Providing comprehensive emergency preparedness training (EPT) to care providers is important to the future success of disaster operations in the US. Few EPT programs possess both competency-driven goals and metrics to measure performance during a multi-patient simulated disaster.

Methods: A 1-day (8-hour) EPT course for care providers was developed to enhance provider knowledge, skill, and comfort necessary to save lives during a simulated disaster. Nine learning objectives, 18 competencies, and 34 performance objectives were developed. During the 2-year demonstration of the curriculum, 24 fourth-year medical students and 17 Veterans Hospital Administration (VHA) providers were recruited and volunteered to take the course (two did not fully complete the research materials). An online pre-test, two post-tests, course assessment, didactic and small group content, and a 6-minute clinical casualty scenario were developed. During the scenario, trainees working in teams were confronted with three human simulators and 10 actor patients simultaneously. Unless appropriate performance objectives were met, the simulators “died” and the team was exposed to “anthrax.” After the scenario, team members participated in a facilitator-led debriefing using digital video and then repeated the scenario.

Results: Trainees (N = 39) included 24 (62%) medical students; seven (18%) physicians; seven (18%) nurses; and one (3%) emergency manager. Forty-seven percent of the VHA providers reported greater than 16 annual hours of disaster training, while 15 (63%) of the medical students reported no annual disaster training. The mean (SD) score for the pre-test was 12.3 (3.8), or 51% correct, and after the training, the mean (SD) score was 18.5 (2.2), or 77% ($P < .01$). The overall rating for the course was 96 out of 100. Trainee self-assessment of “Overall Skill” increased from 63.3 out of 100 to 83.4 out of 100 and “Overall Knowledge” increased from 49.3 out of 100 to 78.7 out of 100 ($P < .01$). Of the 34 performance objectives during the disaster scenario, 23 were completed by at least half of the teams during their first attempt. All teams except one (8 of 9) could resuscitate two simulators and all teams (9 of 9) helped prevent anthrax exposure during their second scenario attempt.

Conclusions: The 1-day EPT course for novice and experienced care providers recreated a multi-actor clinical disaster and enhanced provider knowledge, comfort level, and EPT skill. A larger-scale study, or multi-center trial, is needed to further study the impact of this curriculum and its potential to protect provider and patient lives.

Scott LA, Swartzentruber D, Davis CA, Maddux PT, Schnellman J, Wahlquist AE. Competency in chaos: lifesaving performance of care providers utilizing a competency-based, multi-actor emergency preparedness training curriculum. *Prehosp Disaster Med.* 2013;28(4):322-333.

Received: August 9, 2012
Revised: December 14, 2012
Accepted: January 20, 2013

Online publication: April 26, 2013

doi:10.1017/S1049023X13000368

Introduction

The lack of emergency preparedness training (EPT) for patient care providers—including clinicians, hospital workers, mental health providers, public safety and law enforcement officials, community volunteers, EMS, HazMat and Fire personnel—poses significant risks to both patients and providers. During the 1995 Tokyo sarin gas attacks, for example, most patients bypassed first responders and reported directly to hospitals, where staff was exposed to sarin due to inadequate personal protective equipment (PPE) and training.¹ Furthermore, during Hurricane Katrina, EPT deficits were cited as significant factors contributing to adverse patient outcomes.²⁻⁷

Providing comprehensive EPT for medical trainees is important to the future success of emergency preparedness operations, yet few medical schools have defined and implemented EPT core competencies for health professionals.⁸⁻²⁴ Recent reviews suggest that health care worker EPT programs lack clarity, objectivity, competency-driven goals, scientific rigor, prospective validation, and consistency across medical specialties.²⁵⁻²⁸

A prior study revealed how medical students can value and adopt core EPT elements via a novel addition to an existing curriculum.^{19,29} A significant limitation of the study was the relatively simplistic measurement of EPT performance. A new course from the Center for Health Professional Training and Emergency Response (CHPTER) was developed to create loud and chaotic, multi-actor, clinical disaster scenes (combining more than 10 human actors with human simulators at one time) that would enhance provider knowledge, skills, and comfort necessary to save lives during a simulated disaster. This study presents the performance data of trainees participating in the CHPTER EPT course in 2011 and 2012, including trainee self-assessment, course assessment, precognitive and postcognitive performance, small-group scenario performance, and clinical mass-casualty scenario team performance.

Methods

Curriculum Development—Organization

In 2009, CHPTER was formed as South Carolina's first collaborative EPT center for care providers (www.musc.edu/chpter). Over the course of several months, a CHPTER task force established learning objectives for a 1-day EPT course utilizing existing evaluative frameworks, coursework, and data.³⁰⁻³⁹ Using a modified Delphi process, dozens of competencies were consolidated into 18, and subsequently assigned to five competency domains (Table 1). The task force then developed performance objectives to match competency objectives, and these guided content development for the didactic, small group, and simulation course components. The project was approved by the Medical University of South Carolina Institutional Review Board (IRB).

Small-Group Exercise Development

Didactic Module 1 was developed as an introductory lecture to the training course. Didactic modules 2, 3, and 4 were developed to include a small group exercise designed to prepare students for the afternoon disaster scenario. Following a short slide-based lecture, trainees were randomly assigned to teams of four to six and were assessed by CHPTER instructors who served as small group facilitators.

The team building exercise (module 2) consisted of seven performance components and four fictional disaster scenarios printed on 34-piece puzzles. The unsolved puzzles were presented

to teams in large envelopes, and teams were instructed to complete at least one disaster scenario. Instructors evaluated the teams based on their ability to complete the puzzle and verbalize appropriate responses to threats presented. The communications exercise (module 3) consisted of three scenarios (bus crash, factory explosion, and chemical leak) presented to team members on preprinted handouts, and teams were instructed to complete at least one scenario. Scenarios were discussed in small groups, guided by a facilitator who measured group effectiveness in communicating clinical disaster information. The triage exercise (module 4) consisted of a tabletop triage exercise using 60 small toys; each toy represented a simulated patient and was imprinted with clinical information. Groups were asked to classify the 60 simulated patients according to Simple Triage and Rapid Treatment (START). Facilitators evaluated trainees based on their ability to work as a team to rapidly assess and accurately triage several patients at once.

Human Simulation and Actor Scenario Development

Over six months, CHPTER worked with a university-based human simulation center to develop a series of fictional clinical disasters that combined up to six SimMan high-fidelity and low-fidelity patient simulators (Laerdal, Wappinger Falls, New York, USA) and up to 15 trained live “actors” to simulate a clinical disaster. One of the center's larger observation rooms resembled a small emergency waiting room, and a door to the outside of the simulation center simulated an emergency department (ED) entrance. Storyboards and a stage map (Figure 1) were developed for the simulated exercise. On the stage map, three zones were identified, with an observer assigned to each zone (O1, O2, O3). The letter A designated a human actor and the letter S designated a human simulator. On the color stage map (available online), blue indicated an observer, while green, red, yellow and black indicated a patient's START triage classification. If a chair appeared empty, with no patient (no letters, no color), then it was simply an empty chair.

Members of the CHPTER task force were trained to operate simulation center equipment and software. Observers were assigned to different geographical areas (zones 1, 2 and 3) of the scene so that they could more easily focus on performance objectives during the chaotic movement of patients and actors during the scenario.

Two months prior to training, 15 actors were recruited—including trained patient actors from the medical university—and provided pre-scripted roles. To enhance reliability and validity of performance objectives expected to be achieved by trainees, actors were taught how to follow specific behaviors when confronted with trainees in a chaotic environment.

Multi-Actor Clinical Disaster Scenario: “Influenza-like Illness”

The clinical casualty scenario developed by the curriculum task force involved the acute presentation of cruise-line tourists complaining of cough and shortness of breath. During the 6-minute scenario (“Influenza-like Illness”) both actors and simulators presented with various levels of respiratory complaints, several of them outside awaiting access to the ED waiting room. Working in teams of four to six, trainees were told they were health care providers from different areas of the hospital called down to mitigate the complex and chaotic scene using skills they had learned during didactic and small group lessons. The responders were not aware that some of patients were suffering

Learning Objectives
A. Define a health care disaster and the components of emergency preparedness as it applies to care providers. ^a
B. Understand ethical implications of a health care disaster and its impact on the community. ^b
C. Differentiate between National Incident Management System (NIMS), Incident Command System (ICS/HICS) and the operational disaster/emergency preparedness plan for a health care facility. ^a
D. Identify functional roles (and appreciate your individual limits) of care providers during a disaster. ^{a,b}
E. Define and demonstrate ability to function within the chain of command during a patient care disaster scene. ^{a,c}
F. Define and respond to vulnerabilities and security risks facing providers, health care workers, and health care facilities during a disaster scenario. ^{a,c}
G. Summarize components of teamwork, communication, and triage that are essential to an effective response during a health care disaster and list specific actions to take and to avoid during a health care disaster. ^a
H. Demonstrate effective teamwork, communication and triage to protect care providers and save lives during a disaster scenario. ^c
I. Achieve a greater comfort level with knowledge and skill to provide effective patient care during a clinical disaster. ^b
Competency Domains and Performance Objective Categories
<i>Mobilization: Pre-stage planning and team development</i>
Define and recognize a disaster. (A, F) ^d
Stop: Establish a safety plan. (D, F, G, H)
Develop clinical disaster team (ie, establish leadership, roles and duties). (D, F, G, H)
Select and don appropriate personal protective equipment (PPE) and supplies. (F, G, H)
<i>Clinical disaster operations and communications</i>
Establish operations command (ie, assume team roles). (C, D, E, H)
Establish communications with health care authority and activate Healthcare Incident Command System (HICS). (C, D, E, H)
Appropriately report scene information and needs. (ie, maintain situational awareness). (C, D, E, H)
Optimize teamwork and coordinate tasks. (C, D, E, H)
<i>Protect and preserve human life and continuity of health care facility during a disaster</i>
Ensure personal safety. (C, D, E, F, G, H)
Ensure safety of patients, families, and staff. (D, E, F, G, H)
Accurately assess, reassess and care for patients. (D, E, F, G, H)
Ensure continuity of patient care operations. (D, E, F, G, H)
Preserve integrity and conservation of the physical plant. (D, E, F, G, H)
Perform maneuvers to save simulated patients during a disaster scenario. (F, H)
<i>Demobilization</i>
Ensure effective demobilization of health care resources. (A, C, F, G)
<i>Clinical disaster and emergency preparedness awareness</i>
Understand role of provider to support patients and the community. (B,D)
Understand ethical implications of patient care during a disaster. (B, H)
Self-assess capabilities and limits as a provider during a disaster. (B, D)

Scott © 2013 Prehospital and Disaster Medicine

Table 1. Course Learning Objectives, Competency Domains, and Performance Objective Categories^aDiscrete Knowledge/Cognitive Objective.^bAttitudinal/Affective Objective.^cPerformance/Skill Objective.^dReferences are to learning objectives, above.




Figure 1. Simulator and Actor Staging in Human Performance Lab
 Blue, observer trainer; green, triage green; yellow, triage yellow; red, triage red; black, triage black
 A, actor; S, human simulator; O, observer trainer; 1, zone 1; 2, zone 2; 3, zone 3
 Example: 2S1 (red) is a high-fidelity human simulator patient who is triage red in zone 2

from inhalation anthrax. They also were unaware that one of the patients was carrying two bags of powder, presumed to be anthrax.

Of the four human simulators used in this scenario, two were unstable and required active airway maneuvers (i.e., simple jaw thrusts) in order to open their airways. If teams recognized patient acuity and acted within four minutes, the simulators' vital signs normalized. If not, CHAPTER observers allowed the physiologic parameters of the simulators to deteriorate irreversibly, resulting in cardiac arrest.

Trainees were also confronted with several actors demanding care who were trained to escalate their behavior during the scenario unless appropriate performance measures were met. Once escorted to the Green Triage (this coincides with the “Team Staging Area and BLS/PPE Equipment” area in Figure 1) waiting area, patients with minor complaints changed costumes and presented to the ED entrance as new patients. One actor (representing the terrorist disguised as a tourist) utilized a distraction caused by other actors to bypass security and enter the ED waiting room. If within four minutes the teams recognized that their resources were outstripped and closed the ED door, the patient surge (and terrorist threat) was averted. If not, the actor entered the ED and held up two bags of powder, exposing all patients and all providers to a simulated lethal dose of anthrax.

An educational intervention was developed for all teams participating in the mass-casualty scenario. Immediately after completing their first attempt at the scenario, team members participated in a facilitator-led debriefing using digital video (and split-screen technology) from four different camera viewpoints. Center for Health Professional Training and Emergency Response instructors reinforced key components of the learning objectives and competencies for the course without prompting trainees with specifics about how to remedy the disaster scene. Per study design, teams repeated the 6-minute scenario after the debriefing. Team membership and scenario content did not change.

Research Metrics

The task force developed an online pre-test and two post-tests for trainees utilizing a pool of questions developed by the task force to meet the learning objectives and competencies of the course. The pre-test consisted of two parts. In the first part, Likert-scale self-assessment questions measured trainees' sense of personal capability and comfort level in handling a disaster. The second part included 23 discrete multiple-choice questions.

The post-test consisted of three parts. The first part was identical to the first part of the pre-test (Likert-scale self-assessment questions). The second part of the post-test contained 23 discrete

multiple-choice questions from the question pool. In some cases, slightly modified questions from the question pool were selected. For example, one post-test question described the same patient triage scenario in the pre-test but assigned different vital signs to the patient so the triage answer was “Red” instead of “Yellow”. The third part of the post-test consisted of Likert-scale and open-ended questions for trainees to evaluate course content and instructors. Trainees were asked to complete the same post-test four to six months following the training.

Performance objectives during the mass-casualty scenario were measured and recorded by three CHAPTER observers (zones 1, 2 and 3 observers). Observers recorded whether performance objectives were “met” based on their observations of the team during the scenario. Observers developed a single list of “met” performance objectives for each team and, with the help of digital video playback, were able to complete any part of the checklist they did not have time to finish. Any data missing from an observer’s checklist was considered an “unmet” performance objective.

Descriptive statistics (means, standard deviations, and medians) were used to describe elements of the EPT training, as appropriate. Wilcoxon rank sum tests were used to compare the pre- and post-test values within the cohorts of medical students and VHA trainees. Because many of the pre- and post-tests were completed by the same individuals, and because the test was completed anonymously (as required by the IRB and, unfortunately, making it impossible to conduct paired testing), the *P* values obtained from the Wilcoxon rank sum tests are conservative estimates. In other words, if the pre- and post-test scores were linked and a paired analysis performed, the resulting *P* values would have been smaller than what was observed in the independent testing done by the Wilcoxon rank sum test.⁴⁰

Results

Participants

On February 28, 2011, 10 fourth-year medical students received approximately nine hours of training at the university simulation center. The students volunteered for the course after receiving e-mail from CHAPTER and the College of Medicine. On March 13, 2011, 17 participants from the Veterans Health Administration (VHA) Strategic Health Care Group (EMSHG) were provided the same 9-hour course. The VHA clinicians volunteered for the course after receiving e-mail from CHAPTER and the local VHA medical center. In the second year of the course, 14 fourth-year medical students were trained on April 2, 2012. With the exception of a small percentage of actors and course instructors, recruitment methods and course content did not change in the second year of the course. Trainee descriptions are shown in Table 2, along with medical student and VHA demographic data. Trainees (*N* = 39) included 24 (62%) medical students; seven (18%) physicians; seven (18%) nurses; and one (3%) emergency manager. Of the VHA providers, seven (47%) reported greater than 16 annual hours of disaster training while 63% (*n* = 15) of the medical students reported no annual disaster training. One medical student erroneously self-identified as a mental health provider. After confirming that this was an error, the student’s responses were included in the analysis. In addition, two of the VHA clinicians completed the course but did not complete all components of the online assessments, including the demographic data survey. The responses from these clinicians were excluded from the analysis.

Discrete Knowledge: Before and After EPT

Pre- and post-test data for medical students and VHA clinicians are depicted in Table 3 for before and after EPT as well as at a point four to six months after EPT.

Self-Assessment of EPT Knowledge and Skills

Both medical students and VHA clinicians were asked to assess their EPT skills and comfort levels with performance during a clinical disaster utilizing a 1 to 100 analog scale with higher scores indicating increased comfort levels. Data for medical student skill ratings are depicted in Table 3.

EPT Course Assessment

Trainees were asked to rate the course using an analog scale from 1 (did not meet expectations) to 100 (exceeded expectations). Higher scores indicated positive responses for the evaluation of the training components. These data are presented in Table 3.

Small-Group Exercise Performance

Results of the small-group exercise were as follows:

1. *Team Building: Module 2*—For the seven performance components in the team building exercise, 89% (8 of 9) of the teams “identified a team leader” and 78% (7 of 9) “pre-huddled, team leader assigned roles, and team members understood roles” whereas 69% (6 of 9) of the teams “recognized and discussed the threat.” All teams “completed at least one image puzzle, accurately described scene” and reported a positive learning experience with the team building exercise.
2. *Communication: Module 3*—Eight of nine teams completed all performance components (discussion of chain of command; discussion of relevant verbal communication; and verbalize an accurate, concise and clear report). Teams’ average rating of the “bus crash” scenario during the small group exercise using an analog scale, from 1 (did not meet expectations) to 5 (exceeded expectations), was 4.4 out of 5.
3. *Triage: Module 4*—For the first five performance components (developed team, communicated appropriately with other medical personnel, constructed appropriate scene layout, classified patients using START triage system, and reassessed patients already triaged), 8 of 9 groups completed all. For the last performance component (classified all victims within the time limit), 3 of 9 groups were not able to finish on time. The average number of patients reported by each triage category relative to accurate triage count was red (20.3 vs 21), yellow (17.0 vs 19), green (12.2 vs 12) and black (8.1 vs 8).

Clinical Mass-Casualty Scenario Team Performance

During the 6-minute scenario, nine teams of four to five trainees were asked to mitigate a clinical disaster scene that included three simulated patients and 11 actors. Ten of the actors were patients and one of the actors served as the ED security guard. The zone 1 observer was responsible for rating team performance for all teams (Table 4).

Likert Scale Assessment of Team Performance

There were nine categories of team performance (recognition of disaster, external communication, internal communication, utilization of resources, patient triage and care, patient counseling, personal and

Variable	VHA Clinicians (n = 15)		Medical Students (n = 24)		Combined (N = 39)	
	n	%	n	%	n	%
Male	10	67	12	50	22	56
Non-Hispanic White	9	60	20	83	29	74
Age group, y						
25 or under	0	0	4	17	4	10
26-40	2	13	20	83	22	56
41-55	9	60	0	0	9	23
56+	4	27	0	0	4	10
Educational degree						
Vocational	1	7	0	0	1	3
Bachelor's	2	13	16	67	18	46
Master's	4	27	1	4	5	13
Doctoral	0	0	3	13	3	8
Professional	8	53	4	17	12	31
Occupation						
Nurse	7	47	0	0	7	18
Physician	7	47	21	88	28	72
Emergency Management	1	7	0	0	1	3
Mental health	0	0	1	4	1	3
Other	0	0	2	8	2	5
Years working, y						
Still in training	0	0	22	92	22	56
3-5	1	7	2	8	3	8
6-10	3	20	0	0	3	8
11-20	4	27	0	0	4	10
> 21	7	47	0	0	7	18
Training required, h						
0	5	33	17	71	22	56
1-2	7	47	2	8	9	23
3-8	2	13	2	8	4	10
9-16	1	7	0	0	1	3

Scott © 2013 Prehospital and Disaster Medicine

Table 2. Trainee Demographics and Hours of Disaster Training Required and Performed per Year (*continued*)

Abbreviation: VHA, Veterans Health Administration.

Variable	VHA Clinicians (n = 15)		Medical Students (n = 24)		Combined (N = 39)	
	n	%	n	%	n	%
Training performed, h						
0	0	0	15	63	15	38
1-2	1	7	3	13	4	10
3-8	4	27	4	17	8	21
9-16	3	20	0	0	3	8
17-25	3	20	0	0	3	8
26+	4	27	0	0	4	10

Scott © 2013 Prehospital and Disaster Medicine

Table 2 (*continued*). Trainee Demographics and Hours of Disaster Training Required and Performed per Year
Abbreviation: VHA, Veterans Health Administration.

staff safety, and facility safety and security), and their assessments are depicted in Table 5. For all categories, teams improved their scores after the debriefing (Table 5).

Disaster Scenario Team Performance

Of the 34 performance objectives during the 6-minute scenario, 23 were completed by at least half of the teams during their first attempt. Thirty-one were completed by at least half of the teams during their second attempt. Of the 13 patients in this disaster scene, two simulated patient (2S1 and 3S2) were suffering from occluded airways and could be saved if teams intervened clinically with an airway maneuver prior to the 4-minute mark. Four of nine teams in zones 2 and 3 were able to resuscitate 2S1 during their first attempt; 5 of 9 teams were able to resuscitate 3S2 during their first attempt. All but one of the teams resuscitated 2S1 and 3S2 after the debriefing. In zone 1, 3 of 9 were able to secure the ED door and thwart an exposure to anthrax within four minutes during their first attempt and all teams were able to secure the ED within four minutes during their second attempt. Team performance (ie, teams' ability to resuscitate 2S1 and 3S2 and prevent anthrax exposure) as a function of declining patient stability and scene safety are presented in Figures 2, 3, and 4.

Discussion

Efforts to foster EPT for medical trainees in a simulated environment date back to the 1950s when researchers assessed the use of movies to augment students' training for military and disaster events.⁴¹ The Institute of Medicine's 1999 report, *To Err is Human: Building a Safer Health System*, spurred new development of the use of human patient simulators to train medical providers.⁴² In the 1990s, lessons learned from the aviation industry's Cockpit Resource Management (CRM) led a group of anesthesiologists to pioneer the Acute Crisis Resource Management (ACRM) training module that uses human patient simulators to help reduce error during emergent clinical events.⁴³⁻⁴⁵ Subsequent research has suggested a potential benefit of human simulators to train care providers for clinical emergencies.⁴⁶⁻⁵⁶

Early applications of simulation to emergency medicine include its efficacy to provide emergency medicine team training,⁵⁷ a pilot

study of simulation to provide Emergency Medicine Crisis Resource Management (EMCRM) training⁵⁸ and the theoretical use of multiple patients during simulation to replicate the chaotic environment in the Emergency Department.⁵⁹ Other descriptions of simulation in emergency and military medicine are available.^{9,13,28,57,60-67} Recently a comprehensive list of disaster core competencies for acute care providers was published, but it provides little guidance as to how to incorporate simulation into the disaster classroom.⁶⁸

A literature search in the PubMed database was performed. Additional online searches were completed to capture government reports from, for example, AHRQ (US Agency for Healthcare Research and Quality, www.ahrq.gov). The search yielded approximately 350 articles and reports. Several evidence-based studies were found that established the efficacy of human simulation to train care providers for a disaster.^{12,33,63,69-72} Only one study utilizing a multi-actor clinical disaster scenario to measure the lifesaving performance of medical trainees was found. Wallace quantified the disparity between the times required to resuscitate simulators and actors during a simulated disaster drill. Six simulators and six actors were presented to triage teams in waves of three patients at a time.⁷³ Because the maximum number of patients during an individual triage encounter was limited in the Wallace study, it is not clear whether the tested curricula would have an impact on trainees confronted with several patients at once.

The Center for Health Professional Training and Emergency Response's 1-day EPT course for care providers recreated a loud and chaotic clinical disaster scene (more than 10 human actors with human simulators at one time) and enhanced care provider knowledge, skills, and comfort level necessary to save lives during a simulated disaster. Discrete knowledge of trainees improved significantly across the board after taking the course in 2011 and 2012. Statistically significant improvements were noteworthy for both the medical students and the experienced disaster medical providers from the VHA, suggesting that this course could be suitable to both novice and experienced audiences. Trainee self-assessment of their EPT knowledge and skill also dramatically improved after taking the course. The overall rating for the course

	VHA Clinicians				Medical Students			
	Pre-test	Post-test	P Value	Follow-up	Pre-test	Post-test	P Value	Follow-up
	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	Mean (SD)		Mean (SD)
Cognitive Learning								
Overall scores (out of 24)	15.1 (3.0)	19.4 (2.3)	<.01	18.5 (1.6)	10.6 (3.2)	17.8 (2.0)	<.01	15.3 (2.0)
Percent correct (%)	63.0 (12.0)	81.0 (10.0)	<.01	77.0 (7.0)	44.0 (13.0)	74.0 (8.0)	<.01	64.0 (8.0)
Self Assessments								
Mobilization	69.1 (20.9)	87.5 (9.3)	<.01	82.8 (12.5)	40.6 (23.5)	77.7 (14.0)	<.01	80.2 (11.7)
Team	74.7 (21.7)	88.4 (9.0)	.02	86.2 (12.0)	26.5 (16.5)	74.4 (12.1)	<.01	81.7 (5.5)
Communications	69.9 (23.6)	85.2 (10.1)	.04	76.7 (6.6)	36.2 (17.5)	75.8 (12.8)	<.01	81.0 (6.1)
Triage	69.7 (12.7)	89.1 (7.1)	<.01	88.5 (4.5)	33.3 (19.9)	82.7 (11.9)	<.01	83.3 (11.9)
Incident Command System	71.6 (16.9)	87.6 (10.0)	<.01	87.5 (10.5)	22.5 (19.9)	71.0 (13.8)	<.01	73.0 (9.7)
Overall Knowledge	73.5 (16.7)	83.5 (12.8)	.04	83.0 (9.3)	24.6 (15.2)	71.7 (12.2)	<.01	76.3 (6.0)
Overall Skill	77.8 (18.4)	82.8 (17.3)	.4	88.7 (5.6)	31.7 (15.8)	75.9 (13.5)	<.01	79.0 (3.6)
Course Assessments		Median	Min	Max		Median	Min	Max
Recommend course		92.5	65	100		98.0	78	100
One-day course feasible		82.5	38	100		84.0	67	100
Performance		84.5	60	100		95.0	70	100
Precourse material		81.0	19	100		75.0	50	100
Lectures		90.0	42	100		88.0	71	100
Small group sessions		94.5	71	100		82.0	68	100
Mass casualty		90.0	74	100		93.0	70	100
Presenters		94.0	67	100		100.0	62	100
Facilities		95.0	58	100		97.5	66	100
Overall evaluation		94.5	76	100		100.0	61	100

Scott © 2013 Prehospital and Disaster Medicine

Table 3. Trainee Cognitive Results Before and After Training and Self Assessments, and Course Assessment Results

was 96 out of 100, which represents one of the highest course evaluations completed at the medical university.

The human performance measured during the three small group exercises (team building, communication and triage) demonstrated that nearly all of the groups were able to complete performance components accurately and on time. During the disaster scenario in the afternoon session, trainees were faced with a multi-actor simulated disaster. Overall, teams performed well; two out of three performance objectives were met during the first attempt and teams improved their performance after participating in a short debriefing. Teams successfully identified and performed rescue airway maneuvers to resuscitate two simulated patients who otherwise would have expired. They also identified security threats and implemented measures to prevent exposure to anthrax.

While none of the components of the curriculum are individually unique or novel, the curriculum is competency-based with measurable performance outcomes and potentially applicable to a wide range of medical providers. With the exception of Wallace, this study is unique in that it demonstrates medical trainees' ability to recognize and resuscitate human simulators during a multi-actor simulated clinical disaster. Based on a review of the literature, this is the first published curriculum to use high-fidelity simulation and multi-actor scenarios to measure the lifesaving performance of care providers confronted with more than 10 patients at once.

Limitations

The power assessment of this study (N = 39) is low and the external validity of this curriculum has not been established.

Performance Objective	Patient	Scenario Attempt	
		First (%)	Second (%)
Zone 1			
1. Established roles and responsibilities	All	6 (67)	9 (100)
2. Selected appropriate equipment	All	8 (89)	7 (78)
3. Discussed safety plan prior to action	All	5 (56)	9 (100)
4. Selected and donned appropriate PPE	All	8 (89)	9 (100)
5. Established communication link with EOC	All	8 (89)	8 (89)
6. Effectively communicated needs on scene to EOC	All	4 (44)	7 (78)
7. Followed assigned roles	All	6 (67)	9 (100)
8. Followed chain of command	All	7 (78)	6 (67)
9. Reassessed patients and scene	All	1 (11)	5 (56)
10. Secured doors to ER after recognizing disaster	All	3 (33)	9 (100)
Zone 2			
1. Correctly assigned yellow	2A1	8 (89)	9 (100)
2. Attempted to reassure and calm	2A1	8 (89)	8 (89)
3. Reassessed (no change)	2A1	2 (22)	2 (22)
4. Performed airway maneuver	2S1	4 (44)	8 (89)
5. Assigned red	2S1	4 (44)	7 (78)
6. Reassessed and changed triage to yellow	2S1	0 (0)	3 (33)
7. Reassured and calmed after performing airway maneuver	2S1	0 (0)	2 (22)
8. Assigned green	2A2	7 (78)	9 (100)
9. Reassured and calmed	2A2	8 (89)	8 (89)
10. Reassessed (no change) and sent to green triage staging facility (no change)	2A2	7 (89)	9 (100)
Zone 3			
1. Performed airway maneuver	3S1	6 (67)	7 (78)
2. Assigned black	3S1	7 (78)	9 (100)
3. Reassessed (no change)	3S1	3 (33)	6 (67)
4. Discontinued treatment	3S1	3 (33)	7 (78)
5. Performed airway maneuver	3S2	5 (56)	9 (100)
6. Assigned red	3S2	5 (56)	8 (89)
7. Reassessed	3S2	2 (22)	8 (89)
8. Reassured and calmed	3S2	3 (33)	8 (89)
9. Assigned green	3A1	6 (67)	8 (89)

Scott © 2013 Prehospital and Disaster Medicine

Table 4. Performance Objectives Completed by Nine Teams During Clinical Mass-Casualty Scenario, by Zone (*continued*)
Abbreviations: ER, emergency room; EOC, emergency operations center; PPE, personal protective equipment.

Performance Objective	Patient	Scenario Attempt	
		First (%)	Second (%)
10. Reassured and calmed	3A1	7 (78)	7 (78)
11. Reassessed and sent to green triage staging facility	3A1	5 (56)	8 (89)
12. Assign green	3A2	6 (67)	9 (100)
13. Reassured and calmed	3A2	6 (67)	8 (89)
14. Reassessed and sent to green triage	3A2	7 (78)	8 (89)


Scott © 2013 Prehospital and Disaster Medicine

Table 4 (continued). Performance Objectives Completed by Nine Teams During Clinical Mass-Casualty Scenario, by Zone
Abbreviations: ER, emergency room; EOC, emergency operations center; PPE, personal protective equipment.

Team Assessment Categories	Teams' First Attempt	Teams' Second Attempt
	Mean (SD)	Mean (SD)
Recognition of disaster	5.5 (1.9)	8.1 (1.5)
External communication	6.6 (2.4)	8.0 (2.1)
Internal communication	5.3 (1.8)	7.9 (1.5)
Utilization of resources	6.3 (1.3)	7.6 (1.0)
Patient triage and care	6.5 (1.1)	7.8 (1.6)
Patient counseling	6.4 (2.1)	7.2 (1.7)
Personal and staff safety	5.3 (2.0)	7.4 (2.3)
Facility safety and security	3.4 (2.5)	6.8 (2.4)


Scott © 2013 Prehospital and Disaster Medicine

Table 5. Clinical Mass-Casualty Scenario Team Assessment
1-10 Scale; 1 = did not meet expectations and 10 = exceeded expectations.



Scott © 2013 Prehospital and Disaster Medicine

Figure 2. Median Time to Meet Performance Objectives in Zone 1, by Team Attempt and Safety of Disaster Team



Scott © 2013 Prehospital and Disaster Medicine

Figure 3. Median Time to Meet Performance Objectives in Zone 2, by Team Attempt and Stability of Patient 2S1




Figure 4. Median Time to Meet Performance Objectives in Zone 3, by Team Attempt and Stability of Patient 3S2

A larger-scale study, or preferably a multi-center trial, would allow further analysis of the impact and validity of the curriculum. The use of trained professional actors (who played key patients) as well as trained volunteers (who played patients in the crowd)

References

1. Tokuda Y, Kikuchi M, Takahashi O, Stein GH. Prehospital management of sarin nerve gas terrorism in urban settings: 10 years of progress after the Tokyo subway sarin attack. *Resuscitation*. 2006;68(2):193-202.
2. Barkemeyer BM. Practicing neonatology in a blackout: the University Hospital NICU in the midst of Hurricane Katrina: caring for children without power or water. *Pediatrics*. 2006;117(5 Pt 3):S369-S374.
3. Currier M, King DS, Wofford MR, Daniel BJ, Deshazo R. A Katrina experience: lessons learned. *Am J Med*. 2006;119(11):986-992.
4. Edwards TD, Young RA, Lowe AF. Caring for a surge of Hurricane Katrina evacuees in primary care clinics. *Ann Fam Med*. 2007;5(2):170-174.
5. Hamm LL. Personal observations and lessons from Katrina. *Am J Med Sci*. 2006;332(5):245-250.
6. Kline DG. Inside and somewhat outside Charity. *J Neurosurg*. 2007;106(1):180-188.
7. Leder HA, Rivera P. Six days in Charity Hospital: two doctors' ordeal in Hurricane Katrina. *Compr Ther*. 2006;32(1):2-9.
8. Ciraulo DL, Frykberg ER, Feliciano DV, et al. A survey assessment of the level of preparedness for domestic terrorism and mass casualty incidents among Eastern Association for the Surgery of Trauma members. *J Trauma*. 2004;56(5):1033-1039; discussion 1039-1041.
9. Bartley BH, Stella JB, Walsh LD. What a disaster?! Assessing utility of simulated disaster exercise and educational process for improving hospital preparedness. *Prehosp Disaster Med*. 2006;21(4):249-255.
10. Galante JM, Jacoby RC, Anderson JT. Are surgical residents prepared for mass casualty incidents? *J Surg Res*. 2006;132(1):85-91.
11. Martini SD, Bush AC, Lynch JA. A national survey of terrorism preparedness training among pediatric, family practice, and emergency medicine programs. *Pediatrics*. 2006;118(3):e620-e626.
12. Miller GT, Scott JA, Issenberg SB, et al. Development, implementation and outcomes of a training program for responders to acts of terrorism. *Prehosp Emerg Care*. 2006;10(2):239-246.
13. Scott JA, Miller GT, Issenberg SB, et al. Skill improvement during emergency response to terrorism training. *Prehosp Emerg Care*. 2006;10(4):507-514.
14. Sklar DP, Richards M, Shah M, Roth P. Responding to disasters: academic medical centers' responsibilities and opportunities. *Acad Med*. 2007;82(8):797-800.
15. Coico R, Kachur E, Lima V, Lipper S. Guidelines for preclerkship bioterrorism curricula. *Acad Med*. 2004;79(4):366-375.
16. Waacklerle JF, Seamans S, Whiteside M, et al. Executive summary: developing objectives, content, and competencies for the training of emergency medical technicians, emergency physicians, and emergency nurses to care for casualties resulting from nuclear, biological, or chemical incidents. *Ann Emerg Med*. 2001;37(6):587-601.
17. Association of American Medical Colleges. Training future physicians about weapons of mass destruction. In: *Report of the Expert Panel on Bioterrorism Education*. Washington, DC: Association of American Medical Colleges; 2008.
18. National Research Council. *Hospital-Based Emergency Care: At the Breaking Point*. Washington, DC: The National Academies Press; 2007.
19. Scott LA, Carson DS, Greenwell IB. Disaster 101: a novel approach to disaster medicine training for health professionals. *J Emerg Med*. 2010;39(2):220-226.
20. Association of American Medical Colleges. Number of US medical schools teaching selected topics 2004-2005. http://services.aamc.org/curdir/section2/04_05hottopic.pdf. Accessed November 1, 2008.
21. Rico E, Trepka M, Zhang G, et al. Knowledge and attitudes about bioterrorism and smallpox: a survey of physicians and nurses. Miami, FL: Miami-Dade County Health Department, Office of Epidemiology and Disease Control. *Epi Monthly Report*. 2002;3(7):1-7.
22. Lanzilotti SS, Galanis D, Leoni N, Craig B. Hawaii medical professionals assessment. *Hawaii Med J*. 2002;61(8):162-173.
23. Gershon RR, Canton AN, Magda LA, DiMaggio C, Gonzalez D, Dul MW. Web-based training on weapons of mass destruction response for emergency medical services personnel. *Am J Disaster Med*. 2009;4(3):153-161.
24. Chen J, Wilkinson D, Richardson RB, Waruszynski B. Issues, considerations and recommendations on emergency preparedness for vulnerable population groups. *Radiat Prot Dosimetry*. 2009;134(3-4):132-135.
25. Subbarao I, Lyznicki JM, Hsu EB, et al. A consensus-based educational framework and competency set for the discipline of disaster medicine and public health preparedness. *Disaster Med Public Health Prep*. 2008;2(1):57-68.
26. Williams J, Nocera M, Casteel C. The effectiveness of disaster training for health care workers: a systematic review. *Ann Emerg Med*. 2008;52(3):211-222, 22 e1-e2.
27. Improving Patient Safety Through Simulation Research. Agency for Healthcare Research and Quality (US) web site. <http://www.aahrq.gov/legacy/qual/simul-proj11.htm>. Accessed March 23, 2013.
28. Hsu EB, Jenckes MW, Catlett CL, et al. Training to hospital staff to respond to a mass casualty incident. *Evid Rep Technol Assess*. (Summ) 2004(95):1-3.
29. Scott LA. Disaster 101: a novel approach to health care students' disaster medicine and emergency preparedness training. *Disaster Med Public Health Prep*. 2009;3(3):139-140.
30. Scott L, Ross A, Schnellmann J, Wahlquist A. Surge Capacity: CHAPTER and the South Carolina Healthcare Worker Preparedness. *The Journal of the South Carolina Medical Society*. 2011;107:74-77.
31. Barbera JA, Yeatts DJ, Macintyre AG. Challenge of hospital emergency preparedness: analysis and recommendations. *Disaster Med Public Health Prep*. 2009; 3(2 Suppl):S74-S82.

was utilized to minimize variations in human behavior, improving but not perfecting the study's reliability.

The study sought to measure the impact of EPT on discrete knowledge both immediately after and four to six months after the training day. Unfortunately, these results were not statistically significant, largely because several trainees were lost to follow-up during the first year of the course. A more robust predesign and postdesign with appropriate follow-up could sufficiently power this comparison. In addition, trainees were not tested prior to the disaster scenario (before the training day), primarily because of time and financial constraints. This limits the study's ability to ascertain a performance baseline for the trainees.

Future iterations of CHAPTER curricula will need to include a cost assessment to demonstrate the feasibility of this curriculum for target audiences lacking human simulation capability. A version of the curriculum that uses low-cost, low-fidelity simulators and actors is currently under production by CHAPTER.

Conclusion

This one-day EPT course improved care provider knowledge, skill, and comfort necessary to save lives during a simulated disaster. Trainees were able to resuscitate two unstable simulated patients and prevent anthrax exposure to the hospital during a chaotic and loud disaster scenario. A larger-scale study, or preferably a multi-center trial, is needed to further study the impact of this curriculum and its potential to protect provider and patient lives.

32. The George Washington University Institute for Crisis Disaster and Risk Management. VHA-EMA Certification Program, Healthcare Emergency Management Professional Certification Program, Final Program Recommendations. http://www.gwu.edu/~icdr/publications/PDF/CertificationProgramRecommendations_FinalReport.pdf. Accessed March 23, 2013.
33. Subbarao I, Bond WF, Johnson C, Hsu EB, Wasser TE. Using innovative simulation modalities for civilian-based, chemical, biological, radiological, nuclear, and explosive training in the acute management of terrorist victims: A pilot study. *Prehosp Disaster Med.* 2006;21(4):272-275.
34. Agency for Healthcare Research and Quality. Tool for Evaluating Core Elements of Hospital Disaster Drill. AHRQ Publication No 08-0019. Rockville, Maryland USA; 2008.
35. King HB, Battles J, Baker DP, et al. Team STEPPS: Team Strategies and Tools to Enhance Performance and Patient Safety (Performance and Tools). http://www.ncbi.nlm.nih.gov/books/NBK43686/pdf/advances-king_1.pdf. Accessed March 23, 2013.
36. Markenson D, DiMaggio C, Redlener I. Preparing health professions students for terrorism, disaster, and public health emergencies: core competencies. *Acad Med.* 2005;80(6):517-526.
37. Smith P, Ragan T. Instructional design, 3rd ed. New York: John Wiley & Sons; 2005.
38. Krathwohl D, Bloom B, Masia B. Taxonomy of educational objectives; the classification of educational goals. New York: Longman, Green; 1964.
39. Bloom B, Engelhart M, Furst E, Hill W, Krathwohl D. Taxonomy of educational objectives: the classification of educational goals. New York: Longmans, Green; 1956.
40. Diehr P, Martin D, Koepsell T, Cheadle A. Breaking the matches in a paired /-test for community interventions when the number of pairs is small. *Statistics in Medicine.* 1995;14(13):1491-1504.
41. Ruhe DS, Byfield GV. Audiovisual aids for disaster and military medicine in the medical schools. *J Med Educ.* 1954;29(81):59-62.
42. Kohn LT, Corrigan J, Donaldson MS. To err is human: building a safer health system. Washington, DC: National Academy Press; 2000.
43. Howard SK, Gaba DM, Fish KJ, Yang G, Sarnquist FH. Anesthesia crisis resource management training: teaching anesthesiologists to handle critical incidents. *Aviat Space Environ Med.* 1992;63(9):763-770.
44. Wiener EL, Kanki BG, Helmreich RL. Cockpit resource management. San Diego: Academic Press; 1993.
45. Murray DJ, Boulet JR, Kras JF, Woodhouse JA, Cox T, McAllister JD. Acute care skills in anesthesia practice: a simulation-based resident performance assessment. *Anesthesiology.* 2004;101(5):1084-1095.
46. Wayne DB, Didwania A, Feinglass J, Fudala MJ, Barsuk JH, McGaghie WC. Simulation-based education improves quality of care during cardiac arrest team responses at an academic teaching hospital: a case-control study. *Chest.* 2008; 133(1):56-61.
47. Wayne DB, Butter J, Siddall VJ, et al. Mastery learning of advanced cardiac life support skills by internal medicine residents using simulation technology and deliberate practice. *J Gen Intern Med.* 2006;21(3):251-256.
48. Rosenthal ME, Adachi M, Ribaldo V, Mueck JT, Schneider RF, Mayo PH. Achieving housestaff competence in emergency airway management using scenario based simulation training: comparison of attending vs housestaff trainers. *Chest.* 2006;129(6):1453-1558.
49. Tsai TC, Harasym PH, Nijssen-Jordan C, Jennett P. Learning gains derived from a high-fidelity mannequin-based simulation in the pediatric emergency department. *J Formos Med Assoc.* 2006;105(1):94-98.
50. Shavit I, Keidan I, Hoffmann Y, et al. Enhancing patient safety during pediatric sedation: the impact of simulation-based training of nonanesthesiologists. *Arch Pediatr Adolesc Med.* 2007;161(8):740-743.
51. DeVita MA, Schaefer J, Lutz J, Wang H, Dongilli T. Improving medical emergency team (MET) performance using a novel curriculum and a computerized human patient simulator. *Qual Saf Health Care.* 2005;14(5):326-331.
52. Franc-Law J, Ingrassia P, Ragazzoni L, Corte F. The effectiveness of training with an emergency department simulator on medical student performance in a simulated disaster. *CJEM.* 2010;12(1):27-32.
53. Gordon JA, Shaffer DW, Raemer DB, Pawlowski J, Hurford WE, Cooper JB. A randomized controlled trial of simulation-based teaching versus traditional instruction in medicine: a pilot study among clinical medical students. *Adv Health Sci Educ Theory Pract.* 2006;11(1):33-39.
54. Morgan PJ, Cleave-Hogg D, Desousa S, Lam-McCulloch J. Applying theory to practice in undergraduate education using high fidelity simulation. *Med Teach.* 2006;28(1):e10-e15.
55. Ten Eyck RP, Tews M, Ballester JM. Improved medical student satisfaction and test performance with a simulation-based emergency medicine curriculum: a randomized controlled trial. *Ann Emerg Med.* 2009;54(5):684-691.
56. Wong G, Jenkins C, Yao TJ, Irwin M. A trend toward improved learning of cardiovascular pathophysiology in medical students from using a human patient simulator: results of a pilot study. *Adv Physiol Educ.* 2007;31(4):372.
57. Small SD, Wuerz RC, Simon R, Shapiro N, Conn A, Setnik G. Demonstration of high-fidelity simulation team training for emergency medicine. *Acad Emerg Med.* 1999;6(4):312-323.
58. Reznek M, Smith-Coggins R, Howard S, et al. Emergency medicine crisis resource management (EMCRM): pilot study of a simulation-based crisis management course for emergency medicine. *Acad Emerg Med.* 2003;10(4):386-389.
59. Kobayashi L, Shapiro MJ, Gutman DC, Jay G. Multiple encounter simulation for high-acuity multipatient environment training. *Acad Emerg Med.* 2007;14(12): 1141-1148.
60. Andreatta PB, Maslowski E, Petty S, et al. Virtual reality triage training provides a viable solution for disaster-preparedness. *Acad Emerg Med.* 2010;17(8):870-876.
61. Eaves RH, Flagg AJ. The U.S. Air Force pilot simulated medical unit: a teaching strategy with multiple applications. *J Nurs Educ.* 2001;40(3):110-115.
62. Decker SI, Galvan TJ, Sridaromont K. Integrating an exercise on mass casualty response into the curriculum. *J Nurs Educ.* 2005;44(7):339-340.
63. Vincent DS, Burgess L, Berg BW, Connolly KK. Teaching mass casualty triage skills using iterative multimakin simulations. *Prehosp Emerg Care.* 2009;13(2):241-246.
64. Coule PL, Schwartz RB. The national disaster life support programs: a model for competency-based standardized and locally relevant training. *J Public Health Manag Pract.* 2009;15(2 Suppl):S25-S30.
65. Fritz PZ, Gray T, Flanagan B. Review of mannequin-based high-fidelity simulation in emergency medicine. *Emerg Med Australas.* 2008;20(1):1-9.
66. LeRoy Heinrichs W, Youngblood P, Harter PM, Dev P. Simulation for team training and assessment: case studies of online training with virtual worlds. *World J Surg.* 2008;32(2):161-170.
67. Schumacher J, Runte J, Brinker A, Prior K, Heringlake M, Eichler W. Respiratory protection during high-fidelity simulated resuscitation of casualties contaminated with chemical warfare agents. *Anaesthesia.* 2008;63(6):593-598.
68. Schultz CH, Koenig KL, Whiteside M, Murray R. Development of national standardized all-hazard disaster core competencies for acute care physicians, nurses, and EMS professionals. *Ann Emerg Med.* 2012;59(3):196-208.e1.
69. Kyle RR, Via DK, Lowy RJ, Madsen JM, Marty AM, Mongan PD. A multidisciplinary approach to teach responses to weapons of mass destruction and terrorism using combined simulation modalities. *J Clin Anesth.* 2004;16(2):152-158.
70. Lerner E, Schwartz R, Coule P, Pirralo R. Use of SALT triage in a simulated mass-casualty incident. *Prehosp Emerg Care.* 2010;14:21-25.
71. Summerhill EM, Mathew MC, Stipho S, et al. A simulation-based biodefense and disaster preparedness curriculum for internal medicine residents. *Med Teach.* 2008;30(6): Schultz CH, Koenig KL, Whiteside M, Murray R. Development of national standardized all-hazard disaster.
72. Gillett B, Peckler B, Sinert R, et al. Simulation in a disaster drill: comparison of high-fidelity simulators versus trained actors. *Acad Emerg Med.* 2008;15(11):1144-1151.
73. Wallace D, Gillett B, Wright B, Stetz J, Arquilla B. Randomized controlled trial of high fidelity patient simulators compared to actor patients in a pandemic influenza drill scenario. *Resuscitation.* 2010;81(7):872-876.