

# Heritability and short-term effects of inbreeding in the progenetic trematode *Coitocaecum parvum*: is there a need for the definitive host?

C. LAGRUE\* and R. POULIN

Department of Zoology, University of Otago, P.O. Box 56, Dunedin 9054, New Zealand

(Received 23 September 2008; revised 20 October 2008; accepted 22 October 2008; first published online 18 December 2008)

## SUMMARY

Self-fertilization (or selfing), defined as the fusion of male and female reproductive cells originating from the same individual, is the most extreme case of inbreeding. Although most hermaphroditic organisms are in principle able to self-fertilize, this reproductive strategy is commonly associated with a major disadvantage: inbreeding depression. Deleterious effects due to the loss of genetic diversity have been documented in numerous organisms including parasites. Here we studied the effects of inbreeding depression on the offspring of the progenetic trematode *Coitocaecum parvum*. The parasite can use 2 alternative life-history strategies: either it matures early, via progenesis, and produces eggs by selfing in its second intermediate host, or it waits and reproduces by out-crossing in its definitive host. We measured various key parameters of parasite fitness (i.e. hatching and multiplication rates, infectivity, survival) in offspring produced by both selfing and out-crossing. Altogether, we found no significant difference in the fitness of offspring from progenetic (selfing) and adult (out-crossing) parents. In addition, we found no evidence that either strategy (progenesis or the normal three-host cycle) is heritable, i.e. the strategy adopted by offspring is independent of that used by their parents. Although it is unclear why both reproductive strategies are maintained in *C. parvum* populations, our conclusion is that producing eggs by selfing has few, if any, negative effects on parasite offspring. Inbreeding depression is unlikely to be a factor acting on the maintenance of the normal three-host life cycle, and thus out-crossing, in *C. parvum* populations.

Key words: *Coitocaecum parvum*, inbreeding depression, progenesis, selfing, out-crossing, abbreviated life cycle.

## INTRODUCTION

The mode of reproduction of a species is a major factor influencing its genetic structure and therefore its evolutionary trajectory (Lymbery *et al.* 1997). In hermaphroditic organisms, individuals can either reproduce by self-fertilization (i.e. selfing: fusion of male and female gametes from the same individual) or cross-fertilization (i.e. out-crossing). These two different reproductive strategies can have drastic effects on the genetic structure of populations. For example, studies of plants have shown that, compared to species using out-crossing, predominantly selfing species show significant heterozygote deficiencies and linkage disequilibrium (Loveless and Hamrick, 1984; Heywood, 1991). Hermaphroditism is also widespread in animals but the frequency with which self-fertilization is preferred over cross-fertilization is unknown for most hermaphrodite species (Jarne, 1995).

While selfing can be seen as advantageous by removing the need for a sexual partner, it is also a severe case of inbreeding: both male and female reproductive cells originate from the same parent.

The main cost of this strategy is thought to be genetic deficiencies arising from inbreeding depression and inducing fitness costs in offspring produced by self-fertilization (Charlesworth and Charlesworth, 1987). Because selfing increases homozygosity, inbreeding depression results from the accumulation of deleterious recessive alleles and/or the lower performance of homozygotes compared to heterozygote individuals (Trouvé *et al.* 1996). Genetically homogeneous offspring may also be less likely to survive fluctuating environments (Christen and Milinski, 2003). Thus, the lower fitness of inbred progeny should strongly select against self-fertilization and for the maintenance of reproduction via mating (Milinski, 2006). Whatever the causes of inbreeding depression, it is possible that its effects have played a part in the evolution of breeding systems avoiding, or at least reducing the occurrence of self-fertilization in natural populations. The benefits of out-crossing, as opposed to selfing, are thought to have even wider implications for life-history strategies of both plants and animals. For example, they may have led to the evolution and maintenance of complex life cycles in parasites (Brown *et al.* 2001; Christen *et al.* 2002).

From simple beginnings, parasitic helminths have evolved complex life cycles so challenging that they

\* Corresponding author: Tel: +64 3 479 7964. Fax: +64 3 479 7584. E-mail: lagcl981@student.otago.ac.nz

impose strong selective pressures on these parasite species (Poulin and Cribb, 2002; Poulin, 2007). For example, the typical life cycle of digenean trematodes involves 3 different hosts and as many transmission events. First, eggs are released by adult worms within the definitive host and later hatch into free-swimming larvae (miracidia) which must find and penetrate the mollusc first intermediate host. Second, miracidia develop and multiply into structures (sporocysts) producing free-living cercariae that emerge from the mollusc host to infect the next intermediate host where they encyst as metacercariae (Combes *et al.* 2002). Third, metacercariae must be ingested, along with their intermediate host, by the appropriate definitive host, which is always a vertebrate, for the life cycle to be completed. Complex life cycles also provide several advantages to the parasites such as longer life span, greater adult body size, higher fecundity (Parker *et al.* 2003) and, more importantly, greater access to sexual partners (Brown *et al.* 2001); facilitating cross-fertilization by concentrating conspecific individuals in the same predator definitive host (Poulin and Cribb, 2002; Rauch *et al.* 2005).

However, some trematodes have developed a radical adaptation with 1 host (sometimes 2), most often the vertebrate definitive host, being dropped from the cycle (Lefebvre and Poulin, 2005*a*). The resulting abbreviated life cycle should be easier to complete as the number of transmission events is decreased (Poulin and Cribb, 2002). Shorter life cycles are likely to be favoured in parasites faced with low or highly variable probabilities of transmission. In fact, life-cycle abbreviations from 3 to 2 hosts have occurred independently in numerous trematode lineages (Poulin and Cribb, 2002; Lefebvre and Poulin, 2005*a,b*). This is usually achieved via progenesis: metacercariae mature precociously while still inside the second intermediate host and produce viable offspring by selfing without the need to reach the definitive host. Given that progenesis could serve as a reproductive insurance against failed transmission or even extinction (Bush and Kennedy, 1994), it is surprising that shorter life cycles are not more widespread among trematodes (Lefebvre and Poulin, 2005*c*). It is possible that fitness costs associated with shorter life cycles offset the increased likelihood of completion. Two obvious costs are associated with offspring quality and quantity. First, fecundity may be reduced in abbreviated life cycles as egg production is likely to be much lower inside a small, short-lived invertebrate host than in the larger and longer-lived vertebrate definitive host (Poulin, 2001; Poulin and Cribb, 2002; Lefebvre and Poulin, 2005*a*). Second, trematodes adopting progenesis can only reproduce by selfing within their intermediate host because they are individually enclosed within a cyst, causing inbreeding. Therefore, shorter life cycles could

lead to the production of offspring with low genetic heterogeneity, potentially less viable than offspring produced by out-crossing. The deleterious effects of homozygosity induced by self-fertilization are commonly believed to explain why progenesis is not more widespread (Lefebvre and Poulin, 2005*a*).

It is also possible that trematode alternative life cycles, and hence reproductive mode, are genetically fixed. The co-existence of both normal and abbreviated cycles in parasite populations could thus reflect an underlying genotypic variation such that different parasite strains, using alternative life cycles, co-exist in the same population (Crossan *et al.* 2007). Because the availability of definitive hosts is likely to vary seasonally, each parasite strain's strategy may be periodically advantageous or detrimental but both could be maintained over time by environmental stochasticity. In this case, offspring of parasites reproducing through abbreviated life cycles should be more likely to adopt the shorter route themselves. Trematode species using both the normal three-host cycle and a shorter cycle as alternative strategies are consequently ideal models to determine the heritability of each strategy and to measure the effects of inbreeding depression on important fitness parameters such as infectivity, growth and virulence of the offspring produced by progenetic parasites (Wedekind *et al.* 1998).

The trematode *Coitocaecum parvum* uses either the typical three-host life cycle or a truncated life cycle in which metacercariae mature precociously (progenesis) inside the amphipod second intermediate host and start producing both eggs and sperm. Progenetic metacercariae from amphipod hosts examined by Holton (1984*a*), although enclosed within their cyst, had active spermatozoa in the seminal vesicle and seminal receptacle and viable eggs are most likely produced by self-fertilization rather than asexually. Both strategies are observed synchronously in *C. parvum* populations with up to 60% of the parasites being progenetic (Lefebvre and Poulin, 2005*c*) and, although environmental factors can influence the parasite strategy (Poulin, 2003; Lagrue and Poulin, 2007), it is still unclear why not all parasites use progenesis as a reproductive insurance against failed transmission. Because progenetic individuals are enclosed within cysts and can only self-fertilize, it has been suggested that their offspring could be of lower quality (Poulin, 2001). However, the viability and infectivity of *C. parvum* offspring produced by both progenetic individuals inside the second intermediate host and adult worms in the definitive host are yet to be compared. The adoption of either strategy in *C. parvum* could also be under genetic influence: offspring from progenetic parents simply adopting progenesis as an inherited feature and therefore reproducing by selfing, and vice-versa for individuals that out-crossed in the

definitive host. Once again, this heritability hypothesis remains to be tested.

In this study, we used *C. parvum* to test for the effects of alternative life strategies on the fitness of offspring produced by selfing via the abbreviated cycle and by adult worms in the definitive host. Fitness parameters such as viability and/or infectivity to the next host, used in previous studies on inbreeding depression (Christen *et al.* 2002; Christen and Milinski, 2003; Milinski, 2006), were measured as an estimate of offspring fitness in *C. parvum*. We compared the hatching rate of eggs from both origins, infection success of free-living larvae (miracidia and cercariae) and asexual multiplication within the snail first intermediate host. Furthermore, testing whether trematode life cycles, and hence reproductive strategies, are genetically fixed or environmentally adaptable is an important step towards the understanding of the evolution of complex life cycles in parasites. By experimentally infecting crustacean intermediate hosts, we were also able to compare the proportions of progenetic individuals between metacercariae from selfing and out-crossing parents and consequently determine whether progenesis and life-cycle abbreviation are inherited characters in *C. parvum*.

#### MATERIALS AND METHODS

##### *Life cycle of Coitocaecum parvum*

*Coitocaecum parvum* (Trematode, Opencolidae) is a common parasite of freshwater fish in New Zealand, mostly the common bully (*Gobiomorphus cotidianus*; Macfarlane, 1939; Holton, 1984*b*). Eggs produced by adult worms inside the fish gut are released in host faeces and hatch into free-swimming miracidia. These penetrate mud snails (Hydrobiidae, *Potamopyrgus antipodarum*) in which they mature as a mother sporocyst before asexually producing daughter sporocysts. Each daughter sporocyst then asexually produces cercariae that actively leave the snail host and enter the amphipod *Paracalliope fluviatilis* where they encyst as metacercariae in the body cavity. At this stage, metacercariae can either await ingestion by a fish where they will mature and reproduce, or keep growing and reach maturity while still inside the amphipod. Worms that reach maturity in the crustacean intermediate host reproduce by selfing and lay eggs that are contained in their cyst (Holton, 1984*a*; Poulin, 2001); eggs are released after host death and decomposition. Both alternative life cycles occur simultaneously in natural *C. parvum* populations.

##### *Animal collection*

Snails, amphipods and fish were all collected in Lake Waihola, South Island, New Zealand during spring

and summer 2006. Amphipods and snails were captured by dragging a dip net (mesh size 500  $\mu\text{m}$ ) through macrophytes (*Myriophyllum triphyllum*). All amphipods and snails were returned alive to the laboratory and kept separated in aerated lake water with strands of *M. triphyllum* as food source. Fish were captured using a seine net and kept in groups of 10 in 25-litre containers filled with aerated lake water. Fish were fed exclusively with live amphipods from Lake Waihola for several weeks to increase the intensity of infection with *C. parvum*. Natural prevalence of *C. parvum* infection in bullies is usually 100% and mating pairs are regularly observed during dissection. However, the average infection intensity is low (between 5 and 10 worms per fish; Poulin, 2001; Lefebvre and Poulin, 2005*b*), although some naturally infected fish can contain over 30 and up to 70 *C. parvum* adults (Lefebvre and Poulin, 2005*b*; personal observations). Bullies experimentally fed on amphipods contained between 60 and 100 adult *C. parvum* with an average of around 75. Adult *C. parvum* gather in the rectal region of the fish gut and, even in naturally and artificially heavily infected hosts, we did not observe displacement of individuals to other areas of the digestive tract of the host that might be expected with overcrowding (personal observations). Therefore, by increasing the encounter rate between individual parasites, high parasite densities should enhance the opportunities for cross-fertilization without modifying parasite behaviour. In fact, all laboratory fed bullies contained mating pairs of parasites at the time of dissection and, on average, 5 mating pairs were found per fish (personal observations). However, this only represents a snapshot of the situation as these parasites can store sperm after mating. In such high densities, although only a low proportion of worms are actually observed mating, most if not all individuals have had access to a mating partner and possibly stored sperm for cross-fertilization. The number of eggs produced by selfing in these worms should therefore be much lower than in progenetic metacercariae (Lefebvre and Poulin, 2005*b*).

##### *Hatching success of eggs*

*C. parvum* eggs were obtained from both progenetic metacercariae and adult worms. Because the prevalence of *C. parvum* is usually low (<10%) in second intermediate hosts, amphipods were screened under a dissecting microscope and only individuals judged as infected were dissected (Lefebvre and Poulin, 2005*b*). In total, 150 amphipods were found to contain a progenetic metacercaria. Approximately 35 eggs per metacercaria were taken and transferred into a Petri dish containing 10 ml of filtrated lake water. Eggs from adult *C. parvum* in fish were obtained by dissecting the experimentally fed bullies. Adult worms found in fish guts were kept for 24 h in lake

water to induce them to release their eggs. After that time, all eggs liberated by the parasites were transferred to another Petri dish filled with water. Typically, only few eggs are released daily by adult parasites (Lefebvre and Poulin, 2005b) and, in our study, we obtained 400 adult *C. parvum* that released around 3 eggs each. The large number of parasites used as sources of eggs for both categories should reduce the potential confounding factor of individual variation in egg quality. The 2 pools of eggs (from progenetic metacercariae and from adult worms; i.e. 'progenetic' and 'normal' eggs) were separated into groups of 20–25 and transferred to 96-well microtitre plates, each well filled with 200  $\mu$ l of lake water. The number of hatched eggs was counted everyday until all were hatched or no more were seen to be hatching. At the end of the experiment, 450 eggs of each origin (progenetic or normal) were haphazardly chosen and measured, 300 that hatched and 150 that failed to hatch. Assuming a regular ellipsoid shape, individual egg volume was calculated as  $V_{\text{egg}} = (\pi * L * W^2) / 6$ , where L and W are egg length and width, respectively. The proportions of hatched eggs from progenetic metacercariae and adult worms were compared using a Fisher's exact test. A two-way ANOVA with egg type (progenetic or normal) and hatching status (hatched or failed) as independent factors, and egg volume as the dependent variable, was also used to test for any size differences between each type of egg. Egg volume was log-transformed before analysis to normalize the data.

#### *Infectivity of miracidia to the snail first intermediate host*

Eggs were again obtained from both progenetic individuals and adults as described above. They were then transferred by groups of 10 to Petri dishes containing 4 ml of lake water and 1 uninfected snail per dish was later added. Eggs were protected from the snail by a wire-netted circular cage (10 mm diameter; mesh size 0.5 mm) glued to the bottom of the container. Uninfected snails used in this experiment were selected by size and shell shape. Altered shell shape being characteristic of infection by *C. parvum*, any individual displaying unusual shells were discarded (Lagrue *et al.* 2007a). Additionally, 100 snails judged as uninfected were dissected to verify the accuracy of our screening technique and none were found to be infected by *C. parvum*. Only immature snails (2 mm < shell length < 3 mm at the time of experimental infections) were used for this experiment as mature snails are less likely to get infected (Zakikhani and Rau, 1999; personal observation). Petri dishes were checked everyday until 1 egg hatched. At that time, the remaining eggs were taken out of the Petri dish; each snail was therefore exposed to only 1 miracidium. Snails were kept individually in lake water and fed with strands of

macrophytes for the rest of the experiment. A total of 600 snails were exposed to 1 *C. parvum* larva each, half of those to miracidia from progenetic eggs and the other half from eggs laid by adult worms. After 10 weeks, half of the snails in each treatment were measured to assess for possible differences in shell growth and their shell volume was calculated using the formula  $V = 1/3(\pi r^2 h)$  where V is the volume, r the radius (half the width of whorl 1) and h the vertical height (total length; see Lagrue *et al.* 2007a for details). Snails were then dissected to assess for infection by *C. parvum*. When infections were found, the number of sporocysts was counted. The remaining snails were used to obtain cercariae for subsequent experimental infections of amphipods. These were also measured to assess for possible differences in shell growth, and dissected 10 months post-infection and again, both the proportion of *C. parvum* infected snails and the number of sporocysts were recorded. The proportions of snails infected by miracidia from selfed and out-crossed eggs were compared using Fisher's exact tests. The effect of miracidial origin (selfed or out-crossed eggs) on the number of sporocysts inside the snail was tested using a non-parametric test (Mann-Whitney U test).

#### *Infectivity of cercariae to the amphipod second intermediate host and life-history strategy of metacercariae*

Uninfected amphipods used for experimental infections were obtained by inspecting each amphipod under a microscope and discarding all amphipods that showed any sign of infection, i.e. an opaque mass in the body cavity corresponding to a metacercaria. This method allows the selection of only uninfected individuals with an accuracy of about 95% (Lefebvre and Poulin, 2005b). Also, only male amphipods of similar size (mean body length of  $3.0 \pm 0.0$  mm, maximum 3.2 mm and minimum 2.8 mm respectively) were used to reduce the effects of host conditions on the parasite. Snails experimentally infected in the previous experiment were used as sources of cercariae for this part of the study. Cercariae of *C. parvum* were obtained under controlled conditions to ensure that the cercariae used to experimentally infect amphipods had recently emerged and, therefore, were more likely to penetrate the amphipod. For each round of experimental infections, snails were transferred to large Petri dishes filled with 10 ml of filtrated lake water. Snails were incubated at 25 °C for 20 min under constant light, conditions known to induce cercarial release (Hay *et al.* 2005). The Petri dishes were then screened under a microscope and the cercariae transferred to 500  $\mu$ l Eppendorf tubes using a 20  $\mu$ l micropipette. One cercaria was placed in each tube with 2.5  $\mu$ l of filtrated lake water and an uninfected



Table 1. Results of Fisher's exact tests for comparison between offspring from 'progenetic' and 'normal' eggs (Note that all values for fitness measurements correspond to proportions (%) of success.)

	Reproduction origin		Fisher's exact test	
	'Progenetic'	'Normal'	$\chi^2$	P-value
Egg hatching success	70	77.3	26.23	<0.0001
Miracidia infection success at 10 weeks	14.2	16.9	0.43	0.514
Miracidia infection success at 10 months	13.8	15.1	0.08	0.771
Cercariae infection success	65.5	63.6	0.17	0.680
Metacercariae survival at 5 weeks	66.7	65.6	0.03	0.861
Proportion of progenetic metacercariae	23.3	24.4	0.03	0.866

amphipod was then added. Amphipods were left in the tube along with the cercariae for 5 h, a time after which unsuccessful cercariae stop moving and die. Amphipod survival, at this stage, is usually over 99%. For logistical reasons, amphipods were then haphazardly separated into groups of 5 individuals and placed in plastic Falcon tubes filled with 10 ml of lake water and a strand of macrophyte (*Elodea canadensis*) was added for food. After 5 weeks, a period sufficient for *C. parvum* to adopt one or the other strategy (Lagrue and Poulin, 2007), the 412 surviving amphipods were dissected under the microscope to assess their infection status. Metacercariae found in amphipods were recorded as dead (i.e. encapsulated and melanized; Thomas *et al.* 2000; Bryan-Walker *et al.* 2007) or alive (i.e. moving and within a transparent cyst). Live individuals were measured (length and width) and recorded as 'normal' (non egg-producing worm) or 'progenetic' (egg-producing worm); in the case of progenetic parasites, eggs were also counted. These included both eggs released by the worm in its thin-walled cyst and eggs still *in utero*. The body surface of each parasite was then determined and used as a surrogate for body size. This was done using the formula for an ellipsoid,  $(\pi LW)/4$ , where L and W are the length and width of the parasite. The proportion of cercariae, from progenetic and normal descent, successfully penetrating the host and then developing into metacercariae, as well as the proportion of metacercariae adopting progenesis among offspring originating from either reproductive strategy were compared using Fisher's exact tests. The possible effects of parental strategy on metacercarial body size and egg production were assessed using non-parametric tests (Mann-Whitney U test).

## RESULTS

### Hatching success of eggs

A total of 6415 eggs, 5170 from progenetic metacercariae and 1245 from adult worms, were used in this study. Eggs were observed hatching from day 1 to day 35 and no more eggs hatched after that time.

The overall hatching success was 71.4% but there was a significant difference in the hatching success between eggs from progenetic metacercariae (70%) and those from adult worms (77.3%; Table 1). The daily hatching rate of each type of egg was also clearly distinct (Fig. 1): eggs laid by adult worms did not start hatching before day 9 and a large majority (91.7%) hatched between day 12 and day 20. In contrast, eggs from progenetic individuals started hatching from day 1 and a steadier hatching rate was observed during the experiment. Progenetic eggs were also significantly larger (mean  $\pm$  s.e.,  $112.95 \pm 1.1 \times 10^3 \mu\text{m}^3$ ) than normal eggs ( $108.9 \pm 1.1 \times 10^3 \mu\text{m}^3$ ; ANOVA,  $F_{1,896} = 19$ ,  $P < 0.0001$ ; Fig. 2) and, overall, failed eggs were significantly larger than eggs that successfully hatched ( $128.83 \pm 1.5$  and  $101.97 \pm 0.6 \times 10^3 \mu\text{m}^3$  respectively; ANOVA,  $F_{1,896} = 361$ ,  $P < 0.0001$ ). However, there was a significant interaction between the two factors (ANOVA,  $F_{1,896} = 14$ ,  $P = 0.0002$ ): while failed progenetic eggs were much larger than failed normal eggs ( $134.76 \pm 2$  and  $122.89 \pm 2.1 \times 10^3 \mu\text{m}^3$  respectively; Fisher's LSD, D.F. = 896,  $P < 0.0001$ ), there was no significant difference between progenetic and normal eggs that successfully hatched ( $102.05 \pm 0.8$  and  $101.89 \pm 1 \times 10^3 \mu\text{m}^3$  respectively; Fisher's LSD, D.F. = 896,  $P = 0.625$ ; Fig. 2).

### Infectivity of miracidia to the snail first intermediate host

After 10 weeks, none of the 600 experimentally infected snails were producing cercariae. Among the 300 snails dissected at that time, the mortality was very low (1%) and snail growth was minimal. The overall prevalence of *C. parvum* was 15.5% and no difference was found in the proportion of infected snails between individuals exposed to miracidia from progenetic (14.2%) and normal eggs (16.9%; Table 1). After 10 months, the mortality among the 300 remaining snails was 18.4% and *C. parvum* prevalence was 14.5%. The mortality was similar between the two treatment groups (19.4% and 17.4% respectively, Fisher's exact test,  $\chi^2 = 0.21$ ,  $P = 0.648$ ) and again, no difference was found in the proportions




Fig. 1. Daily hatching rate in terms of the proportion (%) of the total number of eggs ('progenetic' or 'normal') that had hatched at the end of the experiment.




Fig. 2. Mean volumes ( $\pm$  S.E.) of 'progenetic' and 'normal' eggs, overall and for eggs that successfully hatched into miracidia or failed to hatch and died. Statistically significant differences ( $P < 0.05$ ) are marked by (\*).

of infected snails between individuals exposed to miracidia from progenetic (13.8%) and normal eggs (15.1%; Table 1). Although there was a slight difference in size (shell volume) between snails exposed to progenetic and normal miracidia ( $3.77 \pm 0.14$  and  $3.30 \pm 0.12$  mm<sup>3</sup> respectively; Mann-Whitney U test,  $Z = 2.406$ ,  $P = 0.016$ ), we found no size difference among snails actually infected by *C. parvum* ( $2.29 \pm 0.23$  and  $2.03 \pm 0.14$  mm<sup>3</sup> respectively; Mann-Whitney U test,  $Z = 0.466$ ,  $P = 0.641$ ). Consequently, any difference in the number of sporocysts between snails infected by progenetic or normal miracidia would not be influenced by snail size. The

reproductive origin of miracidia (progenetic or normal eggs) had no effect on the mean number of sporocysts inside snail hosts, either after 10 weeks ( $6.7 \pm 0.4$  and  $6.8 \pm 0.4$  respectively; Mann-Whitney U test,  $Z = -0.073$ ,  $P = 0.941$ ) or 10 months ( $44.1 \pm 3.6$  and  $41.6 \pm 3.4$  respectively; Mann-Whitney U test,  $Z = 0.449$ ,  $P = 0.653$ ).

#### *Infectivity of cercariae to the amphipod second intermediate host and life-history strategy of metacercariae*

A total of 1030 amphipods were experimentally exposed to *C. parvum*; 511 to cercariae descending from selfed (progenetic) eggs and 519 from eggs obtained from adult worms in fish. After 5 weeks, 412 of the treated amphipods (40%) were still alive (206 in each group); high mortality is commonly observed in laboratory maintained *P. fluviatilis* (Lagrue and Poulin, 2007). Of these 412 individuals, 266 (64.6%) had been successfully infected by *C. parvum* although 33.8% of the cercariae (90 out of 266) that penetrated the amphipod did not survive and were encapsulated by the host immune system. We found no difference in the proportion of cercariae, produced by either progenetic or normal parents, successfully penetrating the host (65.5% and 63.6% respectively; Table 1) and subsequently surviving to develop into metacercariae (66.7% and 65.6% respectively of cercariae that penetrated the host; Table 1). Thus, there was no difference in the probability of a cercaria being encapsulated depending on its parental origin (normal or progenetic: 34.4 and 33.3% respectively; Fisher's exact test,  $\chi^2 = 0.03$ ,  $P = 0.86$ ).

There was no difference in the proportion of metacercariae adopting the progenetic developmental strategy between offspring of progenetic and normal parents (23.3% and 24.4% respectively; Table 1). Furthermore, the parental reproductive strategy (selfing or out-crossing) did not seem to influence the fitness of the offspring, either in terms of metacercariae body size ( $0.054 \pm 0.005$  and  $0.057 \pm 0.005$  mm<sup>2</sup>; respectively; Mann-Whitney U test,  $Z = 0.786$ ,  $P = 0.432$ ) or egg production of progenetic individuals ( $5.3 \pm 1.3$  and  $7.5 \pm 1.5$  respectively; Mann-Whitney U test,  $Z = -0.926$ ,  $P = 0.354$ ).

#### DISCUSSION

Inbreeding depression has long been thought to reduce the genetic value of inbred progeny, with strong deleterious effects on the fitness of plants and animals having been documented (see Charlesworth and Charlesworth, 1987 for review). This includes parasitic organisms: for example, the effects of inbreeding depression on the fitness of the cestode *Schistocephalus solidus* include reduced growth, lower infection and transmission success as well as limited intraspecific competitive ability (Christen *et al.* 2002; Christen and Milinski, 2003; Milinski, 2006). Whatever induces these deleterious effects, inbreeding depression is often strong and should drive the evolution of breeding systems and life-history strategies that avoid inbreeding (Charlesworth and Charlesworth, 1987). However, in this study, we found no evidence of such depression on any fitness parameter of *C. parvum* offspring. Although the expression of inbreeding depression depends largely on the environment in which its effects are tested, our experimental setups should have allowed us to detect some of these potential effects. Still, the progeny of progenetic metacercariae and adult worms were equally successful in terms of infectivity to snail and amphipod intermediate hosts, asexual multiplication inside snails, and survival and growth inside the amphipod host.

The only difference found in our study was in the hatching success of eggs: a significantly higher proportion of eggs produced by adult worms in fish hatched into miracidia. However, this is unlikely to be the effect of inbreeding depression but rather due to the different mechanisms of egg release. Adult *C. parvum* lay their eggs into the gut lumen of their fish definitive host; the eggs are rapidly released to the environment along with host faeces (Poulin and Cribb, 2002). These eggs are then free to hatch when the miracidium has developed. Because progenetic metacercariae are contained inside the body cavity of their amphipod host, eggs can be released only after the host has died and decomposed; waiting for host death is the most common way by which progenetic parasites release their eggs (Lefebvre and Poulin,

2005a). Some trematode species show high levels of virulence to accelerate host death and therefore the liberation of their eggs (Pampoulie *et al.* 2000), although *Coitocaecum parvum* seems to lower its virulence to extend host life and thus to maximize its egg output (Poulin, 2001). While this might be advantageous in terms of total egg production, it also means that eggs are trapped within the host for prolonged periods of time, waiting to reach the external environment to hatch. This could explain both the observed volume difference between progenetic and normal eggs and the peak of progenetic egg hatching on day 1 post-release. Eggs do not hatch within the cysts and those that are retained for longer than the optimal incubation period inside the amphipod should hatch immediately or might not be viable anymore when they eventually reach the external environment, hence the lower hatching success. At the same time, it is possible that eggs of progenetic metacercariae have a slower development when contained within the cyst, waiting to be released from the host to achieve maturation and hatch, hence the longer hatching time observed in this study and the bimodal shape of the daily hatching rate distribution for progenetic eggs. On the other hand, eggs contained within the amphipod host are protected against potential pathogens (fungi or bacteria) and predation; for instance, snails actively graze on the substrate and consume the eggs (personal observations). Eggs produced by adult worms in the fish host are quickly released in the external environment and remain unprotected during the incubation period, on average 15 days and virtually no eggs hatched before 12 days. Progenetic eggs that hatch during that time after host death, almost a quarter (22.6%) in our study, should be less likely to be predated upon or infected by pathogens, therefore compensating for the lower overall hatching success of eggs produced in the amphipod host.

It has been suggested that high levels of inbreeding should purge the population of deleterious alleles and mutations and therefore quickly reduce or eliminate the effects of inbreeding depression (Charlesworth and Charlesworth, 1987). Previous studies on trematodes have shown that some species can survive using self-fertilization without noticeable deleterious effects, at least for a few generations (see Lefebvre and Poulin, 2005a for review). Producing eggs by selfing inside the second intermediate host can be seen as a cheap reproductive insurance against failed transmission (Wang and Thomas, 2002), particularly if there is no cost associated with inbreeding. It is then possible that progenesis in the second intermediate host is actually not an alternative life-history strategy, when transmission to the definitive host fails, but the preferred reproductive strategy used by *C. parvum*; the high rate of selfing in *C. parvum* populations could cause a constant

purging of strongly deleterious alleles, reducing or even eliminating the short-term effects of inbreeding depression (Charlesworth and Charlesworth, 1987).

Indeed, quantitative models have shown that the balance between the advantages of self-fertilization and costs of inbreeding depression should result in populations using either selfing or out-crossing only (Charlesworth *et al.* 1990). In any given population, *C. parvum* should use preferentially one or the other reproductive strategy. However, it has also been predicted that environmental stochasticity could produce evolutionarily stable mixed mating systems in which both strategies co-occur (Cheptou and Dieckmann, 2002). In the case of *C. parvum*, environmental stochasticity in the second intermediate host is likely to be high due to the host itself (host sex, size, age, body condition and stress levels) and/or competition (intra or interspecific) between co-infecting parasites, and could either prevent or induce metacercariae to adopt progenesis (Lagrue and Poulin, 2007). *C. parvum* may use progenesis whenever possible but still have to use the fish definitive host when necessary: limited space and resources in the amphipod host, intra and or interspecific competition is known to limit parasite growth and prevent any egg production (Lagrue and Poulin, 2008b). In this situation, *C. parvum* populations may be mainly inbred with only a few individuals descending from outcrossing parents. Previous genetic studies have detected very low levels of heterozygosity and high linkage disequilibrium in *C. parvum* populations (Lagrue *et al.* 2007a,b; unpublished data); both are characteristics of predominantly self-fertilizing populations (Lymbery *et al.* 1997; Annan *et al.* 2007). However, the presence of heterozygous genotypes in populations of *C. parvum* shows that some outcrossing does occur (Lagrue *et al.* 2007a,b; unpublished data). In fact, within fish dissected for this study, some mating pairs were observed, providing direct evidence for sexual reproduction. Still, even though the relative importance of each reproductive strategy is unknown, progenesis could easily become the main way to produce offspring, especially in situations where fish are scarce or absent, as there are no obvious costs linked to inbreeding depression. It is also possible that adult worms in fish use selfing as well as outcrossing when sexual partners are scarce, thus increasing even further the inbreeding coefficient of the population (Lefebvre and Poulin, 2005a). However, in our study, a substantial number of offspring from adult *C. parvum* should be produced by cross-fertilization as access to mating partners was not limited and mating was indeed seen to happen; a comparison with the purely selfed offspring of progenetic worms should thus detect fitness differences, if there are any.

Finally, the proportion of progenesis observed in the next generation was comparable in offspring from progenetic metacercariae and those from adults. This shows that the reproductive strategy adopted by parents is not strictly heritable, i.e. it has no effect on the strategy adopted by their offspring, at least in the first generation. It suggests that life-cycle truncation in *C. parvum* is a conditional strategy rather than genetically predetermined (Poulin and Cribb, 2002), therefore corroborating our previous hypothesis that the adoption of progenesis by metacercariae depends on the environment provided by the amphipod host in which metacercariae develop (Lagrue and Poulin, 2007, 2008a,b).

In conclusion, inbreeding is unlikely to have any effect on the quality of offspring produced by metacercariae selfing in their second intermediate host. Furthermore, it seems that the progenetic strategy is not inherited from parents. Nevertheless, if there are no genetic costs of producing offspring by selfing and no genetic basis of progenesis, then why don't all metacercariae adopt the shorter life cycle and reproduce by selfing? First, it is possible that progenetic metacercariae in their second intermediate host cannot produce as many eggs as adult worms in the fish (Lefebvre and Poulin, 2005a) and the two strategies may have on average equal offspring output: progenesis insuring the production of at least some eggs whereas reaching the definitive host provides a low probability of higher fecundity (Poulin, 2001). Second, it is also possible that not all *C. parvum* individuals can achieve progenesis in the amphipod intermediate host. The amphipod host might be ingested by a fish definitive host before any egg production occurs, in which case the parasite would have no other solution but to reproduce in the fish. Also, progenesis is characterized by a drastic increase in the size of metacercariae, egg production being positively correlated with parasite size (Lagrue and Poulin, 2007). Therefore, intrinsic host resources and/or intra-host competition between individual parasites (hetero or conspecific) can restrict the growth of metacercariae and prevent the production of eggs, leaving no choice to the parasite but to wait for transmission to the definitive host (Lagrue and Poulin, 2008b). Overall, a multitude of interconnected factors have been documented to influence the life-history strategy of *Coitocaecum parvum* (Poulin, 2001, 2003; Lagrue and Poulin, 2007, 2008a,b). Whatever the reason for the co-occurrence of the two strategies, from our results, it is likely that inbreeding depression has little if any short-term effects on the maintenance of the normal three-host cycle in *Coitocaecum parvum* populations.

This research was supported by a grant from the Marsden Fund (The Royal Society of New Zealand) to R. P.



## REFERENCES

- Annan, Z., Durand, P., Ayala, F. J., Arnathau, C., Awono-Ambene, P., Simard, F., Razakandrainibe, F. G., Koella, J. C., Fontenille, D. and Renaud, F.** (2007). Population genetic structure of *Plasmodium falciparum* in the two main African vectors, *Anopheles gambiae* and *Anopheles funestus*. *Proceedings of the National Academy of Sciences, USA* **19**, 7987–7992.
- Brown, S. P., Renaud, F., Guegan, J. F. and Thomas, F.** (2001). Evolution of trophic transmission in parasites: the need to reach a mating place? *Journal of Evolutionary Biology* **14**, 815–820.
- Bryan-Walker, K., Leung, T. L. F. and Poulin, R.** (2007). Local adaptation of immunity against a trematode parasite in marine amphipod populations. *Marine Biology* **152**, 687–695.
- Bush, A. O. and Kennedy, C. R.** (1994). Host fragmentation and helminth parasites: hedging your bets against extinction. *International Journal for Parasitology* **24**, 1333–1343.
- Charlesworth, D. and Charlesworth, B.** (1987). Inbreeding depression and its evolutionary consequences. *Annual Review of Ecology and Systematics* **18**, 237–268.
- Charlesworth, D., Morgan, M. T. and Charlesworth, B.** (1990). Inbreeding depression, genetic load and the evolution of outcrossing rates in a multi-locus system with no linkage. *Evolution* **44**, 1469–1489.
- Cheptou, P. and Dieckmann, U.** (2002). The evolution of self-fertilization in density-regulated populations. *Proceedings of the Royal Society of London, B* **269**, 1177–1186.
- Christen, M., Kurtz, J. and Milinski, M.** (2002). Outcrossing increases infection success and competitive ability: experimental evidence from a hermaphrodite parasite. *Evolution* **56**, 2243–2251.
- Christen, M. and Milinski, M.** (2003). The consequences of self-fertilization and outcrossing of the cestode *Schistocephalus solidus* in its second intermediate host. *Parasitology* **126**, 369–378.
- Combes, C., Bartoli, P. and Théron, A.** (2002). Trematode transmission strategies. In *The Behavioural Ecology of Parasites* (ed. Lewis, E. E., Campbell, J. F. and Sukhdeo, M. V. K.), pp. 1–12. CAB International, Wallingford, UK.
- Crossan, J., Paterson, S. and Fenton, A.** (2007). Host availability and the evolution of parasite life-history strategies. *Evolution* **61**, 675–684.
- Hay, K. B., Fredensborg, B. L. and Poulin, R.** (2005). Trematode-induced alterations in shell shape of the mud snail *Zeacumantus subcarinatus* (Prosobranchia: Batillariidae). *Journal of the Marine Biological Association, UK* **85**, 989–992.
- Heywood, J. S.** (1991). Spatial analysis of genetic variation in plant populations. *Annual Review of Ecology and Systematics* **22**, 335–355.
- Holton, A. L.** (1984a). Progenesis as a mean of abbreviating life histories in two New Zealand trematodes, *Coitocaecum parvum* Crowfoot, 1945 and *Stegodexamene anguillae* MacFarlane, 1951. *Mauri Ora* **11**, 63–70.
- Holton, A. L.** (1984b). A redescription of *Coitocaecum parvum* Crowfoot, 1945 (Digenea: Allocreadiidae) from crustacean and fish hosts in Canterbury. *New Zealand Journal of Zoology* **11**, 1–8.
- Jarne, P.** (1995). Mating system, bottlenecks and genetic polymorphism in hermaphroditic animals. *Genetic Research* **65**, 193–207.
- Lagrué, C. and Poulin, R.** (2007). Life cycle abbreviation in the trematode *Coitocaecum parvum*: can parasites adjust to variable conditions? *Journal of Evolutionary Biology* **20**, 1189–1195.
- Lagrué, C. and Poulin, R.** (2008a). Lack of seasonal variation in the life-history strategies of the trematode *Coitocaecum parvum*: no apparent environmental effect. *Parasitology* **135**, 1243–1251.
- Lagrué, C. and Poulin, R.** (2008b). Intra- and interspecific competition among helminth parasites: effects on *Coitocaecum parvum* life history strategy, size and fecundity. *International Journal for Parasitology* **38**, 1435–1444.
- Lagrué, C., McEwan, J., Poulin, R. and Keeney, D. B.** (2007a). Co-occurrences of parasite clones and altered host phenotype in a snail-trematode system. *International Journal for Parasitology* **37**, 1459–1467.
- Lagrué, C., Waters, J. M., Poulin, R. and Keeney, D. B.** (2007b). Microsatellite loci for the progenetic trematode *Coitocaecum parvum* (Opecoelidae). *Molecular Ecology Notes* **7**, 694–696.
- Lefebvre, F. and Poulin, R.** (2005a). Progenesis in digenean trematodes: a taxonomic and synthetic overview of species reproducing in their second intermediate hosts. *Parasitology* **130**, 587–605.
- Lefebvre, F. and Poulin, R.** (2005b). Alternative reproductive strategies in the progenetic trematode *Coitocaecum parvum*: comparison of selfing and mating worms. *Journal of Parasitology* **91**, 93–98.
- Lefebvre, F. and Poulin, R.** (2005c). Life history constraints on the evolution of abbreviated life cycles in parasitic trematodes. *Journal of Helminthology* **79**, 47–53.
- Loveless, M. D. and Hamrick, J. L.** (1984). Ecological determinants of genetic structure in plant populations. *Annual Review of Ecology and Systematics* **15**, 65–95.
- Lymbery, A. J., Constantine, C. C. and Thompson, R. C. A.** (1997). Self-fertilization without genomic or population structuring in a parasitic tapeworm. *Evolution* **51**, 289–294.
- MacFarlane, W. V.** (1939). Life cycle of *Coitocaecum anaspidis* Hickman, a New Zealand digenetic trematode. *Parasitology* **31**, 172–184.
- Milinski, M.** (2006). Fitness consequences of selfing and outcrossing in the cestode *Schistocephalus solidus*. *Integrative and Comparative Biology* **46**, 373–380.
- Pampoulie, C., Lambert, A., Rosecchi, E., Crivelli, A. J., Bouchereau, J. L. and Morand, S.** (2000). Host death: A necessary condition for the transmission of *Aphalloides coelomicola* Dollfus, Chabaud, and Golvan, 1957 (Digenea, Cryptogonimidae)? *Journal of Parasitology* **86**, 416–417.
- Parker, G. A., Chubb, J. C., Ball, M. A. and Roberts, G. N.** (2003). Evolution of complex life cycles in helminth parasites. *Nature, London* **425**, 480–484.
- Poulin, R.** (2001). Progenesis and reduced virulence as an alternative transmission strategy in a parasitic trematode. *Parasitology* **123**, 623–630.

- Poulin, R.** (2003). Information about transmission opportunities triggers a life-history switch in a parasite. *Evolution* **57**, 2899–2903.
- Poulin, R.** (2007). *Evolutionary Ecology of Parasites*, 2nd Edn. Princeton University Press, Princeton, USA.
- Poulin, R. and Cribb, T. H.** (2002). Trematode life cycles: short is sweet? *Trends in Parasitology* **18**, 176–183.
- Rauch, G., Kalbe, M. and Reusch, T. B. H.** (2005). How a complex life cycle can improve a parasite's sex life. *Journal of Evolutionary Biology* **18**, 1069–1075.
- Thomas, F., Guldner, E. and Renaud, F.** (2000). Differential parasite (Trematoda) encapsulation in *Gammarus aequicauda* (Amphipoda). *Journal of Parasitology* **86**, 650–654.
- Trouvé, S., Renaud, F., Durand, P. and Jourdane, J.** (1996). Selfing and outcrossing in a parasitic hermaphrodite helminth (Trematoda, Echinostomatidae). *Heredity* **77**, 1–8.
- Wang, C. L. and Thomas, F.** (2002). Egg production by metacercariae of *Microphallus papillorobustus*: a reproductive insurance? *Journal of Helminthology* **76**, 279–281.
- Wedekind, C., Strahm, D. and Schärer, L.** (1998). Evidence for strategic egg production in a hermaphroditic cestode. *Parasitology* **117**, 373–382.
- Zakikhani, M. and Rau, M. E.** (1999). *Plagiorchis elegans* (Digenea: Plagiorchiidae) infections in *Stagnicola elodes* (Pulmonata: Lymnaeidae): host susceptibility, growth, reproduction, mortality, and cercarial production. *Journal of Parasitology* **85**, 454–463.