

## Research Article

**Cite this article:** Raghu BR, Shivashankara KS, Mahadappa P, Lokesh AN, Nandini KS, Rao VK, Prasanna HC, Dutta SK (2024). *Ex-situ* evaluation of curry leaf (*Murraya koenigii* (L.) Spreng) germplasm to unravel the genetic diversity of leaf essential oils. *Plant Genetic Resources: Characterization and Utilization* 1–15. <https://doi.org/10.1017/S1479262124000662>

Received: 31 July 2024

Revised: 19 November 2024

Accepted: 19 November 2024



### Keywords:

chemometric analysis; chemotypes; cultivated types; germplasm; wild types

### Corresponding author:

Borehalli Rangaswamy Raghu;  
Email: [Raghu.r@icar.gov.in](mailto:Raghu.r@icar.gov.in)

# *Ex-situ* evaluation of curry leaf (*Murraya koenigii* (L.) Spreng) germplasm to unravel the genetic diversity of leaf essential oils

Borehalli Rangaswamy Raghu<sup>1</sup> , Kodthalu Seetharamaiah Shivashankara<sup>1</sup>, Priyanka Mahadappa<sup>2</sup>, Ankanahalli Narayanashetty Lokesh<sup>1</sup>, Kebbahalli Shivashankarappa Nandini<sup>1</sup>, Vala Keshava Rao<sup>1</sup>, H.C. Prasanna<sup>1</sup> and Sudip Kumar Dutta<sup>3</sup> 

<sup>1</sup>Indian Institute of Horticultural Research (ICAR-IIHR), Hesaraghatta, Bengaluru, 560089, India; <sup>2</sup>Indian Veterinary Research Institute (ICAR-IVRI), Regional Campus, Hebbal, Bengaluru, 560024, India and <sup>3</sup>ICAR Research Complex for NEH Region, Sikkim Centre, Gangtok, Sikkim, 737 102, India

## Abstract

The present study was conducted with the objective of studying the genetic diversity of essential oils (EOs) in curry leaf (CL) *ex-situ*. Chemometric methods and pattern analysis were employed to assess the genetic diversity of EOs and to characterise diverse sets of CL germplasm into different chemotypes. The study revealed a huge genetic diversity for EO yield and its composition among the tested genotypes. Cultivated types had significantly higher EO yields and showed a greater degree of genetic divergence compared to wild types. In total, 80 different compounds were identified in the EOs of CL and classified into major (6) and minor (74) compounds. The major compounds  $\alpha$ -pinene,  $\gamma$ -terpinene, and  $\alpha$ -selinene and 14 minor compounds were highly variable among the tested genotypes. They may play an important role in the formation of different chemotypes. Other important compounds, such as trans-caryophyllene and  $\alpha$ -humulene, were more widely distributed among the tested genotypes and indicated their predominant occurrence in the EOs of CL. Some major compounds, such as valencene and  $\gamma$ -terpinene, showed a significant regional correlation, indicating the role of geographic factors in the evolution of different chemotypes. Furthermore, some compounds such as  $\alpha$ -pinene, bornyl acetate, and camphene had significantly higher concentrations in wild types compared to cultivated types, indicating the influence of domestication through human selection on the composition of EOs in CL. A total of 4 major chemotypes were characterised, of which three new chemotypes are being reported for the first time in CL.

## Introduction

*Murraya koenigii* (L.) Spreng, commonly known as curry leaf (CL), is a popular aromatic plant, leafy spice, and a familiar medicinal plant belonging to the family Rutaceae (Prakash and Natarajan, 1974). It is native to India and is geographically distributed throughout the tropical and subtropical regions of the Indian subcontinent (Joseph and Peter, 1985; Raghu, 2020). It is a good source of essential oil (EO); the leaves, fruits, and seeds yield EO on hydro-distillation (Rao *et al.*, 2011a). CL is also rich in several nutrients and bioactive compounds such as  $\beta$ -carotene, calcium, iron, vitamin C, and carbazole alkaloids (Philip *et al.*, 1981; Khatoon *et al.*, 2011; Vyas *et al.*, 2015; Raghu *et al.*, 2020; Raghu, 2023). Aside from its regular culinary usage as a spice (Verghese, 1989), it is also widely used in medical and nutraceutical applications (Igara *et al.*, 2016; Poornima *et al.*, 2022; Raghu *et al.*, 2022). Due to its anti-inflammatory, anti-diuretic, memory enhancing, antimicrobial, anti-diabetic, antioxidant, anti-tumour, and wound healing properties, CL is used in the treatment of various diseases and disorders such as piles, inflammation, itching, poison bites, fresh cuts, dysentery, diarrhoea, vomiting, and dropsy (Dastur, 1970; Drury, 1978; Ganesan *et al.*, 2013; Igara *et al.*, 2016). In recent times, it has become increasingly popular as a profitable commercial crop, especially in various regions of southern India, owing to its low input requirements, perennial nature, and steady market demand around the year (Mohan, 2012). Consequently, there is an escalating demand for CL genotypes that possess genetic superiority for commercial farming.

Essential oils (EOs) are the secondary metabolites produced by aromatic plants to protect themselves against several biotic and abiotic stresses (Sahoo *et al.*, 2022). EOs possess several bioactive properties, such as antimicrobial, anti-inflammatory, antioxidant, etc., which make them valuable commercial products in the pharmaceutical, cosmetic, food, and beverage industries (Ray *et al.*, 2018; Kamila *et al.*, 2021). The chemical composition of essential oils (EOs) directly influences their biological activity (Zhang *et al.*, 2015). For example, the EO



derived from CL, which was high in  $\alpha$ -Pinene (49.3% of the total volume), demonstrated significant antibacterial effects against five strains of gram-positive and five strains of gram-negative bacteria (Senthilkumar *et al.*, 2014). Likewise, the EO containing a high proportion of oxygenated monoterpenes (72.15%), including linalool (32.83%), displayed both antibacterial and antioxidant properties (Priya *et al.*, 2013). Moreover, previous studies have indicated that the EOs extracted from the leaves of the CL plant possess antimicrobial (Goutam and Purohit, 1974), antifungal (Deshmukh *et al.*, 1986), and pesticidal (Pathak *et al.*, 1997) activities. Therefore, it is very much essential to identify and characterise the existing CL germplasm into different chemotypes for better industrial applications. Similar to other aromatic species (Masotti *et al.*, 2003; Angioni *et al.*, 2006; Sahoo *et al.*, 2019), the yield and composition of EO in CL are influenced by its genetic makeup and the age of the plant (Raina *et al.*, 2002; Rao *et al.*, 2011a). Other factors, such as agronomic practices, climate, soil type and composition, and genotype  $\times$  environment interactions, also influence the EO composition and yield (Ram *et al.*, 2005; Verma *et al.*, 2009, 2010). Prevailing edaphic and abiotic factors exert selection pressure on underlying genetic differences for the biosynthetic pathway for volatile compounds, thereby leading to the development of different chemotypes in a single species across various regions (Morshedloo *et al.*, 2015). Hence, a proper understanding of the variations in the EO yield and composition is a prerequisite to exploring superior chemotypes.

*In situ* evaluation (on-site evaluation or evaluation in its natural habitat) of CL for its EO yield and composition has been extensively studied at different locations in India, and several chemotypes have been reported so far (Mallavarapu *et al.*, 1999, 2000; Raina *et al.*, 2002; Rao *et al.*, 2011a, 2011b; Syamasundar *et al.*, 2012; Verma *et al.*, 2013). Whereas evaluation of diverse germplasm with proper experimental design under common growing conditions (*ex-situ* evaluation) would accurately estimate genetic differences between tested genotypes and precisely characterise exploitable chemotypes. However, such efforts have yet to be made so far in CL for EO yield and composition. Therefore, it is necessary to evaluate CL germplasm *ex-situ* to quantify genetic variance for EO composition and yield. With this background, diverse sets of CL germplasm and elite genotypes were collected from different parts of the Indian subcontinent, covering the north, south, east, and western regions. A total of 125 accessions were collected and conserved in a field gene bank established at the ICAR- Indian Institute of Horticultural Research (ICAR-IIHR), Bengaluru (Raghu and Nandini, 2021). The present investigation was carried out on selected diverse germplasm accessions with the objective of studying the genetic diversity of EO yield and composition in CL and identifying different chemotypes.

## Materials and method

### Plant materials

Twenty diverse accessions of CL were utilised in the present study (Table 1). They comprise 15 accessions of cultivated types and 4 wild type genotypes collected from three Indian states and Suwasini, a commercially available improved variety (Fig. 1). The present experiment was carried out in RBD design by planting all 20 selected genotypes in three replications with a plant density of 10 plants per 23 square meters. All the recommended agronomic practices were followed, and plants were maintained by pruning one year after planting at regular intervals of three months.

### Sample preparation and extraction of EO

Leaf samples were collected from 3-month-old shoots during June 2022. Matured and pest-free leaves were selected. Fresh leaflets were retained by removing the leaf stalks. Finely chopped leaflet samples (500 g) were hydro-distilled in a Clevenger-type glass apparatus (Clevenger, 1928) for 5 h (Syamasundar *et al.*, 2012). The EO was collected and dried over anhydrous sodium sulphate and stored in the dark at 5°C until further analysis (Syamasundar *et al.*, 2012). The content of EOs (v/w) in the leaves of CL was expressed as a percentage. The experiment was repeated in June 2023.

### Analysis of the chemical composition of EOs in Gas chromatography-mass spectrometry (GC-MS)

GC-MS analyses of the EOs were performed using a Varian-3800 Gas Chromatograph equipped with a Varian-4000 Ion-Trap MS detector. The ion trap, transfer line and ion source temperatures were maintained at 190, 240 and 200°C, respectively. A fused-silica capillary column VF-5 ms from Varian, USA, with 30 m  $\times$  0.25 mm id, 0.25 mm film thickness, was used for the analysis. The injector temperature was set at 260°C and all injections were made in split mode (1:5). Samples were injected under the following conditions: helium was used as a carrier gas at approximately 1 ml/min, pulsed in split mode (20:1), the solvent delay was 3 min, and the injection volume was 1.0  $\mu$ l. The mass spectrometric detector was operated in the external electron ionisation mode operating at 70 eV, with a full mass scan- range of 45–450 amu. The detector temperature was maintained at 270°C and the temperature programme used for the column was as follows: 50°C for 5 min, followed by an increment of 4°C/min till 170°C, held for 2 min; subsequently, increased by 5°C/min till it reached 250°C and then, a constant temperature of 250°C was maintained for 7 min. The total run time was 60 min. Total volatile production was estimated by a sum of all peak areas in the chromatogram, and individual compounds were quantified as relative per cent area. Individual volatile compounds were identified by comparing their retention index (RI), which was determined using a homologous series of n-alkanes (C5 to C32, procured from Sigma-Aldrich) as Standard (Kovats, 1965) and comparing mass spectra with the available two spectral libraries, using Wiley and NIST-2007 (Adams, 1995).

### Statistical analysis

Replicated data derived from two years (2022 and 2023) on the EO yield were arcsine square root-transformed, and the combined analysis of variance was carried out according to the Duncan multiple comparison test at the probability level of  $P < 0.05$  using web-based free statistical software SAS (SAS On Demand for Academics, 'https://welcome.oda.sas.com'). Adjusted mean values or least squares mean (LS-means) were derived for EO yield. Similarly, combined ANOVA for EO components was performed, and adjusted mean values for individual volatile compounds were derived for each genotype. Two sample *T* tests were also performed using SAS software. The adjusted mean values for volatile components were auto-scaled and logarithmically transformed and further subjected to the chemometric analysis through the free statistical software MetaboAnalyst 5.0 (Pang *et al.*, 2021). The chemometric analysis consisted of significance analysis of metabolomics (SAM) based on *F* statistics, pattern analysis based on Spearman rank correlation, principal component analysis (PCA), partial least squares discriminant analysis (PLS-DA), and variable

**Table 1.** Details of curry leaf germplasm used in the present study

Genotype (Collection#)	Region/ State	Latitude	Longitude	Altitude (Mt.)	Biological status	LSM of Essential oil Yld (%)	Salient morphological features
LSR/18/6	KA	17°10' N	76°47' E	413	Cultivated	0.62	Bushy habit, medium and light-green leaves with very high fragrance
LSR/18/7	KA	17°10' N	76°48' E	413	Cultivated	0.56	Tree habit, medium and dark-green leaves with very high fragrance
LSR/18/8	KA	15°45' N	76°12' E	413	Cultivated	0.30	Tree habit, medium and green leaves with medium fragrance
LSR/18/9	KA	15°46' N	76°44' E	413	Cultivated	0.41	Tree habit, medium and green leaves with medium fragrance
LSR/18/18	KA	17°15' N	76°47' E	343	Cultivated	0.35	Bushy habit, very small and light green leaves with high fragrance
LSR/18/75	KA	17°09' N	75°57' E	374	Cultivated	0.30	Tree habit, medium and dark-green leaves with high fragrance
LSR/18/93	KA	15°40' N	75°17' E	530	Cultivated	0.12	Bushy habit, very small and dark-green leaves with medium fragrance
LSR/18/162	KA	15°01' N	76°05' E	446	Cultivated	0.11	Tree habit, medium and dark-green leaves with medium fragrance
LSR/18/175	KA	15°46' N	76°45' E	338	Cultivated	0.30	Bushy habit, medium and green leaves with medium fragrance
Suwasini	KA	15°29' N	74°58' E	678	Cultivated	0.23	Bushy habit, small and dark-green leaves with very high fragrance
RRP/18/4	Od	18°35' N	82°12' E	272	Cultivated	0.33	Tree habit, medium and dark-green leaves with high fragrance
RRP/18/17	Od	18°15' N	82°52' E	158	Cultivated	0.30	Tree habit, medium and dark-green leaves with high fragrance
RRP/18/20	Od	18°15' N	82°01' E	207	Cultivated	0.36	Tree habit, medium and green-leaves with medium fragrance
RRP/18/30	Od	18°17' N	81°48' E	198	Cultivated	0.20	Tree habit, medium and dark-green leaves with medium fragrance
BRR/18/3	TN	11°50' N	77°56' E	351	Cultivated	0.28	Bushy habit, small and dark-green leaves with high fragrance
BRR/18/28	TN	11°43' N	77°58' E	298	Cultivated	0.30	Bushy habit, small and dark-green leaves with high fragrance
BRR/18/8	TN	11°36' N	77°17' E	1300	Wild	0.21	Tree habit, big and dark-green leaves with low fragrance
BRR/18/9	TN	11°15' N	77°01' E	378	Wild	0.27	Bushy habit, big and dark-green leaves with low fragrance
BRR/18/10	TN	11°13' N	77°06' E	288	Wild	0.25	Tree habit, big and dark-green leaves with low fragrance
BRR/18/19	TN	11°43' N	77°28' E	138	Wild	0.22	Tree habit, big and dark-green leaves with low fragrance
Mean						0.30	

KA, Karnataka; Od, Odisha; TN, Tamil Nadu.

of importance in prediction (VIP score), and hierarchical cluster analysis (HCA) and heatmap with the Euclidean distance between the genotypes given by the Ward algorithm.

## Results


### EO yield and composition

Analysis of variance among CL genotypes revealed a significant ( $P < 0.05$ ) variability for EO yield (Table 2). The EO yield in the present study ranged from 0.11–0.62% (v/w), with a mean

yield of 0.3%. The highest EO yield was recorded in LSR/18/06 (0.62%), followed by LSR/18/07 (0.56%). The lowest EO yield was observed in LSR/18/162 (0.11%). Cultivated types of CL showed significantly ( $P < 0.05$ ) higher EO yield as compared to wild types (online Supplementary Table S1).

Eighty distinct compounds were detected in the EOs of 20 different genotypes of CL. Table 3 summarises the identified compounds in terms of their occurrences across the tested genotypes, along with the range and mean values. Each genotype contained between 27 (BRR/18/10) and 45 (RPP/18/4) compounds, contributing to approximately 98.1% (LSR/18/75) to 99.9% (BRR/18/10)





**Figure 1.** Map showing collection sites of CL germplasm in three Indian states.

of the EOs (Fig. 2). Among the 80 compounds identified, 13 compounds –  $\beta$ -cadinene, agarospirol,  $\delta$ -selinene, patchoulene, santalol, trans- $\alpha$ -bisabolene, safranal, cumic alcohol,  $\delta$ -cadinol,  $\alpha$ -santalol,  $\alpha$ -lonol,  $\beta$ -pachoulene, and  $\beta$ -himachalene – are reported for the first time in this study (Table 3). Additional details about the volatile compounds found in each genotype are provided in online Supplementary Table S2.

#### *Occurrence and distribution of EO components*

Univariate analysis of EO components indicated a statistically significant variability among tested genotypes (Table 3). Trans-caryophyllene was the most abundant (mean;  $33.78 \pm 11.8\%$ ) and widely distributed compound (range; 16.2%–65.4%) detected in all 20 tested genotypes.



**Table 2.** Analysis of variance of essential oil yield among curry leaf germplasm

Source	DF	SS	Mean Square	F Value	Pr > F
Genotype	19	0.888	0.046	23.47	<0.0001
Replication	2	0.0008	0.0004	0.20	0.81
Error	38	0.075	0.0019		

Similarly, compounds valencene (mean;  $11.51 \pm 6.90\%$ ),  $\gamma$ -terpinene (mean;  $9.52 \pm 13.6\%$ ),  $\alpha$ -humulene (mean;  $7.23 \pm 2.61\%$ ),  $\alpha$ -pinene (mean;  $6.01 \pm 9.6\%$ ), and  $\alpha$ -selinene (mean;  $4.53 \pm 6.23\%$ ), were also present in significantly ( $P < 0.05$ ) higher quantities. Wherein  $\alpha$ -humulene and  $\alpha$ -pinene were present in all the tested genotypes, whereas valencene,  $\gamma$ -terpinene, and  $\alpha$ -selinene were detected in >90 and 75% of tested genotypes, respectively. All these top six compounds that we found in the present study as major compounds are listed in Table 3, along with their range of distribution.

Whereas spathulenol, along with 73 remaining compounds (Sl. No. 8-80 in Table 3), were detected in small but varying quantities, with the mean values ranging from <0.5 to 2.67%. This set of compounds comprised compounds restricted to a few genotypes as well as compounds found in the majority of the tested genotypes with varying low concentrations. Thus, we have listed all these 74 compounds (Sl. No. 7-80 in Table 3) as minor compounds.

Further, SAM analysis of EO components between wild and cultivated types was performed to understand the differential distribution of EO components among wild and cultivated types (Fig. 3). The results indicated that concentrations of  $\alpha$ -pinene, bornyl acetate, and camphene were significantly high ( $P < 0.05$ ) in wild types. In contrast, trans- $\alpha$ -bergamotene and 1-terpineol were significantly high in cultivated types (online Supplementary Table S3). Although  $\alpha$ -pinene was detected in all 20 tested genotypes, its concentration was higher in wild types (mean of 23.8%) compared to cultivated types (mean of 1.6%). Besides, pattern analysis based on Spearman's rank correlation confirmed the SAM results. It showed  $\alpha$ -pinene, bornyl acetate, and camphene had an increasing concentration in wild types and a decreasing concentration in cultivated types (Fig. 4). In contrast, compound trans-caryophyllene and 21 other minor compounds were in increasing concentrations in cultivated types and decreasing concentrations in wild types.

### Population variance for EO components

PCA was performed to understand the population variance of EO components in CL germplasm. It is an unsupervised estimation of a given population. To better understand the dispersion of genotypes, we presented the graph of the PCA vectors (Fig. 5), the loadings of the variables (compounds) (online Supplementary Table S4), and the scores of the samples (genotypes) (online Supplementary Table S5).

The 3D principal component analysis (Fig. 5) retained the first three main principal components (PCs), which explained 49.7% of data variability. Results demonstrated a greater genetic divergence both within and between different geographical regions of India. Wild CL genotypes were clustered at the top-left of the score plot, with the highest positive values for PC2 and the highest negative values for PC1, indicating a similar chemical profile

among them. Whereas cultivated types were dispersed at the bottom of the score plot with negative values for PC2, indicating their clear chemical distinctness from wild CL genotypes. Further, these cultivated CL genotypes were dispersed in a divergent direction due to positive/or negative values for PC1 and PC3, indicating a greater genetic divergence among cultivated CL genotypes than wild CL genotypes. There was a clear intermixing of cultivated CL genotypes from different regions, indicating the existence of genotypes with similar chemical compositions across the regions.

Further, PCA indicated that among 80 compounds, 17 compounds had the highest loadings ( $\pm$ values) for the first 3 PCs (online Supplementary Table S4). This indicated their greater contribution to the overall population variance. Amongst these 17 compounds, 3 were major ( $\alpha$ -pinene,  $\gamma$ -terpinene, and  $\alpha$ -selinene), and the remaining 14 were minor (sabinene,  $\alpha$ -phellandrene,  $\beta$ -phellandrene, cryptone,  $\beta$ -elemene, trans- $\alpha$ -bergamotene, camphene, bornyl acetate,  $\alpha$ -terpinene,  $\alpha$ -terpineol, 1-terpineol,  $\delta$ -selinene,  $\delta$ -elemene,  $\gamma$ -cadinene). These findings further confirmed that major and minor compounds together play an important role in the formation of different chemotypes in CL.

Meanwhile, the compounds with almost null loading values ( $\leq |0.09|$ ) for all three PCs were placed near the origin in the score plot and had a limited role in imparting population variance. A total of 34 compounds had null values ( $\leq |0.09|$ ) (online Supplementary Table S4). Among them, 3 major compounds (trans-caryophyllene,  $\alpha$ -humulene, and valencene) and 12 minor compounds (spathulenol,  $\delta$ -cadinene, agarospirol, viridiflorol,  $\alpha$ -copaene, and aromadendrene,  $\alpha$ -gurjunene,  $\alpha$ -muurolene, linalool,  $\alpha$ -cubebene,  $\alpha$ -ylangene, and guaiol) were important due to their wider occurrence across genotypes (present in more than 75% of the tested genotypes). These compounds might play a fixative role in EOs and be responsible for constituting the basic characteristic aroma of CL across various chemotypes.

### Influence of origin on genetic divergence of EO components

In order to identify the compound(s) with significant region-specific expression, a supervised multivariate discriminant analysis was performed. We used the presented variable of importance (VIP) in the prediction score for EO components (Fig. 6) following Sahoo *et al.* (2022). The compounds with >1.0 VIP scores are reported to have more influence on population discrimination. Accordingly, we identified 22 compounds with >1.0 VIP scores (Fig. 6). Major compound valencene along with minor compounds such as sabinene,  $\alpha$ -guaiene,  $\beta$ -elemene,  $\delta$ -elemene, agarospirol,  $\beta$ -guaiene,  $\beta$ -eudesmol, juniper camphor, linalool, and  $\beta$ -selinene were present in significantly higher concentrations in the genotypes from Tamil Nadu. Whereas the major compound  $\gamma$ -terpinene, along with minor compounds such as  $\beta$ -cubebene and  $\alpha$ -phellandrene, were present in significantly higher concentrations in the genotypes of Odisha. No major compounds; however, only a few minor compounds such as torreyol, 1-terpineol,  $\gamma$ -cadinene, camphene,  $\alpha$ -terpineol, caryophyllene oxide, cryptone, and  $\beta$ -humulene were present in significantly higher concentrations in the genotypes from Karnataka.

### Association among EO components

Analysis of hierarchical clustering between volatile compounds and CL genotypes was performed with the objective of identifying genotypes with similar chemical profiles (Fig. 7). Results indicated that

**Table 3.** Components of leaf essential oils of curry leaf germplasm


Sl. No	Compound	Number of genotypes reported the compound	RT	KI	% Range (In reported genotypes)		LS Means (Percent) ( $P = 0.05$ )*
					Min	Max	
1	Trans-Caryophyllene	20	26.60	1442	16.2	65.41	33.78 ± 11.8 <sup>a</sup>
2	Valencene	19	29.76	1499	2.30	23.30	11.51 ± 6.90 <sup>b</sup>
3	$\gamma$ -Terpinene	19	10.65	1056	0.02	37.69	9.52 ± 13.6 <sup>c</sup>
4	$\alpha$ -Humulene	20	28.03	1448	3.1	12.84	7.23 ± 2.61 <sup>d</sup>
5	$\alpha$ -Pinene	20	5.57	959	0.01	31.70	6.01 ± 9.6 <sup>de</sup>
6	$\alpha$ -Selinene	15	29.88	1475	0.01	26.05	4.53 ± 6.23 <sup>e</sup>
7	Spathulenol	18	33.30	1536	0.69	6.32	2.57 ± 2.47 <sup>f</sup>
8	$\beta$ -Elemene	18	25.36	1398	0.02	5.05	1.98 ± 1.76 <sup>fg</sup>
9	$\delta$ -Cadinene	19	30.62	1497	0.66	3.19	1.50 ± 0.77 <sup>fg</sup>
10	$\delta$ -3-Carene	17	10.15	1009	0.10	4.78	1.41 ± 1.41 <sup>fg</sup>
11	$\beta$ -Phellandrene	15	10.15	964	0.02	9.59	1.38 ± 2.45 <sup>fg</sup>
12	Juniper-Camphore	16	36.37	1680	0.51	4.41	1.34 ± 1.21 <sup>fg</sup>
13	$\beta$ -Selinene	17	29.44	1483	0.04	9.47	1.10 ± 2.89 <sup>fg</sup>
14	Trans- $\alpha$ -Bergamotene	12	27.10	1434	0.25	3.76	1.06 ± 1.25 <sup>fg</sup>
15	Aromadendrene	19	27.31	1439	0.21	6.94	1.01 ± 1.44 <sup>fg</sup>
16	Sabinene	20	8.83	984	0.01	4.44	1.00 ± 1.34 <sup>fg</sup>
17	Viridiflorol	17	33.53	1588	0.38	1.99	0.86 ± 0.65 <sup>fg</sup>
18	$\beta$ -Eudesmol	17	34.36	1593	0.17	4.23	0.73 ± 1.20 <sup>fg</sup>
19	Agarospirol	19	34.79	1598	0.05	1.59	0.70 ± 0.54 <sup>fg</sup>
20	Cryptone	6	17.13	1148	0.15	3.9	0.67 ± 1.40 <sup>fg</sup>
21	$\alpha$ -Santalol	2	33.29	1454	1.61	9.5	0.65 ± 2.58 <sup>fg</sup>
22	$\alpha$ -Phellandrene	13	8.38	969	0.01	4.26	0.64 ± 1.14 <sup>fg</sup>
23	$\alpha$ -Copaene	20	24.66	1375	0.11	1.92	0.61 ± 0.55 <sup>fg</sup>
24	$\beta$ -Humulene	8	33.57	1754	0.24	6.57	0.51 ± 1.46 <sup>fg</sup>
25	Germacrene-D	10	31.24	1582	0.07	8.07	0.51 ± 0.20 <sup>fg</sup>
26	Guaiol	19	33.91	1591	0.12	0.9	0.37 ± 0.25 <sup>g</sup>
27	$\alpha$ -Muurolene	18	30.43	1495	0.16	1.04	0.37 ± 0.27 <sup>g</sup>
28	4-Terpineol	6	16.74	1182	0.04	4.83	0.37 ± 1.52 <sup>g</sup>
29	$\gamma$ -Cadinene	12	30.43	1471	0.11	1.98	0.37 ± 0.48 <sup>g</sup>
30	$\alpha$ -Cubebene	20	23.44	1345	0.08	0.64	0.30 ± 0.15 <sup>g</sup>
31	Torreyol	14	34.54	1646	0.07	0.99	0.25 ± 0.28 <sup>g</sup>
32	Camphene	11	9.33	947	0.03	3.54	0.25 ± 0.79 <sup>g</sup>
33	d Limonene	2	9.31	1048	1.26	3.3	0.22 ± 0.78 <sup>g</sup>
34	$\beta$ -Cubebene	7	26.94	1400	0.04	1.36	0.21 ± 0.39 <sup>g</sup>
35	$\gamma$ -Elemene	15	22.80	1425	0.01	0.99	0.20 ± 0.31 <sup>g</sup>
36	Bornyl Acetate	11	21.00	1289	0.01	1.03	0.20 ± 0.31 <sup>g</sup>
37	Cubenol	4	32.59	1580	0.20	2.92	0.19 ± 0.65 <sup>g</sup>
38	$\beta$ -Guaiene	18	35.45	1523	0.05	1.39	0.19 ± 0.36 <sup>g</sup>
39	$\alpha$ -Terpineol	12	17.95	1175	0.01	0.63	0.17 ± 0.22 <sup>g</sup>
40	$\delta$ -Elemene	11	22.34	1353	0.01	0.99	0.17 ± 0.29 <sup>g</sup>
41	$\alpha$ -Gurjunene	16	25.35	1419	0.02	1.03	0.17 ± 0.23 <sup>g</sup>

(Continued)

Table 3. (Continued.)


Sl. No	Compound	Number of genotypes reported the compound	RT	KI	% Range (In reported genotypes)		LS Means (Percent) ( $P = 0.05$ )*
					Min	Max	
42	$\beta$ -Cadinene	5	30.62	1497	0.12	2.02	0.16 $\pm$ 0.46 <sup>B</sup>
43	$\gamma$ -Gurjunene	9	32.16	1509	0.07	0.89	0.14 $\pm$ 0.23 <sup>B</sup>
44	Calamenene	7	30.82	1537	0.12	0.81	0.13 $\pm$ 0.23 <sup>B</sup>
45	Phytol	10	50.81	2045	0.06	0.98	0.13 $\pm$ 0.23 <sup>B</sup>
46	Caryophyllene oxide	7	32.01	1507	0.07	0.75	0.12 $\pm$ 0.22 <sup>B</sup>
47	(Z)- $\beta$ -Gurjunene	4	10.17	1044	0.09	1.92	0.11 $\pm$ 0.43 <sup>B</sup>
48	$\alpha$ -Terpinolene	4	11.82	1052	0.19	1.13	0.10 $\pm$ 0.27 <sup>B</sup>
49	Cadinene	7	25.90	1440	0.09	0.41	0.10 $\pm$ 0.16 <sup>B</sup>
50	$\delta$ -Selinene	6	26.06	1481	0.01	0.6	0.10 $\pm$ 0.19 <sup>B</sup>
51	$\alpha$ -Terpinene	7	11.91	1000	0.01	0.9	0.09 $\pm$ 0.22 <sup>B</sup>
52	Germacrene-B	3	32.62	1584	0.23	1.05	0.09 $\pm$ 0.26 <sup>B</sup>
53	$\alpha$ -Ylangene	17	24.38	1396	0.03	0.35	0.09 $\pm$ 0.07 <sup>B</sup>
54	1-Terpineol	10	14.13	1192	0.01	0.41	0.09 $\pm$ 0.13 <sup>B</sup>
55	$\alpha$ -Guaiene	5	35.10	1490	0.08	0.53	0.09 $\pm$ 0.17 <sup>B</sup>
56	$\alpha$ -Lonol	1	34.36	1455	0	1.58	0.07 $\pm$ 0.35 <sup>B</sup>
57	Linalool	15	13.29	1109	0.01	0.37	0.07 $\pm$ 0.09 <sup>B</sup>
58	Alloaromadrene	2	23.98	1386	0.42	1.11	0.07 $\pm$ 0.09 <sup>B</sup>
59	$\beta$ -Himachalene	1	30.29	1528	0	1.32	0.06 $\pm$ 0.29 <sup>B</sup>
60	$\beta$ -Pachoulene	4	27.56	1484	0.17	0.48	0.06 $\pm$ 0.14 <sup>B</sup>
61	Patchoulane	3	20.04	1393	0.20	0.31	0.06 $\pm$ 0.07 <sup>B</sup>
62	Santalol	2	35.42	1668	0.39	0.8	0.05 $\pm$ 0.19 <sup>B</sup>
63	Trans- $\alpha$ -Bisabolene	5	31.53	1500	0.06	0.38	0.05 $\pm$ 0.12 <sup>B</sup>
64	Trans-Ocimene	2	10.12	1055	0.01	1.08	0.05 $\pm$ 0.24 <sup>B</sup>
65	$\beta$ -Bisabolene	2	30.94	1500	0.38	0.6	0.04 $\pm$ 0.15 <sup>B</sup>
66	$\alpha$ -Thujene	7	5.49	902	0.01	0.52	0.04 $\pm$ 0.13 <sup>B</sup>
67	Cumic alcohol	2	22.57	1284	0.27	0.6	0.04 $\pm$ 0.14 <sup>B</sup>
68	Linalyl acetate	5	17.58	1272	0.02	0.5	0.04 $\pm$ 0.12 <sup>B</sup>
69	Tau-Murolene	1	29.53	1435	0	0.69	0.03 $\pm$ 0.16 <sup>B</sup>
70	$\delta$ -Cadinol	1	30.43	1580	0	0.53	0.02 $\pm$ 0.12 <sup>B</sup>
71	$\alpha$ -Cadinol	3	35.69	1632	0.13	0.19	0.02 $\pm$ 0.06 <sup>B</sup>
72	Nerol	2	15.08	1228	0.04	0.28	0.01 $\pm$ 0.06 <sup>B</sup>
73	Borneol	3	16.40	1156	0.01	0.26	0.01 $\pm$ 0.06 <sup>B</sup>
74	Tau-Cadinol	1	29.98	1580	0	0.29	0.01 $\pm$ 0.06 <sup>B</sup>
75	Safranal	2	21.69	1244	0.09	0.11	0.01 $\pm$ 0.03 <sup>B</sup>
76	Cis-Farnesol	1	36.69	1710	0	0.13	0.006 $\pm$ 0.02 <sup>B</sup>
77	Neryl acetate	1	21.12	1352	0	0.09	0.004 $\pm$ 0.02 <sup>B</sup>
78	$\alpha$ -elemene	1	24.91	1373	0	0.05	0.002 $\pm$ 0.01 <sup>B</sup>
79	Geraniol	1	16.70	1228	0	0.01	0.001 $\pm$ 0.03 <sup>B</sup>
80	Camphor	1	14.96	1138	0	0.002	0.001 $\pm$ 0.04 <sup>B</sup>

RT, Retention Time; KI, Kovate Index; \*Data are expressed as Mean  $\pm$  Standard Deviation; Mean values followed by different letters (a–g) at superscript are significantly different at the probability level of  $P < 0.05$  according to the Duncan test.



**Figure 2.** Number and total volume of volatile compounds present in essential oils of CL germplasm.

### SAM Plot for Delta = 0.9



**Figure 3.** Significant compounds identified between wild and cultivated curry leaves by SAM (Significance Analysis of Microarray) (Delta = 0.9). The green-coloured dots indicated compounds with significant differences between the two groups. Compounds with positive *d*-values are placed at the top, indicating higher concentration in wild types (i.e., mean concentration in wild types > mean concentration in cultivated curry leaves), and compounds with negative *d*-values placed at the bottom, indicating higher concentration in cultivated types (i.e., mean concentration in cultivated types > mean concentration in wild curry leaves).


CL germplasm was grouped into two major genetically distinct populations, A and B. Population-A consisted of 8 cultivated genotypes, of which 6 were from Karnataka and 2 from Odisha. Population A had significantly higher concentrations of  $\alpha$ -selinene and  $\gamma$ -terpinene as major compounds, and  $\alpha$ -phellandrene, crypton,  $\alpha$ -terpineol, 1-terpineol, trans-

$\alpha$ -bisabolene, calamenene,  $\alpha$ -cubebene, and  $\alpha$ -copaene as minor compounds. Further, population-A was sub-grouped into A1 and A2 based on the relative concentrations of  $\alpha$ -selinene,  $\gamma$ -terpinene, crypton,  $\alpha$ -phellandrene,  $\alpha$ -cubebene, and  $\alpha$ -copaene.


Population B had higher concentrations of valencene and was comprised of both cultivated ( $n = 8$ ) and wild CL genotypes ( $n =$



### Top 25 compounds correlated with the 1-2




**Figure 4.** Pattern analysis among the top 25 correlated compounds with wild and cultivated curry leaves based on Spearman rank correlation. '1' denotes Cultivated, and '2' denotes wild types. The top 3 compounds show a positive correlation with wild types and a negative correlation with cultivated types, showing upward expression among wild types and downward expression for cultivated types. The bottom 22 compounds have a positive correlation with cultivated types and a negative correlation with wild types, showing upward expression among cultivated types and downward expression for cultivated types.



**Figure 5.** Principal component analysis for EO components in CL germplasm. 3D Score plot of essential compounds of wild and cultivated curry leaf germplasm collected from 3 Indian states: KA, Karnataka; O, Odisha, and TN, Tamil Nadu.

4). Based on the chemical profile of other major and minor compounds, it was further divided into 3 sub-populations, namely, B1, B2, and B3. Subpopulations B1 (LSR/18/8, LSR/18/9, LSR/18/175,

and RPP/18/30) and B2 (BRR/18/3, BRR/18/28, Suwasini, and RPP/18/4) were comprised of only cultivated CL genotypes and had higher concentrations of trans-caryophyllene and



**Figure 6.** Important volatile compounds identified by partial least squares-discriminant analysis (PLS-DA) among cultivated curry leaves from 3 Indian states. Twenty-five top compounds according to the VIP (variable importance in projection) score in Karnataka, Odisha and Tamil Nadu are shown. Coloured boxes indicated the relative concentrations of the corresponding compound between groups (red, higher concentration; green, lowest concentration; grey, moderate concentration).

$\alpha$ -humulene as major compounds. These two subpopulations, B1 and B2, differed with respect to the composition of minor compounds. Sub-population B3 (BRR/18/8, BRR/18/9, BRR/18/10, BRR/18/19) was comprised of only wild CL genotypes. It was predominantly composed of  $\alpha$ -pinene as the major compound and bornyl acetate, camphene, agarospinol,  $\delta$ -selinene, cadinene,  $\delta$ -elemene, and (*Z*-)- $\beta$ -ocimene as the minor compounds.

#### Chemotype classification in CL


Based on the significant distribution of major compounds and minor compounds in the EOs, the tested genotypes were grouped into four major chemotypes, namely, (i) trans-caryophyllene and  $\gamma$ -terpinene dominant, (ii) trans-caryophyllene dominant, (iii) trans-caryophyllene, valencene, and  $\alpha$ -humulene dominant, and (iv) trans-caryophyllene,  $\alpha$ -pinene, and valencene dominant chemotypes (Table 4). The chemotype characterisation of CL germplasm was in line with the results of PCA, PLS-DA and HCA.  $\alpha$ -pinene dominant chemotypes were limited to wild types, and  $\gamma$ -terpinene dominant chemotypes were confined to cultivated types. The valencene dominant chemotypes were found both in wild and cultivated types of CL genotypes; however, all were collected from the Tamil Nadu region. The trans-caryophyllene was

present in all chemotypes, indicating the predominant occurrence of trans-caryophyllene dominant chemotypes in CL.

#### Discussion

The present research has made significant efforts to explore the biochemical diversity and richness that are inherent in this aromatic plant. The germplasm selected for this study was sourced from a diverse array of geographical areas (three Indian states), ensuring an ample number of accessions from each state to examine any possible inter- and intra-regional differences (Fig. 1). A thorough understanding of the genetic variations in CL is essential for developing an effective strategy for future germplasm exploration initiatives and conservation efforts, as well as for breeding traits related to important essential oils (Salgotra and Chauhan, 2023). This study represents a fundamental step toward fully utilising native species, like CL, for pharmaceutical applications (Raghu, 2020).

The essential oil (EO) yield observed in this study ranged considerably (0.11 to 0.62%) with a mean value of 0.3%, suggesting substantial genetic variation for EO yield in CL, which could be further utilised through breeding strategies. We identified two CL genotypes (LSR/18/06 and LSR/18/07) that exhibited



**Figure 7.** Heatmap of EO components according to different chemotypes in CL germplasm. Clusters of highly similar chemical profiles were highlighted in the box.

particularly high EO yields (>0.5%). Both genotypes are cultivated types showcasing an intense fragrance and were collected from a nearby geographical area (Table 1). The EO yield in CL appears to have a significant correlation with geographical origin (Mallavarapu *et al.*, 1999; Raina *et al.*, 2002; Syamasundar *et al.*,

2012), altitude (Verma *et al.*, 2013), and the domestication process influenced by human selection (Rao *et al.*, 2011b), which likely has had a continuous modifying effect on the underlying biosynthetic pathways of EOs (Shamsheer *et al.*, 2022). This was further supported by the notably lower EO yield found in wild



**Table 4.** Chemotype classification of curry leaf germplasm

Chemotype	Major compound	Minor compound	Explained volume	Corresponding Genotype	Fragrance intensity
Trans-Caryophyllene & $\gamma$ -Terpinene dominant	Trans-Caryophyllene (18–42.4%), $\gamma$ -Terpinene (18.1–37.7%), $\alpha$ -Humulene (4.5–8.0%), $\alpha$ -Selinene (1.5–7.8%)	Spathulenol (1.8–9.5%), $\alpha$ -Phellandrene (0.2–4.3%) Cryptone (0.2–3.9%)	71.8–81.8%	LSR/18/06 LSR/18/07 LSR/18/18 LSR/18/75 RRP/18/17	High High High High High
Trans-Caryophyllene, $\alpha$ -Pinene dominant	Trans-Caryophyllene (65.4%), $\alpha$ -Humulene (3.1%)	Spathulenol (2.0–7.3%) $\alpha$ -Phellandrene (0.2–2.7%) $\beta$ -Phellandrene (1.9–9.6%) $\alpha$ -Santalol (9.5%)	57.8–80.1%	LSR/18/93 RRP/18/20 LSR/18/162	Medium Medium Medium
Trans-Caryophyllene, Valencene & $\alpha$ -Humulene dominant	Trans-Caryophyllene (30.0–52.5%), $\alpha$ -Humulene (7.6–12.9%), Valencene, (9.7–22.3%)	Spathulenol (1.4–2.4%), $\delta$ -Cadinene (1.1–3.1%), Viridiflorol (1.8–1.9%), Aromadendrene (0.2–1.4%), Trans- $\alpha$ -bergamotene (1.2–2.8%), Juniper camphore (1.0–4.4%)	66.3–74.9%	BRR/18/3 BRR/18/28 Suwasini RRP/18/4	High High High High
Trans-Caryophyllene, $\alpha$ -Pinene & Valencene dominant	Trans-Caryophyllene (16.2–27.8%), $\alpha$ -Pinene (20.2–31.7%), Valencene, (10.8–23.3%), $\alpha$ -Humulene (3.4–5.9%),	Spathulenol (1.4–2.4%), $\delta$ -Cadinene (1.1–3.1%), Viridiflorol (1.8–1.9%), Aromadendrene (0.2–1.4%), Germacrene-D (0.2–8.0%) Camphene (0.3–3.6%), Bornyl acetate (0.9–1.2%), $\beta$ -Phellandrene (2.2–5.6%), $\beta$ -Elenene (0.4–4.4%), Sabinene (0.1–4.4%)	72.8–85.9%	LSR/18/8 LSR/18/9 LSR/18/175 RRP/18/30 BRR/18/8 BRR/18/9 BRR/18/10 BRR/18/19	Medium Medium Medium Medium Low Low Low Low

types compared to cultivated types, as all wild genotypes consistently displayed reduced fragrance (Table 1). Hence, it is suggested that the variations in fragrance intensity among CL are positively correlated with leaf EO content.

The chemical diversity of the CL is substantial in terms of EO composition (Rao *et al.*, 2011a). Besides inherent genetic differences (Raina *et al.*, 2002), various elements like location (Mallavarapu *et al.*, 1999, 2000), seasonal changes (Verma *et al.*, 2012), habitat (Rana *et al.*, 2004; Syamasundar *et al.*, 2012; Verma *et al.*, 2013), and cultivation practices (Rao *et al.*, 2011b) notably influence the biochemical properties of CL. Nevertheless, there is currently no research available that directly compares the components of EO by categorising them into major or minor compounds within CL. To enable meaningful direct comparisons, CL oils must be extracted under identical conditions while maintaining all other variables constant. This method would accurately delineate the chemotypes and genuine heritable variation (Rao *et al.*, 2011a). Consequently, in this research, we conducted a statistical analysis of the EO profiles from 20 CL genotypes, all extracted under uniform conditions.

A total of 80 compounds were found across 20 CL accessions, with each genotype having between 27 and 48 compounds, which together constituted more than 98% of the total EO volume. Out of the 80 compounds, only six were identified to be in significantly higher concentrations among the different genotypes; these six compounds – trans-caryophyllene, valencene,  $\gamma$ -terpinene,  $\alpha$ -humulene,  $\alpha$ -pinene, and  $\alpha$ -selenine – were categorised as major volatile compounds. The remaining 74 compounds (Sl. No. 7-80 in Table 3) were detected in substantially lower concentrations and were thus labelled as minor compounds. This classification of EO components into major and minor categories based on statistical differences is being reported in CL for the first time. However, earlier research has identified higher concentrations of  $\alpha$ -pinene, sabinene,  $\beta$ -pinene,  $\delta$ -3-carene, limonene,  $\beta$ -phellandrene, (Z)- $\beta$ -ocimene, lavandulol, terpinen-4-ol, geraniol,  $\alpha$ -copaene, isocaryophyllene, and  $\gamma$ -elemene in EOs of CL (MacLeod and Pieris, 1982; Onayade and Adebajo, 2000; Rao *et al.*, 2011a; Sukkaew *et al.*, 2014; Santhanakrishnan *et al.*, 2023).

Additionally, this study represents the first attempt to examine the differences in EO yield and composition between wild and cultivated types to examine the degree and direction of genetic divergence in CL during the domestication process. Generally, trans-caryophyllene emerged as the primary component in cultivated types, while other significant compounds like  $\gamma$ -terpinene,  $\alpha$ -humulene, and  $\alpha$ -selenine showed quantitative variation among the cultivated genotypes. Conversely, wild types primarily contained  $\alpha$ -pinene as the main constituent of their EOs. This chemical composition variation underscores the essential connection between the culinary and medicinal preference for trans-caryophyllene and the genetic alterations resulting from human selection during the domestication of CL. Therefore, trans-caryophyllene could explain the enduring aroma found in CL plant leaves and their popularity as a spice (Onayade and Adebajo, 2000).

In the current study, multivariate analysis such as PCA was employed to describe population variance for EO composition with a few key compounds (Fig. 4). PCA is a powerful dimension-reduction technique that is frequently employed in large amounts of biochemical data to describe overall variation with the smallest possible number of components. In curry leaf (Santhanakrishnan *et al.*, 2023) and other aromatic crops (Ray *et al.*, 2019; Kamila *et al.*, 2021; Sahoo *et al.*, 2022), the chemical diversity and

chemotype characterisation have been reported based on PCA. In the present study, the PCA indicated that  $\alpha$ -pinene,  $\gamma$ -terpinene, and  $\alpha$ -selenine were contributing the most to population variance due to the highest variable loadings (online Supplementary Table S4). Santhanakrishnan *et al.* (2023) reported that  $\alpha$ -pinene,  $\alpha$ -fenchene,  $\beta$ -pinene, chloral hydrate, and  $\alpha$ -caryophyllene are distributed distinctly. They were found to be prominent components for the distribution of the 11 curry leaf accessions they examined. Although trans-caryophyllene and  $\alpha$ -humulene were detected as the major components of EOs, they contribute less to population variance due to null variable loading values (online Supplementary Table S4), and thus, they were the most widely distributed and commonly available volatile compounds in EOs of CL. In other studies, trans-caryophyllene was reported to be the main component of EOs in CL genotypes from Sri Lanka, Nigeria, and Thailand (Onayade and Adebajo, 2000; Rao *et al.*, 2011b).

In the current study, hierarchical clustering was performed to classify the genotypes based on similar chemical profiles and identify the chemotypes (Fig. 6). The HCA classification was in line with the PCA results. The whole population was divided into two major genetically distinct populations, A and B, which were further sub-grouped into 5 sub-populations (Fig. 6). Grouping of genotypes was not in accordance with their origin, indicating the existence of similar chemotypes across locations. The  $\alpha$ -selenine and  $\gamma$ -terpinene major cluster consisted only of cultivated types (Fig. 6). In contrast, a predominance of valencene, trans-caryophyllene, and  $\alpha$ -humulene formed the second major cluster, which consisted of both wild and cultivated types. Variety Suwasini grouped in the second cluster and had higher concentrations of valencene, trans-caryophyllene, and  $\alpha$ -humulene and a lower concentration of  $\alpha$ -pinene. Interestingly, Santhanakrishnan *et al.* (2023) reported a higher concentration of  $\alpha$ -pinene in the leaf volatiles of Suwasini based on thermal desorption gas chromatography-mass spectroscopy.

Rao *et al.* (2011b) summarised several previous studies in CL. They described 14 chemotypes under three major categories (monoterpenoid, sesquiterpenoid, and mono- and sesquiterpenoid predominant oils) based on leaf EO chemical profiles. However, these studies were characterised by the *in situ* evaluation of a few genotypes, and the identification of chemotypes was simply based on predominant compounds rather than the significant contribution of the compounds based on statistical differences (Rao *et al.*, 2011a). The present CL germplasm was classified into four major chemotypes based on HCA results (Table 4). Notably, trans-caryophyllene was present in all the chemotypes, indicating its predominance in the EOs of CL. Previously,  $\beta$ -caryophyllene dominant (Onayade and Adebajo, 2000; Raina *et al.*, 2002) and  $\beta$ -caryophyllene and  $\alpha$ -pinene dominant chemotypes (Syamasundar *et al.*, 2012) were reported in CL. In the present study, three new chemotypes have been identified: trans-caryophyllene and  $\gamma$ -terpinene dominant; trans-caryophyllene, valencene, and  $\alpha$ -humulene dominant; and trans-caryophyllene,  $\alpha$ -pinene, and valencene dominant. Two superior CL genotypes, LSR/18/06 and LSR/18/07 that have been identified in the current study for the higher EO content belong to the newly reported chemotype, i.e., trans-caryophyllene and  $\gamma$ -terpinene dominant chemotype.

This research highlights the heredity aspects of biochemical variation observed in the EOs of CL. It sets a benchmark for the future management of plant genetic resources and targeted breeding programs in CL. This lays the groundwork for more comprehensive research in the future to uncover the biochemical mechanisms involved and to identify the key genes that improve

EO yield and quality in CL, which can also be applied to other aromatic plants. Additionally, two newly identified superior genotypes of CL (LSR/18/06 and LSR/18/07) from this study have immediate relevance in both the agriculture and essential oil sectors.

## Conclusion

In the current study, we made pioneer efforts to establish an *ex-situ* field bank and then to understand the genetic diversity of EO composition in cultivated and wild types of CL germplasm. We employed statistically robust techniques, such as cluster analysis, PCA, PLS-DA, SAM, and pattern analysis, for chemotype characterisation in CL for the first time. A greater degree of genetic divergence was observed for EO yield and composition among tested genotypes, indicating the extent of chemical divergence that is inherent in CL. The current study has identified and classified a large number of volatile compounds of EOs as major and minor compounds for the first time in CL. Cultivated types yielded substantially higher amounts of EO as compared to wild types, and their chemical profile was genetically distinct from wild types.

Further, a greater degree of chemical divergence was observed in cultivated types as compared to wild types, reflecting the impact of human selection and domestication in CL for a variety of leaf fragrances. In addition, the study has unravelled three new chemotypes in CL. The information generated from the present study will be useful for the pharmaceutical and EO industries and also for planning future exploration trips and subsequent conservation programs in CL. This will support breeding initiatives focused on improving both EO yield and quality in upcoming cultivars, such as the two superior genotypes (LSR/18/06 and LSR/18/07) mentioned in this study.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262124000662>

**Acknowledgements.** The authors are thankful to Dr M R Dinesh, former Director of ICAR – Indian Institute of Horticultural Research and current Director – for providing basic infrastructure and necessary support.

**Author contributions.** B.R. Raghu: Conceptualization, Methodology, Writing- Original draft preparation; K.S. Shivashankara: Supervision, Manuscript editing; Priyanka Mahadappa: Software-data analysis, Manuscript editing; A.N. Loksha: Methodology-GCMS analysis; K. S. Nandini: Methodology-Essential oils extraction; V. K. Rao: Methodology-Essential oils extraction; H.C. Prasanna: Manuscript editing; Sudip Kumar Dutta: Manuscript editing.

**Competing interests.** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- Adams RP (eds) (1995) *Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy*, 2nd Edn. Carol Stream, IL: USA: Allured Publishing Corporation.
- Angioni A, Barra A, Coroneo V, Dessi S and Cabras P (2006) Chemical composition, seasonal variability, and antifungal activity of *Lavandula stoechas* L. ssp. *stoechas* essential oils from stem/leaves and flowers. *Journal of Agricultural and Food Chemistry* **54**, 4364–4370.
- Clevenger JF (1928) Apparatus for the determination of volatile oil. *Journal of the American Pharmaceutical Association* **17**, 346–349.
- Dastur JF (ed.) (1970) *Medicinal Plants of India and Pakistan*, 3rd Edn. Bombay: DB Taraporevala Sons & Co. Private Ltd.
- Deshmukh SK, Jain PC and Agarwal SC (1986) Antimicrobial activity of the essential oil of the leaves of *Murraya koenigii* (Linn.) Spreng (Indian curry leaf). *Fitoterapia* **57**, 295–297.
- Drury HC (ed.) (1978) *The Useful Plants of India*, 2nd Edn. London: William H Allen & Co.
- Ganesan P, Phaiphon A, Murugan Y and Baharin BS (2013) Comparative study of bioactive compounds in curry and coriander leaves: an update. *Journal of Chemical and Pharmaceutical Research* **5**, 590–594.
- Goutam MP and Purohit RM (1974) Antimicrobial activity of essential oils of the leaves of *Murraya koenigii* (L.) Spreng. (Indian curry leaf). *Indian Journal of Pharmacology* **36**, 11.
- Igara C, Omoboyowa D, Ahuchaogu A, Orji N and Ndukwe M (2016) Phytochemical and nutritional profile of *Murraya Koenigii* (Linn) Spreng leaf. *Journal of Pharmacognosy and Phytochemistry* **5**, 7–9.
- Joseph S and Peter KV (1985) Curry leaf (*Murraya koenigii*), perennial nutritious leafy vegetable. *Economic Botany* **39**, 68–73.
- Kamila PK, Ray A, Jena S, Sahoo A, Kar SK, Nayak S and Panda PC (2021) Intraspecific variability in yield and chemical composition of essential oil of the endemic species *Hypericum gaitii* from different natural habitats of Eastern India. *Plant Biosystems* **156**, 1167–1176.
- Khatoun J, Verma A, Chacko N and Sheikh S (2011) Utilisation of dehydrated curry leaves in different food products. *Indian Journal of Natural Products and Resources* **2**, 508–511.
- Kovats E (1965) Gas chromatographic characterisation of organic substances in the retention index system. *Advances in Chromatography* **1**, 229–247.
- Macleod AJ and Pieris NM (1982) Analysis of the volatile essential oils of *Murraya koenigii* and *Pandanus latifolius*. *Phytochemistry* **21**, 1653–1657.
- Mallavarapu GR, Ramesh S, Syamasundar KV and Chandrasekhara RS (1999) Composition of Indian curry leaf oil. *Journal of Essential Oil Research* **11**, 176–178.
- Mallavarapu GR, Rao L and Ramesh S (2000) Volatile constituents of the leaf and fruit oils of *Murraya koenigii* Spreng. *Journal of Essential Oil Research* **12**, 766–768.
- Masotti V, Juteau F, Bessière JM and Viano J (2003) Seasonal and phenological variations of the essential oil from the narrow endemic species *Artemisia molinieri* and its biological activities. *Journal of Agricultural and Food Chemistry* **51**, 7115–7121.
- Mohan RS (2012) Curry leaf campaign. *Spice India* **25**, 10–12.
- Morshedloo MR, Ebadi A, Maggi F, Fattahi R, Yazdani D and Jafari M (2015) Chemical characterisation of the essential oil compositions from Iranian populations of *Hypericum perforatum* L. *Industrial Crops and Products* **76**, 565–573.
- Onayade OA and Adebajo AC (2000) Composition of the leaf volatile oil of *Murraya koenigii* growing in Nigeria. *Journal of Herbs, Spices & Medicinal Plants* **7**, 59–66.
- Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, Gauthier C, Jacques P, Li S and Xia J (2021) Metaboanalyst 5.0: narrowing the gap between raw spectra and functional insights. *Nucleic Acids Research* **49**, 388–396.
- Pathak N, Yadav TD, Jha AN, Vasudevan P and Pathak N (1997) Contact and fumigant action of volatile oil of *Murraya koenigii* against *Callosobruchus chinensis*. *Indian Journal of Entomology* **59**, 198–202.
- Philip J, Peter KV and Gopalakrishnan PC (1981) Curry leaf a mineral packed vegetable. *Indian Horticulture* **25**, 2–3.
- Poornima KN, Nandini KS and Raghu BR (2022) Curry leaf a medicinal boon. *Agriculture world* **8**, 24–27.
- Prakash V and Natarajan CP (1974) Studies on curry leaf (*Murraya koenigii* L.). *Journal of Food Science and Technology* **11**, 285–286.
- Priya RM, Blessed BP and Nija S (2013) Chemical composition, antibacterial and antioxidant profile of essential oil from *Murraya koenigii* (L.) leaves. *Avicenna Journal of Phytomedicine* **4**, 200–214.
- Raghu BR (2020) Diversity and distribution of curry leaf in India. *Journal of Horticultural Sciences* **15**, 1–8.
- Raghu BR (2023) Curry leaf. In Dubey RK and Singh J (eds), *Production Technology of Underexploited Vegetable Crops*. New Delhi: Kalyani Publishers, pp. 72–85.
- Raghu BR and Nandini KS (2021) Standardisation of seed propagation method in curry leaf. In International Conference on Vegetable Research



- and Innovations for Nutrition, Entrepreneurship and Environment (ICVEG-21), ICAR-IIVR, Varanasi (UP), India, 14–16th December 2021, pp. 445–446.
- Raghu BR, Aghora TS and Dhananjaya MV** (2020) Genetic improvement of curry leaf in India: challenges and future prospects. *Indian Horticulture* **65**, 127–130.
- Raghu BR, Aghora TS and Dhananjaya MV** (2022) Curry leaf Improvement in India. In Compendium for winter School on “underexploited vegetables: unexplored treasure trove for food, nutritional and economic security, ICAR-IIVR, Varanasi (UP), India, 02–11 February, 2022, pp. 183–186.
- Raina VK, Lal RK, Tripathi S, Khan M, Syamasundar KV and Srivastava SK** (2002) Essential oil composition of genetically diverse stocks of *Murraya koenigii*. *Flavour and Fragrance Journal* **17**, 144–146.
- Ram P, Kumar B, Naqvi AA, Verma RS and Patra NK** (2005) Post-harvest storage effect on quality and quantity of rose-scented geranium (*Pelargonium* sp. cv. Bourbon) oil in Uttaranchal. *Flavour and Fragrance Journal* **20**, 666–668.
- Rana VS, Juyal JP, Rashmi and Blazquez MA** (2004) Chemical constituents of the volatile oil of *Murraya koenigii* leaves. *International Journal of Aromatherapy* **14**, 23–25.
- Rao BRR, Rajput DK and Mallavarapu GR** (2011a) Chemotype categorisation of curry leaf plants (*Murraya koenigii* (L.) Spreng.). *Journal of Essential Oil-Bearing Plants* **14**, 1–10.
- Rao BRR, Rajput DK and Mallavarapu GR** (2011b) Chemical diversity in curry leaf (*Murraya koenigii*) essential oils. *Food Chemistry* **126**, 989–994.
- Ray A, Jena S, Dash B, Kar B, Halder T, Chatterjee T, Ghosh B, Panda PC, Nayak S and Mahapatra N** (2018) Chemical diversity, antioxidant and antimicrobial activities of the essential oils from Indian populations of *Hedychium coronarium* Koen. *Industrial Crops and Products* **112**, 353–362.
- Ray A, Jena S, Haldar T, Sahoo A, Kar B, Patnaik J, Ghosh B, Panda PC, Mahapatra N and Nayak S** (2019) Population genetic structure and diversity analysis in *Hedychium coronarium* populations using morphological, phytochemical and molecular markers. *Industrial Crops and Products* **32**, 118–133.
- Sahoo A, Kar B, Jena S, Dash B, Ray A, Sahoo S and Nayak S** (2019) Qualitative and quantitative evaluation of rhizome essential oil of eight different cultivars of *Curcuma longa* L. (Turmeric). *Journal of Essential Oil-Bearing Plants* **22**, 239–247.
- Sahoo C, Champati BB, Dash B, Jena S, Ray A, Panda PC, Nayak S and Sahoo A** (2022) Volatile profiling of *Magnolia champaca* accessions by gas chromatography mass spectrometry coupled with chemometrics. *Molecules* **27**, 7302.
- Salgotra RK and Chauhan BS** (2023) Genetic diversity, conservation, and utilisation of plant genetic resources. *Genes* **14**, 174.
- Santhanakrishnan VP, Shoba N, Varun E, Mohankumar S and Raveendran M** (2023) Aromatic profiling of *Murraya koenigii* leaves by Thermal Desorption Gas chromatography-Mass Spectroscopy (TD-GC-MS). *Heliyon* **9**, e17832.
- Senthilkumar A, Gopalakrishnan B, Jayaraman M and Venkatesalu V** (2014) Chemical composition and antibacterial activity of essential oil from the leaves of *Murraya koenigii* (L.) Spreng. *Journal of Experimental Sciences* **5**, 1–4.
- Shamsheer B, Riaz N, Yousaf Z, Hyder S, Aftab A, Iqbal R, Rahman MH, Al-Ashkar IF, Almutairi K and El Sabagh A** (2022) Genetic diversity analysis for wild and cultivated accessions of *Cymbopogon citratus* (D.C.) Stapf using phytochemical and molecular markers. *PeerJ* **10**, e13505.
- Sukkaew S, Pripdeevech P, Thongpoon C, Machan T and Wongchuphan R** (2014) Volatile constituents of *Murraya koenigii* fresh leaves using head-space solid phase microextraction – gas chromatography – mass spectrometry. *Natural Product Communications* **9**, 1783–1786.
- Syamasundar KV, Srinivasulu B, Ananda PLG, Ramesh S and Rao RR** (2012) Chemo variations of wild curry leaf (*Murraya koenigii* Spreng.) from Western Ghats of Indian. *Journal of Pharmacognosy* **3**, 126–130.
- Vergheze J** (1989) Indian curry leaf. *Perfumer & Flavorist* **14**, 69–70.
- Verma RS, Verma RK, Chauhan A and Yadav AK** (2009) Changes in the essential oil content and composition of *Eucalyptus citriodora* hook during leaf ontogeny and leaf storage. *India Perfumer* **53**, 22–25.
- Verma RS, Rahman L, Verma RK, Chanotiya CS, Chauhan A, Yadav A, Yadav AK and Singh A** (2010) Changes in the essential oil content and composition of *Origanum vulgare* L. during annual growth from Kumaon Himalaya. *Current Science* **98**, 1010–1012.
- Verma RS, Padalia RC, Arya V and Chauhan A** (2012) Aroma profiles of the curry leaf, *Murraya koenigii* (L.) Spreng. Chemotypes variability in north India during the year. *Industrial Crops and Products* **36**, 343–348.
- Verma RS, Chauhan A, Padalia RC, Jat SK, Thul S and Sundaresan V** (2013) Phytochemical diversity of *Murraya koenigii* (L.) Spreng. from western Himalaya. *Chemistry & Biodiversity* **10**, 628–641.
- Vyas VG, Kandoliya UK, Vidhani SI, Parmar HJ, Bhalani VM and Golakiya BA** (2015) Heavy metal deposition and phytochemical characterisation of curry leaves (*Murraya koenigii*). *International Journal of Current Microbiology and Applied Sciences* **4**, 839–843.
- Zhang DY, Yao XH, Duan MH, Wei FY, Wu GH and Li L** (2015) Variation of essential oil content and antioxidant activity of *Lonicera* species in different sites of China. *Industrial Crops and Products* **77**, 772–779.