

Sleeping-tree fidelity of the spider monkey shapes community-level seed-rain patterns in continuous and fragmented rain forests

Arturo González-Zamora^{*,1}, Víctor Arroyo-Rodríguez[†], Federico Escobar[‡], Ken Oyama^{†,§}, Filippo Aureli^{**} and Kathryn E. Stoner^{††}

* División de Posgrado, Instituto de Ecología A.C., Xalapa, Veracruz, Mexico

† Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Morelia, Michoacán, Mexico

‡ Red de Ecoetología, Instituto de Ecología A.C., Xalapa, Veracruz, Mexico

§ Escuela Nacional de Estudios Superiores, Universidad Nacional Autónoma de México, Morelia, Michoacán, Mexico

** Instituto de Neuroetología, Universidad Veracruzana, Xalapa, Mexico, and Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John Moores University, Liverpool, UK

†† Department of Fish, Wildlife, and Conservation Ecology, New Mexico State University, Las Cruces, New Mexico, USA

(Received 7 February 2014; revised 4 May 2015; accepted 4 May 2015)

Abstract: Repeated use of sleeping trees (STs) by frugivores promotes the deposition and aggregation of copious amounts of seed, thus having key implications for seed dispersal and forest regeneration. Seed-rain patterns produced by this behaviour likely depend on the frequency of use of these sites, yet this hypothesis has been poorly tested. We evaluated community-level seed-rain patterns produced by the spider monkey (*Ateles geoffroyi*) over 13 mo in latrines located beneath 60 STs in the Lacandona rain forest, Mexico. Because this primate is increasingly 'forced' to inhabit fragmented landscapes, we tested whether sleeping-tree fidelity (STF) differed among sites and between continuous and fragmented forests. We also tested whether seed-rain patterns were associated with STF within each site and forest type. STF was highly variable among STs (average = 7 mo, range = 1–12 mo), but did not differ among study sites or forest types. STF was positively associated with seed abundance, species diversity and species turnover. Nevertheless, STF tended to be negatively related to seed community evenness. These results are likely due to the most frequently used STs being in areas with greater food density. Our results demonstrate that site fidelity shapes community-level seed-rain patterns and thus has key ecological implications.

Key Words: aggregated seed dispersal, *Ateles*, frugivory, forest fragmentation, Lacandona, primates, sleeping site

INTRODUCTION

The repeated use of perches, roosting sites, reproductive sites and sleeping sites over time by frugivorous vertebrates promotes the deposition and aggregation of copious amounts of seed in these sites (Russo & Augspurger 2004, Wenny 2001). These behaviours have key implications for seed dispersal and forest regeneration (Jordano & Schupp 2000), particularly in tropical forests where up to 94% of woody plant species are dispersed by frugivorous animals (Jordano 1992). The importance of this spatially aggregated pattern of seed deposition for seed dispersal is largely dependent on site fidelity, that is, on the frequency of use of these sites over time (Russo

& Augspurger 2004, Russo *et al.* 2006); yet, empirical evidence about this relationship is scarce.

Site fidelity is particularly common in territorial animals (Börger *et al.* 2008), such as Geoffroy's spider monkey (*Ateles geoffroyi*) (Chapman *et al.* 1995). This primate is a highly specialized frugivore (González-Zamora *et al.* 2009, Russo *et al.* 2005) that defends stable home ranges, within which the spider monkey concentrates its activities in areas of higher quality (i.e. the so-called 'core areas'; Asensio *et al.* 2012a). As multi-central place foragers (*sensu* Chapman *et al.* 1989), they feed on several plants located near sleeping trees (STs), and return to the same or different STs after their feeding excursions. An important fraction of seeds swallowed by these primates are defecated in latrines located beneath these STs (González-Zamora *et al.* 2014, Russo & Augspurger 2004, Russo *et al.* 2006), particularly beneath in those located within core areas,

1 Corresponding author. Email: toztlan@yahoo.com.mx

Current address: Instituto de Investigaciones Biológicas, Universidad Veracruzana, Xalapa, Veracruz, Mexico

and hence, variation in sleeping-tree fidelity (STF) is expected to alter patterns of seed rain within latrines.

Here, we evaluated the frequency of use of 60 STs by *A. geoffroyi* and the seed-rain patterns produced by this primate during a 13-mo period in two continuous forest sites (CF) and three forest fragments (FF) in the fragmented Lacandona rain forest, Mexico. Because the potential variation between continuous and fragmented forests in STF has not been previously evaluated, we first tested whether STF (operationally defined as the number of months each ST is used) differed among the five study sites and between forest types (continuous and fragmented forests). For each study site and forest type, we assessed the association between STF and six community-level attributes of seed assemblages (Tuomisto 2010): abundance of seeds, species diversity of seeds (i.e. species richness, exponential of Shannon's entropy and inverse Simpson concentration), community evenness, and seed species turnover (β -diversity) between STs. This information has critical ecological implications, as *A. geoffroyi* are increasingly forced to inhabit fragmented landscapes (Ramos-Fernández & Wallace 2008), but we do not know if their seed-dispersal services may be altered in forest fragments (but see Chaves *et al.* 2011, González-Zamora *et al.* 2014).

STF of spider monkeys is expected to be higher in forest fragments (González-Zamora *et al.* 2012), as the home ranges of these primates in fragments are smaller than in continuous forest sites (Chaves *et al.* 2012). Although it is reasonable to expect that the abundance and diversity of seeds increase with STF, such associations largely depend on food availability within the home range. Because the STs most frequently used are expected to function as key nodes within foraging networks routes that connect core areas (Di Fiore & Suarez 2007, Suarez *et al.* 2014), we predicted STF to be positively associated with seed abundance and with all four diversity metrics. Thus, differences between STs in re-use are expected to result in significant differences in species turnover (i.e. increasing β -diversity) between STs. Moreover, because the spider monkey shows high selectivity towards the consumption of a few genera of plants (Chapman 1988, Milton 1980, Russo *et al.* 2005) and a large number of species are used opportunistically (González-Zamora *et al.* 2009, Nunes 1998), we predicted STF to be negatively associated with community evenness (i.e. with higher STF the seed rain will be dominated by fewer species).

METHODS

Study sites

The Lacandona rain forest constitutes the south-western sector of the Mayan forest in Mexico, and represents a

priority area for biodiversity conservation in Mexico and Mesoamerica (Myers *et al.* 2000). The area is located in the north-eastern portion of the state of Chiapas, and is delimited by the Guatemalan border on the south and east, and by the Chiapas highlands on the north and west. Annual precipitation averages 2850 mm, and average monthly temperatures range between 24°C and 26°C. This region was originally covered by over 1.4 million ha of rain forest, but deforestation between 1960 and 1990 resulted in the loss of 70% of the original forest cover (Arizpe *et al.* 1993).

Within this region the Montes Azules Biosphere Reserve (MABR) was created in 1978 to protect biodiversity. Adjacent to the southern extreme of MABR, the Marqués de Comillas Region (MCR) was colonized by humans about 40 y ago and since then MCR has suffered the rapid loss and fragmentation of the original rain forest (Mora 2008). Currently, MCR is dominated by different-sized rain forest patches, embedded in a matrix of cattle pastures, agricultural lands (e.g. corn, oil palm, rubber) and human settlements. The study was conducted in MCR (eastern side of the Lacantún River; 2039 km²) and MABR (western side; 3312 km²).

Experimental design

Based on a recent study on the density and spatial distribution of sleeping sites, STs and latrines of the spider monkey (*Ateles geoffroyi*) in the region (González-Zamora *et al.* 2012, 2014), we selected 60 sleeping sites, each composed by one single ST, in two continuous-forest sites within MABR and three forest fragments in MCR (12 STs per study site). To increase the independence between the two continuous-forest sites, they were separated by 5 km from each other. The forest fragments FF1, FF2 and FF3 have an area of 1125, 33 and 30 ha, respectively, and were isolated ≥ 24 y ago and were immersed in an anthropogenic matrix of pastures and agricultural lands. The average distance between two fragments was 4.2 km (see further details in Gonzalez-Zamora *et al.* 2012, 2014). We do not have accurate information on the home range, population size/density and core area size of each primate's community. Yet a recent study in the same study sites reports that FF1 is occupied by a community of approximately 40 spider monkeys using a home range of 63 ha, whereas the communities from FF2 and FF3 have 30 and 39 individuals, respectively, and use the entire fragment area (Chaves *et al.* 2012). The average home-range size and community size in the continuous forest of the Lacandona rain forest is 68.4 ha and 40 individuals (Chaves *et al.* 2012). Unfortunately, we do not have data about the size of core areas in the study sites, but in Santa Rosa National Park, Costa Rica, Asensio *et al.* (2012b) estimated that spider monkeys use five core areas of 9.2

ha on average (range = 3.4–19.2 ha) that totalled 46.1 ha out of the 304 ha of the entire home range.

Seed-rain patterns

Each selected ST had a single latrine (Gonzalez-Zamora *et al.* 2012, 2014). The STs had a mean crown projection of approximately $21 \pm 21 \text{ m}^2$ (mean \pm SD). This indicates that each trap covered *c.* 3% of the tree crown. Nevertheless, because we located the trap in the centre of the latrine, which had an average diameter of 1.5 m (González-Zamora *et al.* 2014), each trap covered approximately 60% of the latrine's area. Traps consisted of a circular 1.5-m-diameter PVC frame supporting a 0.5-m-deep open-topped nylon mesh bag suspended 1 m above the ground on three steel posts to prevent the possible predation of seeds by terrestrial vertebrates. In fact, we did not detect signs of seed predation (i.e. open husks, teeth marks) in the dispersed seed pool. Although seed traps also captured some fruits and seeds dispersed by wind or gravity, and would also capture seed dispersed by bird and bats, we only considered seeds completely immersed within monkey faeces. These seeds were easily identified in the field based on their typically stained appearance and characteristic adhesion of faecal matter (González-Zamora *et al.* 2014), thus limiting greatly the erroneous inclusion of seeds dispersed by wind, gravity, or other animals. Furthermore, we did not consider seeds < 5 mm in length, which are known to be dispersed by a wider variety of seed dispersers, including small-to-medium terrestrial mammals, bats and birds (Janson 1983, Stoner *et al.* 2007), whereas seeds ≥ 5 mm in length are frequently dispersed by primates (Arroyo-Rodríguez *et al.* 2015, Chaves *et al.* 2011, González-Zamora *et al.* 2014).

During a 13-mo period (1 February 2011–28 February 2012) each trap was emptied once monthly, and the seeds were collected and afterwards washed for subsequent identification in the laboratory. All seeds were counted and identified to the species level based on (1) our own experience with the local flora (from seeds to adults) (Chaves *et al.* 2011, 2012; González-Zamora *et al.* 2012); (2) assistance from a botanical expert familiar with the flora of the region (G. Ibarra-Manríquez, Universidad Nacional Autónoma de México, Morelia, Mexico) and a local parataxonomist; and (3) information from seed catalogues (Ibarra-Manríquez & Cornejo-Tenorio 2010).

Sleeping-tree fidelity

Following Reichard (1998), STF was defined as the number of months each ST was used during the study period. Given that all traps were checked at the end of

each month, the presence of faeces in the traps was used as an indicator of sleeping-tree use by spider monkeys during that month. Since all traps were completely emptied each month, we considered both fresh and old faeces as indicators of sleeping-tree use, thus avoiding the possibility of overestimating STF. Although our definition of STF does not reflect accurately the intensity of use of STs (i.e. it would be better to assess STF on a daily or weekly basis), our results suggest however that our assessment of STF is an appropriate approximation of the intensity of use, as all community attributes of the seed rain were significantly related to STF.

Sample completeness

To assess the sample completeness within each ST, we employed the coverage estimator (\hat{C}) suggested by Chao & Jost (2012), which is a less biased estimator of sample completeness:

$$\hat{C} = 1 - \frac{f_1}{n} \left[\frac{(n-1)f_1}{(n-1)f_1 + 2f_2} \right]$$

where f_1 and f_2 are the number of species represented by one (singletons) and two (doubletons) individuals in the sample, respectively, and n is the total number of individuals in the sample. Sample completeness did not differ between sites (Kruskal–Wallis test, $H = 6.7$, $P = 0.14$; mean \pm SE: $99\% \pm 1\%$, range: 93–100% per ST). This indicates that the seed sample was accurate with our sampling effort, and that our results are not biased by differences in sample completeness among sites (Chao & Jost 2012).

Composition and structure of seed assemblages

Based on our hierarchically nested sampling design (i.e. 60 STs in five sites within two forest types), we calculated the total abundance and species diversity of seeds for each ST during the 13-mo period. Patterns of species diversity were analysed using true diversity measures (i.e. number equivalents, qD). This analytical approach has been recognized as the most appropriate for diversity comparisons (Chao *et al.* 2012, Jost 2006, 2007; Tuomisto 2010). The formulae are detailed elsewhere (Chao *et al.* 2012, Jost 2006). We considered true diversities 0D (species richness), 1D (exponential of Shannon's entropy) and 2D (inverse Simpson concentration). 0D is not sensitive to species abundances and so gives disproportionate weight to rare species (Jost 2006, Tuomisto 2010). 1D weights each species according to its abundance in the community, and therefore can be interpreted as the number of 'common' (or typical) species in the community (Chao *et al.* 2012).

Finally, 2D favours very abundant species, and can be interpreted as the number of ‘dominant’ species in the community (Chao *et al.* 2012, Jost 2010). These three diversity measures were calculated using raw estimators with the ‘entropart’ package for R (v. 3.1.1) (R Core Team Development).

We also assessed differences in species dominance among latrines using the evenness factor proposed by Jost (2010): $EF_{0.2} = {}^2D/{}^0D$. We used this measure because: (1) it is calculated from true diversity measures; (2) it is independent of the number of species in the sample; and (3) it is very easy to interpret. It ranges between 1 (when all species are equally common) and nearly $1/{}^0D$ (when the community is totally dominated by one species), and can be interpreted as the proportion of dominant species in the community (Jost 2010).

Statistical analyses

To assess whether STs can be considered independent samples, we used Mantel tests to correlate the distance between STs with the difference in STF (i.e. differences in the re-use of different STs). The P-values were calculated using the distribution of the R coefficients obtained from 10000 permutations. Because the Mantel tests did not detect a significant spatial autocorrelation of datasets within the continuous forest sites (CF1: $r = -0.23$, $P = 0.05$; CF2: $r = -0.16$, $P = 0.17$), nor within the forest fragments (FF1: $r = 0.14$, $P = 0.24$; FF2: $r = 0.02$, $P = 0.87$; FF3: $r = 0.10$, $P = 0.41$), we considered the STs as replicates in the following analyses.

We first tested for differences in STF among sites and between forest types using analyses of deviance (ANODEV) with generalized linear models (GLM). As suggested for count dependent variables (i.e. number of months), we used a Poisson error and a log link function, and corrected for data overdispersion (Crawley 2002). We then used analyses of covariance (ANCOVA) with GLMs to assess the association between each community-level variable (abundance of seeds, 0D , 1D , 2D , and EF ; the dependent variable) and forest type (categorical factor) and STF (continuous cofactor). The interaction of these two explanatory variables (forest type \times STF) was included in the models to test if the slope of the association between each response variable and STF differed between forest types. We also used Mantel tests to evaluate if seed species turnover between STs within each sampling site was related to the differences between STs in STF. The species turnover between STs was calculated using multiplicative diversity decomposition of Hill numbers: ${}^qD_\beta = {}^qD_\gamma/{}^qD_\alpha$ (Jost 2007). These metrics were calculated using the ‘entropart’ package for R (<<http://CRAN.R-project.org/package=entropart>>) to implement a function to construct a matrix containing β -

Table 1. Effect of sleeping-tree fidelity (STF) of *Ateles geoffroyi* on five community-level seed-rain patterns produced by this primate below sleeping trees located in two forest types (continuous forest and forest fragments) in the Lacandona rain forest, Mexico. The effect of STF, forest type and the interaction between the two was tested with Generalized Linear Models.

| Response variable/factors | df | χ^2 | P |
|-------------------------------|----|----------|--------|
| Abundance of seeds | | | |
| STF | 1 | 31.3 | <0.001 |
| Forest type | 1 | 3.91 | 0.048 |
| Forest type \times STF | 1 | 0.74 | 0.389 |
| Species richness | | | |
| STF | 1 | 40.4 | <0.001 |
| Forest type | 1 | 0.01 | 0.891 |
| Forest type \times STF | 1 | 0.36 | 0.547 |
| Exponential Shannon entropy | | | |
| STF | 1 | 19.4 | <0.001 |
| Forest type | 1 | 0.27 | 0.602 |
| Forest type \times STF | 1 | 0.002 | 0.963 |
| Inverse Simpson concentration | | | |
| STF | 1 | 11.8 | <0.001 |
| Forest type | 1 | 1.42 | 0.233 |
| Forest type \times STF | 1 | 0.44 | 0.506 |
| Evenness factor | | | |
| STF | 1 | 10.6 | 0.001 |
| Forest type | 1 | 3.33 | 0.068 |
| Forest type \times STF | 1 | 0.002 | 0.966 |

diversity values of each pairwise comparison within each sampling site. Thus, for each ST pair, ${}^qD_\gamma$ was the total (gamma) diversity of the two STs, and ${}^qD_\alpha$ was the average alpha diversity per ST. This beta can be interpreted as ‘effective number of completely distinct communities’ (Jost 2007). In our case, it ranged between 1 (when both STs showed identical seed rains) and 2 (when the two STs were completely different from each other).

RESULTS

Sleeping-tree fidelity in continuous and fragmented forests

Overall, STF of spider monkeys was highly variable among sites. It ranged from 3 to 10 mo in continuous forest sites, and from 1 to 12 mo in forest fragments, but it did not differ among sites (GLM; $\chi^2 = 5.55$, $df = 4$, $P = 0.23$; Figure 1a) or between forest types ($\chi^2 = 0.56$, $df = 1$, $P = 0.45$; Figure 1b).

Association between STF and seed-rain patterns

The generalized linear models showed that STF was positively related to all community attributes of the seed rain (Table 1; Figure 2). The interaction in the models (forest type \times STF) was not significant, indicating that




Figure 1. Sleeping-tree fidelity of *Ateles geoffroyi* in continuous (CF) and fragmented (FF) forests in the Lacandona region, Mexico. Differences among sites (a) and between forest types (b) are shown.

Table 2. Correlations between species β -diversity of seeds between sleeping trees of *Ateles geoffroyi* and inter-tree differences in sleeping tree fidelity in two continuous forest sites (CF) and three forest fragments (FF) in the Lacandona rain forest, Mexico. The correlation coefficients were calculated with Mantel tests (* $P < 0.05$; ** $P < 0.001$, after applying a stringent Bonferroni correction to reduce the likelihood of type I statistical errors). ${}^qD_\beta$ represents the species turnover between sleeping trees, considering three orders q (0, 1 and 2), which determine the sensitivity of each β -diversity component to the relative abundances.

| β -diversity component | CF1 | CF2 | FF1 | FF2 | FF3 |
|------------------------------|--------|--------|---------|---------|---------|
| ${}^0D_\beta$ | -0.001 | 0.182 | 0.374** | 0.542** | 0.400** |
| ${}^1D_\beta$ | 0.003 | -0.052 | 0.244* | 0.385** | 0.453** |
| ${}^2D_\beta$ | 0.083 | -0.125 | 0.175 | 0.302* | 0.523** |

this positive association was similar in both continuous and fragmented forests (Figure 2).

The Mantel tests showed that, in the smallest forest fragments (FF2 and FF3), species turnover of seeds among STs increased with increasing inter-ST differences in STF (Table 2). This association was also significant in FF1, but only when considering β -diversity of order 0 and 1 (i.e. ${}^0D_\beta$ and ${}^1D_\beta$). Yet, the Mantel tests did not detect a significant correlation between inter-ST differences in STF and β -diversity for all order q (Table 2).

DISCUSSION

Contrary to our prediction, STF was similar in all study sites and did not differ between continuous and fragmented forests. This was due to the large variation among STs. Although STs were used by spider monkeys on average for 7 out of 13 mo, there was a substantial variation across trees (range = 1–12 mo), which was

relatively higher in forest fragments than in continuous forest sites (Figure 1). Previous studies on the spider monkey (Chapman *et al.* 1989, Russo & Augspurger 2004, Russo *et al.* 2006) and other primate species (*Brachyteles arachnoides*: Bueno *et al.* 2013; *Hylobates lar*: Reichard 1998; *Gorilla gorilla*: Rogers *et al.* 1998; *Lagothrix lagothricha*: Stevenson 2000) also indicated that the use and re-use of STs can be highly variable, with some trees used for long periods, while others are only used occasionally (Anderson 1984, 2000; Reichard 1998, Teichroeb *et al.* 2012). In the case of the spider monkey, this pattern results from its temporal and spatial foraging behaviour in different areas with a high concentration of food trees within the home range (Asensio *et al.* 2012b, Ramos-Fernández *et al.* 2013), and then routinely returning at night to the same or different STs located in proximity to these areas with greater food density (i.e. multiple central-place foraging; *sensu* Chapman *et al.* 1989). This behaviour allows the monkey to monitor the resources for future usage (Asensio *et al.* 2012a) and to move back to its STs using route-based mental maps (Di Fiore & Suarez 2007, Ramos-Fernández *et al.* 2004, Suarez *et al.* 2014), thus minimizing travel time (Asensio *et al.* 2012b, Chapman 1989, Teichroeb *et al.* 2012). Yet, during territorial defence (Chapman *et al.* 1995, Wallace 2008), or exploration to monitor feeding sites at a great distance (Di Fiore & Suarez 2007, Ramos-Fernández *et al.* 2004, Valero & Byrne 2007) males of *A. geoffroyi* can use STs for short periods because they usually do not return to the same ST (Ramos-Fernández *et al.* 2004), possibly explaining the low fidelity found for some STs in our study. Therefore, as reported for other primate species (Heymann 1995, Pontes & Soares 2005, Reichard 1998, Sigg & Stolba 1981, Silva Júnior *et al.* 2009, Smith *et al.* 2007), the high variation in STF in the Lacandona rain forest is most likely related to the spatial and temporal




Figure 2. Effects of sleeping-tree fidelity of *Ateles geoffroyi* on five community-level attributes of seed assemblages in continuous (CF, open dots) and fragmented forests (FF, black dots) in the Lacandona region, Mexico. Abundance of seeds (a), Species richness (b), Exponential Shannon's entropy (c), Inverse Simpson concentration (d) and Evenness factor (e).

changes in the concentration and distribution of food resources and sex differences in the use of space by the spider monkey.

The relatively higher variation in STF in forest fragments than in continuous forest sites can be related to food scarcity in forest fragments (Arroyo-Rodríguez & Mandujano 2006, Chaves *et al.* 2012). In particular, fruit availability is known to be lower in forest fragments than in continuous forest because of the combination of both a lower density of large food trees (dbh > 60 cm), which represent larger fruit patches (Chapman *et al.* 1992), and smaller home-range sizes in fragments (Chaves *et al.* 2012). Thus, the spatial variation in resource availability is expected to be higher in forest fragments than in continuous forest. As a consequence, in forest fragments, the STs that are located in areas with higher availability of resources would be used relatively more often than in continuous forest sites, whereas the STs that are located in areas with lower availability of resources would be used relatively less intensively than in continuous forest sites. Consistent with this idea, we found that in forest fragments 15 out of 36 STs were used for 9–12 mo and five were only used for 1–3 mo. In continuous forest, however, all STs were used for between 3 and 10 mo (Figure 1).

The lower variation in STF in continuous than in fragmented forest can explain why seed β -diversity between STs did not increase with increasing inter-tree differences in STF in continuous forest sites. It is well known that the lack of variation in explanatory variables (STF in our case) result in weaker associations between explanatory and response variables (species turnover in our case) (Eigenbrod *et al.* 2011). As discussed above, the lower variation in STF in continuous forest sites may be associated with higher availability of resources in this sites when compared with the smallest fragments, which in turn can contribute to reduce β -diversity between STs within the continuous forest. In contrast, higher spatial variations in the availability of food resources may contribute to increase the differences in STF in fragments, and also to significant compositional differentiations in seed assemblages between sleeping trees (González-Zamora *et al.* 2014).

In agreement with our predictions, STF was positively related to the abundance and species diversity of seeds in all sites, indicating that the more often a ST is used, the more seeds from a larger variety of species are deposited under it. Such associations can be explained by the changes in the frugivorous diet along the year (Chaves *et al.* 2012, González-Zamora *et al.* 2014) and by the spatial location of the STs in relation to foraging routes used by spider monkeys (Di Fiore & Suarez 2007, Suarez 2003, Suarez *et al.* 2014), resulting in an increase in the abundance and species diversity of seeds. In conclusion, our results suggest that the spatial configuration of seed

communities deposited in latrines depends on the specific location of STs in foraging paths (Di Fiore & Suarez 2007), and by the spatial and temporal variation in territory quality (Asensio *et al.* 2012a, b).

STF was negatively related to community evenness in all study sites. As expected, this finding can be related to the fact that the seeds in the latrines below more frequently used STs reflect the common feeding pattern of the species, i.e. there is a high selectivity towards the consumption of a few genera of plants (Chapman 1988, Milton 1980, Russo *et al.* 2005) and the opportunistic use of a large number of other fruit species (Dew 2008, González-Zamora *et al.* 2009, Link *et al.* 2012, Nunes 1998, Russo *et al.* 2005), thus reducing the number of common and dominant species, as well as the seed community evenness.

Conclusions and implications for seed dispersal

Our study demonstrates that STF is a key factor shaping the seed-rain patterns produced by the spider monkey below STs. Therefore, changes in STF (e.g. those associated with logging or habitat fragmentation) could have important implications for seed dispersal and forest regeneration. Of course, this constitutes a very important avenue for future research, as we still do not have data on seed germination, and seedling recruitment and growth in and around spider monkey latrines differing in re-use.

Although we found no differences in STF between continuous and fragmented forests, further studies in highly fragmented landscapes are required to accurately test if STF can be altered in more fragmented landscapes. Deforestation in the study region is moderate (c. 40% of remaining forest cover), and forest remnants still maintain most of their original vegetation composition and structure (Hernández-Ruedas *et al.* 2014), which may contributed to the apparent lack of differences in STF between forest types.

If STF is higher in fragmented forests, our results indicate that the effectiveness of the spider monkey as a seed disperser (*sensu* Schupp 1993) would change as a result of the increased aggregation of seeds in latrines found in our study. Nevertheless, this will depend on the impact that such an increase in seed/seedling aggregation has on seed germination and seedling recruitment. We can anticipate two alternative scenarios. First, because we found that STF is negatively related to community evenness, higher STF would result in a seed rain dominated by a few seed species. Thus, based on the Janzen–Connell model (Connell 1971, Janzen 1970), we would expect higher seed predation (e.g. by rodents, insects, and/or pathogens) toward the most dominant species, which could reduce predation pressure on less abundant species allowing the recruitment of rare species

in latrines that are more frequently used. Second, an alternative scenario suggests that the low rate of seed arrival in STs less frequently used may allow seeds to avoid the presence of biotic mortality agents, offsetting the lower seed/seedling survival that is expected in such conditions (Bravo 2012, Russo & Augspurger 2004). Since the seed community structure and composition can be decisive for the initial stages of recruitment (Russo & Augspurger 2004, Schupp *et al.* 2010, Wang & Smith 2002), our results call for further studies to assess the persistence of seeds and recruitment of seedlings under different levels of evenness in the seed bank of latrines and to determine which of these two scenarios occurs with increasing STF.

ACKNOWLEDGEMENTS

We are grateful to Sabrina Russo and two anonymous reviewers for the constructive comments on an earlier version of this manuscript. This research was funded by the Consejo Nacional de Ciencia y Tecnología, CONACyT grants (CB-2005-51043 and CB-2006-56799) and the Dirección General de Asuntos del Personal Académico, DGAPA, Universidad Nacional Autónoma de México, UNAM (project IA-203111). AGZ thanks the scholarship (Apoyos Complementarios para la Consolidación Institucional de Grupos de Investigación) provided by CONACyT. KS thanks the UC MEXUS project. We thank Rafael Lombera and Ana M. González-Di Pierro for their invaluable help in the field. The Instituto de Investigaciones Biológicas, UV, and Instituto de Investigaciones en Ecosistemas y Sustentabilidad, UNAM, provided logistical support. This study would not have been possible without the collaboration of Natura y Ecosistemas Mexicanos A.C. and the local people in Chajul, Reforma Agraria and Zamora Pico de Oro ejidos. We are grateful to M. Lobato, H. Ferreira, A. Valencia and A. Lopez for technical support and to G. Ibarra-Manriquez for help with seed identification.

LITERATURE CITED

- ANDERSON, J. R. 1984. Ethology and ecology of sleep in monkeys and apes. *Advances in the Study of Behavior* 14:166–229.
- ANDERSON, J. R. 2000. Sleep-related behavioural adaptations in free ranging anthropoid primates. *Sleep Medicine Reviews* 4:355–373.
- ARIZPE, L., PAZ, F. & VELÁZQUEZ, M. 1993. Cultura y cambio global: percepciones sociales sobre la deforestación en la Selva Lacandona. Pp. 69–89 in Miguel, A. P. (ed.). *La selva Lacandona: tierra de nadie, tierra de todos*. Centro Regional de Investigaciones Multidisciplinarias, Universidad Nacional Autónoma de México, Mexico City.
- ARROYO-RODRÍGUEZ, V. & MANDUJANO, S. 2006. Forest fragmentation modifies habitat quality for *Alouatta palliata*. *International Journal of Primatology* 27:1079–1096.

- ARROYO-RODRÍGUEZ, V., ANDRESEN, E., BRAVO, S. P. & STEVENSON, P. R. 2015. Seed dispersal by howler monkeys: current knowledge, conservation implications and future directions. Pp. 111–139 in Kowalewski, M., Garber, P. A., Cortés-Ortiz, L., Urbani, B. & Youlatos, D. (eds.). *Howler monkeys: behavior, ecology and conservation*. Springer, New York.
- ASENSIO, N., SCHAFFNER, C. M. & AURELI, F. 2012a. Variability in core areas of spider monkeys (*Ateles geoffroyi*) in a tropical dry forest in Costa Rica. *Primates* 53:147–156.
- ASENSIO, N., LUSSEAU, D., SCHAFFNER, C. M. & AURELI, F. 2012b. Spider monkeys use high-quality core areas in a tropical dry forest. *Journal of Zoology* 287:250–258.
- BÖRGER, L., DALZIEL, B. D. & FRYXELL, J. M. 2008. Are there general mechanisms of animal home range behaviour? A review and prospects for future research. *Ecology Letters* 11:637–650.
- BRAVO, S. P. 2012. The impact of seed dispersal by black and gold howler monkeys on forest regeneration. *Ecological Research* 27:311–321.
- BUENO, R. S., GUEVARA, R., RIBEIRO, M. C., CULOT, L., BUFALO, F. S. & GALETTI, M. 2013. Functional redundancy and complementarities of seed dispersal by the last Neotropical megafrugivores. *PLoS ONE* 8:e56252.
- CHAPMAN, C. A. 1988. Patterns of foraging and range use by three species of Neotropical primates. *Primates* 29:177–194.
- CHAPMAN, C. A. 1989. Spider monkey sleeping sites: use and availability. *American Journal of Primatology* 18:53–60.
- CHAPMAN, C. A., CHAPMAN, L. J. & MCLAUGHLIN, R. L. 1989. Multiple central place foraging by spider monkeys: travel consequences of using many sleeping sites. *Oecologia* 79:506–511.
- CHAPMAN, C. A., CHAPMAN, L. J., WRANGHAM, R., HUNT, K., GEBU, D. & GARDNER, L. 1992. Estimators of fruit abundance of tropical trees. *Biotropica* 24:527–531.
- CHAPMAN, C. A., WRANGHAM, R. W. & CHAPMAN, L. J. 1995. Ecological constraints on group size: an analysis of spider monkey and chimpanzee subgroups. *Behavioral Ecology and Sociobiology* 36:59–70.
- CHAO, A. & JOST, L. 2012. Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology* 93:2533–2547.
- CHAO, A., CHIU, C. H. & HSIEH, T. 2012. Proposing a resolution to debates on diversity partitioning. *Ecology* 93:2037–2051.
- CHAVES, O. M., STONER, K. E. & ARROYO-RODRÍGUEZ, V. 2011. Effectiveness of spider monkeys (*Ateles geoffroyi vellerosus*) as seed dispersers in continuous and fragmented rainforests in southern Mexico. *International Journal of Primatology* 32:177–192.
- CHAVES, O. M., STONER, K. E. & ARROYO-RODRÍGUEZ, V. 2012. Differences in diet between spider monkey groups living in forest fragments and continuous forest in Lacandona, Mexico. *Biotropica* 44:105–113.
- CRAWLEY, M. 2002. *Statistical computing: an introduction to data analysis using S-Plus*. John Wiley & Sons, Chichester. 772 pp.
- CONNELL, J. H. 1971. On the role of natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees. Pp. 298–312 in den Boer, P. J. & Gradwell, G. R. (eds.). *Dynamics of populations*. Center for Agricultural Publication and Documentation, Wageningen.
- DEW, J. L. 2008. Spider monkeys as seed dispersers. Pp. 155–182 in Campbell, C. J. (ed.). *Spider monkeys. The biology, behavior and ecology of the genus Ateles*. Cambridge University Press, New York.
- EIGENBROD, F., HECNAR, S. J. & FAHRIG, L. 2011. Sub-optimal study design has major impacts on landscape-scale inference. *Biological Conservation* 144:298–305.
- DI FIORE, A. & SUAREZ, S. A. 2007. Route-based travel and shared routes in sympatric spider and woolly monkeys: cognitive and evolutionary implications. *Animal Cognition* 10:317–329.
- GONZÁLEZ-ZAMORA, A., ARROYO-RODRÍGUEZ, V., CHAVES, O. M., SÁNCHEZ-LÓPEZ, S., STONER, K. E. & RIBA-HERNÁNDEZ, P. 2009. Diet of spider monkeys (*Ateles geoffroyi*) in Mesoamerica: current knowledge and future directions. *American Journal of Primatology* 71:8–20.
- GONZÁLEZ-ZAMORA, A., ARROYO-RODRÍGUEZ, V., OYAMA, K., SORK, V., CHAPMAN, C. A. & STONER, K. E. 2012. Sleeping sites and latrines of spider monkeys in continuous and fragmented rainforests: implications for seed dispersal and forest regeneration. *PLoS ONE* 7:e46852.
- GONZÁLEZ-ZAMORA, A., ARROYO-RODRÍGUEZ, V., ESCOBAR, F., RÖS, M., OYAMA, K., IBARRA-MANRÍQUEZ, G. & STONER, K. E. 2014. Contagious deposition of seeds in spider monkeys' sleeping trees limits effective seed dispersal in fragmented landscapes. *PLoS ONE* 9(2): e89346.
- HERNÁNDEZ-RUEDAS, M. A., ARROYO-RODRÍGUEZ, V., MEAVE, J. A., MARTÍNEZ-RAMOS, M., IBARRA-MANRÍQUEZ, I., MARTÍNEZ, E., JAMANGAPE, J., MELO, F. P. L. & SANTOS, B. A. 2014. Conserving tropical tree diversity and forest structure: the value of small rainforest patches in moderately-managed landscapes. *PLoS ONE* 9(6):e98931.
- HEYMANN, E. W. 1995. Sleeping habits of tamarins, *Saguinus mystax* and *Saguinus fuscicollis* (Mammalia; Primates; Callitrichidae), in north-eastern Peru. *Journal of Zoology* 237:211–226.
- IBARRA-MANRÍQUEZ, G. & CORNEJO-TENORIO, G. 2010. Diversidad de frutos de los árboles del bosque tropical perennifolio de México. *Acta Botánica Mexicana* 90:51–104.
- JANSON, C. H. 1983. Adaptation of fruit morphology to dispersal agents in a Neotropical forest. *Science* 219:187–189.
- JANZEN, D. H. 1970. Herbivores and the number of tree species in tropical forests. *American Naturalist* 104:501–528.
- JORDANO, P. 1992. Fruits and frugivory. Pp. 105–156 in Fenner, M. (ed.). *Seeds: the ecology of regeneration in plant communities*. CAB International, Wallingford.
- JORDANO, P. & SCHUPP, E. W. 2000. Seed disperser effectiveness: the quantity component and patterns of seed rain for *Prunus mahaleb*. *Ecological Monographs* 70:591–615.
- JOST, L. 2006. Entropy and diversity. *Oikos* 113:363–375.
- JOST, L. 2007. Partitioning diversity into independent alpha and beta components. *Ecology* 88:2427–2439.
- JOST, L. 2010. The relation between evenness and diversity. *Diversity* 2:207–232.
- LINK, A., GALVIS, N., MARQUEZ, M., GUERRERO, J., SOLANO, C. & STEVENSON, P. R. 2012. Diet of the critically endangered brown spider monkey (*Ateles hybridus*) in an inter-Andean lowland

- rainforest in Colombia. *American Journal of Primatology* 74:1097–1105.
- MILTON, K. 1980. *The foraging strategy of howler monkeys: a study in primate economics*. Columbia University Press, New York. 165 pp.
- MORA, F. 2008. Caracterización de la cobertura forestal en el Corredor Biológico Mesoamericano-México: patrones espaciales en la pérdida y fragmentación de bosques. Pp. 55–83 in Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO). *Importancia del capital ecológico de la región Corredor Biológico Mesoamericano-México*. Centro de Investigación en Geografía y Geomática, CentroGEO, Mexico City.
- MYERS, N., MITTERMEIER, R. A., MITTERMEIER, C. G., DA FONSECA, G. A. & KENT, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403:853–858.
- NUNES, A. 1998. Diet and feeding ecology of *Ateles belzebuth* at Maracá ecological station, Roraima, Brasil. *Folia Primatologica* 69:61–76.
- PONTES, A. R. & SOARES, M. 2005. Sleeping sites of common marmosets (*Callithrix jacchus*) in defaunated urban forest fragments: a strategy to maximize food intake. *Journal of Zoology* 26:55–63.
- RAMOS-FERNÁNDEZ, G. & WALLACE, R. B. 2008. Spider monkey conservation in the twenty-first century: recognizing risks and opportunities. Pp. 351–372 in Campbell, J. C. (ed.). *Spider monkeys. The biology, behavior and ecology of the genus Ateles*. Cambridge University Press, New York.
- RAMOS-FERNÁNDEZ, G., MATEOS, J. L., MIRAMONTES, O., LARRALDE, H., COCHO, G. & AYALA-OROZCO, B. 2004. Lévy walk patterns in the foraging movements of spider monkeys (*Ateles geoffroyi*). *Behavioral Ecology and Sociobiology* 55:223–230.
- RAMOS-FERNÁNDEZ, G., SMITH AGUILAR, S. E., SCHAFFNER, C. M., VICK, L. G. & AURELI, F. 2013. Site fidelity in space use by spider monkeys (*Ateles geoffroyi*) in the Yucatan Peninsula, Mexico. *PLoS ONE* 8:e62813.
- REICHARD, U. 1998. Sleeping sites, sleeping places, and presleep behavior of gibbons (*Hylobates lar*). *American Journal of Primatology* 46:35–62.
- ROGERS, M. E., VOYSEY, B. C., MCDONALD, K. E., PARNELL, R. J. & TUTIN, C. E. G. 1998. Lowland gorillas and seed dispersal: the importance of nest sites. *American Journal of Primatology* 45:45–68.
- RUSSO, S. E. & AUGSPURGER, C. K. 2004. Aggregated seed dispersal by spider monkeys limits recruitment to clumped patterns in *Virola calophylla*. *Ecology Letters* 7:1058–1067.
- RUSSO, S. E., CAMPBELL, C. J., DEW, L., STEVENSON, P. R. & SUAREZ, S. A. 2005. A multi-forest comparison of dietary preferences and seed dispersal by *Ateles* spp. *International Journal of Primatology* 26:1017–1037.
- RUSSO, S. E., PORTNOY, S. & AUGSPURGER, C. K. 2006. Incorporating animal behavior into seed dispersal models: implications for seed shadows. *Ecology* 87:3160–3174.
- SCHUPP, E. W. 1993. Quantity, quality and the effectiveness of seed dispersal by animals. *Vegetatio* 107:15–29.
- SCHUPP, E. W., JORDANO, P. & GÓMEZ, J. M. 2010. Seed dispersal effectiveness revisited: a conceptual review. *New Phytologist* 188:333–353.
- SIGG, H. & STOLBA, A. 1981. Home range use and daily march in a hamadryas baboon troop. *Folia Primatologica* 36:40–75.
- SILVA JÚNIOR, W. M., MEIRA-NETO, J. A. A., DA SILVA FLÁVIA, M. C., RODRIGUES DE MELO, F., SANTANA MOREIRA, L., FERREIRA BARBOSA, E., DIAS, L. G. & SILVA PERES, C. A. 2009. Habitat quality of the woolly spider monkey. *Folia Primatologica* 80:295–308.
- SMITH, A. C., KNOGGE, C., HUCK, M., LÖTTKER, P., BUCHANAN-SMITH, H. M. & HEYMANN, E. W. 2007. Long term patterns of sleeping site use in wild saddleback (*Saguinus fuscicollis*) and mustached tamarins (*S. mystax*). *American Journal of Physical Anthropology* 134:340–353.
- STEVENSON, P. R. 2000. Seed dispersal by woolly monkeys (*Lagothrix lagotricha*) at Tinigua National Park, Colombia: dispersal distance, germination rates, and dispersal quantity. *American Journal of Primatology* 50:275–289.
- STONER, K. E., RIBA-HERNÁNDEZ, P., VULINEC, K. & LAMBERT, J. E. 2007. The role of mammals in creating and modifying seedshadows in tropical forests and some possible consequences of their elimination. *Biotropica* 39:316–327.
- SUAREZ, S. A. 2003. *Spatio-temporal foraging skills of white-bellied spider monkeys (Ateles belzebuth belzebuth) in the Yasuni National Park, Ecuador*. Doctoral dissertation, State University of New York at Stony Brook, Stony Brook.
- SUAREZ, S. A., KARRO, J., KIPER, J., FARLER, D., MCELROY, B., ROGERS, B. C., STOCKWELL, B. & TAYLOR, Y. 2014. A comparison of computer-generated and naturally occurring foraging patterns in route-network-constrained spider monkeys. *American Journal of Primatology* 76:460–471.
- TEICHROEB, J. A., HOLMES, J. A. & SICOTTE, P. 2012. Use of sleeping trees by ursine colobus monkeys (*Colobus vellerosus*) demonstrates the importance of nearby food. *Primates* 53:287–296.
- TUOMISTO, H. 2010. A diversity of beta diversities: straightening up a concept gone awry. Part 1. Defining beta diversity as a function of alpha and gamma diversity. *Ecography* 33:2–22.
- VALERO, A. & BYRNE, R. W. 2007. Spider monkey ranging patterns in Mexican subtropical forest: do travel routes reflect planning? *Animal Cognition* 10:305–315.
- WALLACE, R. B. 2008. Towing the party line: territoriality, risky boundaries and male group size in spider monkey fission–fusion societies. *American Journal of Primatology* 70:271–281.
- WANG, B. C. & SMITH, T. B. 2002. Closing the seed dispersal loop. *Trends in Ecology and Evolution* 17:379–385.
- WENNY, D. G. 2001. Advantages of seed dispersal: a re-evaluation of directed dispersal. *Evolutionary Ecology Research* 3:51–74.